

Lawrence Berkeley National Laboratory

Recent Work

Title

HIGH-RESOLUTION STUDY OF THE $C^{12}(p,2p)B^{11}$ REACTION AT 50 MeV

Permalink

<https://escholarship.org/uc/item/1365427q>

Authors

Pugh, H.G.
Hendrie, D.L.
Chabre, Marc
et al.

Publication Date

1965-02-01

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

HIGH-RESOLUTION STUDY OF THE $C^{12}(p, 2p)B^{11}$ REACTION AT 50 MeV

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

HIGH-RESOLUTION STUDY OF THE $C^{12}(p,2p)B^{11}$ REACTION AT 50 MeV

H. G. Pugh, D. L. Hendrie, Marc Chabre, and E. Boschitz

February 1965



HIGH-RESOLUTION STUDY OF THE $C^{12}(p,2p)B^{11}$ REACTION AT 50 MeV*

H. G. Pugh, D. L. Hendrie, Marc Chabre[†], and E. Boschitz[‡]

Lawrence Radiation Laboratory
University of California
Berkeley, California

February 1965

The (p,2p) reaction has been extensively studied at energies between 40 MeV and 440 MeV.¹ These studies have given valuable information on the gross shell structure of nuclei but difficulties in obtaining good energy resolution have limited the use of the reaction for detailed spectroscopic investigations.

We have studied the $C^{12}(p,2p)B^{11}$ reaction at 50 MeV using solid state counters for detection of the outgoing protons and have, for the first time in this type of reaction, obtained energy resolution sufficient to distinguish between individual states of the final nucleus.

A carbon target was bombarded with 50 MeV protons from the Berkeley 88-inch cyclotron. Proton pairs from the (p,2p) reaction were detected in two counter telescopes, each consisting of a pair of lithium-drifted silicon detectors: a 0.024 inch "ΔE" detector and a 0.120 inch "E" detector. The circular collimators defining each solid angle subtended 6.9×10^{-3} steradians at the target. Electronic circuits selected only those events in which the outgoing particles, identified as protons with energies between 9 MeV and 25 MeV, produced a fast coincidence between all four counters. The resolution of the summed energy spectra (the total energy deposited in all four detectors) was better than 350 keV. Figure 1(a) shows a summed energy spectrum which is fairly typical and illustrates that the transition to the ground state of B^{11} is predominant. We have measured angular correlations for this transition with the telescopes placed at equal angles, θ , on opposite sides of the incident beam direction.



Figure 2 shows the differential cross-section for the ground-state transition as a function of θ for events in which $|E_1 - E_2| < 5$ MeV, where E_1 and E_2 are the energies of the two detected protons. This angular correlation differs markedly from those obtained at higher energies:

- a) There are more oscillations, presumably diffraction effects due to localization of the interaction in the nuclear surface.
- b) The cross-section rises quickly at small angles, possibly due to the rapid rise of the proton-proton scattering cross-section at low energies.

Recent distorted wave calculations seem to reproduce the major features of the correlation.²

The inset in Fig. 2 shows three spectra of $(E_1 - E_2)$ for the ground-state transition. The absence of sharp structure in these spectra (which as a consequence of our geometry must be symmetrical about $E_1 = E_2$) indicates that the reaction does not proceed through long-lived proton-unstable excited states of C^{12} .

The $(E_1 - E_2)$ spectra do, however, show slowly-varying structure near the minima of the angular correlation. At these points the cross-section averaged over $|E_1 - E_2| < 5$ MeV is not a good approximation to that for equal energy sharing. There we have drawn smooth lines through the spectra to obtain the cross-section for $E_1 = E_2$; the results are given by the dashed line in Fig. 2. The effects of angular resolution on the angular correlation have not been extracted.

It will be seen from Fig. 1(a) that four excited states of B^{11} are produced with appreciable intensity. The states of spin 5/2- and 7/2- are particularly interesting since, in order for them to be produced by simple removal of a proton from C^{12} , it is necessary for the C^{12} ground state to contain large admixtures of 1f particles. While calculations do indicate the

presence of 1f admixtures in C^{12} it is by no means clear whether the quantity is sufficient.³

Alternatively it is tempting to consider the possibility that these four states may have the character of a $p_{3/2}$ proton hole coupled to the 2+ first excited state of C^{12} at 4.43 MeV. Their formation would then naturally proceed through a double excitation process, i.e., pickup of a $p_{3/2}$ proton accompanied by inelastic scattering of either the incoming or one of the outgoing protons. This mechanism is plausible since the 4.43 MeV state in C^{12} is known to be strongly excited by inelastic proton scattering at these energies.⁴ Double excitation processes of this type have been suggested by Penny and Satchler⁵ and have recently been invoked to interpret an anomalous angular distribution in the $Cr^{52}(d,p)$ reaction.⁶ The production of the 1/2- state at 2.14 MeV would be enhanced by simple removal of a proton from the $(p_{3/2})^{-2}$ $(p_{1/2})^2$ admixture present in the ground state of C^{12} . We might also expect significant configuration mixing between the 3/2- excited state and the ground state.

Using the same counter telescopes as for the $(p,2p)$ reaction we have also studied the $C^{12}(p,d)C^{11}$ reaction with comparable energy resolution, producing analog states in the mirror nucleus. Figure 1(b) shows a spectrum for this reaction. The relative populations of the corresponding states in the two reactions are comparable. In particular the relative weakness of the transition to the 6.34 MeV level in the (p,d) reaction supports our assumption that the unresolved doublet at 6.8 MeV in B^{11} for the $(p,2p)$ reaction consists mainly of the 7/2- state.

The 5/2- and 7/2- states have now been observed in a variety of single-nucleon pickup reactions on C^{12} up to 150 MeV.⁷ Clarification of the role of the double-excitation process in these reactions is necessary to enhance their usefulness as spectroscopic tools.



We wish to thank W. R. Gibson, A. R. Johnston and R. J. Griffiths⁸ for sending us their results prior to publication. There seems to be no direct comparison that can be made between the two sets of results.

REFERENCES

1. H. Tyren, P. Hillman and Th. A. J. Maris, Nucl. Phys. 7, 10 (1958);
G. Tibell, O. Sundberg and P. U. Renberg, Arkiv för Fysik 25, 433 (1963);
M. Riou, Proceedings of the Gatlinburg conference, October 1964, (to be published).
2. K. L. Lim and I. E. McCarthy, Phys. Rev. 133, B1006 (1964) and private communication.
3. A. Goswami and M. K. Pal, Nucl. Phys. 44, 294 (1963).
4. R. W. Peelle, Phys. Rev. 105, 1311 (1957);
T. Stovall and N. M. Hintz, Phys. Rev. 135, B330 (1964).
5. S. K. Penny and G. R. Satchler, Nucl. Phys. 53, 145 (1964).
6. R. Bock, H. H. Dohm, R. Rüdell and R. Stock, Phys. Letters 13, 151 (1964).
7. W. H. Kelly, C. A. Ludemann and C. D. Goodman, Bull. Am. Phys. Soc. 10, 121 (1965);
C. Detraz, Ph.D. Thesis, Orsay (1964);
S. M. Austin, G. L. Salmon, A. B. Clegg, K. J. Foley and D. Newton, Proc. Phys. Soc. (London) 80, 383 (1962);
J. W. Watson, F. Ajzenberg-Selove and R. Middleton, Bull. Am. Phys. Soc. 10, 9 (1965).
8. R. J. Griffiths, A. E. McClatchie, W. R. Gibson, S. J. Hall, A. R. Johnston and K. M. Knight, Congres International de Physique Nucleaire, Paris, 3a(1)/c93 (1964);
W. R. Gibson, A. R. Johnston and R. J. Griffiths, (to be published).

FIGURE CAPTIONS


- Fig. 1(a). A summed energy spectrum ($E_1 + E_2$) for the $C^{12}(p,2p)B^{11}$ reaction at 50 MeV. The positions of known excited states of B^{11} are indicated.
- 1(b). A deuteron energy spectrum from the $C^{12}(p,d)C^{11}$ reaction at 50 MeV. The positions of known excited states of C^{11} are indicated.
- Fig. 2. The differential cross-section $d\sigma/d\Omega_1 d\Omega_2 d(E_1 - E_2)$ for the $C^{12}(p,2p)B^{11}$ g.s. reaction at 50 MeV as a function of θ . The cross-section has been averaged over the region $|E_1 - E_2| < 5$ MeV indicated by the vertical dotted lines on the $(E_1 - E_2)$ spectra shown in the inset. The dashed line is an approximation to the cross-section for equal sharing of energy between the outgoing protons, obtained as described in the text.

FOOTNOTES


* Work performed under the auspices of the U. S. Atomic Energy Commission.

† NATO Fellow on leave from CNRS, Institut Fourier, Grenoble, France.

‡ On leave from NASA Lewis Research Center, Cleveland, Ohio.



MUB-5058



MUB-5057

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report; or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

