

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Physician Characteristics Strongly Predict Patient Enrollment In Hospice

### Permalink

<https://escholarship.org/uc/item/1490c9z6>

### Journal

Health Affairs, 34(6)

### ISSN

0278-2715

### Authors

Obermeyer, Ziad  
Powers, Brian W  
Makar, Maggie  
[et al.](#)

### Publication Date

2015-06-01

### DOI

10.1377/hlthaff.2014.1055

Peer reviewed



# HHS Public Access

Author manuscript

*Health Aff (Millwood)*. Author manuscript; available in PMC 2016 May 02.

Published in final edited form as:

*Health Aff (Millwood)*. 2015 June ; 34(6): 993–1000. doi:10.1377/hlthaff.2014.1055.

## Physician Characteristics Strongly Predict Patient Enrollment in Hospice

**Dr Ziad Obermeyer, MD, MPhil,**

Department of Medicine, Harvard Medical School, Boston, MA

Department of Emergency Medicine, Brigham & Women's Hospital, Boston, MA

Department of Health Care Policy, Harvard Medical School, Boston, MA

**Mr Brian W. Powers, AB,**

Francis Weld Peabody Society, Harvard Medical School, Boston, MA

**Ms Maggie Makar, BS,**

Department of Emergency Medicine, Brigham & Women's Hospital, Boston, MA

**Dr Nancy Keating, MD, MPH, and**

Department of Medicine, Harvard Medical School, Boston, MA

Department of Health Care Policy, Harvard Medical School, Boston, MA

Department of Medicine, Brigham & Women's Hospital, Boston, MA

**Dr David M. Cutler, PhD**

Department of Economics, Harvard University, Cambridge, MA, and National Bureau of Economic Research, Cambridge, MA, USA

### Abstract

Individual physicians are widely believed to play a large role in patients' decisions about end-of-life care, but little empirical evidence supports this view. We developed a novel method for measuring the relationship between physicians' characteristics and hospice enrollment in a nationally representative sample of Medicare patients with poor-prognosis cancer—for whom palliative treatment and hospice would be considered standard of care—who died in the period 2006–11. We found that the proportion of a physician's patients that were enrolled in hospice was a strong predictor of whether or not their other patients would enroll in hospice. The magnitude of this association was larger than that of other known predictors of hospice enrollment that we examined, including patient medical comorbidity, age, race, and sex. Patients cared for by medical oncologists, and those cared for in not-for-profit hospitals, were significantly more likely to enroll in hospice than other patients. These findings suggest that physician characteristics are one of the strongest predictors of whether a patient receives hospice care—which mounting evidence indicates can improve care quality and reduce costs. Interventions geared toward physicians, both by specialty and by previous history of patients' hospice enrollment, may help optimize appropriate hospice use.

There is increasing evidence that hospice care addresses patients' needs and preferences at the end of life, improves care experiences for both patients and caregivers, is associated with decreased health care costs, and even prolongs survival in some populations.[1–4] Although hospice utilization has grown over the past decades, there is substantial variation in its use among patients with similar diagnoses and indications. Many experts and policy makers believe that hospice remains underused.[2,5–7]

A variety of factors are known to predict whether or not patients will enroll in hospice. These include demographic factors such as sex,[6] race,[6,8,9] and age:[8–10] geographical factors:[8,10,11] and health system factors such as the number of physicians and the availability of hospice beds.[12–15]

However, these factors collectively explain only 10 percent of observed variation in hospice use and end-of-life care patterns.[14,15] Furthermore, the available data indicate that patient preferences for the intensity and nature of services have little correlation with the use of hospice care.[12,16–18] This means that most of the variation in this important aspect of care delivery, quality, and Medicare costs remains unaccounted for.

Many researchers and policy makers believe that individual physicians matter a great deal in shaping their patients' choices regarding end-of-life care. While this view seems intuitive, it has been difficult to substantiate empirically. Small survey-based studies have shown correlations between physician specialty, board certification, and beliefs about hospice with patients' decisions about hospice.[19–21] However, the generalizability of these results to outcomes at the national level is unknown.

A study in a large, integrated health system found that the health center where patients received their care was significantly associated with their likelihood of receiving hospice care. The contribution of the individual physician was less clear.[22]

In this study, we developed a novel method for measuring how individual physicians affect their patients' hospice enrollment in a large nationally representative sample of Medicare beneficiaries. We studied patients with poor-prognosis cancers—for example, primary tumors with poor prognoses such as those originating in lung, pancreas, or brain; certain hematologic malignancies; and metastatic disease—to focus on people for whom palliative treatment and hospice would be considered the standard of care.

We quantified the relationship between physician characteristics and patient hospice enrollment using logistic regression, adjusting for a range of other factors known to affect hospice use: physician specialty; patient age, sex, race, and comorbidity; geographic region; and year. We also explored the association of hospice enrollment with the profit status of hospitals with which physicians were associated.

## Study Data And Methods

### Study Cohort

Using a nationally representative 20 percent sample of Medicare fee-for-service beneficiaries in the continental United States, we identified people who died in the period 2006–11 after a poor-prognosis cancer diagnosis and who had at least one year of claims data ( $N=198,948$ ). To identify poor-prognosis cancers, we adapted a palliative care screening instrument used at a major US cancer center. The diagnoses used in the algorithm were developed by clinicians treating a wide range of cancer patients to identify those with poor prognoses and lack of options for curative treatments. We further restricted our study population to patients with diagnostic codes observed to have high mortality rates in the year after diagnosis, calculated using previous years of Medicare data. A detailed description of the creation of the study cohort is available in the online Appendix.[23]

### Physician Characteristics

To quantify the impact of physician characteristics and physician-level variability on hospice enrollment, we constructed a measure of an individual patient's likelihood to enroll in hospice if he or she were under treatment by a given physician. This was accomplished by calculating the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice and then evaluating the impact of this measure on a patient's likelihood of enrolling in hospice (see the analysis below).

We excluded physicians who treated fewer than three poor-prognosis cancer patients, and we conducted a sensitivity analysis that excluded physicians who treated five or fewer patients. We also evaluated the impact of additional physician-level variables, such as physician specialty and profit status of the physician's primary hospital, on patient hospice enrollment.

### Attribution Of Patients To Physicians

We identified all face-to-face encounters in which a patient in our sample received care for their poor-prognosis cancer, and we attributed each patient to the physician involved in the highest number of those encounters. We also explored two alternative methods for attributing patients to physicians: using the first or the last physician who treated the patient for his or her poor-prognosis cancer. More details are in the online Appendix.[23]

### Analysis

Our primary outcome, hospice enrollment, was ascertained based on at least one claim for hospice care. Using logistic regression, we modeled hospice enrollment as a function of the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice; patient's age, sex, race, and comorbidity;[24] physician's specialty; profit status of the physician's primary hospital; year; and hospital referral region. To avoid endogeneity when predicting a given patient's hospice enrollment, we removed that patient from the calculation of the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice. Standard errors were clustered at the level of the individual physician.

We conducted additional analyses to estimate the amount of variance in likelihood of hospice enrollment that was explained by individual predictors. This kind of exercise is straightforward when explaining variance in continuous variables (for example, cost, utilization, or proportion of patients using hospice), but it is more difficult with binary variables (such as individual-level hospice enrollment).[25]

We simulated the effect of a given beneficiary's hypothetical movement from the lowest to the highest decile of continuous variables (the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice and patient's comorbidity and age) or moving between categories of binary variables (patient's sex and race and the facility's profit status), holding all other factors constant. Recording the resulting mean variation in likelihood of hospice enrollment in this population allowed us to quantify and compare the explanatory power of key individual variables.

### Limitations

Our study used fee-for-service Medicare claims data to identify patients who died after receiving specific cancer diagnoses. This allowed us to investigate care patterns in people with poor-prognosis cancer who would be suitable candidates for hospice. However, Medicare data have significant limitations in this regard, which could have biased our results. For example, if patients with more or less severe disease clustered in the practices of certain doctors, hospice enrollment patterns would differ because of patient factors, not physician factors.

This is just one of the many potential biases that can affect the validity of retrospective studies of end-of-life care for cancer patients, leading to questions regarding the validity of this approach.[26] To reduce the risk that such biases would influence our results, our inclusion criteria were designed based on a limited set of *International Classification of Diseases*, Ninth Revision (ICD-9), codes (for a complete list of ICD-9 codes used, see the online Appendix)[23] that identified patients with known severe disease.[27]

We verified that this method did identify high-mortality subgroups by applying it to a year of claims data, without first restricting to beneficiaries who died. We calculated one-year mortality rates in patients with these ICD-9 codes and found it to be high (46 percent), which suggests that we did identify patients with prospectively evident poor-prognosis cancers. However, we cannot say for certain that we identified such patients in our own study cohort.

More broadly, this study was observational, meaning that it could quantify associations but not infer causality. If unmeasured variables caused both hospice enrollment and treatment by a given physician, we could have over- or underestimated the true effect of physician characteristics on patient hospice enrollment.

For example, if patients shopped for physicians based on shared preferences regarding care intensity, we would have overestimated the impact on hospice enrollment attributable to the physician (just as that would have been the case if patients with lower rates of mortality clustered within specific physician practices). Our sensitivity analyses with alternative

attribution methods—particularly the analysis using the first physician to treat each patient for their poor-prognosis cancer, since initial assignment to a physician unknown to the patient is unlikely to reflect shared preferences—were attempts to address this possibility.

Furthermore, we included variables such as age, comorbidity, race, sex, and geographic region in our analysis. But we were unable to include other factors that might affect hospice use but that cannot be measured with Medicare claims data: patient preferences, physician beliefs, or various supply-side factors such as geographic access to hospice or other palliative care interventions.

Data on physician specialty from self-reports or claims may not have accurately reflected the physician's residency training or board certification.

Finally, our results are not generalizable outside the defined study population of Medicare fee-for-service beneficiaries with poor-prognosis cancers.

## Study Results

### Patient Characteristics

Our study cohort of 198,948 patients with poor-prognosis cancers had a mean age of seventy-eight; 88 percent were white, and 52 percent were men. The 131,757 patients (66 percent) enrolled in hospice were older, more likely to be female and more likely to be white, and lived in ZIP codes with higher median incomes, compared to the patients not enrolled in hospice.

Use of inpatient, emergency, and home health services did not differ between the two groups. However, patients enrolled in hospice had slightly more clinic visits during the year before hospice enrollment than patients not enrolled had over a similar period before death. Patients in the two groups had the same median comorbidity scores, reflecting a similar burden of disease. A full description of the study cohort is available in the online Appendix. [23]

### Physician Characteristics

We identified 70,073 physicians who cared for patients with poor-prognosis cancers. Patient load was concentrated among a relatively small pool of physicians: the top 10 percent of physicians cared for 47 percent of all patients in the study. Patient load also varied by specialty: Medical and radiation oncologists cared for an average of fifteen and seventeen patients, respectively, while internists, other medical specialists, and surgeons cared for three to five patients on average. Thus, medical and radiation oncologists made up only 19 percent of the physicians in our cohort, but they cared for 57 percent of all patients with poor-prognosis cancer and accounted for 77 percent of all poor-prognosis cancer encounters.

The fraction of a physician's patients with poor-prognosis cancer enrolled in hospice varied substantially across physicians. For example, 1.6 percent of physicians had less than 5 percent of their poor-prognosis cancer patients enrolled in hospice, while 8.3 percent of physicians had more than 95 percent of their patients enrolled (Exhibit 1).

Results were similar for both alternative attribution strategies. More detailed information on physicians is available in the online Appendix.[23]

### **Impact Of Physician And Patient Characteristics On Hospice Enrollment**

After we assigned patients to the physicians who saw them most frequently for poor-prognosis cancer care, we found that several factors were significantly associated with patients' hospice enrollment. There was substantial variation in hospice use among hospital referral regions, but no clear regional patterns (for complete hospital referral region data see the online Appendix).[23] The likelihood of hospice enrollment generally increased over time (Exhibit 2). After we adjusted for geographical and temporal factors, we found that greater medical comorbidity, older age, female sex, and white race were associated with hospice enrollment.

Patients treated by physicians with claims predominantly linked to not-for-profit hospitals were significantly more likely to enroll in hospice, compared to patients treated by physicians linked to for-profit facilities (Exhibit 2). The specialty of a patient's attributed physician was also significantly associated with the patient's hospice enrollment: Compared to patients attributed to medical oncologists, patients attributed to internists or family practitioners, medical subspecialists, or surgeons were significantly less likely to enroll in hospice. The fraction of a physician's patients with poor-prognosis cancer enrolled in hospice was also significantly associated with hospice enrollment.

We compared the amount of variance in the likelihood of hospice enrollment explained by physician characteristics (Exhibit 3). Moving from the lowest to the highest decile of the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice would have increased the mean likelihood of hospice enrollment from 0.58 to 0.73 in our study cohort, a 27 percent relative increase. This effect was greater than the effects of other predictors of enrollment.

### **Sensitivity Analysis**

When we used alternative patient attribution strategies, the fraction of a physician's patients with poor-prognosis cancer enrolled in hospice remained significantly and strongly associated with hospice enrollment by other patients of that physician. The magnitude of the association was smaller in the model that used the first physician to treat the patient for poor-prognosis cancer (odds ratio: 2.26; 95 percent confidence interval: 2.1, 2.38) than in the primary analysis (OR: 2.67; 95 percent CI: 2.53, 2.82). In contrast, the magnitude was larger in the model that used the last physician (OR: 2.81; 95 percent CI: 2.67, 2.96) and when we restricted the analysis to physicians who saw more than five poor-prognosis patients (OR: 4.16; 95 percent CI: 3.87, 4.48). For complete results, see the online Appendix.[23]

### **Discussion**

Despite growing evidence that hospice can improve care quality and reduce costs,[1–4] hospice use among patients with poor-prognosis illness remains suboptimal. We found that physician-level characteristics strongly predicted patient enrollment in hospice in a large, nationally representative sample of Medicare beneficiaries with poor-prognosis cancers. The

magnitude of this effect was larger than other measured predictors of hospice enrollment, including patient age, race, sex, and medical comorbidity. This finding was robust to several different methods of attributing patients to physicians.

The physician-level variation that we observed could have several sources. For example, existing research suggests that physician preference[18] and practice setting[28] both have a significant impact on patient enrollment in hospice.

Together with the findings of other studies,[14,17,18] our results demonstrate that which physician a patient sees is one of the most important predictors of whether or not he or she enrolls in hospice. This has important implications for the development of policies to advance high-quality end-of-life care. Structural changes to Medicare's criteria for hospice eligibility might increase the use of hospice,[29] but they might be insufficient without concurrent efforts to change physicians' behavior. This would require concerted efforts by the payers, providers, and professional societies most involved with the care of patients with advanced illness.

In our study, large numbers of patients with poor-prognosis cancers were concentrated in a relatively small pool of physicians, the top 10 percent of whom saw nearly half of all patients. This suggests that focused interventions to improve care among high-volume providers could have a major impact.

Such interventions can be grouped into three categories. First, efforts to improve training in end-of-life care are needed in residency and fellowship programs for physicians who will care for large numbers of poor-prognosis patients (for example, medical oncologists, but also physicians specializing in nephrology and cardiology). This is particularly important given the known deficiencies in current educational curricula.[30]

Second, improved measurement of the quality of end-of-life care could help incentivize provider behavior change. The American Society for Clinical Oncology's Quality Oncology Practice Initiative[31] measures hospice use at the practice level and is one example of how professional societies can define standards to promote quality improvement.

A related point is that payers should identify and remove any existing elements of quality measures or payment structures that could represent disincentives for discussions about end-of-life care. This would give physicians—regardless of their personal beliefs—more incentive to engage in discussions about hospice care. The public reporting of metrics on end-of-life care could allow patients and caregivers to choose providers whose beliefs and preferences are consistent with their own when deciding on cancer-directed treatment before death is imminent.

Our sensitivity analysis showed that hospice enrollment was more strongly associated with the characteristics of patients' last physician than with those of their first physician (ORs: 2.94 and 2.35, respectively). This may indicate that, over time, patients gravitate toward providers whose practices match their own preferences.



Third, payers and providers could deploy interventions to increase rates of hospice use among specific physicians who underuse hospice, as measured by mean hospice days per patient or fraction of poor-prognosis patients enrolled in hospice. Implementation of accountable care organizations and other provisions of the Affordable Care Act are likely to provide incentives for such population health management strategies.

Few validated interventions exist, however, and previous efforts to improve joint physician-patient decision making for seriously ill patients have produced mixed results. For example, the Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment (SUPPORT) randomized trial found no effect of integrating specially trained nurses into discussions of end-of-life care.[32] A more encouraging trial demonstrated that identifying eligible nursing home patients and asking their providers to discuss hospice care increased the patients' use of hospice.[33]

More research on effective interventions to increase the uptake of hospice use in real practice environments is needed. So are improved predictive analytic methods for identifying patients with poor prognoses, using the limited administrative data sets that are available to most payers and providers.

The specialty of a patient's physician played a key role in their decision to enroll in hospice: Patients cared for by medical oncologists were significantly more likely to enroll in hospice, compared to patients of other physicians. These findings have several possible interpretations: Compared to other providers, oncologists may make better prognostic estimates, or they may be more comfortable or effective in discussing hospice care with patients. In addition, these results could be due to the environment in which specialists practice: again compared to other providers, oncologists may be more likely to work in practices featuring established relationships with hospice providers. Overall, our results further support the need to concentrate care in high-volume, high-quality centers.[34]

We also found that being treated by a physician affiliated with a not-for-profit hospital was associated with a small but significant increase in the likelihood of hospice enrollment. Supply-side factors and system-level characteristics have been shown to affect hospice enrollment,[8,11,13,21] and we believe this correlation reflects similar factors. Previous research has found that not-for-profit hospitals are more likely to have palliative care and hospice programs.[28,35] Also, patients who do not enroll in hospice receive more end-of-life care,[4] which generates additional revenue that may be particularly appealing to for-profit hospitals.

## Conclusion

Continued development and evaluation of programs that target providers and clinical decision making will be necessary to improve decision making about hospice and end-of-life care in general. Given mounting evidence that hospice and palliative care improve quality and reduce costs, we hope that payers and provider organizations will prioritize the development and testing of new provider-oriented models to improve the uptake of high-quality end-of-life care.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

Funding for this research was provided by the National Institutes of Health (Grant No. DP5 OD012161) to Ziad Obermeyer.

## Notes

1. Teno JM, Clarridge BR, Casey V, Welch LC, Wetle T, Shield R, et al. Family perspectives on end-of-life care at the last place of care. *JAMA*. 2004; 291(1):88–93. [PubMed: 14709580]
2. Institute of Medicine. *Dying in America: improving quality and honoring individual preferences near the end of life*. Washington (DC): National Academies Press; 2014.
3. Temel JS, Greer JA, Muzikansky A, Gallagher ER, Admane S, Jackson VA, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med*. 2010; 363(8):733–742. [PubMed: 20818875]
4. Obermeyer Z, Makar M, Abujaber S, Dominici F, Block S, Cutler DM. Association between the Medicare hospice benefit and health care utilization and costs for patients with poor-prognosis cancer. *JAMA*. 2014; 312(18):1888–1896. [PubMed: 25387186]
5. Gazelle G. Understanding hospice—an underutilized option for life's final chapter. *N Engl J Med*. 2007; 357(4):321–324. [PubMed: 17652645]
6. Medicare Payment Advisory Commission. Washington (DC): MedPAC; 2014 Mar. Report to the Congress: Medicare payment policy [Internet]. Available from: [http://www.medpac.gov/documents/reports/mar14\\_entirereport.pdf?sfvrsn=0](http://www.medpac.gov/documents/reports/mar14_entirereport.pdf?sfvrsn=0) [cited 2015 Mar 18]
7. Stevenson DG. Growing pains for the Medicare hospice benefit. *N Engl J Med*. 2012; 367(18):1683–1685. [PubMed: 23113479]
8. Virnig BA, Kind S, McBean M, Fisher E. Geographic variation in hospice use prior to death. *J Am Geriatr Soc*. 2000; 48(9):1117–1125. [PubMed: 10983913]
9. Greiner KA, Perera S, Ahluwalia JS. Hospice usage by minorities in the last year of life: results from the National Mortality Followback Survey. *J Am Geriatr Soc*. 2003; 51(7):970–978. [PubMed: 12834517]
10. Lackan NA, Ostir GV, Freeman JL, Mahnken JD, Goodwin JS. Decreasing variation in the use of hospice among older adults with breast, colorectal, lung, and prostate cancer. *Med Care*. 2004; 42(2):116–122. [PubMed: 14734948]
11. Wennberg JE, Fisher ES, Stukel TA, Skinner JS, Sharp SM, Bronner KK. Use of hospitals, physician visits, and hospice care during last six months of life among cohorts loyal to highly respected hospitals in the United States. *BMJ*. 2004; 328(7440):607. [PubMed: 15016692]
12. Baker LC, Bundorf MK, Kessler DP. Patients' preferences explain a small but significant share of regional variation in Medicare spending. *Health Aff (Millwood)*. 2014; 33(6):957–963. [PubMed: 24889944]
13. Christakis NA, Iwashyna TJ. Impact of individual and market factors on the timing of initiation of hospice terminal care. *Med Care*. 2000; 38(5):528–541. [PubMed: 10800979]
14. Iwashyna TJ, Chang VW, Zhang JX, Christakis NA. The lack of effect of market structure on hospice use. *Health Serv Res*. 2002; 37(6):1531–1551. [PubMed: 12546285]
15. Kelley AS, Ettner SL, Morrison RS, Du Q, Wenger NS, Sarkisian CA. Determinants of medical expenditures in the last 6 months of life. *Ann Intern Med*. 2011; 154(4):235–242. [PubMed: 21320939]
16. Pritchard RS, Fisher ES, Teno JM, Sharp SM, Reding DJ, Knaus WA, et al. Influence of patient preferences and local health system characteristics on the place of death. SUPPORT investigators. Study fo Understand Prognoses and Preferences for Risks and Outcomes of Treatment. *J Am Geriatr Soc*. 1998; 46(10):1242–1250. [PubMed: 9777906]

17. Barnato AE, Herndon MB, Anthony DL, Gallagher PM, Skinner JS, Bynum JPW, et al. Are regional variations in end-of-life care intensity explained by patient preferences? A study of the US Medicare population. *Med Care*. 2007; 45(5):386–393. [PubMed: 17446824]
18. Cutler, D.; Skinner, JS.; Stern, AD.; Wennberg, D. Cambridge (MA): National Bureau of Economic Research; 2013 Aug. Physician beliefs and patient preferences: a new look at regional variation in health care spending [Internet]. (NBER Working Paper No. 19320). Available from: <http://www.nber.org/papers/w19320.pdf> [cited 2015 Mar 16]
19. Kogan AC, Brumley R, Wilber K, Enguidanos S. Physician factors that influence patient referrals to end-of-life care. *Am J Manag Care*. 2012; 18(11):e416–e422. [PubMed: 23198748]
20. Chinn GM, Liu PH, Klabunde CN, Kahn KL, Keating NL. Physicians' preferences for hospice if they were terminally ill and the timing of hospice discussions with their patients. *JAMA Intern Med*. 2014; 174(3):466–468. Erratum in *JAMA Intern Med*. 2014;174(3):484. [PubMed: 24342971]
21. Bradley EH, Fried TR, Kasl SV, Cicchetti DV, Johnson-Hurzeler R, Horwitz SM. Referral of terminally ill patients for hospice: frequency and correlates. *J Palliat Care*. 2000; 16(4):20–26. [PubMed: 11965930]
22. Keating NL, Herrinton LJ, Zaslavsky AM, Liu L, Ayanian JZ. Variations in hospice use among cancer patients. *J Natl Cancer Inst*. 2006; 98(15):1053–1059. [PubMed: 16882942]
23. To access the Appendix, click on the Appendix link in the box to the right of the article online.
24. Gagne JJ, Glynn RJ, Avorn J, Levin R, Schneeweiss S. A combined comorbidity score predicted mortality in elderly patients better than existing scores. *J Clin Epidemiol*. 2011; 64(7):749–759. [PubMed: 21208778]
25. Mittlböck M, Schemper M. Explained variation for logistic regression. *Stat Med*. 1996; 15(19): 1987–1997. [PubMed: 8896134]
26. Bach PB, Schrag D, Begg CB. Resurrecting treatment histories of dead patients: a study design that should be laid to rest. *JAMA*. 2004; 292(22):2765–2770. [PubMed: 15585737]
27. Earle CC, Ayanian JZ. Looking back from death: the value of retrospective studies of end-of-life care. *J Clin Oncol*. 2006; 24(6):838–840. [PubMed: 16484691]
28. Goldsmith B, Dietrich J, Du Q, Morrison RS. Variability in access to hospital palliative care in the United States. *J Palliat Med*. 2008; 11(8):1094–1102. [PubMed: 18831653]
29. Aldridge Carlson MD, Barry CL, Cherlin EJ, McCorkle R, Bradley EH. Hospices' enrollment policies may contribute to underuse of hospice care in the United States. *Health Aff (Millwood)*. 2012; 31(12):2690–2698. [PubMed: 23213153]
30. Buss MK, Lessen DS, Sullivan AM, Von Roenn J, Arnold RM, Block SD. Hematology/oncology fellows' training in palliative care: results of a national survey. *Cancer*. 2011; 117(18):4304–4311. [PubMed: 21365618]
31. American Society of Clinical Oncology. Alexandria (VA): ASCO; The Quality Oncology Practice Initiative [Internet]. Available from: <http://qopi.asco.org/program.html> [cited 2015 Mar 18]
32. A controlled trial to improve care for seriously ill hospitalized patients. The Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment (SUPPORT). The SUPPORT Principal Investigators. *JAMA*. 1995; 274(20):1591–1598. Erratum in *JAMA*. 1996;275(16):1232. [PubMed: 7474243]
33. Casarett D, Karlawish J, Morales K, Crowley R, Mirsch T, Asch DA. Improving the use of hospice services in nursing homes: a randomized controlled trial. *JAMA*. 2005; 294(2):211–217. [PubMed: 16014595]
34. Hillner BE, Smith TJ, Desch CE. Hospital and physician volume or specialization and outcomes in cancer treatment: importance in quality of cancer care. *J Clin Oncol*. 2000; 18(11):2327–2340. [PubMed: 10829054]
35. Morrison RS, Maroney-Galin C, Kralovec PD, Meier DE. The growth of palliative care programs in United States hospitals. *J Palliat Med*. 2005; 8(6):1127–1134. [PubMed: 16351525]

## Biographies


Ziad Obermeyer (zobermeyer@partners.org) is an assistant professor of emergency medicine and health care policy at Harvard Medical School and an emergency physician at the Brigham and Women's Hospital.

Brian W. Powers is an MD candidate at Harvard Medical School.

Maggie Makar is a research assistant in the Department of Emergency Medicine at the Brigham and Women's Hospital.

Nancy L. Keating is a professor of health care policy and medicine at Harvard Medical School and an internist at the Brigham and Women's Hospital.

David Cutler is the Otto Eckstein Professor of applied economics at Harvard University.



**EXHIBIT 1.**

Distribution of Physician Revealed Preference for Hospice

SOURCE: Authors' analysis of Medicare data.

Notes: Physician revealed preference is reported as  $Pr(Hos|MD)$ , the proportion of a physician's patients enrolled in hospice before death.

**EXHIBIT 2**

## Physician Revealed Preference for Hospice According to Self-Reported Specialty

|                               | Number (%)   | Average Pr(hos  MD) (SD) |
|-------------------------------|--------------|--------------------------|
| <b>Primary Care (PC)</b>      |              |                          |
| Internal Medicine             | 22804 (28.1) | 0.66 (0.24)              |
| Family Medicine               | 8769 (10.8)  | 0.65 (0.27)              |
| Geriatric Medicine            | 1329 (1.6)   | 0.65 (0.25)              |
| Phys. Medicine & Rehab.       | 918 (1.1)    | 0.62 (0.25)              |
| <b>Non-Primary Care (NPC)</b> |              |                          |
| Hematology-Oncology           | 6476 (8.0)   | 0.65 (0.16)              |
| Surgery                       | 4802 (5.9)   | 0.64 (0.25)              |
| Specialist                    | 4272 (5.2)   | 0.64 (0.23)              |
| Pulmonary Disease             | 3774 (4.7)   | 0.61 (0.22)              |
| Gastroenterology              | 3463 (4.2)   | 0.68 (0.24)              |
| Radiation Oncology            | 3321 (4.1)   | 0.69 (0.15)              |
| Neurological Surgery          | 2025 (2.5)   | 0.67 (0.23)              |
| Critical Care Medicine        | 1936 (2.4)   | 0.61 (0.23)              |
| Medical Oncology              | 1890 (2.3)   | 0.66 (0.16)              |
| Neurology                     | 1260 (1.6)   | 0.69 (0.26)              |
| Thoracic Surgery              | 1201 (1.5)   | 0.58 (0.23)              |
| Infectious Disease            | 1030 (1.3)   | 0.61 (0.25)              |
| Nephrology                    | 850 (1.1)    | 0.57 (0.27)              |

SOURCE: Authors' analysis of Medicare data.

Notes: Self-reported specialties for all specialties representing >1 percent of the physicians caring for patients in the study cohort. See eTable4 for a description of the self-reported specialty taxonomy. Pr (hos | MD) is a measure of a physicians revealed preference for hospice which represented the number of patients enrolled in hospice as a fraction of all patients attributed to the physician for cancer care. SD represents standard deviation.

**EXHIBIT 3**

Association between Patient and Physician Characteristics and the Likelihood of Hospice Enrollment.

|                                  | OR (CI)              | Likelihood of Enrollment – Lowest decile <sup>†</sup> or absence of risk factor <sup>‡</sup> | Likelihood of Enrollment – Highest decile <sup>†</sup> or presence of risk factor <sup>‡</sup> | % Change (CI)           |
|----------------------------------|----------------------|--|--|-------------------------|
| Age <sup>†</sup>                 | 1.01<br>(1.01, 1.01) | 0.61   | 0.67   | 10.43<br>(10.41, 10.44) |
| Male sex <sup>‡</sup>            | 0.80<br>(0.79, 0.82) | 0.61   | 0.61   | -7.65<br>(-7.66, -7.64) |
| White race <sup>‡</sup>          | 1.41<br>(1.37, 1.44) | 0.57   | 0.64   | 14.44<br>(14.42, 14.46) |
| Comorbidity <sup>†</sup>         | 1.05<br>(1.05, 1.06) | 0.57   | 0.71   | 27.58<br>(27.55, 27.62) |
| Pr (hos   MD) <sup>†</sup>       | 2.92<br>(2.78, 3.07) | 0.54   | 0.71   | 33.38<br>(33.34, 33.41) |
| For-profit facility <sup>‡</sup> | 0.94<br>(0.90, 0.97) | 0.64   | 0.62   | -2.46<br>(-2.46, -2.45) |

SOURCE: Authors' analysis of Medicare data.

Notes: The first column shows odds ratios (OR). ORs of less than 1 mean a negative relationship, and an odds ratio of greater than 1 means a positive relationship, between the variable and the likelihood of a patient to enroll in hospice. The second column shows estimates for the probability of hospice enrollment in the lowest decile of continuous variables (denoted by †: (Pr(Hos|MD), comorbidity, and age), or the absence of binary risk factors (denoted by ‡: male sex, white race, treatment at a for-profit facility). The third column shows the probability of hospice enrollment in the highest decile or in the presence of binary risk factors. The final column shows the % change in likelihood of enrolling in hospice when moving from the second to third column. Geographic region (HRR) and year were controlled for in the analysis but are not presented in this Exhibit. Pr (hos | MD) is a measure of a physicians revealed preference for hospice which represented the number of patients enrolled in hospice as a fraction of all patients attributed to the physician for cancer care. CI represents the lower and upper bounds of the 95% confidence interval. All associations were significant at p<0.05. Please see eTable7 for complete results of the logistic regression.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript