

UC Irvine

UC Irvine Previously Published Works

Title

Cytokines IL-17 and IL-22 in the host response to infection

Permalink

<https://escholarship.org/uc/item/14x64023>

Journal

Pathogens and Disease, 74(9)

ISSN

0928-8244

Authors

Valeri, Maria
Raffatellu, Manuela

Publication Date

2016-12-01

DOI

10.1093/femspd/ftw111

Peer reviewed

MINIREVIEW

Cytokines IL-17 and IL-22 in the host response to infection

Maria Valeri^{1,2} and Manuela Raffatellu^{1,2,*}

¹Department of Microbiology and Molecular Genetics, University of California Irvine School of Medicine, Irvine, CA 92697-4025, USA and ²Institute for Immunology, University of California Irvine School of Medicine, Irvine, CA 92697-4025, USA

*Corresponding author: Department of Microbiology and Molecular Genetics and Institute for Immunology, University of California, Irvine, Irvine, CA 92697-4025, USA. Tel: 949-824-0359; Fax: 949-824-8598; E-mail: manuelar@uci.edu

One sentence summary: The review highlights the pivotal roles of IL-17 and IL-22 in host defense against microbes.

Editor: Brooke Napier

ABSTRACT

Cytokines IL-17 and IL-22 play pivotal roles in host defense against microbes and in the development of chronic inflammatory diseases. These cytokines are produced by cells that are often located in epithelial barriers, including subsets of T cells and innate lymphoid cells. In general, IL-17 and IL-22 can be characterized as important cytokines in the rapid response to infectious agents, both by recruiting neutrophils and by inducing the production of antimicrobial peptides. Although each cytokine induces an innate immune response in epithelial cells, their functional spectra are generally distinct: IL-17 mainly induces an inflammatory tissue response and is involved in the pathogenesis of several autoimmune diseases, whereas IL-22 is largely protective and regenerative. In this review, we compare IL-17 and IL-22, describing overlaps and differences in their cellular sources as well as their regulation, signaling, biological functions and roles during disease, with a focus on the contribution of these cytokines to the gut mucosal barrier during bacterial infection.

Keywords: IL-17; IL-22; mucosal immunity

INTRODUCTION

Leukocytes constitute the second largest class of cells found within the intestine, second only to epithelial cells (Artis 2008; Ma et al. 2008). Within the intestinal mucosa, leukocyte types are segregated into two distinct anatomic areas: the lamina propria (LP) and the epithelium (Hooper and Macpherson 2010; Maynard et al. 2012). The LP harbors adaptive immune cells such as T cells and B cells, as well as innate immune cells, including dendritic cells (DCs), macrophages and eosinophils. In contrast, specific subsets of cells called intraepithelial lymphocytes (IELs) are found associated with the epithelial layer, in particular at the basement membrane between enterocytes (Cheroutre, Lamblez and Mucida 2011). Gut IELs are almost exclusively T cells, as originally estimated based on histological sections

(Darlington and Rogers 1966). In addition, CX3C-chemokine receptor 1 (CX3CR1)⁺ macrophages and DCs reside within the intestinal epithelial layer and have the capacity to sample antigens in the gut and promote appropriate T-cell responses (Bogunovic et al. 2009; Schulz et al. 2009).

The gastrointestinal (GI) tract is home to trillions of microbes, comprising thousands of species that reside within the lumen (Mowat and Agace 2014). These microbes, collectively referred to as the gut microbiota, vary in composition and number along the length of the GI tract (Mowat and Agace 2014). Although most intestinal microbes are beneficial and live symbiotically with the host, disruption of the epithelial barrier or invasion by pathogenic microbes elicits a complex immune response involving the intestinal epithelium and leukocytes. Restoration

of epithelial integrity and eradication of invading microbes are orchestrated through the activity of cytokines and chemokines, which enable epithelial cells and leukocytes to communicate, and promote responses such as cell migration, differentiation, replication or activation of cell-intrinsic defenses (Matthews, Weight and Parkos 2014).

Intestinal cytokine responses comprise complex signaling networks, wherein cytokines regulate one another (Garlanda, Dinarello and Mantovani 2013; Manzanillo, Eidenschenk and Ouyang 2015). As such, disrupting one particular cytokine response can lead to different outcomes depending on the cell types and the intestinal regions involved, as well as on the gut microbiota. Recognizing the context of these signaling networks can help in understanding the various roles of intestinal cytokines under different conditions. In this review, we discuss recent findings on the interleukin (IL-) 17 and IL-22 pathways, focusing on their roles in intestinal homeostasis and in the host response to infection. Furthermore, we describe the various cell types that produce IL-17 and IL-22 during bacterial infection, and we discuss how these responses are activated and regulated by pattern recognition receptor (PRR)-dependent pathways. Finally, we examine the importance of IL-17 and IL-22 responses in mediating gut immunity to bacterial-induced colitis.

IL-17 and IL-22 are related, but distinct cytokines

Cytokines are secreted proteins that play an important role in intercellular communication. In addition to orchestrating the immune response to pathogens, cytokines also regulate wound healing, angiogenesis and physiological and pathological tissue reorganization (Zhu and Emerson 2002; Leoni et al. 2015; Kreuger and Phillipson 2016). Cytokines elicit biological effects by binding to the extracellular moiety of specific transmembrane receptor proteins on the outer membrane of cells. This binding triggers a signaling pathway, which leads to functional changes in these cells. Cytokines are grouped into families based on similarities in genome location, gene structure, structure of the secreted protein and the receptor(s) the cytokine engages. IL-17 and IL-22 are leukocyte-derived cytokines that primarily impact epithelial cells in tissues such as the gut, the lung and the skin (Blaschitz and Raffatellu 2010; Qu et al. 2013).

IL-17 belongs to the IL-17 cytokine family, whereas IL-22 is a member of the IL-10 cytokine family (Park et al. 2005). The *in vivo* effector functions of IL-17 and IL-22 are crucial to maintaining mucosal immunity against specific pathogens and include induction of antimicrobial proteins, recruitment of neutrophils to sites of bacterial invasion and enhancement of mucosal barrier repair and maintenance by stimulating epithelial cell proliferation and tight junction protein production (Ye et al. 2001a; Liang et al. 2006; Aujla et al. 2008; Raffatellu et al. 2008; Zheng et al. 2008; Conti et al. 2009; Pickert et al. 2009).

There are six members of the IL-17 family: IL-17A (commonly referred to as IL-17), IL-17B, IL-17C, IL-17D, IL-17E (also known as IL-25) and IL-17F (Kolls and Linden 2004; Iwakura et al. 2011). These cytokines are often co-expressed and can form homo- and heterodimers. In 1995, IL-17 receptor A (IL-17RA) was identified as a new cytokine receptor for IL-17A and was later found to be part of a new cytokine receptor family (Yao et al. 1995). The IL-17 receptor family now consists of five members (IL-17RA, RB, RC, RD and RE), all of which, like their ligands, share sequence homology (Aggarwal and Gurney 2002). IL-17RA is expressed on a wide range of tissues and cell types. Upon stimu-

lation with IL-17, IL-17RA initiates the activation of downstream signaling pathways to induce the production of proinflammatory molecules. However, IL-17RA alone is insufficient to mediate IL-17 signaling. Further evaluation revealed that IL-17 signals through a heterodimeric receptor complex composed of IL-17RA and IL-17RC (Toy et al. 2006; Rickel et al. 2008; Song et al. 2011). It is proposed that the binding of ligand to the first IL-17 receptor subunit alters the affinity and specificity of the second binding event, thereby promoting the formation of a heterodimeric, rather than a homodimeric, receptor complex (Ely, Fischer and Garcia 2009; Liu et al. 2013). IL-17 receptors work through a pathway that depends on ACT1s (also referred to as CIKS) and activates NF- κ B and MAP kinases for the induction of proinflammatory mediators such as IL-6, IL-8, granulocyte-macrophage colony-stimulating factor (GM-CSF), some chemokines (CXCL1, CXCL10, CCL2, CCL7, CCL20) and matrix metalloproteinase-3 (MMP3) and MMP13 (Fossiez et al. 1996; Gaffen 2009).

IL-22 signals through a heterodimeric receptor comprised of the broadly expressed IL-10R2 subunit and the more restricted IL-22R1 subunit (Dumoutier, Louahed and Renaud 2000; Dumoutier et al. 2000; Xie et al. 2000). IL-22R1 is mainly expressed on intestinal and respiratory epithelial cells, on keratinocytes and on hepatocytes, but not on cells of hematopoietic origin (Dudakov, Hanash and van den Brink 2015). IL-22 binds first to IL-22R1, and then the IL-22-IL-22R1 complex binds IL-10R2 to propagate downstream signals (Logsdon et al. 2002; Li et al. 2004). IL-22 signals through the JAK-STAT pathway, inducing phosphorylation of the kinases JAK1 and TYK2, as well as of the transcription factors STAT1, STAT3 and STAT5 (Dumoutier, Louahed and Renaud 2000; Dumoutier et al. 2000; Xie et al. 2000; Pestka et al. 2004). Triggering this signaling pathway allows IL-22 to induce the expression of various tissue-specific genes including those that encode proteins involved in tissue inflammation, immunosurveillance and homeostasis (Wolk et al. 2004; Liang et al. 2006, 2010; Zheng et al. 2008). By eliciting various innate defense mechanisms from epithelial cells, IL-22 is essential for host defense at mucosal surfaces against extracellular pathogens such as bacteria and yeast (Zheng et al. 2007, 2008; Aujla et al. 2008; Sonnenberg et al. 2012). In general, IL-22 acts to strengthen epithelial barrier functions and is involved in tissue homeostasis as well as in tissue repair and wound healing. However, excessive or prolonged production of IL-22 can cause pathology, such as psoriasis-like skin inflammation (Ma et al. 2008).

IL-17 was initially found to be secreted by a subset of CD4⁺ T cells termed Th17 cells, which also secrete the cytokines IL-22 and IL-21 (Harrington et al. 2005; Langrish et al. 2005; Park et al. 2005; Korn et al. 2007; Zheng et al. 2007). However, recent studies identified several other cell types that contribute to IL-17 and IL-22 production (Wolk et al. 2002, 2011; Kondo et al. 2009; Ortega et al. 2009), including activated CD4⁺ T cells (Harrington et al. 2005; Park et al. 2005; Wolk et al. 2011), CD8⁺ T cells (Kondo et al. 2009; Ortega et al. 2009), as well as various innate lymphoid cells (ILCs) such as natural killer (NK) cells (Hughes et al. 2010; Pandya et al. 2011), NKT cells (Rachitskaya et al. 2008), lymphoid tissue inducer (LTI) cells (Cupedo et al. 2009; Crellin et al. 2010) and LTI-like cells (Luci et al. 2009; Kim et al. 2011) (Table 1). The role of these cells in secreting IL-17 and IL-22 is discussed in detail below.

Increasing evidence shows that IL-17 and IL-22 can be protective against infections, largely by the following two mechanisms. The first involves production of antimicrobial peptides, which is largely dependent on the synergistic action of IL-17 and IL-22 on epithelial cells (Wolk et al. 2004; Liang et al. 2006;

Table 1. Cell types that secrete IL-17 and/or IL-22.

Cell type	Cytokine secreted	Transcription factor	Surface phenotype	Human/mouse studies	Ref
Th17	IL-17, IL-22	ROR γ t	CD4 ⁺ , CCR4 ⁺ , CCR6 ⁺ , CXCR3 ⁻ , CD161 ⁺ , IL23R ⁺	Human and mouse studies	Cosmi et al. (2008), Kleinschek et al. (2009); Park et al. (2005)
Th22	IL-22	Unknown (AHR?)	CD4 ⁺ , CCR4 ⁺ , CCR6 ⁺ , CCR10 ⁺	Human	Eyerich et al. (2009); Trifari et al. (2009)
Tc22	IL-22	Unknown	CD4 ⁺ , CD8 ⁺ , IL-21R ⁺	Human	Nograles et al. (2009)
Tc17	IL-17	Unknown	CD8 ⁺ , CD45RB ⁺ , CD38 ⁺ , CD103 ⁺ , CCR6 ⁺	Mouse	Yen et al. (2009)
$\gamma\delta$ cells	IL-17, IL-22	AP1	CD27 ⁻ , CD28 ⁻ , NKP80 ⁺ , CD45RA ⁺ , CD158 ⁺	Human and mouse	Peng et al. (2008), Riera-Sans and Behrens (2007), Shibata et al. (2008)
NKT	IL-17, IL-22	ROR	CD3 ⁺ , CD56 ⁺	Mouse	Rachitskaya et al. (2008)
LTI	IL-17, IL-22	RORC	CD3 ⁻ , CD56 ⁻ , CD161 ⁺ , NKp44 ⁻ , CD117 ⁺ , CD127 ⁺	Human	Cupedo et al. (2009)
NKp46 ⁺	IL-17, IL-22	RORC	CD3 ⁻ , CD56 ⁺ , NKp44 ⁺ NKp46 ⁺ , NKG2D ⁺ , CD117 ⁺ , CD127 ⁺	Mouse	Satoh-Takayama et al. (2008)
NK22	IL-22	Unknown	CD3 ⁻ , CD56 ⁺ , NKp44 ⁺ CD117 ⁺ , CD127 ⁺	Mouse	Norian et al. (2009)

Zheng et al. 2008; Feng et al. 2009). The second mechanism involves IL-17 and IL-22 inducing gut and lung epithelial cells to express chemokines that attract granulocytes, particularly neutrophils, to sites of infection (Albanesi et al. 2000; Nograles et al. 2008). Despite their protective role against extracellular pathogens, IL-17 and IL-22 may also be detrimental when they are not tightly regulated. Indeed, these cytokines contribute to the pathology observed in several autoimmune and chronic inflammatory diseases, including psoriasis, asthma and inflammatory bowel disease (Teunissen et al. 1998; Andoh et al. 2005; Chan et al. 2006; Wolk et al. 2007; McKinley et al. 2008).

IL-17 and IL-22 are regulated by PRR-dependent pathways

The innate immune system is crucial for controlling infectious agents. By using a wide range of innate immune sensors termed pattern recognition receptors (PRRs), the host recognizes general pathogen-associated molecular patterns (PAMPs), initiating a response that limits the pathogen and alerts the adaptive immune system (Fig. 1). Two of the most important PRR protein families are the membrane-bound Toll-like receptors (TLRs) and the cytosolic Nod-like receptors (NLRs). Mammals have many TLRs, each of which recognizes certain PAMPs. For example, TLR4 recognizes lipopolysaccharide, a major component of the Gram-negative bacterial cell wall (Lu, Yeh and Ohashi 2008). TLR2, which is also located on the cell membrane, recognizes bacterial surface lipoproteins (Kang et al. 2009), whereas TLR9, which is located in the membrane of cellular organelles known as endosomes, recognizes non-methylated CpG sequences present in bacterial DNA (Latz et al. 2004). When a TLR binds to its ligand, host-cell adaptor proteins, such as Myeloid differentiation primary response gene 88 (MyD88), associate with the cytoplasmic part of the receptor, and downstream signaling is initiated (Medzhitov et al. 1998) (Fig. 1).

TLR-dependent pathways are important for host defense against bacterial pathogens (Charrel-Dennis et al. 2008; Kawai

and Akira 2011; Wang and Liu 2016), including pathways that control IL-17 and IL-22 responses (Abt et al. 2016) (Fig. 1). For example, it was recently shown that TLR7 stimulation with the synthetic ligand resiquimod (R848) induces IL-23 expression followed by a burst of IL-22 secretion by ILCs, leading to expression of the antimicrobial lectin REG3 γ and restoration of colonization resistance against vancomycin-resistant *Enterococcus* (VRE) (Abt et al. 2016).

Interleukin-23 is a heterodimeric cytokine comprised of the subunits p19 and p40 (Hunter 2005), and is a crucial upstream regulator of IL-17 and IL-22 expression *in vivo* (Dubin and Kolls 2007; Godinez et al. 2009; Gasse et al. 2011; Cox et al. 2012). In a mouse model of *Klebsiella pneumoniae* lung infection, bacterial stimulation of TLR4 on DCs results in IL-23 production (Happel et al. 2003). IL-23, in turn, triggers rapid production of IL-17 and IL-22 by T cells (Ye et al. 2001a,b; Happel et al. 2005; Aujla et al. 2008), which is required for efficient neutrophil recruitment in this model. Similarly, injection of *Escherichia coli* into the peritoneal cavity of naïve mice triggers IL-23 production in a TLR4-dependent manner, and the resulting secreted IL-17 largely originates from $\gamma\delta$ T cells (Shibata et al. 2007). TLR4 also modulates IL-23 expression in mouse bone marrow-derived conventional DCs during *Salmonella enterica* serovar Enteritidis infection (Siegemund et al. 2007). In the presence of intestinal inflammation, IL-23 stimulates T cells in the murine intestinal mucosa to produce IL-17 and IL-22, promoting neutrophil recruitment in response to *S. enterica* serovar Typhimurium infection (Godinez et al. 2009). The TLR/IL-23/IL-17 axis is also an important player in vaccine-induced protection, wherein TLR4 signaling is required for vaccine-induced immunity to *Bordetella pertussis* in mice (Higgins et al. 2006). In this model, production of IL-23 and IL-1 β by DCs contributed to Th17 cell differentiation, while increases in IL-17 levels led to enhanced bacterial clearance and macrophage bactericidal activity in a dose-dependent fashion (Higgins et al. 2006).

All TLRs except TLR3 require MyD88 as an adaptor to mediate signal transduction (Gay et al. 2014). MyD88 appears to be crucial for inducing maximal expression of IL-17 and IL-22 in




Figure 1. Bacterium-induced signaling in antigen-presenting cells (APCs) leads to IL-23 production and subsequent activation of IL-17 and IL-22-producing cells. APCs, for instance DCs and macrophages, express PRRs on their surface, such as TLRs, and intracellular receptors in their cytosol, such as NLRs. These PRRs recognize conserved microbial-associated molecular motifs and PAMPs, including LPS, lipoproteins, and CpG oligodeoxynucleotides. When a TLR binds to its cognate PAMP, host-cell adaptor molecules such as MyD88, TIRAP and TRIF are recruited, and downstream signaling is initiated. The resulting activation of nuclear factor- κ B (NF- κ B) promotes transcription of a range of genes coding for proinflammatory cytokines, including IL-23, which trigger the production of IL-17 and IL-22 in cell subsets including Th17, $\gamma\delta$ T and ILC3.

the gut and the lung (Lebeis et al. 2007; Gibson et al. 2008; Zhang et al. 2009; Keestra et al. 2011). In the *S. Typhimurium* colitis model, *Myd88*^{-/-} mice exhibit a delayed mucosal inflammatory response and blunted IL-17 and IL-22 levels when compared to wild-type controls (Keestra et al. 2011). During gut infection with *Citrobacter rodentium*, *Myd88*^{-/-} mice display increased mortality, intestinal pathology and systemic bacterial spread; whether this is associated with IL-17 and IL-22 responses has yet to be elucidated (Lebeis et al. 2007; Gibson et al. 2008). In the lung, *Myd88*^{-/-} mice exhibit a reduced innate mucosal IL-17 response during infection with *Chlamydia muridarum* (Zhang et al. 2009). Moreover, in a T-cell transfer model of colitis, *Myd88*^{-/-} CD4⁺ T cells express lower levels of IL-17 than wild-type CD4⁺ T cells, indicating a direct role for MyD88 signaling in modulating T-cell effector function (Fukata et al. 2008).

Whether MyD88-dependent regulation of T-cell-induced colitis is due to TLR signaling or IL-1 family cytokine signaling remains unclear. Fukata et al. (2008) showed that CD4⁺CD45Rb^{high} T cells express significant levels of TLRs, suggesting a role for TLR signaling by T cells in Th17 differentiation and in the development of inflammatory bowel disease. On the other hand, Chang et al. (2013) demonstrated that MyD88 mediates IL-1 signaling in CD4⁺ T cells for the upregulation of IL-23R by IL-23, which expands committed Th17 cells via mTOR activation.

The cytosolic NLRs are another class of PRRs activated during enteric infections, recognizing bacterial PAMPs in the cytosol (Fig. 1) (Inohara and Nunez 2001). NLRs are classified into several subfamilies on the basis of their amino-terminal domain (Ting et al. 2008), including the CARD-containing nucleotide-binding oligomerization domain proteins (e.g. NOD1, NOD2 and NLRP3), the pyrin domain-containing proteins (e.g. NLRP3) and the BIR-containing proteins (also known as NALPs or BIRs). NOD receptors recognize fragments of peptidoglycan; specifically, NOD1 recognizes γ -D-glutamyl-meso-diaminopimelic acid (iE-DAP), a molecule produced by Gram-negative bacteria (Chamaillard et al. 2003; Girardin et al. 2003a,b), whereas NOD2 recognizes muramyl dipeptide (MDP), which is present in both Gram-positive and Gram-negative bacteria (Girardin et al. 2003a,b). Several NLRs have been shown to have a role in inflammatory diarrhea. Mice deficient in *Nod1* and *Nod2* exhibit a delayed intestinal inflammatory response in colitis models employing *S. Typhimurium* or *C. rodentium* (Geddes et al. 2011). In the cecal tissue of these models, when compared to wild-type mice, absence of both *Nod1* and *Nod2* results in reduced amounts of IL-17 and IL-22, as well as the antimicrobial proteins RegIII γ and lipocalin 2 (Geddes et al. 2011).

NLRP3 and NLRP4 trigger the formation of cytosolic protein complexes called inflammasomes, which in turn regulate the maturation of the cytokines IL-1 β and IL-18 (Latz, Xiao and Stutz

2013). Inflammasomes are activated in response to a variety of signals that indicate injury to the host, including tissue damage, metabolic stress and infection. NLRP3 responds to a wide range of PAMPs and DAMPs, including bacterial messenger RNA, bacterial DNA:RNA hybrids, MDP, DNA and RNA viruses, fungi, protozoa, ATP, uric acid crystals, silica, aluminum hydroxide, asbestos and bee venom (Kanneganti et al. 2006; Dostert et al. 2008; Hornung et al. 2008; Marina-Garcia et al. 2008; Sander et al. 2011; Franchi et al. 2014; Kailasan Vanaja et al. 2014; Sha et al. 2014). The NLR4 inflammasome is activated by flagellin (Franchi et al. 2006; Miao et al. 2006) and the inner rod proteins of type III secretion systems (Miao et al. 2010). The inflammasome is a key regulator of immune responses in the gut and inflammasome-dependent IL-17 production has been implicated in protection from bacterial and fungal infection. IL-1 β produced by NLRP3 inflammasome activity has been shown to mediate a protective Th17 response against the *E. coli* heat-labile enterotoxin in mice, indicating that NLRP3 activation could regulate IL-17 responses (Brereton et al. 2011). Inflammasome-mediated IL-1 β plays a critical role in promoting Ag-specific Th17 cells and in generating shielding immunity against *B. pertussis* infection (Dunne et al. 2010). It has also been demonstrated that caspase-1 and ASC are important in the defense against *Candida* through IL-1 β and IL-18 expression and consequent induction of antifungal Th1 and Th17 responses (van de Veerdonk et al. 2011). Moreover, two recent studies (Levy et al. 2015; Nowarski et al. 2015) demonstrated that epithelial inflammasome activation and IL-18 secretion can control intestinal homeostasis or induce autoinflammation. ILC3 cells are triggered to secrete IL-22, regulating IL-18 expression in epithelial cells, in turn modulating homeostasis and inflammation. Finally, the IL-18 pathway seems to take part in the host's defense against *C. rodentium* infection downstream of IL-22 (Munoz et al. 2015).

IL-17 and IL-22-producing cells in response to acute infection

T helper cells

During the immune response to infection, IL-17A/F and IL-22 are often produced at the same time and at high levels in inflamed tissues. In general, a broad variety of T cells and other leukocytes secrete IL-17 and/or IL-22 (Table 1). The most characterized IL-17/IL-22-secreting cells are T helper 17 (Th17) cells (Korn et al. 2009). Th17 cells were first described as peripheral CD4⁺ T cells that differentiate into a distinct lineage in a GATA-3 and T-bet-independent fashion (Harrington et al. 2005; Park et al. 2005). This T-cell population is abundant at homeostasis in gut-associated tissues, particularly in the LP, and is defined by expression of the transcription factor ROR γ t (Ivanov et al. 2006, 2008). Th17-cell differentiation depends on IL-6 and on transforming growth factor β (TGF- β) (Bettelli et al. 2006; Ivanov et al. 2006; Mangan et al. 2006). Signal transduction downstream of IL-6 and TGF- β , including STAT3 activation downstream of the IL-6 receptor, induces expression of ROR γ t, which promotes transcription of *Il17a* and *Il17f* (Ivanov et al. 2006; Harris et al. 2007; Yang et al. 2007).

In addition to IL-6 and TGF β , commensal microbes are necessary for the development of Th17 cells (Atarashi et al. 2008; Hall et al. 2008), as Th17 cells are absent in germ-free mice (Ivanov et al. 2006). Monoassociation of germ-free mice with commensal microbes of the class Clostridiales known as 'Candidatus Arthromitus' or 'segmented filamentous bacteria' (SFB) is sufficient to induce Th17-cell development (Gaboriau-Routhiau et al.

2009; Ivanov et al. 2009; Wu et al. 2010). Colonization of mice with SFB results in rapid upregulation of serum amyloid A (SAA) proteins in intestinal epithelial cells, which is both necessary and sufficient to directly promote proliferation of ROR γ t⁺ CD4⁺ T cells and IL-17 production (Sano et al. 2015). Moreover, IL-22 derived from group 3 innate lymphoid cells (ILC3) seems to potentiate expression of SAAs by intestinal epithelial cells (Atarashi et al. 2015; Sano et al. 2015). CD11c⁺ myeloid cells and adhesion-elicited epithelial cell production of reactive oxygen species also contribute to SAA-dependent production of Th17-promoting cytokines (Atarashi et al. 2015).

TLR, inflammasome and dectin signaling each appear to be required for SFB-mediated Th17-cell development, suggesting that there may be redundant innate signals (Ivanov et al. 2009). Additionally, a recent study suggests that intimate adhesion of SFB to the epithelium is important for Th17 development (Atarashi et al. 2015). As SFB have not been identified in the human gut, it is not clear yet whether microbes contribute to Th17 development in humans. Nevertheless, Atarashi et al. (2015) showed that Th17 cells develop in mice upon infection with the human pathogen Enterohemorrhagic *E. coli* O157:H7, as well as after administration of a mixture of adherent human commensal strains belonging to different species, including *Clostridium*, *Bifidobacterium*, *Ruminococcus* and *Bacteroides* isolated from fecal samples of a patient with ulcerative colitis.

The Th17 response is a crucial arm of mucosal immunity against bacterial pathogens in the lung and the intestine. These cells mediate protection in a number of models of lung infection, including *K. pneumoniae*, *Pseudomonas aeruginosa*, *Shigella flexneri* and several *Mycobacterium* species (Khader et al. 2007; Umemura et al. 2007; Aujla et al. 2008; Priebe et al. 2008). In the GI tract, Th17 cells increase host resistance against *Helicobacter pylori*, *C. rodentium* and *S. Typhimurium* (Wu et al. 2007; Raffatellu et al. 2008; Ivanov et al. 2009; Velin et al. 2009; Sellge et al. 2010). For instance, in a simian immunodeficiency virus infection model in macaques, depletion of gut CD4⁺ Th17 cells led to increased systemic dissemination of *S. Typhimurium* (Raffatellu et al. 2008). Similarly, depletion of CD3⁺ T cells in the streptomycin-treated mouse model of *S. Typhimurium* colitis resulted in a blunted innate IL-17 response and an associated decrease in mucosal protection against the pathogen (Raffatellu et al. 2008; Godinez et al. 2009). A subset of LP CD4⁺ T cells that express IL-17 and IL-22 during the innate phase of infection with *S. Typhimurium* and *C. rodentium* has subsequently been identified and termed innate Th17 (iT17) cells (Geddes et al. 2011). Finally, two novel subsets of Th17 cells have recently been identified: a proinflammatory subset that produces GM-CSF (Codarri et al. 2011; El-Behi et al. 2011) and a regulatory subset that produces IL-10 (Esplugues et al. 2011). The role of these cell subsets in response to infection is unknown.

Beyond the mucosa, SFB also trigger development of systemic Th17 cells, which can be detrimental; for example, by triggering spontaneous arthritis in a mouse model (Wu et al. 2010). Similarly, monoassociation of mice with SFB exacerbates the course of experimental autoimmune encephalomyelitis (Lee et al. 2011), a paralytic autoimmune disease induced by immunization with myelin protein and mediated by Th17 cells. Thus, depending on the setting, Th17 cells can be protective, but they can also initiate or exacerbate autoimmunity.

In addition to Th17 cells, a second T helper cell population was recently described as a major source of IL-22 in human tissue. Named Th22 cells, this population expresses the CCR4 and CCR10 chemokine receptors, and is enriched for during inflammatory skin diseases (Nograles et al. 2008; Duhon et al. 2009;

Eyerich et al. 2009; Trifari et al. 2009). In contrast to Th17 cells, which are able to secrete both IL-17 and IL-22, Th22 cells do not secrete IL-17. A third group of adaptive immune cells that produce IL-17 and/or IL-22 are CD8⁺ T cells (Kondo et al. 2009; Ortega et al. 2009). Similar to Th17 cells, a population of CD8⁺ T cells produces IL-17, but it is unclear whether these cells coproduce IL-22 and whether they represent a functionally distinct subset (Tc17 cells) of CD8⁺ T cells (Shin et al. 1999). Furthermore, there is evidence for a subgroup of CD8⁺ T cells that produces IL-22, but not IL-17 (Tc22 cells) (Nogales et al. 2009). The discovery of these distinct lineages of leukocytes that produce different subsets of cytokines emphasizes that production of IL-17 and IL-22 is regulated by distinct processes, underlining the potential complexity of these responses in different models of infection and autoimmunity.

$\gamma\delta$ T cells

$\gamma\delta$ T cells are an innate immune cell population that plays important roles at the mucosal barrier. These cells are predominantly found in the intestinal epithelial lymphocyte compartment of the intestinal mucosa and they can be classified into an IFN- γ -producing subset or an IL-17 and IL-22-producing subset (Cua and Tato 2010). CD27⁻ ROR γ t⁺ $\gamma\delta$ T cells secrete IL-17 and/or IL-22 when stimulated with IL-23, IL-1 β , or with a cocktail of phorbol 12-myristate-13-acetate (PMA) and ionomycin (Jensen et al. 2008; Ribot et al. 2009). Relatedly, IL-23 is sufficient to induce IL-17A production in $\gamma\delta$ T cells purified from the peritoneum of infected mice (Shibata et al. 2007). Of particular note, $\gamma\delta$ T cells can rapidly produce IL-17A upon TLR and/or cytokine stimulation. Early production of IL-17A was observed during *E. coli* infection, where peritoneal IL-17A was found as early as 1 h after infection (maximal production at 6 h), and large amounts of neutrophils were recruited by 24 h post-infection (Shibata et al. 2007). Similarly, IL-17A production by peritoneal $\gamma\delta$ T cells is rapidly induced after injection of heat-killed mycobacteria (Martin et al. 2009) or after sepsis due to cecal puncture (Flierl et al. 2008).

In the skin, IL-17A-producing $\gamma\delta$ T cells can promote the formation of abscesses and granulomas to help mediate containment of microbial infections (Cho et al. 2010). In cutaneous *Staphylococcus aureus* infections, IL-17A-producing skin cells play a critical role in the immune response against this pathogen. In this setting, IL-17A helps to recruit neutrophils into skin abscesses, to limit the size of abscesses and to reduce the number of *S. aureus* bacteria contained in these abscesses (Cho et al. 2010).

In *Listeria monocytogenes* infections, IL-17A-producing $\gamma\delta$ T cells contribute to bacterial clearance by containing the bacteria within granulomas in the liver and by promoting the recruitment of neutrophils and other myeloid cells (Hamada et al. 2008). In the absence of $\gamma\delta$ T cells or IL-17A, bacterial numbers in the liver are more than 100-fold higher than in wild-type mice (Hamada et al. 2008). Moreover, *L. monocytogenes* infection in $\gamma\delta$ T-cell-deficient mice is associated with large inflammatory lesions in the liver with necrotic hepatocytes (Mombaerts et al. 1993; Fu et al. 1994) that are similar to those seen in IL-17A-deficient mice (Hamada et al. 2008). IL-17RA and IL-23 are required to control *L. monocytogenes* systemic infections and $\gamma\delta$ T cells are the principal source of IL-17A early in infection (Meeks et al. 2009), although CD4⁻8⁻ $\alpha\beta$ T cells also produce IL-17A in the peritoneum (Riol-Blanco et al. 2010).

$\gamma\delta$ T cells can also be pathogenic. As with many T-cell populations, inappropriate or sustained production of a proinflam-

matory cytokine can have negative consequences. Several autoimmune disease models have identified IL-17A-producing $\gamma\delta$ T cells as being important in disease progression. For example, in a model of collagen-induced arthritis, oligoclonal IL-17A-producing $\gamma\delta$ T cells accumulate in the lymph nodes and joints of collagen-injected mice (Roark et al. 2007). Depletion of V γ 4 $\gamma\delta$ T cells greatly reduces disease severity consistent with a significant reduction in pathogenic anticollagen IgG2a (Roark et al. 2007). Moreover, in different models of neuroinflammation, $\gamma\delta$ T cells play an important role in early activation and recruitment of cells through the release of IL-17A. In experimental autoimmune encephalomyelitis (EAE), a mouse model of multiple sclerosis, $\gamma\delta$ T cells producing IL-17A accumulate in the brain (Jensen et al. 2008; Sutton et al. 2009). Although $\gamma\delta$ cells alone are not sufficient to reconstitute disease ($\alpha\beta$ T cells are also required), $\gamma\delta$ -deficient mice exhibit reduced EAE severity (Sutton et al. 2009). In addition to producing IL-17A, $\gamma\delta$ T cells also inhibit CD4⁺ Treg responses in EAE and reverse Treg suppression of $\alpha\beta$ T cells in vitro (Petermann et al. 2010).

The involvement of IL-17A-producing $\gamma\delta$ T cells in models of microbial, autoimmune and inflammatory diseases underscores their importance in the initiation and resolution of acute inflammatory responses in a variety of settings. Despite their small numbers, $\gamma\delta$ T cells can have large effects on the type of immune response and the disease outcome.

Innate lymphoid cells

The importance of a non-T-cell source for IL-22 was first demonstrated in a *C. rodentium* infection model, in which wild-type mice and mice that lack B and T cells (*Rag2*^{-/-}) exhibited comparable IL-22 production and normal host defense during the early phase of infection (Zheng et al. 2008). These newly identified cells were termed ILCs, then later categorized in human tonsils and intestinal LP as ILC3 based on their production of IL-17 and IL-22 (Montaldo et al. 2014).

ILC3s can secrete IL-17 and IL-22 in response to IL-1 β and/or IL-23 due to their constitutive expression of IL-1R and/or IL-23R (Spits and Cupedo 2012). ILCs, like B and T cells, are derived from a common lymphoid progenitor. However, they represent a very heterogeneous group of cells. All IL-17/IL-22-producing ILCs are dependent on ROR γ t for their development, including LTi cells (Takatori et al. 2009), LTi-like ILCs (Buonocore et al. 2010), NKp46⁺ ILCs (Satoh-Takayama et al. 2008; Cella et al. 2009; Luci et al. 2009; Sanos et al. 2009), invariant NKT cells (Doisne et al. 2011) and mucosal-associated invariant T cells (Le Bourhis et al. 2011). Unlike Th17 cells, antigenic priming is not required for activation of LTi, LTi-like, NKp46⁺ ILCs and probably $\gamma\delta$ T cells; IL-23 stimulation is often sufficient for inducing IL-17 and IL-22 secretion by these cell types (Cua and Tato 2010; Cua and Sherlock 2011; Le Bourhis et al. 2011).

The lineage of NKp46⁺ cells was first identified as a major source of IL-22 in the intestine (Satoh-Takayama et al. 2008; Cella et al. 2009; Luci et al. 2009). These cells do not co-express IL-17, lack typical NK cell effector functions, and may or may not express NK1.1. Interestingly, some of these NKp46⁺ cells also express CD11c, an integrin highly expressed by DCs. Surprisingly, mice deficient in NKp46 are still capable of secreting IL-22 and are fully resistant to *C. rodentium* infection (Satoh-Takayama et al. 2009). In the spleen and in the mesenteric lymph nodes, another cell type, CD4⁺ LTi-like cells, produce IL-17 and IL-22 upon IL-23 stimulation (Takatori et al. 2009; Sonnenberg et al. 2011). LTi cells are also the building blocks of cryptopatches, which are lymphoid structures found in both the small and

large intestine. During *C. rodentium* infection, LT α cells are major sources of IL-22 in the mesenteric lymph nodes (Sonnenberg et al. 2011). In contrast to NKp46⁺ ILCs, the development of LT α cells is regulated by the lymphotoxin (LT) pathway (Eberl et al. 2004; Satoh-Takayama et al. 2011). LT is a member of the TNF core family and plays a critical role in regulation of mucosal immune responses (Fu and Chaplin 1999; Tumanov, Christiansen and Fu 2007) and contributes to effector immune responses (Ware 2005; Wang et al. 2010). LT also regulates DCs and CD4⁺ T-cell homeostasis in the steady state, as well as determines the functions of these cells during pathogenic challenge (Lewis et al. 2011; Upadhyay and Fu 2013).

Consistent with a role for LT α cells during *C. rodentium* infection, mice deficient in the LT pathway are also more susceptible (Spahn et al. 2004; Ota et al. 2011). Specifically, wild-type mice treated with either an anti-IL-22 antibody or with LT β RfC fusion protein show similar degrees of susceptibility to *C. rodentium* infection, suggesting a connection between the LT and IL-22 pathways. Indeed, IL-22 induction in the colon during *C. rodentium* infection is blocked when the LT pathway is deficient (Tumanov et al. 2011). Taken together, these studies suggest a redundant role for different ILC subsets during *C. rodentium* infection, and the specific contribution of each cell type needs to be further elucidated.

In addition to the aforementioned cell types, a pathogenic population of Thy1⁺Sca1⁺ ILCs has been identified in the inflamed colon of mice in a *H. hepaticus*-induced colitis model (Buonocore et al. 2010). These cells are negative for both NKp46 and CD4, but they are able to produce IL-17 and IL-22 upon stimulation with IL-23, and drive the development of colitis (Buonocore et al. 2010).

NON-LYMPHOID SOURCES OF IL-17 AND IL-22

Other potential sources of IL-17 and IL-22 include neutrophils that are recruited to the site of infection (Zindl et al. 2013; Taylor et al. 2014; Lee et al. 2015). In the dextran sodium sulfate colitis model, IL-22-producing neutrophils were shown to be protective (Zindl et al. 2013). Intestinal pathology was alleviated in DSS-treated IL22^{-/-} mice transplanted with neutrophils from wild-type mice, whereas neutrophils from IL22^{-/-} mice had little effect on disease progression (Zindl et al. 2013). During fungal infection, IL-17-producing neutrophils are recruited to the site of infection (Taylor et al. 2014), but whether neutrophils produce IL-17 or not appears to be dependent on the infection model (Huppler et al. 2015). Proposed mechanisms for activation of IL-17 and IL-22 production by neutrophils include stimulation with cytokines such as IL-6 and IL-23, activation of dectin-1 and 2, and through regulation of ROR γ t and arylhydrocarbon receptor (AhR) via activation of mTOR (Werner et al. 2011; Zindl et al. 2013; Taylor et al. 2014; Chen et al. 2016). Future studies are needed to further elucidate whether IL-17 and IL-22 production by neutrophils contributes to the host response to infection.

HOST DEFENSE AND PROTECTIVE FUNCTION OF IL-17 AND IL-22 DURING INTESTINAL INFECTION

In the gut mucosa, IL-17 and IL-22 have synergistic effects on the induction of antimicrobial proteins by GI epithelial cells, which

have an important role in limiting dissemination of pathogens and of commensal bacteria that could penetrate a disrupted epithelial barrier (Zheng et al. 2008; Ismail, Behrendt and Hooper 2009). IL-17 and IL-22 act in concert to orchestrate the mucosal barrier, but the biological effects of these cytokines differ in many aspects. IL-22 appears to be a novel type of immune mediator that increases the immune defenses of tissue cells, protects tissues from damage and enhances their regeneration, whereas IL-17A and IL-17F are typical proinflammatory mediators. Below we discuss the effects of IL-17 and IL-22 primarily in the context of gut infection.

IL-17

The most prominent function of IL-17 is the recruitment of neutrophils to the site of infection, which is necessary for clearing microorganisms (Ishigame et al. 2009). IL-17 stimulates target cells to produce neutrophil chemoattractants such as the CXC chemokines IL-8 (CXCL8), CXCL1 and CXCL2, as well as growth factors, including G-CSF and GM-CSF (Fig. 2). Neutrophil recruitment is required for a rapid response (i.e. within 4–8 h) against bacterial and fungal pathogens such as *Citrobacter rodentium*, *Salmonella Typhimurium* and *Candida albicans* (Raffatellu et al. 2008; Saijo et al. 2010; Geddes et al. 2011). IL-17 also synergizes with other cytokines, such as IL-1, IL-6 and TNF- α , which together promote activation of tissue infiltrating neutrophils to effectively eliminate extracellular pathogens (Awane et al. 1999; Dong 2008). Mice infected with *C. rodentium* mount protective IL-17 responses, and disruption of IL-17 or its receptor leads to exacerbated bacterial burden and dissemination, increased disease susceptibility resulting from defective induction of CXC chemokines and impaired neutrophil recruitment (Happel et al. 2005; Mangan et al. 2006). Importantly, administration of recombinant IL-17 into IL-17-deficient mice infected with *C. rodentium* restored neutrophilia at sites of inoculation (Happel et al. 2005), demonstrating the critical importance of the IL-23/IL-17 axis in host defense against extracellular bacterial infections. Neutrophils are also an important first line of mucosal defense against *S. Typhimurium*. In patients affected by primary neutrophil immunodeficiency (e.g. chronic granulomatous disease), *S. Typhimurium* often disseminates from the gut, resulting in bacteremia (Mouy et al. 1989; Winkelstein et al. 2000). In accordance with these observations, IL17ra^{-/-} mice exhibit reduced neutrophil accumulation in the gut, and increased translocation of *S. Typhimurium* to the mesenteric lymph nodes and spleen (Raffatellu et al. 2008).

In addition to neutrophil recruitment, the IL-17 cytokine family targets epithelial cells to induce antimicrobial responses against extracellular pathogens and to promote tissue remodeling (Ouyang, Kolls and Zheng 2008). An important mechanism of host defense in the intestine against bacterial pathogens is the presence of tight junctions that maintain the integrity of the intestinal epithelium, thereby preventing bacterial translocation (Macdonald and Monteleone 2005). In the gut epithelium, IL-17 stimulates the secretion of claudin 1 and claudin 2, proteins that are involved in the formation of tight junctions, ultimately forming a large, interconnecting network and maintaining intestinal integrity (Moran et al. 2009; Dungan and Mills 2011). Although IL-17 is largely considered proinflammatory, the cytokine may also promote restoration of the mucosal barrier and further control microbial translocation through the induction of tight junction proteins.




Figure 2. Functions of IL-17 and IL-22 at epithelial surfaces. IL-17 and IL-22 are upregulated in response to infection with a variety of pathogens. These cytokines are produced by subsets of cells, including $\gamma\delta$ T cells, ILC3 and Th17 cells, and elicit innate responses from mucosal epithelial cells. These responses include production of mucins by goblet cells, upregulation of genes involved in wound healing (e.g. Bcl-2, c-Myc, cyclin D1), the secretion of antimicrobial molecules (e.g. S100A8, S100A9, REG proteins, lipocalin-2) and the induction of proinflammatory mediators (e.g. TNF- α , IL-6, CXCL1, G-CSF) that contribute neutrophil recruitment at the site of infection. The functions of both cytokines are overlapping and in part synergistic. While IL-22 is more potent in tissue protection, repair and induction of antimicrobial peptides, IL-17 mediates stronger inflammatory responses by inducing other proinflammatory cytokines and the recruitment of neutrophils.

IL-22

The beneficial role of IL-22 during host defense has been studied in several models of intestinal infection, including *C. rodentium* (Ota et al. 2011) and *Clostridium difficile* (Hasegawa et al. 2014), as well as during infection with VRE, a gut commensal bacterium that causes bacteremia and endocarditis (Abt et al. 2016).

The functions of IL-22 during host defense against pathogens can be summarized into three major categories (Fig. 2). First, by promoting epithelial proliferation, IL-22 helps to maintain and restore epithelial barrier function after infection. Second, IL-22, together with other cytokines such as IL-17 or TNF- α , induces the expression of antimicrobial proteins involved in host defense in the skin, the airways and the intestine. Some antimicrobial proteins induced by IL-22 include lipocalin-2 and calprotectin, which is a heterodimer of S100A8 and S100A9 (Behnsen et al. 2014). These antimicrobial proteins sequester essential metal ions from pathogens (Corbin et al. 2008; Liu et al. 2012), a process known as nutritional immunity (Kehl-Fie and Skaar 2010). Therefore, IL-22-mediated induction of metal sequestration is an essential process in limiting growth and translocation of commensal microbes from the gut (Flo et al. 2004; Berger et al. 2006; Raffatellu et al. 2009). IL-22 also promotes the production of inflammatory mediators such as IL-1 β , SAA and LPS-binding protein (Wolk et al. 2007; Dalmas et al. 2014; Sano et al. 2015). During *Klebsiella pneumoniae* infection, IL-22 is essential for the release of chemokines such as CXCL1, CXCL5 and CXCL9, as well as IL-6 and G-CSF from airway epithelial cells (Aujla et al. 2008). Overall, the primary function of IL-22 is to promote mucosal barrier integrity, thus limiting bacterial replication and dissemination.

During *C. rodentium* infection, IL-22 stimulates the production of protective mucins from goblet cells (Sugimoto et al. 2008; Turner, Stockinger and Helmsby 2013) and induces the release of the C-type lectins REG3 β and REG3 γ from intestinal epithelial cells in the colon (Cash et al. 2006; Zheng et al. 2008). REG3 γ is a soluble C-type lectin produced by Paneth cells and epithelial cells, which exerts its antimicrobial activity by interacting directly with the bacterial cell wall (Cash et al. 2006). Mice lacking IL-17 and IL-22 have reduced amounts of antimicrobial peptides and are highly susceptible to *C. rodentium* infection; however, they can be rescued by treatment with recombinant REG3 γ (Zheng et al. 2008). Microbiota-derived LPS maintains basal expression of REG3 γ in intestinal epithelial cells and Paneth cells. Indeed, REG3 γ is not detected in germ-free mice (Cash et al. 2006; Mukherjee et al. 2014), and even short-term antibiotic treatment impairs its expression, rendering mice susceptible to VRE infection, a defect that can be reverted by oral administration of LPS (Brandl et al. 2008). Similarly, flagellin from commensal bacteria sensed by TLR5 expressed on CD103⁺ CD11b⁺ DCs found in the LP contributes to the maintenance of REG3 γ expression. Upon TLR5 activation, DCs produce IL-23, promoting IL-22 release by ILCs, and therefore REG3 γ expression in intestinal epithelial cells (Kinnebrew et al. 2010; Kinnebrew et al. 2012). The importance of the IL-22-REG3 γ axis against bacterial infections has been recently demonstrated by Abt et al. (2016). Briefly, these authors showed that Resiquimod (R848), a synthetic ligand for TLR-7 that stimulates antiviral innate immune defenses, induces IL-22 expression, restoring REG3 γ and reestablishing colonization resistance against VRE in antibiotic-treated mice.

Another mechanism by which IL-22 controls opportunistic gut pathogens is by facilitating the binding of complement component C3 to bacteria (Hasegawa et al. 2014). When the intestinal mucosa is infected with *C. difficile*, this organism is capable of translocating to other tissues, contributing to *C. difficile*-induced mortality. However, IL-22 induces an increase of C3 expression and secretion in the liver, which is mediated by the STAT3 pathway. This process leads to a substantial increase in systemic serum levels of C3, which in turn mediates enhanced opsonization of translocated commensals and pathogens, and their subsequent killing and clearance by neutrophils and macrophages (Hasegawa et al. 2014).

Although IL-22 plays a beneficial role in many infection models, recent studies have shown that IL-22 production may also be detrimental to the host. For instance, during *S. Typhimurium* infection, IL-22 does not play a protective role, but is instead exploited by the pathogen to colonize the gut to high levels (Behnsen et al. 2014). IL-22 favors *S. Typhimurium* colonization by inducing antimicrobial proteins that sequester metal ions. As the pathogen is able to overcome metal starvation, it outcompetes related commensals such as *Escherichia coli* (Behnsen et al. 2014). During *Helicobacter pylori* infection, IL-22 also contributes to the development of gastric pathology. Both *H. pylori*-infected humans and mice exhibit an overabundance of IL-22 (Zhuang et al. 2015), which stimulates gastric epithelial cells to secrete CXCL2. CXCL2 then recruits myeloid-derived suppressor cells that produce the proinflammatory proteins S100A8 and S100A9 and inhibit Th1 cell responses, thereby contributing to the development of *H. pylori*-associated gastritis (Zhuang et al. 2015).

CONCLUSIONS

The intestinal tract is home to commensal bacteria and fungi that have coevolved with the mammalian host. In order to maintain intestinal homeostasis, the mucosal immune system must balance between an appropriate response to dangerous pathogens and an inappropriate response to commensal microbiota that breach the epithelial barrier. IL-17 and IL-22 have been shown to play a critical role in maintaining barrier homeostasis against intestinal pathogens and commensal bacteria. It is becoming increasingly apparent that bacterial pathogens trigger a rapid IL-17 and IL-22-dependent innate defense in the gut mucosa. In turn, intestinal production of these cytokines seems to be regulated homeostatically by interactions between the host and the intestinal microbiota (e.g. SFB).

IL-17 and IL-22 enhance basic innate barrier defenses at mucosal surfaces, such as antimicrobial peptide production and neutrophil recruitment, i.e. rapidly occurring events that precede adaptive immunity. Although the functional spectra of these cytokines overlap with regard to inducing an innate immune response in epithelial cells, they are generally different. IL-17 is largely proinflammatory and can be pathogenic, whereas IL-22 primarily exhibits regenerative and protective roles. Nevertheless, IL-22's proinflammatory effects might contribute to pathology, as could be the case in psoriasis (Zheng et al. 2007) or IBD (Brand et al. 2006). Altogether, cytokines IL-17 and IL-22 provide barrier integrity against extracellular pathogens by (i) instructing innate immune responses in tissue cells, (ii) inducing the recruitment of adaptive immune cells via epithelial-derived chemokines and (iii) inducing regeneration of epithelial surfaces after inflammation.

The IL-17 and IL-23 (i.e. the cytokine upstream of both IL-17 and IL-22) pathways are clinical targets for multiple inflammatory diseases. For example, the antibody ustekinumab, which neutralizes the p40 subunit of IL-23 and IL-12, is approved to treat psoriasis and is showing benefit in clinical trials for Crohn's disease (Sandborn et al. 2008). Clinical data on the use of antibodies specific to the IL-23 p19 subunit are beginning to emerge in IBD and also show efficacy (Sandborn et al. 2008). IL-23-specific antibodies are also highly efficacious in psoriasis, suggesting that the efficacy achieved with p40 inhibition is largely a result of IL-23 inhibition (Leonardi et al. 2008; Papp et al. 2008; Croft, Benedict and Ware 2013). Antibodies to IL-17A or IL-17RA are also highly effective in psoriasis (Papp et al. 2012; Langley et al. 2014), but surprisingly, they exacerbated Crohn's disease in a subset of patients (Hueber et al. 2012). Thus, although IL-23 is important for production and maintenance of IL-17A and IL-17F, inhibition of IL-17A or IL-17RA in patients with Crohn's disease does not phenocopy the effects observed with a p40 or p19 inhibitor. As such, the mechanism(s) underlying the differential efficacy observed in Crohn's disease with inhibition of IL-17 versus IL-23 is an important unanswered question.

ACKNOWLEDGEMENTS

We would like to acknowledge the Raffatellu lab for helpful suggestions, and Sean-Paul Nuccio for his help with editing the manuscript.

Work in the MR lab is supported by Public Health Service Grants AI126277, AI121928, AI114625, AI105374, and DK058057. MR holds an Investigator in the Pathogenesis of Infectious Disease Award from the Burroughs Wellcome Fund.

Conflict of interest. None declared.

REFERENCES

- Abt MC, Buffie CG, Susac B et al. TLR-7 activation enhances IL-22-mediated colonization resistance against vancomycin-resistant enterococcus. *Sci Transl Med* 2016;**8**: 327ra25.
- Aggarwal S, Gurney AL. IL-17: prototype member of an emerging cytokine family. *J Leukocyte Biol* 2002;**71**:1-8.
- Albanesi C, Scarponi C, Cavani A et al. Interleukin-17 is produced by both Th1 and Th2 lymphocytes, and modulates interferon-gamma- and interleukin-4-induced activation of human keratinocytes. *J Invest Dermatol* 2000;**115**: 81-7.
- Andoh A, Zhang Z, Inatomi O et al. Interleukin-22, a member of the IL-10 subfamily, induces inflammatory responses in colonic subepithelial myofibroblasts. *Gastroenterology* 2005;**129**:969-84.
- Artis D. Epithelial-cell recognition of commensal bacteria and maintenance of immune homeostasis in the gut. *Nat Rev Immunol* 2008;**8**:411-20.
- Atarashi K, Nishimura J, Shima T et al. ATP drives lamina propria T(H)17 cell differentiation. *Nature* 2008;**455**:808-12.
- Atarashi K, Tanoue T, Ando M et al. Th17 cell induction by adhesion of microbes to intestinal epithelial cells. *Cell* 2015;**163**:367-80.
- Aujla SJ, Chan YR, Zheng M et al. IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nat Med* 2008;**14**:275-81.
- Awane M, Andres PG, Li DJ et al. NF-kappa B-inducing kinase is a common mediator of IL-17-, TNF-alpha-, and IL-1-

- beta-induced chemokine promoter activation in intestinal epithelial cells. *J Immunol* 1999;162:5337–44.
- Behnsen J, Jellbauer S, Wong CP et al. The cytokine IL-22 promotes pathogen colonization by suppressing related commensal bacteria. *Immunity* 2014;40:262–73.
- Berger T, Togawa A, Duncan GS et al. Lipocalin 2-deficient mice exhibit increased sensitivity to *Escherichia coli* infection but not to ischemia-reperfusion injury. *P Natl Acad Sci USA* 2006;103:1834–9.
- Bettelli E, Carrier Y, Gao W et al. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 2006;441:235–8.
- Blaschitz C, Raffatellu M. Th17 cytokines and the gut mucosal barrier. *J Clin Immunol* 2010;30:196–203.
- Bogunovic M, Ginhoux F, Helft J et al. Origin of the lamina propria dendritic cell network. *Immunity* 2009;31:513–25.
- Brand S, Beigel F, Olszak T et al. IL-22 is increased in active Crohn's disease and promotes proinflammatory gene expression and intestinal epithelial cell migration. *Am J Physiol Gastro* 2006;290:G827–38.
- Brandl K, Plitas G, Mihu CN et al. Vancomycin-resistant enterococci exploit antibiotic-induced innate immune deficits. *Nature* 2008;455:804–7.
- Brereton CF, Sutton CE, Ross PJ et al. *Escherichia coli* heat-labile enterotoxin promotes protective Th17 responses against infection by driving innate IL-1 and IL-23 production. *J Immunol* 2011;186:5896–906.
- Buonocore S, Ahern PP, Uhlig HH et al. Innate lymphoid cells drive interleukin-23-dependent innate intestinal pathology. *Nature* 2010;464:1371–5.
- Cash HL, Whitham CV, Behrendt CL et al. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science* 2006;313:1126–30.
- Cella M, Fuchs A, Vermi W et al. A human natural killer cell subset provides an innate source of IL-22 for mucosal immunity. *Nature* 2009;457:722–5.
- Chamaillard M, Hashimoto M, Horie Y et al. An essential role for NOD1 in host recognition of bacterial peptidoglycan containing diaminopimelic acid. *Nat Immunol* 2003;4:702–7.
- Chan JR, Blumenschein W, Murphy E et al. IL-23 stimulates epidermal hyperplasia via TNF and IL-20R2-dependent mechanisms with implications for psoriasis pathogenesis. *J Exp Med* 2006;203:2577–87.
- Chang J, Burkett PR, Borges CM et al. MyD88 is essential to sustain mTOR activation necessary to promote T helper 17 cell proliferation by linking IL-1 and IL-23 signaling. *P Natl Acad Sci USA* 2013;110:2270–5.
- Charrel-Dennis M, Latz E, Halmen KA et al. TLR-independent type I interferon induction in response to an extracellular bacterial pathogen via intracellular recognition of its DNA. *Cell Host Microbe* 2008;4:543–54.
- Chen F, Cao A, Yao S et al. mTOR mediates IL-23 induction of neutrophil IL-17 and IL-22 production. *J Immunol* 2016;196:4390–9.
- Cheroutre H, Lambolez F, Mucida D. The light and dark sides of intestinal intraepithelial lymphocytes. *Nat Rev Immunol* 2011;11:445–56.
- Cho JS, Pietras EM, Garcia NC et al. IL-17 is essential for host defense against cutaneous *Staphylococcus aureus* infection in mice. *J Clin Invest* 2010;120:1762–73.
- Codarri L, Gyulveszi G, Tosevski V et al. ROR γ drives production of the cytokine GM-CSF in helper T cells, which is essential for the effector phase of autoimmune neuroinflammation. *Nat Immunol* 2011;12:560–7.
- Conti HR, Shen F, Nayyar N et al. Th17 cells and IL-17 receptor signaling are essential for mucosal host defense against oral candidiasis. *J Exp Med* 2009;206:299–311.
- Corbin BD, Seeley EH, Raab A et al. Metal chelation and inhibition of bacterial growth in tissue abscesses. *Science* 2008;319:962–5.
- Cosmi L, De Palma R, Santarlasci V et al. Human interleukin 17-producing cells originate from a CD161+CD4+ T cell precursor. *J Exp Med* 2008;205:1903–16.
- Cox JH, Kljavin NM, Ota N et al. Opposing consequences of IL-23 signaling mediated by innate and adaptive cells in chemically induced colitis in mice. *Mucosal Immunol* 2012;5:99–109.
- Crellin NK, Trifari S, Kaplan CD et al. Human NKp44+IL-22+ cells and LT α -like cells constitute a stable ROR γ + lineage distinct from conventional natural killer cells. *J Exp Med* 2010;207:281–90.
- Croft M, Benedict CA, Ware CF. Clinical targeting of the TNF and TNFR superfamilies. *Nat Rev Drug Discov* 2013;12:147–68.
- Cua DJ, Sherlock JP. Autoimmunity's collateral damage: Gut microbiota strikes 'back'. *Nat Med* 2011;17:1055–6.
- Cua DJ, Tato CM. Innate IL-17-producing cells: the sentinels of the immune system. *Nat Rev Immunol* 2010;10:479–89.
- Cupedo T, Crellin NK, Papazian N et al. Human fetal lymphoid tissue-inducer cells are interleukin 17-producing precursors to ROR γ + CD127+ natural killer-like cells. *Nat Immunol* 2009;10:66–74.
- Dalmas E, Venteclaf N, Caer C et al. T cell-derived IL-22 amplifies IL-1 β -driven inflammation in human adipose tissue: relevance to obesity and type 2 diabetes. *Diabetes* 2014;63:1966–77.
- Darlington D, Rogers AW. Epithelial lymphocytes in the small intestine of the mouse. *J Anat* 1966;100:813–30.
- Doisne JM, Soulard V, Becourt C et al. Cutting edge: crucial role of IL-1 and IL-23 in the innate IL-17 response of peripheral lymph node NK1.1-invariant NKT cells to bacteria. *J Immunol* 2011;186:662–6.
- Dong C. TH17 cells in development: an updated view of their molecular identity and genetic programming. *Nat Rev Immunol* 2008;8:337–48.
- Dostert C, Petrilli V, Van Bruggen R et al. Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science* 2008;320:674–7.
- Dubin PJ, Kolls JK. IL-23 mediates inflammatory responses to mucoid *Pseudomonas aeruginosa* lung infection in mice. *Am J Physiol Lung C* 2007;292:L519–28.
- Dudakov JA, Hanash AM, van den Brink MR. Interleukin-22: immunobiology and pathology. *Annu Rev Immunol* 2015;33:747–85.
- Duhen T, Geiger R, Jarrossay D et al. Production of interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T cells. *Nat Immunol* 2009;10:857–63.
- Dumoutier L, Louahed J, Renaud JC. Cloning and characterization of IL-10-related T cell-derived inducible factor (IL-TIF), a novel cytokine structurally related to IL-10 and inducible by IL-9. *J Immunol* 2000;164:1814–9.
- Dumoutier L, Van Roost E, Colau D et al. Human interleukin-10-related T cell-derived inducible factor: molecular cloning and functional characterization as an hepatocyte-stimulating factor. *P Natl Acad Sci USA* 2000;97:10144–9.
- Dungan LS, Mills KH. Caspase-1-processed IL-1 family cytokines play a vital role in driving innate IL-17. *Cytokine* 2011;56:126–32.
- Dunne A, Ross PJ, Pospisilova E et al. Inflammasome activation by adenylate cyclase toxin directs Th17 responses and

- protection against *Bordetella pertussis*. *J Immunol* 2010;**185**:1711–9.
- Eberl G, Marmon S, Sunshine MJ *et al*. An essential function for the nuclear receptor ROR γ (t) in the generation of fetal lymphoid tissue inducer cells. *Nat Immunol* 2004;**5**:64–73.
- El-Behi M, Ciric B, Dai H *et al*. The encephalitogenicity of T(H)17 cells is dependent on IL-1- and IL-23-induced production of the cytokine GM-CSF. *Nat Immunol* 2011;**12**:568–75.
- Ely LK, Fischer S, Garcia KC. Structural basis of receptor sharing by interleukin 17 cytokines. *Nat Immunol* 2009;**10**:1245–51.
- Esplugues E, Huber S, Gagliani N *et al*. Control of TH17 cells occurs in the small intestine. *Nature* 2011;**475**:514–8.
- Eyerich S, Eyerich K, Pennino D *et al*. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* 2009;**119**:3573–85.
- Feng JY, Rao GZ, Liu YP *et al*. Expression of CD1a and CD207 in condyloma acuminatum epidermis. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 2009;**25**:420–2.
- Flierl MA, Rittirsch D, Gao H *et al*. Adverse functions of IL-17A in experimental sepsis. *FASEB J* 2008;**22**:2198–205.
- Flo TH, Smith KD, Sato S *et al*. Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 2004;**432**:917–21.
- Fossiez F, Djossou O, Chomarat P *et al*. T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines. *J Exp Med* 1996;**183**:2593–603.
- Franchi L, Amer A, Body-Malapel M *et al*. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1 β in salmonella-infected macrophages. *Nat Immunol* 2006;**7**:576–82.
- Franchi L, Eigenbrod T, Munoz-Planillo R *et al*. Cytosolic double-stranded RNA activates the NLRP3 inflammasome via MAVS-induced membrane permeabilization and K⁺ efflux. *J Immunol* 2014;**193**:4214–22.
- Fu YX, Chaplin DD. Development and maturation of secondary lymphoid tissues. *Annu Rev Immunol* 1999;**17**:399–433.
- Fu YX, Roark CE, Kelly K *et al*. Immune protection and control of inflammatory tissue necrosis by gamma delta T cells. *J Immunol* 1994;**153**:3101–15.
- Fukata M, Breglio K, Chen A *et al*. The myeloid differentiation factor 88 (MyD88) is required for CD4⁺ T cell effector function in a murine model of inflammatory bowel disease. *J Immunol* 2008;**180**:1886–94.
- Gaboriau-Routhiau V, Rakotobe S, Lecuyer E *et al*. The key role of segmented filamentous bacteria in the coordinated maturation of gut helper T cell responses. *Immunity* 2009;**31**:677–89.
- Gaffen SL. Structure and signalling in the IL-17 receptor family. *Nat Rev Immunol* 2009;**9**:556–67.
- Garlanda C, Dinarello CA, Mantovani A. The interleukin-1 family: back to the future. *Immunity* 2013;**39**:1003–18.
- Gasse P, Riteau N, Vacher R *et al*. IL-1 and IL-23 mediate early IL-17A production in pulmonary inflammation leading to late fibrosis. *PLoS One* 2011;**6**:e23185.
- Gay NJ, Symmons MF, Gangloff M *et al*. Assembly and localization of Toll-like receptor signalling complexes. *Nat Rev Immunol* 2014;**14**:546–58.
- Geddes K, Rubino SJ, Magalhaes JG *et al*. Identification of an innate T helper type 17 response to intestinal bacterial pathogens. *Nat Med* 2011;**17**:837–44.
- Gibson DL, Ma C, Bergstrom KS *et al*. MyD88 signalling plays a critical role in host defence by controlling pathogen burden and promoting epithelial cell homeostasis during *Citrobacter rodentium*-induced colitis. *Cell Microbiol* 2008;**10**:618–31.
- Girardin SE, Boneca IG, Carneiro LA *et al*. Nod1 detects a unique muropeptide from gram-negative bacterial peptidoglycan. *Science* 2003a;**300**:1584–7.
- Girardin SE, Boneca IG, Viala J *et al*. Nod2 is a general sensor of peptidoglycan through muramyl dipeptide (MDP) detection. *J Biol Chem* 2003b;**278**:8869–72.
- Godinez I, Raffatellu M, Chu H *et al*. Interleukin-23 orchestrates mucosal responses to *Salmonella enterica* serotype Typhimurium in the intestine. *Infect Immun* 2009;**77**:387–98.
- Hall JA, Bouladoux N, Sun CM *et al*. Commensal DNA limits regulatory T cell conversion and is a natural adjuvant of intestinal immune responses. *Immunity* 2008;**29**:637–49.
- Hamada S, Umemura M, Shiono T *et al*. IL-17A produced by gammadelta T cells plays a critical role in innate immunity against *Listeria monocytogenes* infection in the liver. *J Immunol* 2008;**181**:3456–63.
- Happel KI, Dubin PJ, Zheng M *et al*. Divergent roles of IL-23 and IL-12 in host defense against *Klebsiella pneumoniae*. *J Exp Med* 2005;**202**:761–9.
- Happel KI, Zheng M, Young E *et al*. Cutting edge: roles of Toll-like receptor 4 and IL-23 in IL-17 expression in response to *Klebsiella pneumoniae* infection. *J Immunol* 2003;**170**:4432–6.
- Harrington LE, Hatton RD, Mangan PR *et al*. Interleukin 17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol* 2005;**6**:1123–32.
- Harris TJ, Grosso JF, Yen HR *et al*. Cutting edge: An in vivo requirement for STAT3 signaling in TH17 development and TH17-dependent autoimmunity. *J Immunol* 2007;**179**:4313–7.
- Hasegawa M, Yada S, Liu MZ *et al*. Interleukin-22 regulates the complement system to promote resistance against pathogens after pathogen-induced intestinal damage. *Immunity* 2014;**41**:620–32.
- Higgins SC, Jarnicki AG, Lavelle EC *et al*. TLR4 mediates vaccine-induced protective cellular immunity to *Bordetella pertussis*: role of IL-17-producing T cells. *J Immunol* 2006;**177**:7980–9.
- Hooper LV, Macpherson AJ. Immune adaptations that maintain homeostasis with the intestinal microbiota. *Nat Rev Immunol* 2010;**10**:159–69.
- Hornung V, Bauernfeind F, Halle A *et al*. Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal destabilization. *Nat Immunol* 2008;**9**:847–56.
- Hueber W, Sands BE, Lewitzky S *et al*. Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* 2012;**61**:1693–700.
- Hughes T, Becknell B, Freud AG *et al*. Interleukin-1 β selectively expands and sustains interleukin-22⁺ immature human natural killer cells in secondary lymphoid tissue. *Immunity* 2010;**32**:803–14.
- Hunter CA. New IL-12-family members: IL-23 and IL-27, cytokines with divergent functions. *Nat Rev Immunol* 2005;**5**:521–31.
- Huppler AR, Verma AH, Conti HR *et al*. Neutrophils do not express IL-17A in the context of acute oropharyngeal candidiasis. *Pathogens* 2015;**4**:559–72.
- Inohara N, Nunez G. The NOD: a signaling module that regulates apoptosis and host defense against pathogens. *Oncogene* 2001;**20**:6473–81.
- Ishigame H, Kakuta S, Nagai T *et al*. Differential roles of interleukin-17A and -17F in host defense against

- mucoepithelial bacterial infection and allergic responses. *Immunity* 2009;**30**:108–19.
- Ismail AS, Behrendt CL, Hooper LV. Reciprocal interactions between commensal bacteria and gamma delta intraepithelial lymphocytes during mucosal injury. *J Immunol* 2009;**182**:3047–54.
- Ivanov II, Atarashi K, Manel N et al. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 2009;**139**:485–98.
- Ivanov II, Frutos Rde L, Manel N et al. Specific microbiota direct the differentiation of IL-17-producing T-helper cells in the mucosa of the small intestine. *Cell Host Microbe* 2008;**4**:337–49.
- Ivanov II, McKenzie BS, Zhou L et al. The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17+ T helper cells. *Cell* 2006;**126**:1121–33.
- Iwakura Y, Ishigame H, Saijo S et al. Functional specialization of interleukin-17 family members. *Immunity* 2011;**34**:149–62.
- Jensen KD, Su X, Shin S et al. Thymic selection determines gamma delta T cell effector fate: antigen-naive cells make interleukin-17 and antigen-experienced cells make interferon gamma. *Immunity* 2008;**29**:90–100.
- Kailasan Vanaja S, Rathinam VA, Atianand MK et al. Bacterial RNA:DNA hybrids are activators of the NLRP3 inflammasome. *P Natl Acad Sci USA* 2014;**111**:7765–70.
- Kang JY, Nan X, Jin MS et al. Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity* 2009;**31**:873–84.
- Kanneganti TD, Ozoren N, Body-Malapel M et al. Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature* 2006;**440**:233–6.
- Kawai T, Akira S. Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* 2011;**34**:637–50.
- Keestra AM, Godinez I, Xavier MN et al. Early MyD88-dependent induction of interleukin-17A expression during *Salmonella* colitis. *Infect Immun* 2011;**79**:3131–40.
- Kehl-Fie TE, Skaar EP. Nutritional immunity beyond iron: a role for manganese and zinc. *Curr Opin Chem Biol* 2010;**14**:218–24.
- Khader SA, Bell GK, Pearl JE et al. IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell responses after vaccination and during *Mycobacterium tuberculosis* challenge. *Nat Immunol* 2007;**8**:369–77.
- Kim S, Han S, Withers DR et al. CD117(+) CD3(-) CD56(-) OX40Lhigh cells express IL-22 and display an LT α phenotype in human secondary lymphoid tissues. *Eur J Immunol* 2011;**41**:1563–72.
- Kinnebrew MA, Buffie CG, Diehl GE et al. Interleukin 23 production by intestinal CD103(+)CD11b(+) dendritic cells in response to bacterial flagellin enhances mucosal innate immune defense. *Immunity* 2012;**36**:276–87.
- Kinnebrew MA, Ubeda C, Zenewicz LA et al. Bacterial flagellin stimulates Toll-like receptor 5-dependent defense against vancomycin-resistant *Enterococcus* infection. *J Infect Dis* 2010;**201**:534–43.
- Kleinschek MA, Boniface K, Sadekova S et al. Circulating and gut-resident human Th17 cells express CD161 and promote intestinal inflammation. *J Exp Med* 2009;**206**:525–34.
- Kolls JK, Linden A. Interleukin-17 family members and inflammation. *Immunity* 2004;**21**:467–76.
- Kondo T, Takata H, Matsuki F et al. Cutting edge: Phenotypic characterization and differentiation of human CD8+ T cells producing IL-17. *J Immunol* 2009;**182**:1794–8.
- Korn T, Bettelli E, Gao W et al. IL-21 initiates an alternative pathway to induce proinflammatory T(H)17 cells. *Nature* 2007;**448**:484–7.
- Korn T, Bettelli E, Oukka M et al. IL-17 and Th17 Cells. *Annu Rev Immunol* 2009;**27**:485–517.
- Kreuger J, Phillipson M. Targeting vascular and leukocyte communication in angiogenesis, inflammation and fibrosis. *Nat Rev Drug Discov* 2016;**15**:125–42.
- Langley RG, Elewski BE, Lebwohl M et al. Secukinumab in plaque psoriasis—results of two phase 3 trials. *New Engl J Med* 2014;**371**:326–38.
- Langrish CL, Chen Y, Blumenschein WM et al. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J Exp Med* 2005;**201**:233–40.
- Latz E, Schoenemeyer A, Visintin A et al. TLR9 signals after translocating from the ER to CpG DNA in the lysosome. *Nat Immunol* 2004;**5**:190–8.
- Latz E, Xiao TS, Stutz A. Activation and regulation of the inflammasomes. *Nat Rev Immunol* 2013;**13**:397–411.
- Le Bourhis L, Guerri L, Dusseaux M et al. Mucosal-associated invariant T cells: unconventional development and function. *Trends Immunol* 2011;**32**:212–8.
- Lebeis SL, Bommarius B, Parkos CA et al. TLR signaling mediated by MyD88 is required for a protective innate immune response by neutrophils to *Citrobacter rodentium*. *J Immunol* 2007;**179**:566–77.
- Lee YK, Menezes JS, Umesaki Y et al. Proinflammatory T-cell responses to gut microbiota promote experimental autoimmune encephalomyelitis. *P Natl Acad Sci USA* 2011;**108** (Suppl 1):4615–22.
- Lee YS, Yang H, Yang JY et al. Interleukin-1 (IL-1) signaling in intestinal stromal cells controls KC/ CXCL1 secretion, which correlates with recruitment of IL-22-secreting neutrophils at early stages of *Citrobacter rodentium* infection. *Infect Immun* 2015;**83**:3257–67.
- Leonardi CL, Kimball AB, Papp KA et al. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomized, double-blind, placebo-controlled trial (PHOENIX 1). *Lancet* 2008;**371**:1665–74.
- Leoni G, Neumann PA, Sumagin R et al. Wound repair: role of immune-epithelial interactions. *Mucosal Immunol* 2015;**8**:959–68.
- Levy M, Thaiss CA, Zeevi D et al. Microbiota-modulated metabolites shape the intestinal microenvironment by regulating NLRP6 inflammasome signaling. *Cell* 2015;**163**:1428–43.
- Lewis KL, Caton ML, Bogunovic M et al. Notch2 receptor signaling controls functional differentiation of dendritic cells in the spleen and intestine. *Immunity* 2011;**35**:780–91.
- Li J, Tomkinson KN, Tan XY et al. Temporal associations between interleukin 22 and the extracellular domains of IL-22R and IL-10R2. *Int Immunopharmacol* 2004;**4**:693–708.
- Liang SC, Nickerson-Nutter C, Pittman DD et al. IL-22 induces an acute-phase response. *J Immunol* 2010;**185**:5531–8.
- Liang SC, Tan XY, Luxenberg DP et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* 2006;**203**:2271–9.
- Liu JZ, Jellbauer S, Poe AJ et al. Zinc sequestration by the neutrophil protein calprotectin enhances *Salmonella* growth in the inflamed gut. *Cell Host Microbe* 2012;**11**:227–39.

- Liu S, Song X, Chrnyk BA et al. Crystal structures of interleukin 17A and its complex with IL-17 receptor A. *Nat Commun* 2013;4:1888.
- Logsdon NJ, Jones BC, Josephson K et al. Comparison of interleukin-22 and interleukin-10 soluble receptor complexes. *J Interf Cytok Res* 2002;22:1099–112.
- Lu YC, Yeh WC, Ohashi PS. LPS/TLR4 signal transduction pathway. *Cytokine* 2008;42:145–51.
- Luci C, Reynders A, Ivanov II et al. Influence of the transcription factor RORgammat on the development of NKp46+ cell populations in gut and skin. *Nat Immunol* 2009;10:75–82.
- Ma HL, Liang S, Li J et al. IL-22 is required for Th17 cell-mediated pathology in a mouse model of psoriasis-like skin inflammation. *J Clin Invest* 2008;118:597–607.
- Macdonald TT, Monteleone G. Immunity, inflammation, and allergy in the gut. *Science* 2005;307:1920–5.
- McKinley L, Alcorn JF, Peterson A et al. TH17 cells mediate steroid-resistant airway inflammation and airway hyperresponsiveness in mice. *J Immunol* 2008;181:4089–97.
- Mangan PR, Harrington LE, O'Quinn DB et al. Transforming growth factor-beta induces development of the T(H)17 lineage. *Nature* 2006;441:231–4.
- Manzanillo P, Eidenschenk C, Ouyang W. Deciphering the crosstalk among IL-1 and IL-10 family cytokines in intestinal immunity. *Trends Immunol* 2015;36:471–8.
- Marina-Garcia N, Franchi L, Kim YG et al. Pannexin-1-mediated intracellular delivery of muramyl dipeptide induces caspase-1 activation via cryopyrin/NLRP3 independently of Nod2. *J Immunol* 2008;180:4050–7.
- Martin B, Hirota K, Cua DJ et al. Interleukin-17-producing gamma-delta T cells selectively expand in response to pathogen products and environmental signals. *Immunity* 2009;31:321–30.
- Matthews JD, Weight CM, Parkos CA. Leukocyte-epithelial interactions and mucosal homeostasis. *Toxicol Pathol* 2014;42:91–8.
- Maynard CL, Elson CO, Hatton RD et al. Reciprocal interactions of the intestinal microbiota and immune system. *Nature* 2012;489:231–41.
- Medzhitov R, Preston-Hurlburt P, Kopp E et al. MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol Cell* 1998;2:253–8.
- Meeks KD, Sieve AN, Kolls JK et al. IL-23 is required for protection against systemic infection with *Listeria monocytogenes*. *J Immunol* 2009;183:8026–34.
- Miao EA, Alpuche-Aranda CM, Dors M et al. Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1beta via Ipaf. *Nat Immunol* 2006;7:569–75.
- Miao EA, Mao DP, Yudkovsky N et al. Innate immune detection of the type III secretion apparatus through the NLRC4 inflammasome. *P Natl Acad Sci USA* 2010;107:3076–80.
- Mombaerts P, Arnoldi J, Russ F et al. Different roles of alpha beta and gamma delta T cells in immunity against an intracellular bacterial pathogen. *Nature* 1993;365:53–6.
- Montaldo E, Teixeira-Alves LG, Glatzer T et al. Human RORgammat(+)/CD34(+) cells are lineage-specified progenitors of group 3 RORgammat(+) innate lymphoid cells. *Immunity* 2014;41:988–1000.
- Moran EM, Mullan R, McCormick J et al. Human rheumatoid arthritis tissue production of IL-17A drives matrix and cartilage degradation: synergy with tumour necrosis factor-alpha, Oncostatin M and response to biologic therapies. *Arthritis Res Ther* 2009;11:R113.
- Mouy R, Fischer A, Vilmer E et al. Incidence, severity, and prevention of infections in chronic granulomatous disease. *J Pediatr* 1989;114:555–60.
- Mowat AM, Agace WW. Regional specialization within the intestinal immune system. *Nat Rev Immunol* 2014;14:667–85.
- Mukherjee S, Zheng H, Derebe MG et al. Antibacterial membrane attack by a pore-forming intestinal C-type lectin. *Nature* 2014;505:103–7.
- Munoz M, Eidenschenk C, Ota N et al. Interleukin-22 induces interleukin-18 expression from epithelial cells during intestinal infection. *Immunity* 2015;42:321–31.
- Nogralas KE, Zaba LC, Guttman-Yassky E et al. Th17 cytokines interleukin (IL)-17 and IL-22 modulate distinct inflammatory and keratinocyte-response pathways. *Brit J Dermatol* 2008;159:1092–102.
- Nogralas KE, Zaba LC, Shemer A et al. IL-22-producing "T22" T cells account for upregulated IL-22 in atopic dermatitis despite reduced IL-17-producing TH17 T cells. *J Allergy Clin Immunol* 2009;123:1244–52.e2.
- Norian LA, Rodriguez PC, O'Mara LA et al. Tumor-infiltrating regulatory dendritic cells inhibit CD8+ T cell function via L-arginine metabolism. *Cancer Res* 2009;69:3086–94.
- Nowarski R, Jackson R, Gagliani N et al. Epithelial IL-18 equilibrium controls barrier function in colitis. *Cell* 2015;163:1444–56.
- Ortega C, Fernandez AS, Carrillo JM et al. IL-17-producing CD8+ T lymphocytes from psoriasis skin plaques are cytotoxic effector cells that secrete Th17-related cytokines. *J Leukocyte Biol* 2009;86:435–43.
- Ota N, Wong K, Valdez PA et al. IL-22 bridges the lymphotoxin pathway with the maintenance of colonic lymphoid structures during infection with *Citrobacter rodentium*. *Nat Immunol* 2011;12:941–8.
- Ouyang W, Kolls JK, Zheng Y. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* 2008;28:454–67.
- Pandya AD, Al-Jaderi Z, Högglund RA et al. Identification of human NK17/NK1 cells. *PLoS One* 2011;6:e26780.
- Papp KA, Langley RG, Lebwohl M et al. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 52-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 2). *Lancet* 2008;371:1675–84.
- Papp KA, Leonardi C, Menter A et al. Brodalumab, an anti-interleukin-17-receptor antibody for psoriasis. *New Engl J Med* 2012;366:1181–9.
- Park H, Li Z, Yang XO et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nat Immunol* 2005;6:1133–41.
- Peng MY, Wang ZH, Yao CY et al. Interleukin 17-producing gamma delta T cells increased in patients with active pulmonary tuberculosis. *Cell Mol Immunol* 2008;5:203–8.
- Pestka S, Krause CD, Sarkar D et al. Interleukin-10 and related cytokines and receptors. *Annu Rev Immunol* 2004;22:929–79.
- Petermann F, Rothhammer V, Claussen MC et al. gamma-delta T cells enhance autoimmunity by restraining regulatory T cell responses via an interleukin-23-dependent mechanism. *Immunity* 2010;33:351–63.
- Pickert G, Neufert C, Leppkes M et al. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J Exp Med* 2009;206:1465–72.

- Priebe GP, Walsh RL, Cederroth TA et al. IL-17 is a critical component of vaccine-induced protection against lung infection by lipopolysaccharide-heterologous strains of *Pseudomonas aeruginosa*. *J Immunol* 2008;**181**:4965–75.
- Qu N, Xu M, Mizoguchi I et al. Pivotal roles of T-helper 17-related cytokines, IL-17, IL-22, and IL-23, in inflammatory diseases. *Clin Dev Immunol* 2013;**2013**:968549.
- Rachitskaya AV, Hansen AM, Horai R et al. Cutting edge: NKT cells constitutively express IL-23 receptor and RORgammat and rapidly produce IL-17 upon receptor ligation in an IL-6-independent fashion. *J Immunol* 2008;**180**:5167–71.
- Raffatellu M, George MD, Akiyama Y et al. Lipocalin-2 resistance confers an advantage to *Salmonella enterica* serotype Typhimurium for growth and survival in the inflamed intestine. *Cell Host Microbe* 2009;**5**:476–86.
- Raffatellu M, Santos RL, Verhoeven DE et al. Simian immunodeficiency virus-induced mucosal interleukin-17 deficiency promotes *Salmonella* dissemination from the gut. *Nat Med* 2008;**14**:421–8.
- Ribot JC, deBarros A, Pang DJ et al. CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat Immunol* 2009;**10**:427–36.
- Rickel EA, Siegel LA, Yoon BR et al. Identification of functional roles for both IL-17RB and IL-17RA in mediating IL-25-induced activities. *J Immunol* 2008;**181**:4299–310.
- Riera-Sans L, Behrens A. Regulation of alphabeta/gammadelta T cell development by the activator protein 1 transcription factor c-Jun. *J Immunol* 2007;**178**:5690–700.
- Riol-Blanco L, Lazarevic V, Awasthi A et al. IL-23 receptor regulates unconventional IL-17-producing T cells that control bacterial infections. *J Immunol* 2010;**184**:1710–20.
- Roark CL, French JD, Taylor MA et al. Exacerbation of collagen-induced arthritis by oligoclonal, IL-17-producing gamma delta T cells. *J Immunol* 2007;**179**:5576–83.
- Saijo S, Ikeda S, Yamabe K et al. Dectin-2 recognition of alpha-mannans and induction of Th17 cell differentiation is essential for host defense against *Candida albicans*. *Immunity* 2010;**32**:681–91.
- Sandborn WJ, Feagan BG, Fedorak RN et al. A randomized trial of Ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with moderate-to-severe Crohn's disease. *Gastroenterology* 2008;**135**:1130–41.
- Sander LE, Davis MJ, Boekschoten MV et al. Detection of prokaryotic mRNA signifies microbial viability and promotes immunity. *Nature* 2011;**474**:385–9.
- Sano T, Huang W, Hall JA et al. An IL-23R/IL-22 circuit regulates epithelial serum amyloid A to promote local effector Th17 responses. *Cell* 2015;**163**:381–93.
- Sanos SL, Bui VL, Mortha A et al. RORgammat and commensal microflora are required for the differentiation of mucosal interleukin 22-producing NKp46+ cells. *Nat Immunol* 2009;**10**:83–91.
- Satoh-Takayama N, Dumoutier L, Lesjean-Pottier S et al. The natural cytotoxicity receptor NKp46 is dispensable for IL-22-mediated innate intestinal immune defense against *Citrobacter rodentium*. *J Immunol* 2009;**183**:6579–87.
- Satoh-Takayama N, Lesjean-Pottier S, Sawa S et al. Lymphotoxin-beta receptor-independent development of intestinal IL-22-producing NKp46+ innate lymphoid cells. *Eur J Immunol* 2011;**41**:780–6.
- Satoh-Takayama N, Voshenrich CA, Lesjean-Pottier S et al. Microbial flora drives interleukin 22 production in intestinal NKp46+ cells that provide innate mucosal immune defense. *Immunity* 2008;**29**:958–70.
- Schulz O, Jaensson E, Persson EK et al. Intestinal CD103+, but not CX3CR1+, antigen sampling cells migrate in lymph and serve classical dendritic cell functions. *J Exp Med* 2009;**206**:3101–14.
- Sellge G, Magalhaes JG, Konradt C et al. Th17 cells are the dominant T cell subtype primed by *Shigella flexneri* mediating protective immunity. *J Immunol* 2010;**184**:2076–85.
- Sha W, Mitoma H, Hanabuchi S et al. Human NLRP3 inflammasomes sense multiple types of bacterial RNAs. *P Natl Acad Sci USA* 2014;**111**:16059–64.
- Shibata K, Yamada H, Hara H et al. Resident Vdelta1+ gammadelta T cells control early infiltration of neutrophils after *Escherichia coli* infection via IL-17 production. *J Immunol* 2007;**178**:4466–72.
- Shibata K, Yamada H, Nakamura R et al. Identification of CD25+ gamma delta T cells as fetal thymus-derived naturally occurring IL-17 producers. *J Immunol* 2008;**181**:5940–7.
- Shin HC, Benbernou N, Esnault S et al. Expression of IL-17 in human memory CD45RO+ T lymphocytes and its regulation by protein kinase A pathway. *Cytokine* 1999;**11**:257–66.
- Siegmund S, Schutze N, Freudenberg MA et al. Production of IL-12, IL-23 and IL-27p28 by bone marrow-derived conventional dendritic cells rather than macrophages after LPS/TLR4-dependent induction by *Salmonella* Enteritidis. *Immunobiology* 2007;**212**:739–50.
- Song X, Zhu S, Shi P et al. IL-17RE is the functional receptor for IL-17C and mediates mucosal immunity to infection with intestinal pathogens. *Nat Immunol* 2011;**12**:1151–8.
- Sonnenberg GF, Monticelli LA, Alenghat T et al. Innate lymphoid cells promote anatomical containment of lymphoid-resident commensal bacteria. *Science* 2012;**336**:1321–5.
- Sonnenberg GF, Monticelli LA, Elloso MM et al. CD4(+) lymphoid tissue-inducer cells promote innate immunity in the gut. *Immunity* 2011;**34**:122–34.
- Spahn TW, Maaser C, Eckmann L et al. The lymphotoxin-beta receptor is critical for control of murine *Citrobacter rodentium*-induced colitis. *Gastroenterology* 2004;**127**:1463–73.
- Spits H, Cupedo T. Innate lymphoid cells: emerging insights in development, lineage relationships, and function. *Annu Rev Immunol* 2012;**30**:647–75.
- Sugimoto K, Ogawa A, Mizoguchi E et al. IL-22 ameliorates intestinal inflammation in a mouse model of ulcerative colitis. *J Clin Invest* 2008;**118**:534–44.
- Sutton CE, Lalor SJ, Sweeney CM et al. Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells, amplifying Th17 responses and autoimmunity. *Immunity* 2009;**31**:331–41.
- Takatori H, Kanno Y, Watford WT et al. Lymphoid tissue inducer-like cells are an innate source of IL-17 and IL-22. *J Exp Med* 2009;**206**:35–41.
- Taylor PR, Roy S, Leal SM et al. Activation of neutrophils by autocrine IL-17A-IL-17RC interactions during fungal infection is regulated by IL-6, IL-23, RORgammat and dectin-2. *Nat Immunol* 2014;**15**:143–51.
- Teunissen MB, Koomen CW, de Waal Malefyt R et al. Interleukin-17 and interferon-gamma synergize in the enhancement of proinflammatory cytokine production by human keratinocytes. *J Invest Dermatol* 1998;**111**:645–9.
- Ting JP, Lovering RC, Alnemri ES et al. The NLR gene family: a standard nomenclature. *Immunity* 2008;**28**:285–7.

- Toy D, Kugler D, Wolfson M et al. Cutting edge: interleukin 17 signals through a heteromeric receptor complex. *J Immunol* 2006;**177**:36–9.
- Trifari S, Kaplan CD, Tran EH et al. Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat Immunol* 2009;**10**:864–71.
- Tumanov AV, Christiansen PA, Fu YX. The role of lymphotoxin receptor signaling in diseases. *Curr Mol Med* 2007;**7**:567–78.
- Tumanov AV, Koroleva EP, Guo X et al. Lymphotoxin controls the IL-22 protection pathway in gut innate lymphoid cells during mucosal pathogen challenge. *Cell Host Microbe* 2011;**10**:44–53.
- Turner JE, Stockinger B, Helmby H. IL-22 mediates goblet cell hyperplasia and worm expulsion in intestinal helminth infection. *PLoS Pathog* 2013;**9**:e1003698.
- Umemura M, Yahagi A, Hamada S et al. IL-17-mediated regulation of innate and acquired immune response against pulmonary *Mycobacterium bovis* bacille Calmette-Guerin infection. *J Immunol* 2007;**178**:3786–96.
- Upadhyay V, Fu YX. Lymphotoxin signalling in immune homeostasis and the control of microorganisms. *Nat Rev Immunol* 2013;**13**:270–9.
- van de Veerdonk FL, Joosten LA, Shaw PJ et al. The inflammasome drives protective Th1 and Th17 cellular responses in disseminated candidiasis. *Eur J Immunol* 2011;**41**:2260–8.
- Velin D, Favre L, Bernasconi E et al. Interleukin-17 is a critical mediator of vaccine-induced reduction of *Helicobacter* infection in the mouse model. *Gastroenterology* 2009;**136**:2237–46 e1.
- Wang Y, Koroleva EP, Kruglov AA et al. Lymphotoxin beta receptor signaling in intestinal epithelial cells orchestrates innate immune responses against mucosal bacterial infection. *Immunity* 2010;**32**:403–13.
- Wang Y, Liu L. The membrane protein of severe acute respiratory syndrome coronavirus functions as a novel cytosolic pathogen-associated molecular pattern to promote beta interferon induction via a Toll-like-receptor-related TRAF3-independent mechanism. *MBio* 2016;**7**:e01872–15.
- Ware CF. Network communications: lymphotoxins, LIGHT, and TNF. *Annu Rev Immunol* 2005;**23**:787–819.
- Werner JL, Gessner MA, Lilly LM et al. Neutrophils produce interleukin 17A (IL-17A) in a dectin-1- and IL-23-dependent manner during invasive fungal infection. *Infect Immun* 2011;**79**:3966–77.
- Winkelstein JA, Marino MC, Johnston RB et al. Chronic granulomatous disease. Report on a national registry of 368 patients. *Medicine (Baltimore)* 2000;**79**:155–69.
- Wolk K, Kunz S, Asadullah K et al. Cutting edge: immune cells as sources and targets of the IL-10 family members? *J Immunol* 2002;**168**:5397–402.
- Wolk K, Kunz S, Witte E et al. IL-22 increases the innate immunity of tissues. *Immunity* 2004;**21**:241–54.
- Wolk K, Warszawska K, Hoeflich C et al. Deficiency of IL-22 contributes to a chronic inflammatory disease: pathogenetic mechanisms in acne inversa. *J Immunol* 2011;**186**:1228–39.
- Wolk K, Witte E, Hoffmann U et al. IL-22 induces lipopolysaccharide-binding protein in hepatocytes: a potential systemic role of IL-22 in Crohn's disease. *J Immunol* 2007;**178**:5973–81.
- Wu HJ, Ivanov II, Darce J et al. Gut-residing segmented filamentous bacteria drive autoimmune arthritis via T helper 17 cells. *Immunity* 2010;**32**:815–27.
- Wu Q, Martin RJ, Rino JG et al. IL-23-dependent IL-17 production is essential in neutrophil recruitment and activity in mouse lung defense against respiratory *Mycoplasma pneumoniae* infection. *Microbes Infect* 2007;**9**:78–86.
- Xie MH, Aggarwal S, Ho WH et al. Interleukin (IL)-22, a novel human cytokine that signals through the interferon receptor-related proteins CRF2-4 and IL-22R. *J Biol Chem* 2000;**275**:31335–9.
- Yang XO, Panopoulos AD, Nurieva R et al. STAT3 regulates cytokine-mediated generation of inflammatory helper T cells. *J Biol Chem* 2007;**282**:9358–63.
- Yao Z, Fanslow WC, Seldin MF et al. Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 1995;**3**:811–21.
- Ye P, Garvey PB, Zhang P et al. Interleukin-17 and lung host defense against *Klebsiella pneumoniae* infection. *Am J Resp Cell Mol* 2001a;**25**:335–40.
- Ye P, Rodriguez FH, Kanaly S et al. Requirement of interleukin 17 receptor signaling for lung CXC chemokine and granulocyte colony-stimulating factor expression, neutrophil recruitment, and host defense. *J Exp Med* 2001b;**194**:519–27.
- Yen HR, Harris TJ, Wada S et al. Tc17 CD8 T cells: functional plasticity and subset diversity. *J Immunol* 2009;**183**:7161–8.
- Zhang X, Gao L, Lei L et al. A MyD88-dependent early IL-17 production protects mice against airway infection with the obligate intracellular pathogen *Chlamydia muridarum*. *J Immunol* 2009;**183**:1291–300.
- Zheng Y, Danilenko DM, Valdez P et al. Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* 2007;**445**:648–51.
- Zheng Y, Valdez PA, Danilenko DM et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat Med* 2008;**14**:282–9.
- Zhu J, Emerson SG. Hematopoietic cytokines, transcription factors and lineage commitment. *Oncogene* 2002;**21**:3295–313.
- Zhuang Y, Cheng P, Liu XF et al. A pro-inflammatory role for Th22 cells in *Helicobacter pylori*-associated gastritis. *Gut* 2015;**64**:1368–78.
- Zindl CL, Lai JF, Lee YK et al. IL-22-producing neutrophils contribute to antimicrobial defense and restitution of colonic epithelial integrity during colitis. *P Natl Acad Sci USA* 2013;**110**:12768–73.