UC Davis

UC Davis Previously Published Works

Title

Lipidomic Analysis of Oxidized Fatty Acids in Plant and Algae Oils.

Permalink

<https://escholarship.org/uc/item/15m9d7pd>

Journal

Journal of agricultural and food chemistry, 65(9)

ISSN

0021-8561

Authors

Richardson, Christine E
Hennebelle, Marie
Otoki, Yurika
et al.

Publication Date

2017-03-01

DOI

10.1021/acs.jafc.6b05559

Peer reviewed

Lipidomic Analysis of Oxidized Fatty Acids in Plant and Algae Oils

Christine E. Richardson,[†] Marie Hennebelle,[‡] Yurika Otoki,^{‡,§} Daisy Zamora,^{||} Jun Yang,[⊥]
Bruce D. Hammock,[⊥] and Ameer Y. Taha^{*,†,‡,§}[†]Graduate Group in Nutritional Biology, College of Agriculture and Environmental Sciences; [‡]Department of Food Science and Technology, College of Agriculture and Environmental Sciences; and [⊥]Department of Entomology and UC Davis Comprehensive Cancer Center, University of California, Davis, California 95616, United States[§]Food and Biodynamic Chemistry Laboratory, Graduate School of Agricultural Science, Tohoku University, Sendai 981-8555, Japan^{||}Department of Psychiatry, University of North Carolina—Chapel Hill, North Carolina 27516, United States

S Supporting Information

ABSTRACT: Linoleic acid (LA) and α -linolenic acid (ALA) in plant or algae oils are precursors to oxidized fatty acid metabolites known as oxylipins. Liquid chromatography tandem mass spectrometry was used to quantify oxylipins in soybean, corn, olive, canola, and four high-oleic acid algae oils at room temperature or after heating for 10 min at 100 °C. Flaxseed oil oxylipin concentrations were determined in a follow-up experiment that compared it to soybean, canola, corn, and olive oil. Published consumption data for soybean, canola, corn, and olive oil were used to estimate daily oxylipin intake. The LA and ALA fatty acid composition of the oils was generally related to their respective oxylipin metabolites, except for olive and flaxseed oil, which had higher LA derived monohydroxy and ketone oxylipins than other oils, despite their low LA content. Algae oils had the least amount of oxylipins. The change in oxylipin concentrations was not significantly different among the oils after short-term heating. The estimated oxylipin intake from nonheated soybean, canola, corn, and olive oil was 1.1 mg per person per day. These findings suggest that oils represent a dietary source of LA and ALA derived oxylipins and that the response of oils to short-term heating does not differ among the various oils.

KEYWORDS: linoleic acid, α -linolenic acid, oxidized fatty acids, oxylipins, lipidomics, plant oils, soybean, corn, canola, flaxseed, olive, algae oils, mass spectrometry

■ INTRODUCTION

Plant oils are a major dietary source of the essential polyunsaturated fatty acids, linoleic acid (LA, 18:2n-6) and α -linolenic acid (ALA, 18:3n-3). LA and ALA are required for infant development and supporting optimal nutrition. The estimated requirements are 0.5–1% of calories for ALA and 1–2% for LA.^{1,2} Currently, however, LA accounts for approximately 7% of caloric intake in the US due to the widespread consumption of high-LA oils such as soybean and corn.³ The estimated US intake of ALA is 0.72% energy,³ provided mainly through soybean oil and, to a lesser extent, canola oil.³

LA and ALA are precursors to bioactive oxidized fatty acid metabolites known as oxylipins. In vivo, LA derived oxylipins are involved in inflammatory cascades, pain perception, and skin barrier integrity.^{4–7} Little is known about the role of ALA derived oxylipins in mammals, although plants produce them enzymatically to regulate root development and defense against pathogens.^{8,9}

Oxylipins can be synthesized *in vivo* from their precursor fatty acids via oxygenase enzymes,^{10–12} or obtained through food or dietary oils.^{13–16} Ingested hydroxy, epoxy, and dihydroxy oxylipins are bioavailable.^{17–21} LA derived hydroperoxides have been shown to degrade in the stomach into aldehydes, which accumulate in liver.^{22,23}

The oxidation of fatty acids in oils is nonenzymatic,²⁴ although enzymatic oxidation via lipoxygenase enzymes occurs during the extraction of oils from seed sources.²⁵ Studies reported the

formation of LA derived hydroxy, hydroperoxy, ketone, or epoxy metabolites in oils heated at 40, 100, or 180 °C for 10–264 h or up to 156 days.^{13–15} Oils with higher LA content, such as safflower oil, produced more LA derived oxylipins as compared to low-LA oils.^{13–15} In these studies, oxylipins were detected after prolonged heating of the oils but not at room temperature, likely because the analytical methods involving high performance liquid chromatography with ultraviolet detection, gas chromatography coupled to a flame ionization detection (GC-FID) or nuclear magnetic resonance lacked sensitivity.^{13–15} Fankhauser-Noti et al., however, detected the presence of LA derived monoepoxy and diepoxy fatty acids with GC-FID in fresh (nonheated) and 4-year-old olive and sunflower oil.²⁶ We are not aware of studies that reported on ALA derived oxylipins in oils.

The advent of lipidomic analysis with ultrahigh pressure liquid chromatography tandem mass spectrometry (UPLC-MS/MS) has enabled the separation, detection, and quantitation of many oxylipin species at picomolar concentrations.²⁷ To gain detailed analytical insight into the type and quantity of oxylipins in oils, the aim of the present study was to measure LA and ALA derived oxylipins in plant and algae oils containing varying amounts of LA and ALA at room temperature or after heating for 10 min with UPLC-MS/MS.²⁷ Algae oils were tested because they are an

Received: December 12, 2016

Revised: January 30, 2017

Accepted: February 3, 2017

Published: February 3, 2017

emerging and potentially sustainable source of dietary fat.²⁸ Like plants, algae synthesize unsaturated fatty acids and contain oxygenase enzymes that may contribute to lipid oxidation during the extraction.²⁹ Oxylipins were measured in oils at room temperature or after heating for 10 min at 100 °C to simulate low-heat cooking or simmering conditions. This allowed us to test whether highly unsaturated oils were more vulnerable to oxidation compared to less saturated oils. Daily intake of LA and ALA derived oxylipins was also estimated based on published consumption data of commonly consumed plant oils in the US.³

We hypothesized that concentrations of LA and ALA derived oxylipins at room temperature will be proportional to their precursor fatty acid concentrations in oils, and that high LA or ALA oils will produce more oxylipins following short-term heating for 10 min compared to low LA or ALA oils. Understanding the dietary contribution of oils to potentially bioactive oxylipins is likely to have significant implications for human health and disease.

MATERIALS AND METHODS

Chemicals and Reagents. Methanol, ethyl acetate, chloroform, toluene, and hexane were obtained from Fisher Scientific (Hampton, NH). Methanol was LC-MS grade, whereas all other solvents were HPLC grade. Acetic acid, butylated hydroxytoluene, sodium carbonate, glycerol, triphenylphosphine, ethylenediaminetetraacetic acid (EDTA), and hydrochloric acid were purchased from Sigma-Aldrich (St. Louis, MO). Oxylipin standards were purchased from Cayman Chemicals (Cayman Chemical, Ann Arbor, MI) or synthesized by Dr. Bruce Hammock's lab at UC Davis. The synthetic standards included the following ALA derived oxylipins: 15(16)-EpODE, 12(13)-EpODE, 9(10)-EpODE, 15,16-DiHODE, 12,13-DiHODE and 9,10-DiHODE. Fatty acid standards were purchased from NuCheck Prep (Elysian, MN). Oils were purchased from local stores.

General Study Design. Two experiments were carried out to test the aforementioned hypothesis. In the first experiment, oxylipins and fatty acids were measured in soybean, corn, canola and olive oil as well as 4 algae oils provided by TerraVia Holdings, Inc. that were high in oleic acid, low in LA and ALA, and either lacked or contained 1000 ppm of mixed tocopherols (FORTIUM MT70 IP Liquid from Kemin IA, USA) as an antioxidant. The algae oils tested were a High Stability Algae Oil (HSAO), HSAO without added antioxidants (w/AO), Ultra Omega-9 Algae oil ("Thrive"), and "Thrive" w/AO. Oxylipins were measured at room temperature or after heating for 10 min at 100 °C. Heat was applied at 100 °C (212 °F), because it is below the smoke point range of the oils used in the present study (140–244 °C; 280–471 °F).^{30,31}

In the second experiment, we confirmed oxylipin concentrations in off-the-shelf oils tested in [Experiment 1](#) (corn, canola, soybean and olive oil), and compared them to flaxseed (linseed) oil, which is prone to oxidation due to its high ALA content (54% of total fatty acids).³² Estimated oxylipin intake levels in the US diet were then calculated from published consumption data for soybean, corn, canola and olive oil³ and measured oxylipins in these oils. The fatty acid composition of the oils in both experiments was confirmed with GC-FID.

Experiment 1 oils were purchased from local supermarkets in San Francisco. **Experiment 2** oils were purchased from local supermarkets in Davis.

Experiment 1. Five mL of each of the soybean, corn, canola, olive and 4 algae oils were added to 8 mL glass vials (17 mm diameter, 64 mm height) and heated, uncapped at 100 °C on a heating block for 10 min. Temperature was measured each minute. Samples were staggered 1 min apart to allow time for temperature recording ([Supporting Information Figure 2](#)). Two 10 μ L samples were taken from each oil at baseline (room temperature) and after 10 min of heating for fatty acid and oxylipin analysis (methodological details below). There were three replicates per oil in total, and each was performed on a separate day (i.e., all 8 oils heated and sampled on day 1 and then on days 2 and 3). Oils

were stored in a 4 °C fridge and thawed prior to aliquoting samples each day.

Experiment 2. Soybean, corn, canola, olive and flaxseed oils were purchased from various supermarket outlets in Davis (CA, USA). [Supporting Information Table 1](#) shows the oil brand, store they were purchased from, date they were purchased and expiry date. For each oil, 5 bottles from the same brand but with different lot numbers were purchased from the various stores, except for flaxseed oil, because we were not able to find the same brand in different stores. Therefore, 3 different brands of flaxseed oil were purchased, with one brand being purchased in duplicate from the same store to increase the sample size to 4 ([Supporting Information Table 1](#)). The reason we aimed to keep the oil company source consistent was to confirm the validity of our methods on the same oil brand. Oxylipins were extracted and measured within a week after purchasing the oils. Fatty acid composition of the oils was confirmed a few weeks later.

Oxylipin Analysis with UPLC-MS/MS. A total of 8 LA and 8 ALA oxylipins were measured by targeted UPLC-MS/MS ([Supporting Information Table 2](#)). Ten μ L of oil sample were mixed with 200 μ L of methanol containing 0.1% acetic acid and 0.1% butylated hydroxytoluene (BHT) after adding 10 μ L surrogate standard solution (purity $\geq$ 95%) containing 5 pmol (per sample) of d11–11(12)-EpETrE, d11–14,15-DiHETrE, d4–6-keto-PGF1 α , d4–9- HODE, d4-LTB4, d4-PGE2, d4-TXB2, d6–20-HETE and d8–5-HETE in methanol¹²⁷ and 10 μ L of antioxidant solution (0.2 mg/mL BHT, ethylenediaminetetraacetic acid (EDTA), and triphenylphosphine (TPP) in water/methanol (1:1 v/v)). While the test oils did not contain arachidonic and docosahexaenoic acid, d4–6-keto-PGF1 α , d4-LTB4, d4-PGE2 and d4-TXB2, which are standards for arachidonate or docosahexaenoate derived oxylipins, constituted part of our routinely used surrogate standard mix.^{10,11} They were included in the assay but were not used in any of the LA or ALA oxylipin calculations. It should be noted that the oil aliquots in methanol (containing acetic acid, antioxidant and surrogate standard) were stored in –80 °C until they were hydrolyzed on the same day, 1–3 days after the heating experiment (as described in the following paragraph).

Oils were hydrolyzed by adding 200 μ L of 0.25 M sodium carbonate solution (1.13 g in 21.3 mL water and 21.3 mL methanol) to 10 μ L oil aliquot containing antioxidant and surrogate standard.^{10,33,34} The samples were vortexed and heated for 30 min at 60 °C under constant shaking. The samples were allowed to cool to room temperature and 25 μ L acetic acid and 1575 μ L of Millipore water were added to each sample. Litmus paper was used to confirm that the pH was below 7 after adding the acetic acid.

Oxylipins were extracted using solid phase extraction (SPE) Waters Oasis HLB columns (60 mg, 3 cm cartridges; Waters, Milford, MA). The SPE columns were washed with 4 mL ethyl acetate and twice with 4 mL of methanol and preconditioned twice with 4 mL of SPE wash buffer containing 95/5/0.1 v/v/v Millipore water/methanol/acetic acid. The hydrolyzed oil samples were loaded onto the column and washed with 4 mL SPE buffer twice. The SPE filter was dried under high vacuum for 20 min. Oxylipins were then eluted from the filter with 0.5 mL methanol and 1.5 mL ethyl acetate into collection tubes containing 6 μ L of 30% glycerol (in methanol). Samples were dried with vacuum centrifugation and reconstituted in 50 μ L methanol containing 200 nM 1-cyclohexyl ureido, 3-dodecanoic acid (CUDA) as a surrogate recovery standard. The reconstituted samples were filtered using Ultrafree-MC VV Centrifugal Filter (0.1 μ m; EMD Millipore, Bedford, MA, USA) tubes and transferred to LC-MS/MS vials.

A surrogate standard mix containing 10, 50, or 200 nM of d11–11(12)-EpETrE, d11–14,15-DiHETrE, d4–6-keto-PGF1 α , d4–9-HODE, d4-LTB4, d4-PGE2, d4-TXB2, d6–20-HETE and d8–5-HETE was spiked with 200 nM CUDA and used to calculate accuracy and sample percent recovery. The accuracy was determined by dividing the observed concentrations of the surrogate standards using CUDA, by the expected concentrations and multiplying by 100 ($n = 3$). The observed concentration of the surrogate standards was calculated as follows:

$$\begin{aligned} &\text{Observed surrogate concentration} \\ &= (\text{Surrogate standard area/CUDA area}) \\ &\quad \times (1/\text{slope of surrogate standard curve}) \times 200 \text{ nM CUDA} \end{aligned}$$

The percent recovery in the oil samples was calculated as follows:

$$\begin{aligned} \text{Percent recovery} &= (\text{Surrogate area in oil sample} \\ &\quad / \text{CUDA area in oil sample}) \\ &\quad \times (1/\text{slope of surrogate standard curve}) \\ &\quad \times 200 \text{ nM CUDA} / 100 \text{ nM surrogate concentration} \times 100 \end{aligned}$$

The slope of surrogate standard curve was derived from a plot of the standard surrogate concentration/CUDA concentration versus surrogate area/CUDA area,

Oxylipins were analyzed on an Agilent 1200 SL LC series UPLC system (Agilent Corporation, Palo Alto, CA, USA) connected to a 4000 QTRAP tandem mass spectrometer (Applied Biosystems Instrument Corporation, Foster City, CA, USA) equipped with an electrospray source (Turbo V). The system was operated in negative electrospray ionization mode and used optimized multiple reaction monitoring (MRM) conditions.²⁷ Oxylipins were separated on an Agilent Eclipse Plus C-18 reverse-phase column (2.1 × 150 mm, 1.8 μm particle size). The autosampler temperature was kept constant at 4 °C and the column at 50 °C. Mobile phase A contained Millipore water containing 0.1% glacial acetic acid, and mobile phase B contained acetonitrile/methanol (80/15 v/v) with 0.1% glacial acetic acid. The flow rate was 250 μL/min. Solvent B was held at 35% for 0.25 min, and then increased to 45% between 0.25 and 1 min, 55% from 1 to 3 min, 66% from 3 to 8.5 min, 72% from 8.5 to 12.5 min, 82% from 12.5 to 15 min and 95% from 15 to 16.5 min. It was maintained at 95% to 18 min, lowered to 35% from 18 to 18.1 min and held at 35% between 18.1 and 21 min. The retention time, MRM conditions, collision energy, limits of quantitation and surrogate standard used for each oxylipin are presented in [Supporting Information Table 3](#). The limits of detection were set at 3 times the signal-to-noise ratio, whereas the limits of quantitation were set at 10 times the signal-to-noise ratio.

Oil Hydrolysis under Air or Nitrogen. To ensure that heat applied during the hydrolysis process described above does not cause oxylipin artifacts, we performed the hydrolysis reaction described above with soybean oil samples (10 μL per sample) capped under air (n = 3) or flushed with nitrogen prior to capping (n = 3) at 60 °C for 30 min. The hydrolyzed oxylipins were subjected to SPE and measured on an Agilent 1290 Infinity UPLC system interfaced to a 6460 triple-quadrupole mass spectrometer with electrospray ionization LC-MS/MS. The same oxylipin method was adapted from the 4000 QTRAP tandem mass spectrometer system (Applied Biosystems) to the Agilent 6460 tandem mass spectrometer. The data shown in [Supporting Information Table 4](#) confirm no differences between oxylipins hydrolyzed under air or nitrogen.

Fatty Acid Analysis with Gas Chromatography. Ten microliters of oil sample was nonadecanoic acid (19:0) ethyl ester in chloroform/methanol (2:1 v/v; [Experiment 1](#)) or free heptadecanoic acid in methanol ([Experiment 2](#)) as internal standards. Fatty acids were analyzed according to the method of Ichihara et al.³⁵ Four hundred μL of toluene, 3 mL methanol and 600 μL of 8% hydrochloric acid in methanol solution were added to each sample before placing vials on a dry heating block at 90 °C for 60 min. The samples were allowed to cool at room temperature for approximately 10 min. One mL of hexane and 1 mL of water were added to each sample. The samples were vortexed and the hexane and water layer were allowed to separate by leaving the sample undisturbed for 15 min. Six hundred μL of the upper hexane layer were transferred to a microcentrifuge tube containing sodium sulfate as a drying agent. The hexane layer containing fatty acid methyl esters (FAMES) was transferred into new microcentrifuge tubes and stored at −80 °C until analysis.

FAMES were analyzed on a Varian 3800 gas-chromatography system equipped with a DB-23 fused silica capillary column (30 m × 0.25 mm

inner diameter, 0.25 μm film thickness; Agilent Technologies, Santa Clara, CA, USA). The injector and detector temperature were set at 250 and 300 °C, respectively. The initial oven temperature was held at 50 °C for 2 min, and was increased by 10 °C/min to 180 °C, held at 180 °C for 5 min, increased to 240 °C at 5 °C/min and held at 240 °C for 5 min. The total run time was 37 min. The carrier gas was helium, which was maintained at a flow rate of 1.3 mL/min. A custom mix of 31 fatty acid methyl ester standards (NuChek Prep, Elysian, MN, USA) was used to identify the individual fatty acids. Retention times of the main fatty acids in oils are presented in [Supporting Information Table 5](#). Fatty acid concentrations were determined by comparison of the GC peak areas to the internal standard area. Data were expressed as percent of total identified fatty acid peaks or absolute concentrations.

Estimation of Dietary LA and ALA Derived Oxylipin Intake Levels. US oil intake data reported by Blasbalg et al. (2011)³ was used to derive the mean amount of oxylipins consumed in the US diet. The amount of soybean, corn, canola and olive oil consumed in grams per person per day based on the Blasbalg et al. study is 31.8, 2.2, 2.2 and 1.9, respectively ([Supporting Information Table 6](#)). This amount was multiplied by the measured mean concentration of oxylipins in each oil, to estimate LA and ALA oxylipin intake levels, after correcting for oil density.

Statistical Analysis. Statistical analysis was performed on GraphPad Prism v. 6.05 (La Jolla, CA, USA). For [Experiment 1](#), the effect of heat on the rise in temperature of the 8 oils over time was determined using a two-way repeated measures analysis of variance (ANOVA) followed by Tukey's multiple comparison test. The differences in fatty acid and oxylipin concentrations at baseline, and in the change in oxylipins after heating (relative to baseline) amongst the different oils were assessed using a Kruskal-Wallis one-way ANOVA. No post-hoc tests were performed in [Experiment 1](#) because the sample size was too small (n = 3 per oil) to provide accurate comparisons without risking a type I or II statistical error. Data from [Experiment 1](#) are presented as median and range of the lowest and highest points.

Data for [Experiment 2](#) were analyzed by Kruskal-Wallis test, followed by Dunn's multiple comparison test. The total sample size of 24 oils (n = 4–5 per oil) was sufficient to allow meaningful comparisons. Fatty acid and oxylipin data for [Experiment 2](#) are presented as median and interquartile ranges (25th and 75th percentiles). Spearman's correlation analysis was used to correlate LA and ALA concentrations (μM) to LA and ALA derived oxylipins (nM).

Oxylipin intake data are expressed as mean without standard deviation, because it is a calculated value so true variability cannot be established. Hence, no statistical comparisons were done for calculated oxylipin intake data.

Statistical significance was set at $p < 0.05$.

RESULTS

Experiment 1. Standard Recovery. The mean accuracy for d11-11(12)-EpETrE, d11-14,15-DiHETrE, d4-9-HODE, and d8-5-HETE, the surrogates used to quantify LA and ALA derived oxylipins, was 82%, 81%, 92%, and 82%, respectively. The correlation coefficient reflecting the linearity of the standard curve was above 0.99 for all measured oxylipins. The mean percent recovery of d11-11(12)-EpETrE, d11-14,15-DiHETrE, d4-9-HODE, and d8-5-HETE, the surrogates used to quantify LA and ALA derived oxylipins, was 39%, 70%, 55%, and 48%, respectively. The low percent recovery of the surrogate standards in the oils is likely due to their partial degradation during base hydrolysis and ion suppression.^{36,37} However, the use of the surrogate standards added prior to base hydrolysis and SPE corrects for these losses.³⁶ Although a reduced percent recovery may decrease sensitivity, only peaks with a signal-to-noise ratio above 10 were included in the analysis to minimize the risk of quantitating noise. A representative UPLC-MS/MS chromatogram of peaks detected in soybean oil before and after 10 min of heating is shown in [Supporting Information Figure 1](#).

Figure 1. Oil fatty acid percent composition.

Temperature. Two-way repeated measures ANOVA showed a significant effect of time on oil temperature ($F(10,160) = 3204$; $P < 0.0001$) during the 10 min heating period. Oil temperature increased significantly within the first minute compared to baseline and reached a steady-state level of 100 °C within 7 min (Supporting Information Figure 2).

Baseline Oil Fatty Acid Composition. Oil fatty acid percent composition is shown in Figure 1. Concentration data are presented in Supporting Information Figure 3. Differences in fatty acid percent composition among the oils were reflected in the concentration data. Therefore, statistical analysis of the percent composition data will be discussed in this section.

Kruskal–Wallis test showed significant differences among the groups for myristic acid (14:0; $p = 0.014$), palmitic acid (16:0; $p = 0.003$), stearic acid (18:0; $p = 0.007$), oleic acid (18:1n-9; $p = 0.003$), vaccenic acid (18:1n-7; $p = 0.013$), LA ($p = 0.002$), and ALA ($p = 0.01$). Myristate, stearate, and vaccinate were generally low in composition (<5%) or not detected in some oils. Palmitate was highest in olive (12%), corn (11.5%), and soybean oils (11%), followed by canola oil (4%), HSAO (neat and w/AO; ~3.5%), and “Thrive” (neat and w/AO; ~1.5%). Oleate was highest in the 4 algae oils (~91%), followed by olive (74%), canola (63.5%), soybean (21%), and corn (29%) oil.

LA composition was highest in corn and soybean oil (55%), followed by canola (19%), olive (7%), “Thrive” (neat and w/AOs; ~3.9%), and HSAO (neat and w/AO; ~1.5%). ALA was highest in canola oil (7.2%), followed by soybean oil (6.5%), corn oil (0.8%), and olive oil (0.4%). It was negligible or not detected in the 4 algae oils.

Baseline Oxylipin Concentrations. The LA and ALA oxylipin data are presented in Figures 2A and 2B, respectively. Overall, high LA or ALA oils had high concentrations of their respective LA or ALA metabolites, except for olive oil, which had comparable levels of LA and ALA monohydroxylated products compared to corn oil despite being low in LA (7%) and ALA (0.4%). The presence or absence of α -tocopherol in the algae oils did not appear to affect oxylipin concentrations.

Results from the one-way ANOVA indicated that LA derived oxylipin median values differed significantly among the groups for 9-HODE ($p = 0.006$), 13-HODE ($p = 0.006$), 9-oxo-ODE ($p = 0.003$), 13-oxo-ODE ($p = 0.007$), 9,10-DiHOME ($p = 0.006$),

and 12,13-DiHOME ($p = 0.006$) (Figure 2A). Monohydroxylated metabolites (9-HODE and 13-HODE) were 19 to 24 times higher in olive and corn oil than soybean and canola oil, which were both 3 to 4 times higher than the 4 algae oils. LA dihydroxylated metabolites (9,10-DiHOME and 12,13-DiHOME) were at least 11 times higher in corn and soybean oil as compared to other oils. No significant differences were observed for epoxy-metabolites of LA (9(10)-EpOME and 12(13)-EpOME).

One-Way ANOVA also showed statistically significant differences among the groups in ALA derived monohydroxy (9-HOTrE, $p = 0.019$; 13-HOTrE, $p = 0.005$), epoxy (9(10)-EpODE, $p = 0.04$; 15(16)-EpODE, $p = 0.035$), and dihydroxy metabolites (9,10-DiHODE, $p = 0.008$; 12,13-DiHODE, $p = 0.004$; 15,16-DiHODE, $p = 0.004$) (Figure 2B). Monohydroxylated ALA metabolites (9- and 13-HOTrE) were highest in olive oil, followed sequentially by corn, soybean, and canola oil, compared to algae oils. Epoxidized ALA metabolites (9(10)-EpODE, 15(16)-EpODE), and the dihydroxylated ALA metabolite 15,16-DiHODE were 5- to 22- fold higher in canola and soybean oil compared to the other oils. Dihydroxylated ALA derived 9,10-DiHODE was 4-fold higher in soybean oil compared to corn oil, which was 4- to 19-fold higher compared to other oils. 12,13-DiHODE was detected in corn, canola, and soybean oils, but was negligible or undetected in the remaining oils.

Effect of Oil Type on the Change in Oxylipin and Fatty Acid Concentrations after Short-Term Heating. Heat was applied at 100 °C for 10 min to test whether oxylipins increased more in oils with higher levels of LA and ALA compared to less unsaturated oils such as algae. Kruskal–Wallis one-way ANOVA found no significant differences among the oils in the change (from baseline) in LA or ALA oxidized metabolites (Supporting Information Figure 4A and 4B). There were no significant differences among the oils in the change in fatty acid concentrations or percent composition from baseline (data not shown).

Experiment 2. Fatty Acid Composition of Various off-the-Shelf Oils. The fatty acid percent composition data for olive, corn, canola, soybean, and flaxseed oils obtained from 4 to 5 stores (one oil per store) are presented in Table 1. Fatty acid

Figure 2. Oil LA (A) or ALA (B) derived oxylipin concentrations at room temperature.

Table 1. Fatty Acid Composition of the Different Oils (Soybean, Corn, Canola, Olive, and Flaxseed)^a

	Soybean Oil	Corn Oil	Canola Oil	Olive Oil	Flaxseed Oil
16:0	10.2 (10.0–10.5) ^{ab}	11.9 (11.8–11.9) ^{ac}	4.0 (4.0–4.2) ^b	13.7 (12.5–14.3) ^a	5.3 (5.2–5.5) ^{bc}
18:0	3.8 (3.7–4.0) ^a	1.5 (1.5–1.6) ^b	1.6 (0.0–1.7) ^{ab}	2.6 (2.6–3.1) ^{ab}	3.7 (3.4–4.1) ^{ab}
18:1 n-9	22.5 (22.4–22.8) ^{ac}	28.5 (28.1–28.7) ^{bc}	61.5 (60.9–63.1) ^{ab}	71.3 (67.4–73.2) ^b	20.2 (18.9–21.4) ^c
18:1 n-7	1.2 (1.1–1.2) ^{ab}	0.0 (0.0–0.0) ^a	1.6 (1.5–2.2) ^b	1.6 (1.3–1.8) ^b	0.0 (0.0–0.0) ^a
18:2 n-6	55.0 (54.7–55.0) ^{ac}	56.9 (56.8–57.3) ^a	19.7 (19.6–19.8) ^{ab}	8.6 (7.2–11.2) ^b	15.0 (14.5–15.5) ^{bc}
18:3 n-3	7.1 (7.0–7.1) ^{ab}	1.0 (1.0–1.1) ^{ac}	8.9 (8.7–8.9) ^{bc}	0.8 (0.8–0.9) ^a	55.7 (54.2–57.0) ^b
18:2 n-6/18:3 n-3	7.8 (7.6–7.9) ^{abc}	54.3 (52.3–61.6) ^a	2.2 (2.2–2.3) ^{bc}	12.0 (8.6–12.5) ^{ac}	0.3 (0.3–0.3) ^b

^aData (% of total detected fatty acids) are expressed as median and interquartile range (25th and 75th percentiles). Data were analyzed by Kruskal–Wallis test followed by Dunn's multiple comparison posthoc test. For each row, different alphabetical superscripts mean that the oils differed significantly ($P < 0.05$) from each other. Only the main fatty acids present in the oils are presented, namely, palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1 n-9), vaccenic acid (18:1 n-7), linoleic acid (18:2 n-6), and α -linolenic acid (18:3 n-3).

analysis confirmed the composition of the oils.^{13,38} Flaxseed oil contained the highest amount of ALA (56%), followed by canola (9%), soybean (7%), corn (1%), and olive (0.8%) oil. Corn and soybean oil contained the highest amount of LA (55–57%)

followed by canola (20%), flaxseed (15%), and olive (9%) oil.

Significant differences among the various oils are shown in Table

1.

Table 2. Concentrations of Linoleic Acid (LA) and α -Linolenic Acid (ALA) Derived Metabolites in Nanomolar in the Different Oils (Olive, Corn, Canola, Soybean, and Flaxseed)^a

	Soybean Oil	Corn Oil	Canola Oil	Olive Oil	Flaxseed Oil
13-HODE	1570 (1025–2095) ^{ab}	8400 (6085–10205) ^{bc}	147 (101–232) ^b	14300 (6235–17600) ^{ac}	24100 (16900–45925) ^c
9-HODE	333 (260–795.5) ^{ab}	5450 (4270–6065) ^{bc}	215 (165.5–303) ^b	14000 (4915–16000) ^{ac}	68050 (40900–78100) ^c
13-oxo-ODE	277 (147.5–352.5) ^{ab}	446 (330.5–545) ^{bc}	120 (64.4–181) ^b	4420 (2980–5240) ^{ac}	25800 (20275–50000) ^c
9-oxo-ODE	892 (722–1590) ^a	2320 (1470–3485) ^{ab}	679 (444.5–963) ^a	10300 (4850–12250) ^b	2880 (1640–3805) ^{ab}
12(13)-EpOME	5100 (3510–6140) ^{ab}	13000 (11050–16600) ^a	2880 (2445–3210) ^{ab}	1210 (609–1445) ^b	12970 (7905–18575) ^a
9(10)-EpOME	4430 (3135–5345) ^{ab}	17300 (16250–21200) ^a	2550 (2025–2975) ^b	250 (939–2685) ^b	9200 (6368–11950) ^{ab}
12,13-DiHOME	18800 (10870–22700) ^a	9030 (5230–13565) ^a	982 (906–1065) ^{ab}	56.9 (40.85–78.1) ^b	352.5 (159–868.5) ^{ab}
9,10-DiHOME	23900 (14200–28800) ^a	15400 (9130–21050) ^a	357 (314.5–414) ^{ab}	75.5 (59.8–99.5) ^b	444 (158–934) ^{ab}
Total LA-metabolites	56590 (37475–63568)^{ab}	72802 (60932–84871)^a	7922 (6850–8963)^b	46244 (20954–55257)^{ab}	143157 (111198–193904)^a
9-HOTrE	99 (59–140) ^{ab}	231 (208–283) ^{bc}	42 (26–63) ^b	2090 (1915–2955) ^{ac}	50750 (28275–77800) ^c
13-HOTrE	160 (113–243) ^{ab}	42 (28–53) ^a	95 (77–114) ^{ac}	908 (604–1260) ^{bc}	19300 (11890–34975) ^b
15(16)-EpODE	3590 (2525–4095) ^{ab}	2430 (2155–3055) ^{ab}	3840 (3210–4075) ^a	224 (194–235) ^b	248500 (141500–384750) ^a
12(13)-EpODE	574 (353–628) ^{ab}	213 (206–260) ^{ac}	851 (710.5–1024) ^{bc}	85 (64–116) ^a	15050 (9165–21325) ^b
9(10)-EpODE	742 (457–791) ^{ab}	1010 (809–1370) ^{ac}	1030 (899–1375) ^{bc}	178 (122–231) ^a	20650 (13950–27500) ^c
15,16-DiHODE	28000 (16250–32900) ^a	1530 (935–2630) ^b	11200 (9710–12600) ^{ab}	15 (11–27) ^b	7990 (2600–14700) ^{ab}
12,13-DiHODE	720 (415–914) ^a	187 (126–277) ^{ab}	103 (85–114) ^{bc}	14 (10–17) ^b	381 (225–519) ^{ac}
9,10-DiHODE	9160 (5000–11150) ^a	1720 (939–2775) ^{ab}	146 (132–173) ^{bc}	7 (6–13) ^c	306 (180–507) ^{ac}
Total ALA-metabolites	43127 (26067–49923)^{ab}	7073 (6362–9890)^{ac}	17192 (16166–18277)^{bc}	3570 (2942–4810)^c	355205 (279990–497593)^b

^aData are expressed as median and interquartile range (25th and 75th percentiles). HODE, hydroxyoctadecadienoic acid; oxo-ODE, oxo-octadecadienoic acid; EpOME, epoxyoctadecamonoenoic acid; DiHOME, dihydroxyoctadecamonoenoic acid; OxLAM, oxidized linoleic acid metabolites; HOTrE, hydroxyoctadecatrienoic acid; EpODE, epoxyoctadecadienoic acid; DiHODE, dihydroxyoctadecadienoic acid; ALA, α -linolenic acid. Data were analyzed by Kruskal–Wallis test followed by Dunn's multiple comparison tests. For each row, different alphabetical superscripts mean that the oils differed significantly ($P < 0.05$) from each other.

Oxylipin Concentrations of Various off-the-Shelf Oils.

Oxylipin concentrations of five off-the-shelf oils are shown in Table 2. Kruskal–Wallis one-way ANOVA followed by Dunn's multiple comparison test revealed that LA- and ALA-oxidized metabolite concentrations differed significantly among the different oils.

With regard to LA metabolites, concentrations of LA derived monohydroxy (9- and 13-HODE) and ketone metabolites (9- and 13-oxo-ODE) were highest in flaxseed and olive oil relative to corn, soybean, and canola oil. 9- and 13-HODE and 13-oxo-ODE were significantly higher by 15–316-fold in flaxseed oil as compared to canola ($p < 0.001$) and soybean oil ($p < 0.05$). These metabolites were also significantly higher by 37–97-fold in olive oil relative to canola oil ($p < 0.05$). 9-oxo-ODE was significantly higher in olive oil as compared to both canola ($p < 0.001$) and soybean oils ($p < 0.05$) by 15- and 12-fold, respectively.

Flaxseed and corn oil had the highest concentration of LA derived epoxides (9(10)- and 12(13)-EpOME). Both of these oils had an 11-fold higher concentration of 12(13)-EpOME relative to olive oil ($p < 0.01$). The 9(10)-EpOME concentration was significantly higher than canola ($p < 0.01$) and olive oil ($p < 0.001$) by 7- and 69-fold, respectively.

Dihydroxy products of LA derived epoxides (9,10-DiHOME and 12,13-DiHOME) were significantly higher in soybean ($p < 0.001$) and corn oil ($p < 0.01$) by 159–330-fold and 200–317-fold, respectively, as compared to olive oil.

With regard to ALA metabolites, flaxseed oil had the highest concentration of monohydroxylated ALA derived metabolites, followed by olive oil (Table 2). 9- and 13-HOTrE were significantly higher in flaxseed oil than soybean, corn, and canola oil by 121–1208-fold. 9-HOTrE was significantly higher in olive oil than canola oil by 510-fold, whereas 13-HOTrE was significantly higher than corn oil by 22-fold.

Epoxy-ALA metabolites (15(16)-, 12(13)-, and 9(10)-EpODE) were highest in flaxseed oil, followed by canola oil. All three epoxidized ALA metabolites were 116–1109 and 6–10 times higher in flaxseed and canola oil ($p < 0.05$), respectively, than olive oil, which had the lowest concentration of epoxidized ALA metabolites among the five oils. Soybean, corn, and canola oil had similar concentrations of epoxidized ALA metabolites; however, 12(13)-EpODE was 98% lower in corn oil relative to flaxseed oil ($p < 0.05$) and 9(10)-EpODE was 83% lower in soybean oil than flaxseed oil ($p < 0.05$).

Dihydroxy ALA metabolites (15,16-, 12,13-, and 9,10-diHODEs) were highest in soybean oil and lowest in olive oil. The differences were statistically significant between soybean and olive oil for all three diHODEs, which were lower in olive oil by 98–99.9% ($p < 0.001$). Soybean oil also had a higher concentration of 12,13-DiHODE (+85%; $p < 0.05$) and 9,10-DiHODE (+98%; $p < 0.05$) compared to canola oil.

Reproducibility between Experiments. Supporting Information Table 7 summarizes mean oil oxylipin concentrations from Experiments 1 and 2, and the experiment comparing air to nitrogen capping during hydrolysis ("Experiment 3"). As shown, oxylipin concentrations were similar between Experiments. Epoxides in canola oil, however, differed by 4–7-fold between Experiments 1 and 2, suggesting they are unstable in this particular oil. The oils used came from different outlets (San Francisco or Davis) and likely had different expiration dates. Fankhauser-Noti et al. reported similar fluctuations in epoxides between fresh and old olive and sunflower oil.²⁶

Estimated Daily Oxidized Fatty Acid Intake. Mean daily intake of oxidized LA and ALA metabolites from soybean, corn, canola, and olive oil was estimated, based on published mean consumption (kg/p/year) data on these oils.³ Consumption data for flaxseed oil and the four algae oils were not available.

As shown in Figures 3A and 3B, of the four oils, soybean oil contributed the most of LA and ALA oxylipins in the diet,

Figure 3. Estimated US consumption of LA (A) and ALA (B) derived oxylipins, and total oxylipins (C) from various oils.

because it is the most consumed plant oil in the US (Supporting Information Table 6).³ The majority of oxylipins in the diet were in the form of epoxy and dihydroxy metabolites of LA and ALA.

Estimated daily intake of total oxylipins from olive, corn, canola, and soybean oils averaged 1.1 mg/day (Figure 3C). Of the measured oils, soybean oil contributed most oxylipins in the diet, followed by corn, olive, and canola oil (Figure 3C).

Correlations. Oil LA and ALA concentrations positively correlated with concentrations of their respective metabolites as shown in Table 4. LA correlated positively with LA derived

9(10)- and 12(13)-EpOME and 9,10- and 12,13-DiHOME ($P < 0.05$). It also correlated with ALA derived 9-HOTrE and DiHODEs ($P < 0.05$).

ALA did not correlate with LA derived oxylipins but positively correlated with ALA derived hydroxy, epoxy, and dihydroxy metabolites ($P < 0.05$).

DISCUSSION

The present study reported the presence of LA and ALA derived oxylipin species in plant and algae oils. Concentrations of LA and

Table 4. Spearman's Correlation between Fatty Acid and Oxylipin Concentrations from Experiment 2^a

	Correlation between Fatty Acid and Oxylipin Concentration			
	LA conc		ALA conc	
	r-value	p-value	r-value	p-value
LA-oxylipins				
9-HODE	−0.3191	0.1285	−0.00956	0.9646
13-HODE	−0.2643	0.2119	−0.01565	0.9421
9-oxo-ODE	−0.3148	0.1341	−0.3696	0.0755
13-oxo-ODE	−0.3765	0.0698	0.04174	0.8465
9(10)-EpOME	0.4991	0.0130	0.06174	0.7744
12(13)-EpOME	0.4652	0.0220	0.2565	0.2263
9,10-DiHOME	0.8652	<0.0001	0.1104	0.6075
12,13-DiHOME	0.86	<0.0001	0.06174	0.7744
ALA-oxylipins				
9-HOTrE	−0.4904	0.0150	0.4304	0.0358
13-HOTrE	−0.3852	0.0630	−0.0087	0.9678
9(10)-EpODE	0.01566	0.9421	0.5214	0.0090
12(13)-EpODE	0.04001	0.8527	0.855	<0.0001
15(16)-EpODE	0.1339	0.5327	0.7896	<0.0001
9,10-DiHODE	0.8339	<0.0001	0.2009	0.3466
12,13-DiHODE	0.6551	0.0005	0.492	0.0146
15,16-DiHODE	0.5466	0.0057	0.6458	0.0007

^aTable depicts correlations between total LA and ALA concentration and total LA- and ALA-oxylipin concentrations in all oils. Data was analyzed using Spearman correlation analysis. Significant *p*-values were determined to be <0.05. HODE, hydroxyoctadecadienoic acid; oxo-ODE, oxo-octadecadienoic acid; EpOME, epoxyoctadecamonoenoic acid; DiHOME, dihydroxyoctadecamonoenoic acid; OxLAM, oxidized linoleic acid metabolites; HOTrE, hydroxyoctadecatrienoic acid; EpODE, epoxyoctadecadienoic acid; DiHODE, dihydroxyoctadecadienoic acid; ALA, α -linolenic acid.

ALA derived oxylipins were generally proportional to the concentrations of their precursor fatty acids except for flaxseed and olive oil, which had higher oxylipin concentrations than other oils. Estimated oxylipin intake levels based on available consumption data on soybean, corn, canola, and olive oils averaged 1.1 mg per person per day. Oxylipin concentrations were comparable between experiments, thus confirming the reproducibility of our measurements.

The oxylipins detected in nonheated oils were likely formed by nonenzymatic or enzymatic pathways during the seed extraction process. Nonenzymatic auto-oxidation is known to be influenced by storage or processing conditions.³⁹ Enzymatic oxidation is mediated in part by lipoxygenase enzyme, which is activated when the barrier integrity of the seed, fruit, or algae is compromised by homogenization during the oil extraction process.^{25,29,40} Other plant or algae enzymes involved in oxylipin formation include soluble epoxide hydrolase, cytochrome P450, or pathogen-inducible oxygenases,^{41–44} although their activation during the oil extraction process is not known.

There were no significant differences among the oils in the change in oxylipin concentrations following 10 min of heat relative to baseline (Experiment 1, Supporting Information Figure 4). Previous studies reported the formation of LA derived oxylipins after heating high LA oils at 40, 100, or 180 °C for 10–264 h.^{13–15} We predicted, however, that we might detect changes in oxylipins in the high LA or ALA oils within 10 min of heating at 100 °C using UPLC-MS/MS. The lack of differences between the oils could be due to the short heating duration, as previous

work has demonstrated that prolonged heating is required to oxidize oils.^{13,15} It is also possible that oxylipin products of heat-induced oxidation (HODEs, oxoODEs) were increased in the high LA or ALA oils during the 10 min heating process, but were rapidly degraded into secondary volatile compounds.^{45,46}

The LA and ALA contents of the oils from Experiment 2 were related to the concentration of their respective oxylipin metabolites (Table 4). LA also correlated highly with ALA derived oxylipins. This association was likely driven by olive and flaxseed oil, which unexpectedly contained a high amount of LA derived oxylipins potentially caused by processing, storage, or handling conditions that need to be further investigated. Overall, however, the findings suggest that the precursor fatty acid pool in oils is an important determinant of oxylipin concentrations, consistent with the observation that the low LA and ALA algae oils had the least concentration of LA and ALA derived oxylipins compared to other plant oils, irrespective of vitamin E content (Experiment 1, Figure 2). The presence of vitamin E may not be critical for low polyunsaturated fatty acid oils when heated for a short period of time (10 min) at 100 °C.

In both experiments, olive oil had higher concentrations of LA derived ketones (9- and 13-oxo-ODE), and monohydroxylated LA (9- and 13-HODE) and ALA (9- and 13-HOTrE) metabolites, than soybean, canola and corn oil, despite being low in LA (9%) and ALA (0.8%). This could be due to enhanced lipoxygenase activity upon homogenizing the olive fruit to extract the oil compared to seeds (soybean, canola, or corn).⁴⁰ However, Jarén-Galán et al. reported higher lipoxygenase activity from soybean compared to olives.⁴⁷ It is possible that other oxygenase enzymes that differ in activity between soybeans and olives account for the unexpected high oxylipin concentrations in olive oil. Other factors such as processing or storage conditions may also explain the high monohydroxylated oxylipin metabolites detected in olive oil relative to other higher-LA or ALA oils.

LA derived monohydroxy and ketone metabolites, and ALA derived monohydroxy and epoxy metabolites were highest in flaxseed oil as compared to other oils (Experiment 2). The high concentration of ALA derived oxylipins is expected because flaxseed oil contained the highest concentration of ALA (54%) compared to the other oils tested in this study (2–6%). Flaxseed oil has less LA relative to soybean oil (14% versus 50%), yet LA-metabolites were 3-fold higher in flaxseed oil compared to soybean oil. It is not known whether these metabolites were formed during processing, storage, or flaxseed crushing. The high concentrations of ALA epoxides may have catalyzed the oxidation of LA via electrophilic attack of the allylic carbon next to the double bonds, although this remains to be determined.

The consumption of LA has increased from 2% to 7% of energy over the past century, due to increased consumption of high LA plant oils, such as soybean oil.³ Soybean oil is the most commonly consumed plant oil in the US and the fourth major contributor of total calories.³ It is likely that the consumption of oxidized LA metabolites concomitantly increased over the past few decades with increased soybean oil intake. Approximating the levels of dietary oxylipins will allow future studies evaluating the bioavailability and effect of oxylipins on health and disease to utilize relevant doses.

Oxylipins are bioavailable, and circulating LA derived oxylipins in particular have been associated with atherosclerosis, pain syndromes, and hypertension, consistent with their role in mediating pro-inflammatory signaling in tissues or vasculature.^{4,48,49} A recent meta-analysis reported that fried food

consumption was associated with hypertension and weight gain.⁵⁰ In vivo, the concentration of circulating oxylipins depends on the availability of their precursor fatty acid.^{10,11,51} However, studies have demonstrated that dietary oxylipins are absorbed^{17–20} and incorporated into blood chylomicrons.⁵² The relative plasma contribution of dietary oxylipins compared to endogenously produced oxylipins is not known and merits future evaluation.

Estimated US daily intake of LA and ALA oxylipins from soybean, corn, canola and olive oil was 0.64 and 0.44 mg, respectively, and amounted to a total of 1.1 mg per person per day. These values are likely underestimated, however, because (1) oxylipins in this study were quantified in off-the-shelf oils maintained at room temperature, (2) the oxylipin content of commonly consumed foods such as peanut butter and french fries was not measured,^{53,54} and (3) a targeted UPLC-MS/MS approach was used to quantify oxylipins, which means that other oxylipin species, such as hydroperoxides of LA or ALA or oleic acid derived compounds, were not accounted for in our estimates. Accounting for the amount and type of oxylipins produced during food processing, cooking, or prolonged storage, in relation to water, metal, and antioxidant content, will provide a better estimate of daily oxylipin consumption levels.^{39,55,56} The use of nontargeted mass spectrometry methods may identify other oxylipin species in commonly consumed oils that can be quantified with targeted UPLC-MS/MS as performed in the present study.⁵⁷

Limitations of this study include the low sample size and lack of information on the processing methods used to produce the oils and storage conditions and duration since date of production. The risk of statistical errors associated with the low sample size or number of replicates in [Experiment 1](#) was mitigated by reproducing the measurements of some oils (corn, soybean, olive, and canola) in [Experiment 2](#). Information on processing and storage would require coordination with each of the oil manufacturers in future studies.

In summary, this study quantified LA and ALA oxylipins in various oils and found them to be related to LA and ALA fatty acid composition, with a notable exception being olive and flaxseed oils. The amount of oxylipins derived from commonly consumed plant oils in the US was estimated to be 1.1 mg per person per day, but this value is underestimated because it does not account for oxylipins in commonly consumed oils not measured in this study, or processing, frying, or storage effects. Knowing the amount and type of oxylipins chronically consumed through dietary oils is important for understanding their health implications.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.jafc.6b05559](https://doi.org/10.1021/acs.jafc.6b05559).

Tables and figures related to oil brands tested, oxylipin method details and reproducibility, and data from [Experiment 1](#) (temperature changes over time, fatty acid concentrations, and change in oxylipins after heating for 10 min) ([PDF](#))

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: ataha@ucdavis.edu. Tel: 530-752-7096.

ORCID

Ameer Y. Taha: [0000-0003-4611-7450](https://orcid.org/0000-0003-4611-7450)

Funding

This work was supported by TerraVia Holdings, Inc., USDA National Institute of Food and Agriculture, Hatch/Taha (project #1008787), NIEHS R01 ES002710, NIEHS/Superfund Research Program P42 ES004699, and NIH/NIDDK U24 DK097154.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Mark S. Horowitz is thanked for statistical programming expertise.

■ ABBREVIATIONS USED

ALA, α -linolenic acid; AO, antioxidants; CUDA, 1-cyclohexyl-dodecanoic acid urea; DiHODE, dihydroxyoctadecadienoic acid; DiHOME, dihydroxyoctadecamonoenoic acid; EDTA, triphenylphosphine, ethylenediaminetetraacetic acid; EpODE, epoxyoctadecadienoic acid; EpOME, epoxyoctadecamonoenoic acid; FID, flame ionization detector; GC, gas-chromatography; GC-FID, gas chromatography coupled to a flame ionization detection; HODE, hydroxyoctadecadienoic acid; HOTrE, hydroxyoctadecatrienoic acid; HSAO, high stability algae oil; LA, linoleic acid; LC-MS/MS, liquid chromatography tandem mass spectroscopy; MRM, multiple reaction monitoring; NMR, nuclear magnetic resonance; oxo-ODE, oxo-octadecadienoic acid; PUFAs, polyunsaturated fatty acids; SPE, solid-phase extraction; UV, ultraviolet; w/AO, without added antioxidants

■ REFERENCES

- (1) Hansen, A. E.; Haggard, M. E.; Boelsche, A. N.; Adam, D. J.; Wiese, H. F. Essential fatty acids in infant nutrition. III. Clinical manifestations of linoleic acid deficiency. *Journal of nutrition* **1958**, *66*, 565–76.
- (2) Holman, R. T.; Johnson, S. B.; Hatch, T. F. A case of human linolenic acid deficiency involving neurological abnormalities. *American journal of clinical nutrition* **1982**, *35*, 617–23.
- (3) Blasbalg, T. L.; Hibbeln, J. R.; Ramsden, C. E.; Majchrzak, S. F.; Rawlings, R. R. Changes in consumption of omega-3 and omega-6 fatty acids in the United States during the 20th century. *Am. J. Clin. Nutr.* **2011**, *93*, 950–62.
- (4) Patwardhan, A. M.; Akopian, A. N.; Ruparel, N. B.; Diogenes, A.; Weintraub, S. T.; Uhlson, C.; Murphy, R. C.; Hargreaves, K. M. Heat generates oxidized linoleic acid metabolites that activate TRPV1 and produce pain in rodents. *J. Clin. Invest.* **2010**, *120*, 1617–26.
- (5) Chiba, T.; Thomas, C. P.; Calcutt, M. W.; Boeglin, W. E.; O'Donnell, V. B.; Brash, A. R. The precise structures and stereochemistry of trihydroxy-linoleates esterified in human and porcine epidermis and their significance in skin barrier function: IMPLICATION OF AN EPOXIDE HYDROLASE IN THE TRANSFORMATIONS OF LINOLEATE. *J. Biol. Chem.* **2016**, *291*, 14540.
- (6) Moghaddam, M. F.; Grant, D. F.; Cheek, J. M.; Greene, J. F.; Williamson, K. C.; Hammock, B. D. Bioactivation of leukotoxins to their toxic diols by epoxide hydrolase. *Nat. Med.* **1997**, *3*, 562–6.
- (7) Feldstein, A. E.; Lopez, R.; Tamimi, T. A.; Yerian, L.; Chung, Y. M.; Berk, M.; Zhang, R.; McIntyre, T. M.; Hazen, S. L. Mass spectrometric profiling of oxidized lipid products in human nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *J. Lipid Res.* **2010**, *51*, 3046–54.
- (8) Marcos, R.; Izquierdo, Y.; Vellosillo, T.; Kulasekaran, S.; Cascon, T.; Hamberg, M.; Castresana, C. 9-Lipoxygenase-Derived Oxylipins Activate Brassinosteroid Signaling to Promote Cell Wall-Based Defense and Limit Pathogen Infection. *Plant Physiol.* **2015**, *169*, 2324–34.

- (9) Hamberg, M.; Sanz, A.; Rodriguez, M. J.; Calvo, A. P.; Castresana, C. Activation of the fatty acid alpha-dioxygenase pathway during bacterial infection of tobacco leaves. Formation of oxylipins protecting against cell death. *J. Biol. Chem.* **2003**, *278*, 51796–805.
- (10) Taha, A. Y.; Hennebelle, M.; Yang, J.; Zamora, D.; Rapoport, S. I.; Hammock, B. D.; Ramsden, C. E. Regulation of rat plasma and cerebral cortex oxylipin concentrations with increasing levels of dietary linoleic acid. *Prostaglandins, Leukotrienes Essent. Fatty Acids* **2016**, *10.1016/j.plefa.2016.05.004*
- (11) Ramsden, C. E.; Ringel, A.; Majchrzak-Hong, S. F.; Yang, J.; Blanchard, H.; Zamora, D.; Loewke, J. D.; Rapoport, S. I.; Hibbeln, J. R.; Davis, J. M.; Hammock, B. D.; Taha, A. Y. Dietary linoleic acid-induced alterations in pro- and anti-nociceptive lipid autacoids: Implications for idiopathic pain syndromes? *Mol. Pain* **2016**, *12*, 12.
- (12) Reinaud, O.; Delaforge, M.; Boucher, J. L.; Rocchiccioli, F.; Mansuy, D. Oxidative metabolism of linoleic acid by human leukocytes. *Biochem. Biophys. Res. Commun.* **1989**, *161*, 883–91.
- (13) Marmesat, S.; Velasco, J.; Dobarganes, M. C. Quantitative determination of epoxy acids, keto acids and hydroxy acids formed in fats and oils at frying temperatures. *J. Chromatogr. A* **2008**, *1211*, 129–34.
- (14) Morales, A.; Dobarganes, M. C.; Márquez-Ruiz, G.; Velasco, J.; Marmesat, S. Quantitative analysis of hydroperoxy-, keto- and hydroxydienes in refined vegetable oils. *Journal of Chromatography A* **2012**, *1229*, 190–197.
- (15) Goicoechea, E.; Guillen, M. D. Analysis of hydroperoxides, aldehydes and epoxides by 1H nuclear magnetic resonance in sunflower oil oxidized at 70 and 100 degrees C. *J. Agric. Food Chem.* **2010**, *58*, 6234–45.
- (16) Mubiru, E.; Shrestha, K.; Papastergiadis, A.; De Meulenaer, B. Development and validation of a gas chromatography-flame ionization detection method for the determination of epoxy fatty acids in food matrices. *J. Agric. Food Chem.* **2014**, *62*, 2982–8.
- (17) Ferreira-Vera, C.; Priego-Capote, F.; Mata-Granados, J. M.; Luque de Castro, M. D. Short-term comparative study of the influence of fried edible oils intake on the metabolism of essential fatty acids in obese individuals. *Food Chem.* **2013**, *136*, 576–84.
- (18) Wilson, R.; Lyall, K.; Smyth, L.; Fernie, C. E.; Riemersma, R. A. Dietary hydroxy fatty acids are absorbed in humans: implications for the measurement of 'oxidative stress' in vivo. *Free Radical Biol. Med.* **2002**, *32*, 162–8.
- (19) Wilson, R.; Smith, R.; Wilson, P.; Shepherd, M. J.; Riemersma, R. A. Quantitative gas chromatography-mass spectrometry isomer-specific measurement of hydroxy fatty acids in biological samples and food as a marker of lipid peroxidation. *Anal. Biochem.* **1997**, *248*, 76–85.
- (20) Goicoechea, E.; Brandon, E. F.; Blokland, M. H.; Guillen, M. D. Fate in digestion in vitro of several food components, including some toxic compounds coming from omega-3 and omega-6 lipids. *Food Chem. Toxicol.* **2011**, *49*, 115–24.
- (21) Wilson, R.; Fernie, C. E.; Scrimgeour, C. M.; Lyall, K.; Smyth, L.; Riemersma, R. A. Dietary epoxy fatty acids are absorbed in healthy women. *Eur. J. Clin. Invest.* **2002**, *32*, 79–83.
- (22) Glavind, J.; Sylven, C. Intestinal absorption and lymphatic transport of methyl linoleate hydroperoxide and hydroxyoctadecadienoate in the rat. *Acta Chem. Scand.* **1970**, *24*, 3723–8.
- (23) Kanazawa, K.; Ashida, H. Dietary hydroperoxides of linoleic acid decompose to aldehydes in stomach before being absorbed into the body. *Biochim. Biophys. Acta, Lipids Lipid Metab.* **1998**, *1393*, 349–61.
- (24) Li, H.; Fan, Y. W.; Li, J.; Tang, L.; Hu, J. N.; Deng, Z. Y. Evaluating and predicting the oxidative stability of vegetable oils with different fatty acid compositions. *J. Food Sci.* **2013**, *78*, H633–41.
- (25) Pulvera, Z. M.; Kitamura, K.; Hajika, M.; Shimada, K.; Matsui, K. Oxylipin metabolism in soybean seeds containing different sets of lipoxygenase isozymes after homogenization. *Biosci., Biotechnol., Biochem.* **2006**, *70*, 2598–603.
- (26) Fankhauser-Noti, A.; Biedermann-Brem, S.; Grob, K. Assessment of epoxidized soy bean oil (ESBO) migrating into foods: Comparison with ESBO-like epoxy fatty acids in our normal diet. *Food Chem. Toxicol.* **2006**, *44*, 1279–1286.
- (27) Yang, J.; Schmelzer, K.; Georgi, K.; Hammock, B. D. Quantitative profiling method for oxylipin metabolome by liquid chromatography electrospray ionization tandem mass spectrometry. *Anal. Chem.* **2009**, *81*, 8085–93.
- (28) Paul Abishek, M.; Patel, J.; Prem Rajan, A. Algae oil: a sustainable renewable fuel of future. *Biotechnol. Res. Int.* **2014**, *2014*, 272814.
- (29) Jacquemoud, D.; Pohnert, G. Extraction and Analysis of Oxylipins from Macroalgae Illustrated on the Example *Gracilaria vermiculophylla*. *Methods Mol. Biol.* **2015**, *1308*, 159–72.
- (30) Morgan, D. A. Smoke, fire, and flash points of cottonseed, peanut, and other vegetable oils. *J. Am. Oil Chem. Soc.* **1942**, *19*, 193–98.
- (31) Choi, H.; Lee, E.; Lee, K. G. Quality evaluation of noble mixed oil blended with palm and canola oil. *J. Oleo Sci.* **2014**, *63*, 653–60.
- (32) Douny, C.; Razanakolona, R.; Ribonnet, L.; Milet, J.; Baeten, V.; Rogez, H.; Scippo, M. L.; Larondelle, Y. Linseed oil presents different patterns of oxidation in real-time and accelerated aging assays. *Food Chem.* **2016**, *208*, 111–5.
- (33) Arnold, C.; Markovic, M.; Blosssey, K.; Wallukat, G.; Fischer, R.; Dechend, R.; Konkel, A.; von Schacky, C.; Luft, F. C.; Müller, D. N.; Rothe, M.; Schunck, W. H. Arachidonic acid-metabolizing cytochrome P450 enzymes are targets of {omega}-3 fatty acids. *J. Biol. Chem.* **2010**, *285*, 32720–33.
- (34) Schebb, N. H.; Ostermann, A. I.; Yang, J.; Hammock, B. D.; Hahn, A.; Schuchardt, J. P. Comparison of the effects of long-chain omega-3 fatty acid supplementation on plasma levels of free and esterified oxylipins. *Prostaglandins Other Lipid Mediators* **2014**, *113–115*, 21–9.
- (35) Ichihara, K.; Fukubayashi, Y. Preparation of fatty acid methyl esters for gas-liquid chromatography. *J. Lipid Res.* **2010**, *51*, 635–40.
- (36) Panuwet, P.; Hunter, R. E., Jr.; D'Souza, P. E.; Chen, X.; Radford, S. A.; Cohen, J. R.; Marder, M. E.; Kartavenka, K.; Ryan, P. B.; Barr, D. B. Biological Matrix Effects in Quantitative Tandem Mass Spectrometry-Based Analytical Methods: Advancing Biomonitoring. *Crit. Rev. Anal. Chem.* **2016**, *46*, 93–105.
- (37) Willenberg, I.; Ostermann, A. I.; Schebb, N. H. Targeted metabolomics of the arachidonic acid cascade: current state and challenges of LC-MS analysis of oxylipins. *Anal. Bioanal. Chem.* **2015**, *407*, 2675–83.
- (38) Orsavova, J.; Jarmila Vavra, A.; Robert, V.; Mlcek, J. Fatty Acid composition of vegetable oils and its contribution to dietary energy intake and dependence of cardiovascular mortality on dietary intake of fatty acids. *Int. J. Mol. Sci.* **2015**, *16*, 12871–12890.
- (39) Claxson, A. W.; Hawkes, G. E.; Richardson, D. P.; Naughton, D. P.; Haywood, R. M.; Chander, C. L.; Atherton, M.; Lynch, E. J.; Grootveld, M. C. Generation of lipid peroxidation products in culinary oils and fats during episodes of thermal stressing: a high field 1H NMR study. *FEBS Lett.* **1994**, *355*, 81–90.
- (40) Soldo, B.; Sprung, M.; Musac, G.; Pavela-Vrancic, M.; Ljubenkov, I. Evaluation of Olive Fruit Lipoxygenase Extraction Protocols on 9- and 13-Z,E-HPODE Formation. *Molecules* **2016**, *21*, 506.
- (41) Morisseau, C.; Beetham, J. K.; Pinot, F.; Debernard, S.; Newman, J. W.; Hammock, B. D. Cress and potato soluble epoxide hydrolases: purification, biochemical characterization, and comparison to mammalian enzymes. *Arch. Biochem. Biophys.* **2000**, *378*, 321–32.
- (42) Sanz, A.; Moreno, J. I.; Castresana, C. PLOX, a new pathogen-induced oxygenase with homology to animal cyclooxygenase. *Plant Cell* **1998**, *10*, 1523–37.
- (43) Grausem, B.; Widemann, E.; Verdier, G.; Nosbusch, D.; Aubert, Y.; Beisson, F.; Schreiber, L.; Franke, R.; Pinot, F. CYP77A19 and CYP77A20 characterized from *Solanum tuberosum* oxidize fatty acids in vitro and partially restore the wild phenotype in an *Arabidopsis thaliana* cutin mutant. *Plant, Cell Environ.* **2014**, *37*, 2102–15.
- (44) Petkova-Andonova, M.; Imaishi, H.; Ohkawa, H. CYP92B1, A cytochrome P450, expressed in petunia flower buds, that catalyzes monooxidation of long-chain fatty acids. *Biosci., Biotechnol., Biochem.* **2002**, *66*, 1819–28.
- (45) Goicoechea, E. Volatile compounds generated in corn oil stored at room temperature. Presence of toxic compounds. *Eur. J. Lipid Sci. Technol.* **2014**, *116*, 395–406.

- (46) Katragadda, H. R.; Sidhu, S. Carbonell-Barrachina A.A., Emissions of volatile aldehydes from heated cooking oils. *Food Chem.* **2010**, *120*, 59–65.
- (47) Jaren-Galan, M.; Carmona-Ramon, C.; Minguez-Mosquera, M. I. Interaction between chloroplast pigments and lipoxygenase enzymatic extract of olives. *J. Agric. Food Chem.* **1999**, *47*, 2671–7.
- (48) Leong, X. F.; Salimon, J.; Mustafa, M. R.; Jaarin, K. Effect of repeatedly heated palm olein on blood pressure-regulating enzymes activity and lipid peroxidation in rats. *Malays J. Med. Sci.* **2012**, *19*, 20–9.
- (49) Jaarin, K.; Mustafa, M. R.; Leong, X. F. The effects of heated vegetable oils on blood pressure in rats. *Clinics (Sao Paulo)* **2011**, *66*, 2125–32.
- (50) Sayon-Orea, C.; Carlos, S.; Martinez-Gonzalez, M. A. Does cooking with vegetable oils increase the risk of chronic diseases?: a systematic review. *Br. J. Nutr.* **2015**, *113* (Suppl 2), S36–48.
- (51) Ramsden, C. E.; Ringel, A.; Feldstein, A. E.; Taha, A. Y.; MacIntosh, B. A.; Hibbeln, J. R.; Majchrzak-Hong, S. F.; Faurot, K. R.; Rapoport, S. I.; Cheon, Y.; Chung, Y. M.; Berk, M.; Mann, J. D. Lowering dietary linoleic acid reduces bioactive oxidized linoleic acid metabolites in humans. *Prostaglandins, Leukotrienes Essent. Fatty Acids* **2012**, *87*, 135–41.
- (52) Staprans, I.; Rapp, J. H.; Pan, X. M.; Kim, K. Y.; Feingold, K. R. Oxidized lipids in the diet are a source of oxidized lipid in chylomicrons of human serum. *Arterioscler., Thromb., Vasc. Biol.* **1994**, *14*, 1900–5.
- (53) Storey, M. L.; Anderson, P. A. Contributions of white vegetables to nutrient intake: NHANES 2009–2010. *Adv. Nutr.* **2013**, *4*, 335S–44S.
- (54) Stroehla, B. C.; Malcoe, L. H.; Velie, E. M. Dietary sources of nutrients among rural Native American and white children. *J. Am. Diet. Assoc.* **2005**, *105*, 1908–16.
- (55) Aladedunye, F. A.; Przybylski, R. Degradation and Nutritional Quality Changes of Oil During Frying. *J. Am. Oil Chem. Soc.* **2009**, *86*, 149–156.
- (56) Spiteller, P.; Spiteller, G. Strong dependence of the lipid peroxidation product spectrum whether Fe²⁺/O₂ or Fe³⁺/O₂ is used as oxidant. *Biochim. Biophys. Acta, Lipids Lipid Metab.* **1998**, *1392*, 23–40.
- (57) Cajka, T.; Fiehn, O. Toward Merging Untargeted and Targeted Methods in Mass Spectrometry-Based Metabolomics and Lipidomics. *Anal. Chem.* **2016**, *88*, 524–45.