

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

An odorant-binding protein required for suppression of sweet taste by bitter chemicals.

Permalink

<https://escholarship.org/uc/item/15s9t3pj>

Journal

Neuron, 79(4)

Authors

Jeong, Yong
Shim, Jaewon
Oh, So
et al.

Publication Date

2013-08-21

DOI

10.1016/j.neuron.2013.06.025

Peer reviewed

Published in final edited form as:

Neuron. 2013 August 21; 79(4): 725–737. doi:10.1016/j.neuron.2013.06.025.

An Odorant Binding Protein required for suppression of sweet taste by bitter chemicals

Yong Taek Jeong^{1,*}, Jaewon Shim^{1,*}, So Ra Oh², Hong In Yoon², Chul Hoon Kim², Seok Jun Moon^{1,4}, and Craig Montell^{3,4,5}

¹Department of Oral Biology, BK21 project, Yonsei University College of Dentistry, 250 Seongsanno, Seodaemun-gu, Seoul 120-752, Korea

²Department of Pharmacology, BK21 project, Yonsei University College of Medicine

³Neuroscience Research Institute and Department of Molecular, Cellular and Developmental Biology, University of California Santa Barbara, Santa Barbara, CA, 93110, USA

⁴Department of Biological Chemistry and Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

Summary

Animals are often confronted with the decision as to consume a diet that contains competing attractive and aversive compounds. Here, using the fruit fly, *Drosophila melanogaster*, we describe a mechanism that influences this decision. Addition of bitter compounds to sucrose suppressed feeding behavior, and this inhibition depended on the odorant binding protein, OBP49a. In wild-type flies, bitter compounds suppressed sucrose-induced action potentials, and the inhibition was impaired in *Obp49a* mutants. However, loss of OBP49a did not affect action potentials in sugar- or bitter-activated gustatory receptor neurons (GRNs) when the GRNs were presented with just one type of tastant. OBP49a was expressed in accessory cells, and acted non-cell autonomously to attenuate nerve firings in sugar-activated GRNs when bitter compounds were combined with sucrose. These findings demonstrate an unexpected role for an OBP in taste, and identify a molecular player involved in the integration of opposing attractive and aversive gustatory inputs.

Introduction

Most foods are comprised of complex mixtures of different tastants, such as sweet and bitter compounds. Consequently, animal food preferences are decided by interactions between multiple constituents, many of which modulate the appeal or aversion of the component tastants. Suppression of the attractiveness of sweet by bitter tasting compounds has a strong survival benefit. Many tastants that are perceived as bitter are toxic, and so inhibition of stimulatory feeding behavior by these chemicals is critical.

When deterrent chemicals are present together with phagostimulatory tastants, they inhibit feeding by acting on two types of gustatory receptor cells. Aversive chemicals in foods not

© 2013 Elsevier Inc. All rights reserved.

⁵Correspondence: sjmoon@yuhs.ac or craig.montell@lifesci.ucsb.edu.

*These authors contributed equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

only stimulate deterrent taste cells, but also inhibit taste receptor cells that are activated by aversive compounds. This interaction between bitter and attractive gustatory stimuli has been observed in a wide array of vertebrate and invertebrate animals (Glendinning, 2007). Most studies dealing with the interactions between deterrent and attractive tastants have focused on quinine, a prototypical bitter compound. Electrophysiological recordings in hamsters show that the response to sucrose is inhibited by quinine (Formaker et al., 1997). In the catfish, quinine inhibits the positive gustatory response of several amino acids (Ogawa et al., 1997). Bitter compounds such as quinine are also aversive to flies (Tompkins et al., 1979), and suppress sugar-evoked firings in gustatory receptor neurons (GRNs) (Meunier et al., 2003).

The suppression of the stimulatory effect of attractive tastants by deterrent compounds could take place in the taste receptor cells, or in higher processing central pathways. While both sites might contribute to inhibition of sugar attractiveness by quinine, there is evidence that the afferent taste receptor cells are important for this phenomenon (Formaker et al., 1997; Talavera et al., 2008). Multiple mechanisms have been proposed to account for inhibition of sweet taste by quinine and other bitter compounds within the peripheral region of the gustatory system. The bitter-sweet interaction could be a consequence of lateral inhibition of sugar responsive gustatory receptor cells by bitter-activated neurons, similar to the inhibition of olfactory receptor neurons (ORNs) following activation of neighboring ORNs (Vandenbeuch et al., 2004; Su et al., 2012). Chemical interactions between the sugars and bitter compounds might also inhibit the attractiveness of the sugars. Competition of sugars and bitter chemicals for the same receptor is also plausible. An important advance is the demonstration that the effectiveness of the mammalian TRP channel, TRPM5, which is indirectly activated by sugars, via a G-protein coupled signaling pathway, is inhibited by quinine (Talavera et al., 2008). Thus, TRPM5 may provide one molecular mechanism through which quinine inhibits the attractiveness of sugars.

In *Drosophila* the molecular mechanism underlying the bitter-sweet interaction is unexplored. In view of an electrophysiological analysis, the site of this interaction is likely to be in the gustatory bristles (sensilla), which house the GRNs and accessory cells, and involve the taste receptors (Meunier et al., 2003). In fly GRNs, the largest class of taste receptors are referred to as gustatory receptors (GRs), which are distantly related to olfactory receptors (ORs) (Clyne et al., 1999; Gao and Chess, 1999; Vosshall et al., 1999; Clyne et al., 2000; Robertson et al., 2003; Scott, 2004; Montell, 2009). The ORs are more extensively characterized than the GRs, and are distinct from mammalian olfactory and taste receptors as the fly ORs are cation channels (Sato et al., 2008; Wicher et al., 2008). Thus, ORs provide the framework for many of the studies focusing on GRs, which may also be cation channels (Sato et al., 2011).

The direct ligand for at least one OR (OR67d) may not be the olfactory cue itself. Rather, there is evidence that the ligand for OR67d is an odorant binding protein (OBP), which is an extracellular protein present in the endolymph (Laughlin et al., 2008). The OBP, referred to as Lush, binds *in vitro* to OR67d when Lush is bound to a volatile pheromone (Laughlin et al., 2008). The actual receptor complex appears to be comprised of OR67d and a CD36-related protein, SNMP (Laughlin et al., 2008). However, whether Lush serves as the ligand *in vivo* is unresolved (Gomez-Diaz et al., 2013).

Some OBPs are expressed in gustatory sensilla (McKenna et al., 1994; Pikielny et al., 1994; Ozaki et al., 1995; Galindo and Smith, 2001; Shanbhag et al., 2001; Koganezawa and Shimada, 2002; Sánchez-Gracia et al., 2009; Yasukawa et al., 2010), although the family of 52 OBPs were identified originally in olfactory sensilla and are referred to as “odorant binding proteins” (Vogt and Riddiford, 1981). The roles of most OBPs have not been

reported, even in the olfactory system. Mutations affecting two OBPs that are expressed in taste sensilla (OBP57d/e) have been described. However, the contribution of these two OBPs to gustatory behavior appears to be small (Matsuo et al., 2007; Harada et al., 2008). Thus, the functions of OBPs in the gustatory response are largely unknown.

Here, we report an unexpected role for a *Drosophila* OBP, referred to as OBP49a. Loss of OBP49a had no impact on the production of action potentials in response to any deterrent or attractive compound tested. Rather, OBP49a was expressed in accessory cells and required by sweet activated GRNs for suppression of the attractive sugar responsive by bitter compounds. These findings provide the first molecular handle on the enigmatic phenomenon by which a deterrent compound inhibits the phagostimulatory signal of an attractive tastant in flies.

Results

OBPs are enriched in taste organs

In a previous study, we performed a DNA microarray analysis and identified *Drosophila* genes that were expressed preferentially in gustatory sensilla on the main taste organ, the labellum (Moon et al., 2009). In this analysis, we found that several genes encoding OBPs were the genes that were the most highly enriched in gustatory sensilla. To evaluate the reliability of the microarray data, we performed quantitative PCR. We prepared total RNA from the labella of control flies (*w¹¹¹⁸*), and from a mutant (*poxn*) in which the chemosensory bristles were transformed into mechanosensory bristles (Awasaki and Kimura, 1997). The enrichment profiles of each of the four *Obp* genes examined were similar to the microarray results (Figures 1A–1B). There were no major differences in the expression levels of the *Obps* in males and females.

Generation of mutations in *Obp56g*, *Obp19b*, *Obp49a* and *Obp57c*

To survey the roles of OBPs in the gustatory response, we generated mutations affecting four *Obp* genes. Three of the mutations disrupted *Obp56g*, *Obp19b* and *Obp49a*, which were the genes most enriched in gustatory organs (~100–800-fold; Figure 1A). In addition, we mutated the *Obp57c* gene, which was enriched in taste sensilla to a lower extent (~5-fold; Figure 1A).

To simultaneously create mutations and gene reporters we used ends-out homologous recombination. We generated a modified targeting vector (pw35*loxPGAL4*), which included the *GAL4* reporter juxtaposed to the *mini-white* marker gene (Figure 1C). We flanked both genes with *loxP* sequences to allow for removal of these genes with Cre recombinase. This would provide flexibility in cases in which it would be useful to introduce other reporters in combination with the mutations. We replaced the entire coding regions of *Obp56g*, *Obp19b* and *Obp49a* with the *GAL4* and the *mini-white* coding sequences (Figures 1D and S1A–S1B). To disrupt *Obp57c* expression, we substituted the start codon with a stop codon so that the neighboring, overlapping gene, *Obp57b*, would be minimally affected (Figure S1C). We confirmed each *Obp* knockout by PCR analysis of genomic DNA (Figures 1E and S1A–S1C).

OBP49a deficient flies showed reduced avoidance to sugar/bitter cocktails

To address whether the *Obp* mutations affected gustatory behavior, we performed two-way choice assays. The flies were given a choice between 1 mM sucrose and 5 mM sucrose mixed with either red or blue food coloring. After allowing the flies to feed for 90 minutes, we determined the preference indexes. All four mutant flies showed normal preferences for the higher concentration of sucrose (Figure S2A). If bitter compounds are combined with the

5 mM sucrose, wild-type prefer 1 mM sucrose (Moon et al., 2006; Lee et al., 2009). Three of the *Obp* mutants (*Obp56g¹*, *Obp19b¹* and *Obp57c¹*) showed normal repulsion to each of the bitter tastants tested (Figures 2A–2B and S2B–S2E). In contrast, mutation of *Obp49a* impaired the avoidance to a wide array of bitter compounds including papaverine, berberine, denatonium, quinine, caffeine, and strychnine (Figures 2 and S2B–S2E). The only exception was L-canavanine avoidance, which did not depend on any of the *Obp* mutants tested (Figure S2F). The decreased avoidances to the bitter chemical/5 mM sucrose mixtures were similar to those elicited by mutation of the broadly required gustatory receptor, *Gr33a* (Figures 2A–2B and S2B–S2D). We obtained *Obp49a^D* flies by excising the *GAL4* and *white* genes. *Obp49a^D* animals displayed the same defects in avoidance to the sucrose/aversive compound cocktails as the *Obp49a¹* animals (Figures 2C–2H).

OBP49a is expressed in thecogen cells

To test for expression of *Obp49a* in the major taste organ, the labellum, we took advantage of the *GAL4* reporter inserted in the *Obp49a* locus to drive expression of *UAS-mCD8::GFP*. The GFP was expressed broadly in many cells in the labellum (Figure 3A). To confirm the spatial distribution in the labellum, and to determine the cell type that expressed OBP49a, we raised OBP49a antibodies. The anti-OBP49a recognized a protein of the predicted size (23kDa) in the labellum of wild-type but not in *Obp49a¹* or *Obp49a^D* (Figure 3B). We performed immunocytochemistry and found that the anti-OBP49a signal was associated with all the chemosensory sensilla in wild-type but not *Obp49a¹* labella (Figure 3C). We double labeled labella from *Obp49a¹/UAS-mCD8::GFP* flies with anti-GFP and anti-OBP49a. The anti-OBP49a signal was distributed more broadly than the anti-GFP staining (Figure 3A), indicating that the reporter was expressed in a subset of OBP49a positive cells.

The 31 taste sensilla on each side of the labellum are classified into L-, I-, and S-type depending on their relative position and length (Vosshall and Stocker, 2007; Montell, 2009). L- and S-type sensilla house four GRNs and I-type sensilla contain two GRNs. Each sensillum also has three different types of accessory cells: tricogen (shaft), tormogen (socket), and thecogen (sheath). To address whether OBP49a was expressed in GRNs, we expressed a *UAS-mCD8::GFP* reporter under control of the *Gr5a-GAL4*, which labels GRNs that respond to attractive compounds such as sugars (Thorne et al., 2004; Wang et al., 2004). In addition, we used transgenic flies that expressed GFP in GRNs that are activated by aversive compounds such as caffeine and quinine (*Gr66a-I-GFP*) (Wang et al., 2004). Anti-OBP49a staining did not overlap with either of these markers (Figures 3D–3E) indicating that OBP49a was expressed in other cells in close proximity to cells marked with the *Gr5a* and *Gr66a* reporters.

To determine which non-neuronal cell type expressed OBP49a, we used markers that stained either the tormogen (*ASE5-GFP*) or the thecogen (*nompA-GAL4*) (Barolo et al., 2000; Chung et al., 2001). Anti-OBP49a was distributed in *nompA* positive cells, indicating that OBP49a was in thecogen cells, but not in cells expressing the *ASE5* reporter (Figures 3F–3G). Moreover, the *nompA-GAL4* reporter in combination with *UAS-Obp49a* restored normal aversion to bitter compound/5 mM sucrose mixtures in an *Obp49a^D* mutant background (Figures 2C–2H).

Requirement for OBP49a for suppression of sucrose-induced action potentials by bitter chemicals

In the olfactory system, the OBP referred to as Lush impacts on the activity of a subset of olfactory receptor neurons (Xu et al., 2005). Thus, in the gustatory system, OBP49a might contribute to the decreased attraction to 5 mM sucrose in the presence of a bitter tastant by affecting the activity of GRNs. If so, OBP49a could act on either of two different types of

GRNs. OBP49a could promote the activity of GRNs that respond exclusively to aversive compounds. Alternatively, OBP49a might be needed to suppress GRNs that are activated by sugars.

To address whether the activities of either the sugar or bitter-responsive GRNs were altered in *Obp49a* mutant labella, we performed tip-recordings, which measure nerve firings elicited by tastants. We focused on L-type sensilla to monitor sucrose-induced action potentials, and S-type sensilla to assay the responses to bitter compounds. Application of sucrose to L-type sensilla (L3, L4, L5, and L6), or aversive chemicals to S-type sensilla (S3, S5, S6, and S10), resulted in virtually the same frequencies of action potentials in wild-type and *Obp49a¹* animals (Figure 4).

The preceding data indicated that OBP49a was not required for stimulation of GRNs by either sweet or bitter compounds. Therefore, we explored the possibility that OBP49a was required for inhibition of the sweet response by bitter chemicals. L-type sensilla house GRNs that are activated by sugars, water, low salt, and high salt, but they do not respond to bitter chemicals (Hiroi et al., 2004; Weiss et al., 2011), thereby allowing us to assay inhibition of sucrose-elicited spikes by bitter chemicals. As described above, L-type sensilla from either wild-type or *Obp49a¹* flies displayed robust action potentials in response to 10 mM sucrose. When we exposed wild-type L-type sensilla to 10 mM sucrose combined with bitter chemicals, the responses were inhibited in a dose-dependent manner (Figures 5A–5F). The responses inhibited by bitter chemicals were generated by sugar-activated GRNs rather than water GRNs because we observed the same extent of inhibition by bitter chemicals in flies missing a channel, Pickpocket28 (*Ppk28*), required for water sensitivity (Figures S3A–S3C) (Cameron et al., 2010; Chen et al., 2010).

Of significance here, inhibition of the sucrose-induced action potentials by bitter chemicals was greatly reduced in *Obp49a¹* and *Obp49a^D* flies (Figures 5A–5F). The impairments in inhibition of sucrose-stimulated nerve firings by aversive chemicals were rescued by expression of wild-type OBP49a in the co-receptor cells using the *GAL4/UAS* system (Figures 5A–5F). The contribution of OBP49a to inhibition of the sucrose response by aversive compounds was broad, as the impairments occurred in response to a wide range of aversive tastants. Mutation of *Obp49a* had no impact on action potentials in L-type sensilla when the sucrose was combined with L-canavanine (Figure 5G). This was expected since L-canavanine did not suppress sucrose-induced nerve firings in wild-type (Figures 5G).

The preceding results suggest that OBP49a is required for inhibiting sucrose-responsive GRs, which may be cation channels (Sato et al., 2011). To test whether bitter compounds and OBP49a might affect the activity of another *Drosophila* cation channel, we ectopically expressed TRPA1 in sugar responsive GRN under the control of the *Gr5a-GAL4*. TRPA1 was activated by N-methylmaleimide to the same extent in the presence or absence of either berberine or OBP49a (Figures S3E–S3H).

Since bitter chemicals can inhibit water spikes, we examined whether OBP49a was also involved in the inhibition of water spikes by bitter chemicals (Meunier et al., 2003). Application of 1 mM KCl evoked nerve firings in L-type sensilla from wild-type and *Obp49a¹* flies but not in the Δ *ppk28* mutant, indicating that the spikes were from water responsive GRNs (Figure S3D). Increasing concentrations of bitter chemicals reduced water spikes in L-type sensilla to the same extent in both wild-type and *Obp49a¹* flies (Figure S3D). Thus, OBP49a was dispensable for the suppression of the water response by bitter chemicals.

Non-redundant roles for OBP49a and GR33a in sensing bitter chemicals

The preceding tip recording analysis indicated that OBP49a participated in suppression of the behavioral attraction to sweet compounds by bitter tastants by attenuating action potentials in sucrose-activated GRNs. The *Obp49a¹* behavioral phenotype was the same as that displayed by *Gr33a¹* mutants, even though *Gr33a* functions in the GRNs in S- and I-type sensilla, which are activated by bitter compounds. Therefore, if *Gr33a* and *Obp49a* act on different GRNs, then the *Gr33a¹, Obp49a¹* double mutant should show a more severe phenotype than either the *Gr33a¹* or *Obp49a¹* single mutants. Alternatively, if *Gr33a* and *Obp49a* acted through a common mechanism in the same GRNs, then the phenotypes of the double and single mutants would be expected to be the same.

We found that the defect in avoidance of the aversive chemical/sucrose cocktail was more severe in the *Gr33a¹, Obp49a¹* double-mutant animals than in *Gr33a¹* or *Obp49a¹* flies (Figures 6A–6E). The only exception was with strychnine (Figure 6F), which was consistent with our previous finding that *Gr33a¹* flies did not display a behavioral defect in strychnine avoidance (Moon et al., 2009). These findings support the conclusion that OBP49a and GR33a are involved in bitter chemical sensing through distinct pathways.

Rescue of the *Obp49a* phenotype by tethering OBP49a to GR5a-expressing cells

OBPs are secreted into the extracellular endolymph in chemosensory sensilla, and therefore have the potential to function non-cell autonomously. The finding that mutation of *Obp49a* impaired the suppression of sucrose-induced action potentials by bitter compounds indicated that OBP49a normally acted on sugar-responsive GRNs. To test this proposal, we expressed a membrane-tethered version of OBP49a so that OBP49a would be displayed extracellularly, but remain attached to the expressing cells. To do so, we generated transgenic flies expressing a form of OBP49a that was fused at the C-terminal end to a MYC linker and a transmembrane domain from the platelet derived growth factor receptor (OBP49a-t) (Figure 7A). We used the *GAL4/UAS* system to express *UAS-Obp49a-t* in sugar-activated GRNs (*Gr5a-GAL4*), bitter-activated GRNs (*Gr33a-GAL4*), or thecogen cells (*nompA-GAL4*), which synthesize OBP49a.

We found that *Obp49a-t* restored normal suppression of the sucrose response in *Obp49a^D* animals, but only if it was expressed in sugar-activated GRNs. This included normal behavioral suppression (Figures 7C–7F), and inhibition of sucrose-induced action potentials by bitter compounds such as berberine, denatonium, quinine and caffeine (Figures 8A–8C and S4A). This was not due to a non-specific effect of OBP49a-t on sugar-activated GRNs because expression of *UAS-Obp49a-t* under the control of the *Gr5a-GAL4* did not alter either the behavioral or electrophysiological responses to sucrose (Figures 7B and S4B). Expression of *Obp49a-t* either in GRNs that are activated by bitter compounds or in the thecogen cells did not rescue the *Obp49a^D* phenotype (Figure 7).

Binding of bitter compounds to OBP49a

The requirement for OBP49a for bitter-induced suppression of the sugar response raised the possibility that it binds to aversive tastants. To test for direct interactions of OBP49a with bitter chemicals, we employed surface plasmon resonance. We ectopically expressed *UAS-Obp49a* in compound eyes under the control of the *GMR-GAL4*, purified OBP49a from head extracts, and coupled the protein to sensor chips. We found that berberine, denatonium and quinine bound to OBP49a in a dose-dependent manner (Figures 8D–8F). In contrast, sucrose did not bind to OBP49a (Figure 8G), suggesting that OBP49a specifically interacted with bitter chemicals.

Close association of OBP49a and GR64a

The OBP49a-dependent suppression of the sucrose response by bitter compounds suggested that OBP49a might physically interact with the sucrose receptor. At least two GRs are required for sucrose detection. These include GR64a (Dahanukar et al., 2007; Jiao et al., 2007), as well as GR64f, which is required for sensing nearly all sugars, including sucrose, and may be a co-receptor for sugar-responsive GRs (Jiao et al., 2008).

To test whether OBP49a was in close proximity to GR64a or GR64f (Dahanukar et al., 2007; Jiao et al., 2007) and might therefore associate directly, we employed a YFP-based protein complementation assay. YFP can be split into two complementing fragments, and fluorescence is generated only when the separated parts are brought together. To address whether OBP49a was juxtaposed or interacted with either GR64a or GR64f *in vivo*, we generated *UAS*-transgenes encoding the N-terminal YFP fragment, YFP(1), fused to the N-termini of GR64a and GR64f, and the C-terminal YFP fragment YFP(2) linked to the C-termini of OBP49-t. As a control we used a previously described transgene *UAS-SNMP1:YFP(2)*, which encoded YFP(2) linked to a CD36-related receptor, SNMP1 (Benton et al., 2007). SNMP1 functions in pheromone detection in olfactory receptor neurons (Benton et al., 2007). We expressed these constructs in sugar-responsive GRNs under control of the *Gr5a-GAL4*.

We assayed for YFP-based protein complementation by dissecting labella from the transgenic flies, and performing confocal microscopy. There was no fluorescence visible in labella isolated from flies harboring the transgenes encoding just a single YFP(1) or YFP(2) fusion protein, such as YFP(1):GR64a or OBP49a-t-YFP(2) (Figure 8H). In contrast, co-expression of YFP(1):GR64a and OBP49a-t-YFP(2) in sugar responsive GRNs produced a strong signal (Figure 8I). To address whether a bitter chemical might enhance the fluorescence, we dipped the labella from immobilized flies in berberine before dissecting the labella. However, exposure to this bitter tastant had no impact on the fluorescence (Figure 8I). We did not detect signals when we expressed OBP49a-t-YFP(2) with YFP(1):GR64f (Figure 8J). The combination of YFP(1):GR64a with SNMP1-YFP(2) also did not produce fluorescence (Figure 8K). These findings support the conclusion that OBP49a either interacts with or is adjacent to GR64a.

Discussion

Many bitter-tasting chemicals are toxic (Glendinning, 2007). Therefore, the ability of animals to suppress their attraction to sugars and other nutritious foods that are laced with bitter tastants is critical for survival. Consequently, this avoidance behavior is conserved throughout the animal kingdom. Nevertheless, the molecules and molecular mechanisms through which positive feeding behavior is inhibited by deterrent compounds are poorly unexplored in most animals, such as the fruit fly.

Potentially, there are multiple neural mechanisms that could explain how aversive tastants suppress the otherwise stimulatory effects of sweeteners and other attractive compounds. Animals ranging from flies to humans have separate taste receptor cells devoted to sensing bitter and sweet tastants, and the suppression of sweet by bitter compounds could take place through integration of separate inputs in the brain. There could also be lateral interactions in the periphery between separate sweet and bitter responsive afferent receptor cells. Alternatively, a bitter compound might directly suppress sweet-activated taste receptor cells.

In this study, we unexpectedly identified a mechanism through which an array of bitter compounds inhibited the stimulatory effects of sucrose in flies. This work emerged from a functional analysis of OBPs in *Drosophila* taste, and was motivated by the finding that

multiple *Obp* genes were highly enriched in taste sensilla (McKenna et al., 1994; Pikielny et al., 1994; Ozaki et al., 1995; Galindo and Smith, 2001; Shanbhag et al., 2001; Koganezawa and Shimada, 2002; Sánchez-Gracia et al., 2009; Yasukawa et al., 2010), but their roles in the gustatory system were largely unexplored. We found that OBP49a was required for avoiding bitter-tasting compounds in a standard two-way choice assay consisting of 1 mM sucrose alone versus 5 mM sucrose plus bitter tastants. Because wild-type flies find bitter tastants aversive, they prefer the lower concentration of sucrose, when the higher concentration of sucrose is laced with tastants such as berberine, quinine or denatonium. However, the *Obp49a* mutant animals were impaired in this avoidance behavior. The phenotype was similar to that resulting from elimination of GRs, such as GR33a and GR66a, which are broadly expressed in avoidance GRNs, and are necessary in these GRNs for induction of bitter-induced action potentials (Moon et al., 2006; Lee et al., 2009; Moon et al., 2009). Unlike the *Gr33a* and *Gr66a* mutants, the decreased avoidance in the *Obp49a¹* animals was not due to impairment of aversive tastant-induced action potentials in *Gr66a*-expressing GRNs.

We conclude that the decreased behavioral avoidance to sucrose/aversive chemicals mixture in *Obp49a* mutant flies was due to a deficit in the sugar-activated GRNs that express *Gr5a*, and not due to effects on GRNs activated by bitter compounds. In support of the conclusion, the *Gr5a*- and *Gr66a*-expressing GRNs from *Obp49a¹* sensilla produced normal action potential frequencies in response to sugars and bitter compounds, respectively. However, when sucrose was combined with bitter tastants, the normal inhibition of the sugar-induced nerve firings was strongly impaired. OBP49a is therefore the first molecule shown to promote the inhibition of the sucrose response by aversive chemicals in *Drosophila*.

We propose that OBP49a, which is synthesized and secreted by thecogen cells into the endolymph, acts directly on sugar-activated (*Gr5a*-expressing) cells to inhibit sugar-induced action potentials, in response to bitter compounds. Further supporting this proposal, we rescued the *Obp49a^D* mutant phenotype with a membrane tagged version of OBP49a (OBP49a-t) that was restricted to the external surface of *Gr5a*-expressing cells, but not if OBP49a-t was confined to thecogen cells or *Gr66a*-expressing cells. Even though OBP49a was normally produced in thecogen cells, we were not able to rescue the *Obp49a* mutant phenotype if we expressed OBP49a-t in these accessory cells. Conversely, expression of an untagged version of OBP49a in thecogen cells fully restored normal inhibition of sucrose-induced action potentials by bitter compounds. These findings demonstrate that OBP49a was required non-cell autonomously by sugar-responsive cells.

Our findings indicate that at least one important cellular mechanism through which bitter and sweet taste is integrated, occurs in the taste receptor neurons. However, the current study does not exclude that there may also be integration of bitter and sugar activated signals in the brain. While *Obp49a* mutants showed a significant deficit in suppression of the sugar response by bitter compounds, the inhibition was not eliminated at the highest concentrations of the noxious tastants. Thus, there is an additional mechanism that contributes to inhibition of the sucrose response by bitter chemicals.

Mammals appear to use a similar cellular strategy for suppressing the appeal of sugars, when they are mixed with the bitter compound, quinine. In mammalian taste buds, sugars activate G-protein coupled receptors, and the signaling pathway culminates with activation of the TRPM5 channel (Pérez et al., 2002; Zhang et al., 2003). Quinine has a profound effect on inhibiting TRPM5, thereby reducing the attraction to sugars (Talavera et al., 2008). However, this TRPM5 mechanism may be relatively specific to quinine, since another bitter compound, denatonium, is ~100-fold less effective at inhibiting TRPM5 (Talavera et al.,

2008). By contrast, the OBP49a-dependent mechanism described here participates in the inhibition of the sucrose response to a wide array of aversive tastants.

There are at least two possible molecular mechanisms through which localization of OBP49a at the cell surface of sugar-responsive GRNs inhibits these neurons. OBP49a binds directly to bitter compounds and either interacts with or lies in close proximity to the sucrose receptor, GR64a. According to one possibility, OBP49a might deliver bitter chemicals to the cell surface of sugar-activated GRs, thereby greatly increasing the local concentration of bitter chemicals. The bitter chemicals might then bind to sugar-activated GRs, causing them to change from a high affinity state to low affinity state for sugars. Alternatively, the bitter chemicals might not bind directly to sugar-activated GRs, even at very high concentrations. Rather, once bound to bitter tastants, OBP49a might undergo a conformational change that in turn inhibits the GR64a complex. Since GRs may be cation channels (Sato et al., 2011), OBP49a might provide insects a mechanism by which bitter compounds suppress sugar-activated cation conductances.

Experimental Procedures

Fly stocks

All fly stocks were maintained on conventional cornmeal-agar-molasses medium, 12-h light/12-h dark cycles at 25°C and 60 % humidity. *70FLP, 70I-SceII/CyO, Sco/CyO, P[w⁺, Cre], UAS-mCD8::GFP* flies were obtained from the Bloomington Stock Center. *Gr5a-GAL4* and *Gr66a-/-GFP* were from K. Scott. *ASE5-GFP, nompA-GAL4, and UAS-SNMP1-YFP(2)* were provided by J.W. Posakony, Y.D. Chung, and L. Vosshall, respectively.

Construction of the pw35loxPGAL4 vector

To generate pw35loxPGAL4, we modified the pw35GAL4 vector (Moon et al, 2009) We inserted loxP oligonucleotides into the NotI and Acc65I sites. Each oligonucleotide also included portions of the NotI and Acc65I sites so that these two restriction sites were preserved. The orientation of the loxP sequences were in same orientation so that we could remove the floxed *mini-white* and the *GAL4* coding sequences after genetically introducing the Cre recombinase.

Generation of the *Obp* mutants

To generate the *Obp19b¹*, *Obp49a¹*, and *Obp56g¹* alleles, we PCR amplified 3 kb genomic DNAs encompassing both the 5' and 3' ends of the *Obp* coding sequences from isogenic *w¹¹¹⁸* flies. The genomic fragments were selected to introduce deletions of 930, 759 and 465 bp, respectively. To produce the *OBP57c¹* allele, we PCR amplified from isogenic *w¹¹¹⁸* flies a 3 kb of genomic DNA extending from the 5' end of the start codon, and a 3 kb genomic DNA extending from the 3' side of the start codon. This latter DNA included a stop codon at codon position one. Each homologous arm was subcloned into the pw35loxPGAL4 vector. The transgenic flies were generated by first obtaining random insertions of the transgenes (BestGene Inc., Chino Hills, CA), and then by mobilizing the transgenes and screening for targeted insertions as described previously (Gong and Golic, 2003). Each *Obp* mutation was confirmed by genomic PCR. The primer information used to confirm the deletions are: 5'-ACATCGTCGAGATGGTGCTGAACA-3' and 5'-TCCGCATCTGGAATACACTTCGCT-3' for the *Obp19b¹* deletion. 5'-TGACTTGCACACGCTGTGAATACG-3' (P1), 5'-TACCCACAATCCGTTTCAGT-3' (P2), and 5'-ATACAGCCAGCATTTCCCTTTCGGG-3' (P3) for the *Obp49a¹* deletion. 5'-TGTCOAAGGAAGTGGTGACGGATT-3' and 5'-TGGCTGCCTGGAAGCAGGATAATA-3' for the *Obp56g¹* deletion. 5'-TTCAGAACCGCCTTCTGCGAGTA-3' and 5'-

TGGCCAGGCAATCTGACACGTAAT-3' for the *Obp57c^l* deletion. To generate the *Obp49a^D* allele, the *Obp49a^l* flies were crossed to flies containing the P[*w^t, Cre*] transgene. The mosaic-eyed progeny were collected and crossed to balancer flies, and the white-eyed flies progeny of this latter cross were subjective to genomic PCR analysis using primers P1 and P3.

Creation of flies expressing a membrane tethered OBP49a-t

To generate the *UAS-Obp49a-t* transgenic flies, we first amplified the *Obp49a* coding sequence lacking the translation stop codon from *w¹¹¹⁸* labellar cDNA using the Hifidelity PCR kit (Roche), and cloned the cDNA into the pUAST vector. Sequences encoding the 10 amino acid MYC linker (EQKLISEEDL) and the transmembrane domain from the platelet derived growth factor receptor were amplified from the pDisplay vector (Invitrogen), and cloned in-frame 3' to the coding region for *Obp49a*. We also subcloned the cDNA encoding *Obp49a* with a normal stop codon and without the sequences encoding MYC and the membrane-tethered tag (*UAS-Obp49a*) into the pUAST vector. The transgenic flies were generated by BestGene Inc. (Chino Hills, CA).

Real time PCR analysis

We extracted total RNA from the labella of adult male and female wild-type and *poxn* flies using the Trizol reagent (Invitrogen, Carlsbad, CA), and generated cDNAs from 0.5 µg RNA using the SuperScript III First Strand Synthesis System (Invitrogen). Quantitative PCR was performed using an ABI7500 real-time PCR machine (Applied Biosystems, Foster City, CA) and the ABI SYBR green system. Transcript levels were normalized to *rp49* as an internal control and the $\Delta\Delta CT$ (CT = threshold cycle) method was used to calculate the relative amount of mRNAs. We repeated the experiments at least four times.

Anti-OBP49a antibodies and Western blots

Rabbit polyclonal OBP49a antibodies were raised to a synthetic peptide (CKPPRGPPPSAEDM; amino acid 199–212). 20 labella were dissected from wild-type, *Obp49a^l*, and *Obp49a^D* flies and homogenized in 1X SDS sample buffer with a pellet pestles (Kimble-Kontes). The extracts were subjected to electrophoresis by SDS-PAGE and transferred to PVDF membranes (Millipore). The membranes were probed with primary antibodies against OBP49a (1:1,000) and Tubulin (1:3,000, 12G10 from Hybridoma Bank), and then with peroxidase-conjugated anti-mouse or rabbit IgG secondary antibodies (1:5,000, Sigma).

Immunohistochemistry

Whole mount fly labellar immunostaining was performed as described previously (Moon et al., 2009) using anti-OBP49a (1:400) and mouse anti-GFP (1:400, Molecular Probes) primary antibodies, and anti-mouse-Alexa488 (1:400, Molecular Probes) and anti-rabbit-Alexa568 (1:400, Molecular Probes) secondary antibodies. The stained samples were mounted with Vectashield (Vector Laboratories, Burlingame, CA) and visualized with a Zeiss LSM700 confocal microscope (Jena, Germany).

Chemicals

Sucrose, denatonium, quinine, papaverine, caffeine, strychnine, L-canavanine, sulforhodamine B and KCl were purchased from Sigma-Aldrich (Saint Louis, MO). Berberine sulfate trihydrate, and Brilliant Blue FCF were obtained from Wako Pure Chemical Industries, Ltd (Osaka, Japan).

Two-way choice behavioral tests

The binary food choice assays were performed as described previously (Meunier et al., 2003; Moon et al., 2006). Briefly 3–6 days old flies were starved for 18 hours and then placed in 72 well microtiter dishes. Each alternating well was filled with 1% agarose combined with one of two types of test mixtures. To perform the sucrose test, the wells contained either 5 mM or 1 mM sucrose. The aversion to bitter chemicals was assayed by comparing the preferences for 1 mM sucrose versus 5 mM sucrose plus the indicated concentrations of aversive compounds. To monitor food intake, one test mixture contained blue dye (Brilliant Blue FCF, 0.125 mg/ml) while the other contained red dye (sulforhodamine B, 0.2 mg/ml). After allowing the flies to feed for 90 minutes at room temperature in the dark, the animals were frozen at -20°C . The numbers of flies that were blue (N^{B}), red (N^{R}) or purple (N^{MIX}) were determined under a dissection microscope and the preference index (P.I.) values were calculated according to the following equation: $(N^{\text{R}}+0.5N^{\text{MIX}})/(N^{\text{R}}+N^{\text{B}}+N^{\text{MIX}})$. P.I.s=1.0 and 0 indicated complete preferences for one or the other food alternative. A P.I.=0.5 indicated no preference.

Electrophysiology

Tip recordings (Hodgson et al., 1955; Wieczorek and Wolff, 1989) were performed as described previously (Moon et al., 2006). 1-day-old flies, which were kept on fresh fly food after eclosion, were immobilized by inserting a glass capillary, which was filled with Ringer's solution, into the abdomen through to the head. This electrode also served as a reference electrode. The indicated labellar sensilla were stimulated with a recording electrode (10–20 μm tip diameter) containing the test tastants in 1mM KCl as the electrolyte. The recording electrode was connected to a preamplifier (TastePROBE, Syntech, Hilversum, The Netherlands). The signals were collected and amplified (10 \times) using a signal connection interface box (Syntech) in conjunction with a 100–3000 Hz band-pass filter. The inputs were also linked to a loudspeaker to facilitate audio monitoring. Recordings of action potentials were acquired at a 12-kHz sampling rate and analyzed with Autospike 3.1 software (Syntech). The spikes were sorted based on their amplitude for further quantitative analyses.

OBP49a expression and purification

OBP49a was expressed in fly eyes under the control of the long *GMR-GAL4* (Wernet et al., 2003). The fly heads expressing OBP49a in eyes were separated from bodies by agitation of frozen flies. 10 ml of collected fly heads were homogenized in 25 ml of 10 mM Tris-HCl, pH 7.4, 10 % glycerol, using a motor-driven homogenizer and further homogenized with a Dounce homogenizer. The extracts were centrifuged for 10 min at 3,000 $\times g$, and the supernatants were ultracentrifuged at 4°C for 60 min at 100,000 $\times g$. OBP49a was purified by serial use of the HiTrap SP XL 5 ml and HiTrap Q XL 5 ml columns (GE Healthcare), followed by affinity purification with OBP49a antibodies. The purity of OBP49a was assessed by fractionation of the protein by SDS-PAGE and silver staining (Figures S4C and S4D).

Surface Plasmon Resonance (SPR) binding experiments

SPR was conducted using a BIAcore 3000 (GE Healthcare) at 25°C . Coupling of OBP49a to CM5 chips (GE Healthcare) was performed by injecting 0.1 $\mu\text{g}/\text{ml}$ of protein with 10 mM sodium acetate, pH 4.0 at a 5 $\mu\text{l}/\text{min}$ flow rate, and confirmed by an increase of 10,000 resonance units on the sensor chip. The chemicals were diluted to the indicated concentrations in the continuous flow buffer HBS-P (10 mM HEPES pH 7.4, 150 mM NaCl, 0.005 % Surfactant P20). Each analytic run was performed at 30 $\mu\text{l}/\text{min}$ flow rate. The chip matrix was regenerated using 20 mM NaOH after each binding analysis.

Protein complementation assay (PCA)

To obtain the *UAS-YFP(1):Gr64a*, *UAS-YFP(1):Gr64f* transgenic flies for the PCA, we first generated pUAST-*YFP(1)* by PCR amplifying a 462 base pair *YFP(1)* fragment from pAKAR3EV (Komatsu et al., 2011) that extended from the Kozak sequence. This fragment was subcloned between the EcoRI and KpnI sites of pUAST. We then inserted the coding sequences of *Gr64a* and *Gr64f* into pUAST-*YFP(1)*, so that YFP(1) was linked to the N-termini of the GRs. To produce the *pUAST-OBP49a-t-YFP(2)* construct, we used pUAST-*Obp49a-t* to PCR amplify the *OBP49a-t* coding sequence that lacked the stop codon, and then inserted the fragment into pUAST. We then used pAKAR3EV as the template to PCR amplify two pieces of DNA fragments encoding a 116 amino acid long flexible EV linker and YFP(2), which encoded residues 155 to 237 of YFP. We inserted these DNA fragments adjacent to the 3' end of the *OBP49a-t* coding region. We expressed these transgenes, as well as *UAS-Snmp1-YFP(2)* (Benton et al., 2007), under the control of the *Gr5a-GAL4*. To apply berberine to the sensilla, we immobilized the flies with a glass capillary and dipped the labella into a solution containing 100 μ M berberine for one minute before dissecting the labella. We also immersed labella in 100 μ M berberine/100 mM sucrose solutions, and obtained indistinguishable results from that generated with untreated labella, or labella dipped in berberine only (data not shown). The labella were fixed with 4% paraformaldehyde in PBS-T (0.2% Triton X-100 in PBS) for 20 min. Fixed labella were washed with PBS-T three times and cut in half with a razor blade and then, mounted in VECTASHELD (Vector Laboratories). Fluorescence was viewed in whole mounts of labella using a Zeiss LSM700 confocal microscope.

Data analyses

All error bars represent SEMs. Unpaired Student's *t*-tests were used to compare two sets of data. ANOVA with the Tukey *post hoc* tests were used to compare multiple sets of data. Asterisks indicate statistical significance (* $p < 0.05$, ** $p < 0.01$).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Bloomington Stock Center, and Drs. K. Scott, J.W. Posakony and Y.D. Chung for fly stocks, and Dr. C.Y. Park and Mr. J. Song at Surface Plasmon Resonance Facility, UNIST for technical help. The tubulin monoclonal antibody developed by Drs. Frankel and Nelsen was obtained from the Developmental Studies Hybridoma Bank developed under the auspices of the NICHD and maintained by The University of Iowa, Department of Biology, Iowa City, IA 52242. This work was supported by a grant to C.M. from the NIDCD (DC007864) and by a grant to S.J.M. from the Converging Research Center Program funded by the Ministry of Education, Science, and Technology (2012K001350) and from the Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science, and Technology (2009-0075341 and 2012R1A1A1012081).

References

- Awasaki T, Kimura K. *pox-neuro* is required for development of chemosensory bristles in *Drosophila*. *J. Neurobiol.* 1997; 32:707–721. [PubMed: 9183748]
- Barolo S, Walker RG, Polyanovsky AD, Freschi G, Keil T, Posakony JW. A Notch-independent activity of Suppressor of Hairless is required for normal mechanoreceptor physiology. *Cell.* 2000; 103:957–969. [PubMed: 11136980]
- Benton R, Vannice KS, Voshall LB. An essential role for a CD36-related receptor in pheromone detection in *Drosophila*. *Nature.* 2007; 450:289–293. [PubMed: 17943085]

- Cameron P, Hiroi M, Ngai J, Scott K. The molecular basis for water taste in *Drosophila*. *Nature*. 2010; 465:91–95. [PubMed: 20364123]
- Chen Z, Wang Q, Wang Z. The amiloride-sensitive epithelial Na⁺ channel PPK28 is essential for *Drosophila* gustatory water reception. *J. Neurosci*. 2010; 30:6247–6252. [PubMed: 20445050]
- Chung YD, Zhu J, Han Y, Kernan MJ. *nompA* encodes a PNS-specific, ZP domain protein required to connect mechanosensory dendrites to sensory structures. *Neuron*. 2001; 29:415–428. [PubMed: 11239432]
- Clyne PJ, Warr CG, Carlson JR. Candidate taste receptors in *Drosophila*. *Science*. 2000; 287:1830–1834. [PubMed: 10710312]
- Clyne PJ, Warr CG, Freeman MR, Lessing D, Kim J, Carlson JR. A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron*. 1999; 22:327–338. [PubMed: 10069338]
- Dahanukar A, Lei YT, Kwon JY, Carlson JR. Two *Gr* genes underlie sugar reception in *Drosophila*. *Neuron*. 2007; 56:503–516. [PubMed: 17988633]
- Formaker BK, MacKinnon BI, Hettinger TP, Frank ME. Opponent effects of quinine and sucrose on single fiber taste responses of the chorda tympani nerve. *Brain Res*. 1997; 772:239–242. [PubMed: 9406978]
- Galindo K, Smith DP. A large family of divergent *Drosophila* odorant-binding proteins expressed in gustatory and olfactory sensilla. *Genetics*. 2001; 159:1059–1072. [PubMed: 11729153]
- Gao Q, Chess A. Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence. *Genomics*. 1999; 60:31–39. [PubMed: 10458908]
- Glendinning JI. How do predators cope with chemically defended foods? *Biol Bull*. 2007; 213:252–266. [PubMed: 18083965]
- Gomez-Diaz C, Reina JH, Cambillau C, Benton R. Ligands for pheromone-sensing neurons are not conformationally activated odorant binding proteins. *PLoS Biol*. 2013; 11:e1001546. [PubMed: 23637570]
- Gong WJ, Golic KG. Ends-out, or replacement, gene targeting in *Drosophila*. *Proc. Natl. Acad. Sci. USA*. 2003; 100:2556–2561. [PubMed: 12589026]
- Harada E, Haba D, Aigaki T, Matsuo T. Behavioral analyses of mutants for two odorant-binding protein genes, *Obp57d* and *Obp57e*, in *Drosophila melanogaster*. *Genes Genet. Syst*. 2008; 83:257–264. [PubMed: 18670137]
- Hiroi M, Meunier N, Marion-Poll F, Tanimura T. Two antagonistic gustatory receptor neurons responding to sweet-salty and bitter taste in *Drosophila*. *J. Neurobiol*. 2004; 61:333–342. [PubMed: 15389687]
- Hodgson ES, Lettvin JY, Roeder KD. Physiology of a primary chemoreceptor unit. *Science*. 1955; 122:417–418. [PubMed: 13246649]
- Jiao Y, Moon SJ, Montell C. A *Drosophila* gustatory receptor required for the responses to sucrose, glucose, and maltose identified by mRNA tagging. *Proc. Natl. Acad. Sci. USA*. 2007; 104:14110–14115. [PubMed: 17715294]
- Jiao Y, Moon SJ, Wang X, Ren Q, Montell C. *Gr64f* is required in combination with other gustatory receptors for sugar detection in *Drosophila*. *Curr. Biol*. 2008; 18:1797–1801. [PubMed: 19026541]
- Koganezawa M, Shimada I. Novel odorant-binding proteins expressed in the taste tissue of the fly. *Chem. Senses*. 2002; 27:319–332. [PubMed: 12006372]
- Komatsu N, Aoki K, Yamada M, Yukinaga H, Fujita Y, Kamioka Y, Matsuda M. Development of an optimized backbone of FRET biosensors for kinases and GTPases. *Mol. Biol. Cell*. 2011; 22:4647–4656. [PubMed: 21976697]
- Laughlin JD, Ha TS, Jones DN, Smith DP. Activation of pheromone-sensitive neurons is mediated by conformational activation of pheromone-binding protein. *Cell*. 2008; 133:1255–1265. [PubMed: 18585358]
- Lee Y, Moon SJ, Montell C. Multiple gustatory receptors required for the caffeine response in *Drosophila*. *Proc. Natl. Acad. Sci. USA*. 2009; 106:4495–4500. [PubMed: 19246397]

- Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. Odorant-binding proteins OBP57d and OBP57e affect taste perception and host-plant preference in *Drosophila sechellia*. *PLoS Biol.* 2007; 5:e118. [PubMed: 17456006]
- McKenna MP, Hekmat-Scafe DS, Gaines P, Carlson JR. Putative *Drosophila* pheromone-binding proteins expressed in a subregion of the olfactory system. *J. Biol. Chem.* 1994; 269:16340–16347. [PubMed: 8206941]
- Meunier N, Marion-Poll F, Rospars JP, Tanimura T. Peripheral coding of bitter taste in *Drosophila*. *J. Neurobiol.* 2003; 56:139–152. [PubMed: 12838579]
- Montell C. A taste of the *Drosophila* gustatory receptors. *Curr. Opin. Neurobiol.* 2009; 19:345–353. [PubMed: 19660932]
- Moon SJ, Köttgen M, Jiao Y, Xu H, Montell C. A taste receptor required for the caffeine response in vivo. *Curr. Biol.* 2006; 16:1812–1817. [PubMed: 16979558]
- Moon SJ, Lee Y, Jiao Y, Montell C. A *Drosophila* gustatory receptor essential for aversive taste and inhibiting male-to-male courtship. *Curr. Biol.* 2009; 19:1623–1627. [PubMed: 19765987]
- Ogawa K, Marui T, Caprio J. Quinine suppression of single facial taste fiber responses in the channel catfish. *Brain Res.* 1997; 769:263–272. [PubMed: 9374194]
- Ozaki M, Morisaki K, Idei W, Ozaki K, Tokunaga F. A putative lipophilic stimulant carrier protein commonly found in the taste and olfactory systems. A unique member of the pheromone-binding protein superfamily. *Eur. J. Biochem.* 1995; 230:298–308. [PubMed: 7601113]
- Pérez CA, Huang L, Rong M, Kozak JA, Preuss AK, Zhang H, Max M, Margolskee RF. A transient receptor potential channel expressed in taste receptor cells. *Nat. Neurosci.* 2002; 5:1169–1176. [PubMed: 12368808]
- Pikielny CW, Hasan G, Rouyer F, Rosbash M. Members of a family of *Drosophila* putative odorant-binding proteins are expressed in different subsets of olfactory hairs. *Neuron.* 1994; 12:35–49. [PubMed: 7545907]
- Robertson HM, Warr CG, Carlson JR. Molecular evolution of the insect chemoreceptor gene superfamily in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA.* 2003; 2(100 Suppl):14537–14542. [PubMed: 14608037]
- Sánchez-Gracia A, Vieira FG, Rozas J. Molecular evolution of the major chemosensory gene families in insects. *Heredity.* 2009; 103:208–216. [PubMed: 19436326]
- Sato K, Pellegrino M, Nakagawa T, Nakagawa T, Vossshall LB, Touhara K. Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature.* 2008; 452:1002–1006. [PubMed: 18408712]
- Sato K, Tanaka K, Touhara K. Sugar-regulated cation channel formed by an insect gustatory receptor. *Proc. Natl. Acad. Sci. USA.* 2011; 108:11680–11685. [PubMed: 21709218]
- Scott K. The sweet and the bitter of mammalian taste. *Curr. Opin. Neurobiol.* 2004; 14:423–437. [PubMed: 15321062]
- Shanbhag SR, Park SK, Pikielny CW, Steinbrecht RA. Gustatory organs of *Drosophila melanogaster*: fine structure and expression of the putative odorant-binding protein PBPRP2. *Cell Tissue Res.* 2001; 304:423–437. [PubMed: 11456419]
- Su CY, Menuz K, Reiser J, Carlson JR. Non-synaptic inhibition between grouped neurons in an olfactory circuit. *Nature.* 2012; 492:66–71. [PubMed: 23172146]
- Talavera K, Yasumatsu K, Yoshida R, Margolskee RF, Voets T, Ninomiya Y, Nilius B. The taste transduction channel TRPM5 is a locus for bittersweet taste interactions. *FASEB J.* 2008; 22:1343–1355. [PubMed: 18070821]
- Thorne N, Chromey C, Bray S, Amrein H. Taste perception and coding in *Drosophila*. *Curr. Biol.* 2004; 14:1065–1079. [PubMed: 15202999]
- Tompkins L, Cardosa MJ, White FV, Sanders TG. Isolation and analysis of chemosensory behavior mutants in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA.* 1979; 76:884–887. [PubMed: 106397]
- Vandenbeuch A, Pillias AM, Faurion A. Modulation of taste peripheral signal through interpapillar inhibition in hamsters. *Neurosci. Lett.* 2004; 358:137–141. [PubMed: 15026167]
- Vogt RG, Riddiford LM. Pheromone binding and inactivation by moth antennae. *Nature.* 1981; 293:161–163. [PubMed: 18074618]

- Vosshall LB, Amrein H, Morozov PS, Rzhetsky A, Axel R. A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell*. 1999; 96:725–736. [PubMed: 10089887]
- Vosshall LB, Stocker RF. Molecular architecture of smell and taste in *Drosophila*. *Annu. Rev. Neurosci.* 2007; 30:505–533. [PubMed: 17506643]
- Wang Z, Singhvi A, Kong P, Scott K. Taste representations in the *Drosophila* brain. *Cell*. 2004; 117:981–991. [PubMed: 15210117]
- Weiss LA, Dahanukar A, Kwon JY, Banerjee D, Carlson JR. The molecular and cellular basis of bitter taste in *Drosophila*. *Neuron*. 2011; 69:258–272. [PubMed: 21262465]
- Wernet MF, Labhart T, Baumann F, Mazzoni EO, Pichaud F, Desplan C. Homothorax switches function of *Drosophila* photoreceptors from color to polarized light sensors. *Cell*. 2003; 115:267–279. [PubMed: 14636555]
- Wicher D, Schafer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS. *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature*. 2008; 452:1007–1011. [PubMed: 18408711]
- Wieczorek H, Wolff G. The labellar sugar receptor of *Drosophila*. *J. Comp. Physiol. A*. 1989; 164:825–834.
- Xu P, Atkinson R, Jones DN, Smith DP. *Drosophila* OBP LUSH is required for activity of pheromone-sensitive neurons. *Neuron*. 2005; 45:193–200. [PubMed: 15664171]
- Yasukawa J, Tomioka S, Aigaki T, Matsuo T. Evolution of expression patterns of two odorant-binding protein genes, *Obp57d* and *Obp57e*, in *Drosophila*. *Gene*. 2010
- Zhang Y, Hoon MA, Chandrashekar J, Mueller KL, Cook B, Wu D, Zuker CS, Ryba NJ. Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar signaling pathways. *Cell*. 2003; 112:293–301. [PubMed: 12581520]

Highlights

- Control of sweet/bitter taste interaction by an odorant binding protein.
- Inhibition of sweet-activated gustatory neurons by an odorant binding protein.
- Bitter compounds inhibit the sweet response by binding to OBP49a.




Figure 1. *Obp* genes with enriched expression in the labella, and generation of the *Obp49a* mutant

(A) Relative expression of *Obp* RNAs in the labella of wild-type and *poxn* mutants. RNA expression was analyzed by microarray analysis (blue bars) and real-time PCR (red bars). The real time PCR analysis was performed using RNA prepared from separate males (M) and females (F). The numbers on the Y-axis are arbitrary units, which indicate relative rather than absolute levels of the various RNAs. Shown are the means \pm SEM (n=3, microarray data; n=4–6, real time PCR). (B) Comparison of the enrichment of *Obp* genes as assessed by microarray and RT-PCR analyses using RNA prepared from males (M) and females (F). (C) The *pw35loxPGAL4* vector used to generate the *Obp* mutants. (D)

Schematic of the *Obp49a* locus and the targeting construct used to generate the *Obp49a^l* allele. The *Obp49a^D* allele was derived by removing the floxed *mini-white* and *GAL4* genes. The boxes represent exons, and the orientation of the *mini-white-GAL4* and *loxP* are indicated by the arrows. (E) Confirmation of the deletion in *Obp49a^l* and the excision of the *mini-white* and the *GAL4* in *Obp49a^D*. Genomic DNA was prepared and PCR performed using the primer pairs (P1, P2, and P3) indicated in panel D (see also Figure S1).




Figure 2. *Obp49a* mutants displayed reduced avoidance to sucrose/bitter mixtures

Two-way choice behavioral assays were performed by allowing the flies to choose between 1 mM sucrose versus 5 mM sucrose/aversive chemical cocktails. (A and B) Screen for *Obps* required for gustatory avoidance of papaverine and berberine. *Gr33a¹* was included as a positive control. (C–H) Testing for requirements for *Obp49a* for avoidance of a variety of sucrose/bitter cocktails. Both *Obp49a¹* and *Obp49a^D* showed similar defects in bitter avoidance. To test for rescue of the *Obp49a^D* phenotype, we expressed an *Obp49a⁺* transgene (*UAS-Obp49a*) in the coxigen cells (*nompA-GAL4*). The numbers of behavioral tests are indicated. Shown are the means ± SEM. The asterisks indicate statistically significant differences (***p*<0.01) from wild-type flies (see also Figure S2).




Figure 3. Expression of OBP49a in the labellum

(A) Co-expression of the *Obp49a* reporter and anti-OBP49a. *Obp49a¹/UAS-mCD8::GFP* labella were stained with GFP and OBP49a antibodies. (B) Western blot using OBP49a antibodies. Extracts were prepared from wild-type, *Obp49a¹*, and *Obp49a^D* labella, and the blot was probed with anti-OBP49a. The blot was also probed with anti-tubulin as a loading control. (C) Staining of labella from wild-type (top) and *Obp49a¹* flies with anti-OBP49a (bottom). The arrowhead indicates the position of the thecogen cells in the *Obp49a¹* mutant labellum. (D–G) OBP49a was expressed in thecogen cells. The cellular distribution of OBP49a was examined by comparing the spatial distribution of cell-type-specific markers with anti-OBP49a. (D) *Gr5a-GAL4/UAS-mCD8::GFP* (sweet GRNs). (E) *Gr66a-/-GFP*

(bitter GRNs). (F) *nompA-GAL4/UAS-mCD8::GFP* (thecogen). (G) *ASE5-GFP* (tormogen). Scale bars represent 50 μ M




Figure 4. *Obp49a¹* mutant animals showed normal action potential frequencies evoked by bitter tastants

Tip recordings were performed on the indicated S-type sensilla. The action potential frequencies were based on the number of spikes produced between 50 ms and 550 ms after initiating the stimulation. (A–G) Representative traces of nerve firings from S6 sensilla and quantification of the mean action potentials induced by the indicated concentration of bitter chemicals in S-type sensilla. (A) 1 mM papaverine. (B) 50 μ M berberine. (C) 1 mM quinine. (D) 1 mM denatonium. (E) 5 mM caffeine. (F) 1 mM strychnine. (G) 25 mM L-canavanine. (H) Representative traces of nerve firings from L6 sensilla and quantification of the mean action potentials induced by 10 mM sucrose in L-type sensilla. (I) Dose response analysis of

the action potential frequencies elicited by papaverine, berberine, and quinine from S6 sensilla. Data are the means \pm SEM (n=5—17).




Figure 5. Requirement for *Obp49a* for inhibition of sucrose-induced action potentials by aversive chemicals

Tip recordings of L-type sensilla (L4, L5, and L6) showing representative traces and the mean action potentials induced by 10 mM sucrose combined with aversive chemicals. Each stimulant contained 10 mM sucrose, 1 mM KCl as the electrolyte and the indicated concentrations of bitter chemicals. The tip recordings were performed using the indicated sensilla and the following bitter chemicals. (A) Papaverine. (B) Berberine. (C) Denatonium. (D) Quinine. (E) Caffeine. (F) Strychnine. (G) L-canavanine. Shown are the means \pm SEM (n=5–13). The genotype of the rescue flies was *Obp49a^D,nompA-GAL4/UASObp49a*. The

asterisks denote statistically significant differences (* $p < 0.05$, ** $p < 0.01$) from wild-type flies (see also Figure S3).




Figure 6. Phenotypes of single and double *Obp49a*¹ and *Gr33a*¹ mutants

The two-way choice behavioral assays were performed using 1 mM sucrose versus 5 mM sucrose combined with the indicated concentrations of bitter chemicals. (A) Papaverine. (B) Berberine. (C) Denatonium. (D) Quinine. (E) Caffeine. (F) Strychnine. Data are the means \pm SEM (n=6–11). The asterisks indicate statistically significant differences between *Gr33a*¹ and *Gr33a*¹, *Obp49a*¹ double mutant flies (**p < 0.01).




Figure 7. Membrane tethered OBP49a rescued *Obp49a^D* behavioral phenotype when expressed in sugar sensing GRNs

(A) Structure of the membrane tethered OBP49a (*OBP49a-t*). TMD, transmembrane domain. A 20 amino acid scale bar is shown. (B) Testing for effects of *OBP49a-t* expression on sucrose sensing behavior in sweet sensing GRNs (*Gr5a* GRNs). The flies were given a choice between 1 mM sucrose and 5 mM sucrose. (C–F) Testing for rescue of the impaired avoidance to sucrose/bitter cocktails in *Obp49a^D* flies by expressing *UAS-Obp49a-t* under the control of the *Gr5a-GAL4* (sweet sensing GRNs), *Gr33a-GAL4* (bitter sensing GRNs) or the *nompA-GAL4* (thecogen cells). The flies were given a choice between 1 mM sucrose and 5 mM sucrose mixed with a bitter tastant. (C) Berberine. (D) Denatonium. (E) Quinine.

(F) Caffeine. Shown are the means \pm SEMs. The numbers of assays performed are indicated. The asterisks indicate statistically significant differences between *Obp49a^D/UAS-Obp49a-t* and *Obp49a^D;Gr5a-GAL4/UAS-Obp49a-t* flies (**p<0.01).




Figure 8. Cellular requirement for OBP49a-t, and testing for interactions between OBP49a and bitter tastants and GR64a

(A–C) Tip-recordings were performed using the indicated sensilla and genotypes. Shown are representative traces and the mean numbers of action potentials ($n=8–11$) induced by 10 mM sucrose plus the indicated bitter tastant. (A) 10 μ M berberine. (B) 10 μ M denatonium. (C) 10 μ M quinine. The asterisks indicate statistically significant differences between *Obp49a^D+UAS-Obp49a-t* and *Obp49a^D;Gr5a-GAL4/UAS-Obp49a-t* flies (** $p<0.01$). (D–G) Sensorgrams showing binding between bitter tastants and OBP49a using the BIAcore system. The indicated concentrations of tastants were allowed to interact with the OBP49a, which was bound to CM5 chips. (R.U., relative resonance units). (H–K) Protein

complementation assay using the split YFP approach. Labella were dissected from transgenic flies expressing the indicated YFP(1) and YFP(2) fusion proteins in *Gr5a* GRNs, under control of the *GAL4/UAS* system. In some cases, 100 μ M berberine was applied to sensilla for 1 minute before dissection. The scale bar in (H) represents 10 μ m, and applies to all panels in (H–K) (see also Figure S4).