

Lawrence Berkeley National Laboratory

Recent Work

Title

DIFFERENTIAL CROSS SECTIONS FOR THE PRODUCTION OF POLARIZED POSITONS IN AN EXTERNAL FIELD

Permalink

<https://escholarship.org/uc/item/169240m1>

Author

Johnson, Coates R.

Publication Date

1958-09-16

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

DIFFERENTIAL CROSS SECTIONS FOR THE PRODUCTION
OF POLARIZED POSITONS AND NEGATONS BY POLARIZED
PHOTONS IN AN EXTERNAL FIELD

Coates R. Johnson

September 16, 1958

DIFFERENTIAL CROSS SECTIONS FOR THE PRODUCTION OF POLARIZED POSITONS AND
NEGATONS BY POLARIZED PHOTONS IN AN EXTERNAL FIELD

Coates R. Johnson

Radiation Laboratory
University of California
Berkeley, California

September 16, 1958

ABSTRACT

The differential cross sections for the production of polarized positons and negatons by polarized photons in an external field are calculated in the Born approximation.

* This work was done under the auspices of the U. S. Atomic Energy Commission.

DIFFERENTIAL CROSS SECTIONS FOR THE PRODUCTION OF PAIR PARTICLES BY POLARIZED PHOTONS IN AN EXTERNAL FIELD

Coates R. Johnson

Radiation Laboratory
University of California
Berkeley, California

September 16, 1958

I. INTRODUCTION

This paper reports the evaluation of the differential cross sections for pair production involving polarized particles in an external field, so that anyone who wishes to carry the calculations further may use them for a check. The reason for this particular calculation is the recent interest in shower phenomena involving polarized particles.

II. RELATION OF THE CROSS SECTION TO THE S MATRIX

The cross section for a process which involves only one photon in the initial state is given by

$$d\sigma = (2\pi)^2 \delta(E_f - E_i) \left| (f | m | i) \right|^2, \quad (1)$$

where E_f and E_i denote the final and initial energies and

$$(f | S | i) = \delta(E_f - E_i) (f | m | i). \quad (2)$$

The S matrix, $(f | S | i)$, is the matrix element of the operator S between the initial and final states considered in the particular problem, and the S operator in quantum electrodynamics is given by

$$S = 1 + \sum_{n=1}^{\infty} S_n \quad S_n = \frac{(-i)^n}{n!} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} d\tau_n \dots d\tau_1 P (H(\tau_n) \dots H(\tau_1)), \quad (3)$$

where P is the chronological operator and τ labels the spacelike surfaces

III. EVALUATION OF THE S MATRIX FOR PAIR PRODUCTION

In our case $H(\tau)$ can be split into two parts,

$$H(\tau) = - \int_{\sigma(\tau)} j_{\mu}(x) a^{\mu}(x) d\sigma - \int_{\sigma(\tau)} j_{\mu}(x) \phi^{\mu}(x) d\sigma, \quad (5)$$

where $j_{\mu}(x)$ is the electron current operator, $a^{\mu}(x)$ is the electromagnetic field operator, and $\phi^{\mu}(x)$ is the operator associated with the external electromagnetic field; $\phi^{\mu}(x)$ will be treated as a number (since we are justified in neglecting the quantum properties of this strong external field).

The terms of lowest order in e which contribute to the matrix element when the initial state contains just one photon and the final state one negaton and positron are given by

$$(p^+ p^- | S | k) = \frac{ie^2 mZe}{(2\pi)^2 (2\omega \epsilon_+ \epsilon_-)^{1/2} (2\pi)^{3/2}} \delta(\epsilon_+ + \epsilon_- - \omega) \phi(q) \bar{u}(p_-) Q(p_+ p_-) V(p_+), \quad (6)$$

where

$$Q(p_+ p_-) = \not{\underline{p}}_- \frac{1(\not{\underline{p}}_- - \not{\underline{u}}) - m}{(p_- - k)^2 + m^2} \not{\underline{n}} + \not{\underline{n}} \frac{1(\not{\underline{k}} - \not{\underline{p}}_+) - m}{(p_+ - k)^2 + m^2} \not{\underline{p}}_+,$$

$$\phi^{\mu}(x) = n^{\mu} \frac{Ze}{(2\pi)^3} \int \phi(q) \delta(q^0) e^{iq \cdot x} d^4 q, \quad (7)$$

$$\vec{q} = \vec{k} - \vec{p}_+ - \vec{p}_-.$$

Here p_+ , ϵ_+ , $p_- \epsilon_-$, k , ω refer to the four momentum and energies respectively of the positron, negaton, and photon, while $\bar{u}(p_-)$ and $V(p_+)$ are the spinors associated with the outgoing negaton and positron. The polarization four-vector of the photon is denoted by $\not{\underline{\ell}}$. A line under a vector means multiplication by γ_{μ} (for example, $\not{\underline{\ell}} = \gamma_{\mu} \ell^{\mu}$).

IV. EVALUATION OF THE CROSS SECTION

Substituting Eq. (6) in Eq. (1) one obtains the cross section for pair production,

$$d\sigma = \frac{\alpha Z^2}{(2\pi)^2} \frac{r_0^2 m^2}{\epsilon_+ \epsilon_- \omega} |\phi(q)|^2 X \delta(\epsilon_+ + \epsilon_- - \omega) d^3 p_+ d^3 p_- , \quad (8)$$

where

$$X = (2m)^2 \left| \bar{u}(p_-) Q(p_+ p_-) v(p_+) \right|^2 , \quad (9)$$

r_0 = classical electron radius,

α = fine-structure constant.

Making use of

$$\begin{aligned} \delta(\epsilon_+ + \epsilon_- - \omega) d^3 p_+ d^3 p_- &= p_+^2 \frac{dp_+}{d\epsilon_+} p_-^2 \frac{dp_-}{d\epsilon_-} \delta(\epsilon_+ + \epsilon_- - \omega) d\Omega_+ d\Omega_- d\epsilon_+ d\epsilon_- \\ &= p_+ dp_+ p_- dp_- \epsilon_+ \epsilon_- d\Omega_+ d\Omega_- d\epsilon_+ , \end{aligned} \quad (10)$$

one gets

$$d\sigma = \frac{\alpha Z^2}{(2\pi)^2} r_0^2 m^2 |\phi(\vec{q})|^2 \frac{|p_+| |p_-|}{\omega} d\epsilon_+ d\Omega_+ d\Omega_- X. \quad (11)$$

It is easily shown that

$$X = (2m)^2 \text{Trace} \left[\bar{Q} \sum(p_-) \Lambda(p_-) Q \sum(p_+) \Lambda(p_+) \right] , \quad (12)$$

where

$$\begin{aligned} \Lambda(p_+) &= \frac{\underline{1} p_+ + m}{2m} , & \sum(p_+) &= \frac{\delta_+ \vec{\sigma} \cdot \vec{p}_+ + |p_+|}{2 |p_+|} , \\ \Lambda(p_-) &= \frac{-\underline{1} p_- + m}{2m} , & \sum(p_-) &= \frac{-\delta_- \vec{\sigma} \cdot \vec{p}_- + |p_-|}{2 |p_-|} , \end{aligned}$$

$$\vec{\sigma} \cdot \vec{p} = \underline{1} p_5 \gamma_5 , \quad \gamma_5 = \gamma_1 \gamma_2 \gamma_3 \gamma_4 , \quad (13)$$

$\delta = 1$ spin in the direction of $\vec{p}$,

$\delta = -1$ spin in the direction of $-\vec{p}$.

Tedious evaluation of this trace leads to

$$\begin{aligned}
 -4X = & \left[\frac{A}{(p_-k)^2} + \frac{B}{(p_+k)^2} + \frac{C}{(p_+k)(p_-k)} \right] + \frac{\delta_+ \delta_-}{|\vec{p}_+| |\vec{p}_-|} \left[\frac{A_{+-}}{(p_-k)^2} + \frac{B_{+-}}{(p_+k)^2} + \right. \\
 & \left. + \frac{C_{+-}}{(p_+k)(p_-k)} \right] + \frac{\delta_+ \delta_0}{|\vec{p}_+|} \left[\frac{A_{+0}}{(p_-k)^2} + \frac{B_{+0}}{(p_+k)^2} + \frac{C_{+0}}{(p_+k)(p_-k)} \right] \\
 & + \frac{\delta_- \delta_0}{|\vec{p}_-|} \left[\frac{A_{-0}}{(p_-k)^2} + \frac{B_{-0}}{(p_+k)^2} + \frac{C_{-0}}{(p_+k)(p_-k)} \right], \quad (14)
 \end{aligned}$$

$$A \equiv 2(kp_+)(kp_-) + 4\omega\epsilon_+(kp_-) + 4a(kp_+) - 2d(kp_-) + 8\omega\epsilon_+a - 4(p_+p_-)a - 8\epsilon_+\epsilon_-a + 4m^2a$$

$$B \equiv 2(kp_+)(kp_-) + 4\omega\epsilon_-(kp_+) + 4b(kp_-) - 2d(kp_+) + 8\omega\epsilon_-b - 4(p_+p_-)b - 8\epsilon_+\epsilon_-b + 4m^2b$$

$$C \equiv 4\omega\epsilon_+(kp_-) + 4\omega\epsilon_-(kp_+) - 4\omega^2(p_+p_-) + 4a(kp_+) + 4b(kp_-) + 4m^2\omega^2$$

$$+ d \left[4\omega^2 - 4\omega\epsilon_- - 2(kp_-) - 2(kp_+) - 4\omega\epsilon_+ + 8\epsilon_+\epsilon_- + 4(p_+p_-) - 4m^2 \right],$$

$$\delta_0 A_{-0} = -2i \left[m^2 - \epsilon_+\epsilon_- \right] g_- + 2i(\vec{p}_-\vec{p}_-)h_- + 2i\omega\epsilon_-r_- - 2i(kp_+)(\vec{p}_-\vec{p}_-)(\vec{k}\ell^* \times \ell)$$

$$- 2i\omega\epsilon_-(kp_+)(\vec{p}_-\ell^* \times \ell) - 4i \left[(kp_+) + 2\omega\epsilon_+ \right] g_- - 4i\omega\epsilon_+(\vec{p}_-\vec{p}_-)(\vec{k}\ell^* \times \ell)$$

$$- 4i \left[\omega\epsilon_+(\vec{k}\vec{p}_-) - \omega\epsilon_-(\vec{k}\vec{p}_+) + (\vec{k}\vec{p}_+)(\vec{k}\vec{p}_-) \right] (\vec{p}_-\ell \times \ell^*),$$

$$\delta_0 B_{+0} = 2i \left[m^2 - \epsilon_+\epsilon_- \right] g_+ - 2i(\vec{p}_-\vec{p}_-)h_+ + 2i\omega\epsilon_+r_+ + 2i\omega\epsilon_-(kp_+)(\vec{p}_-\ell^* \times \ell)$$

$$- 2i(kp_+)(\vec{p}_-\vec{p}_-)(\vec{k}\ell \times \ell^*),$$

$$\delta_0 C_{-0} = 2i \left[m^2 - \epsilon_+\epsilon_- \right] g_+ - 2i \left[m^2 - \epsilon_+\epsilon_- \right] g_- - 2i(\vec{p}_-\vec{p}_-)h_+ + 2i(\vec{p}_-\vec{p}_-)h_- - 2i\omega\epsilon_-r_-$$

$$+ 2i\omega\epsilon_+r_+ - 4i \left[m^2(\vec{k}\vec{p}_-) - \omega\epsilon_+(\vec{p}_+\vec{p}_-) \right] (\vec{k}\ell^* \times \ell) - 4i\omega\epsilon_-(\vec{k}\vec{p}_-)(\vec{p}_-\ell \times \ell^*)$$

$$- 4\epsilon_+\epsilon_-(\vec{k}(\vec{k} \times \vec{p}_+) \times \vec{p}_-),$$

-7-

$$\delta_{00} A_{00} = 21 \left[m^2 - \epsilon_+ \epsilon_- \right] h_- - 21 \omega \epsilon_+ r_- - 21 (\bar{p}_+ \bar{p}_+) g_- + 21 \omega \epsilon_+ (k p_-) (\bar{p}_+ \ell^* \times \ell) \\ - 21 (k p_-) (\bar{p}_+ \bar{p}_+) (\bar{k} \ell \times \ell^*) ,$$

$$\delta_{00} B_{+0} = -21 \left[m^2 - \epsilon_+ \epsilon_- \right] h_+ + 21 (\bar{p}_+ \bar{p}_+) g_+ - 21 \omega \epsilon_+ r_+ - 21 (k p_-) (\bar{p}_+ \bar{p}_+) (\bar{k} \ell^* \times \ell) \\ - 21 \omega \epsilon_+ (k p_-) (\bar{p}_+ \ell \times \ell^*) - 41 \left[(k p_-) + 2 \omega \epsilon_- \right] h_+ - 41 \omega \epsilon_- (\bar{p}_+ \bar{p}_+) (\bar{k} \ell^* \times \ell) \\ - 41 \left[\omega \epsilon_- (\bar{k} \bar{p}_+) - \omega \epsilon_+ (\bar{k} \bar{p}_-) + (\bar{k} \bar{p}_-) (\bar{k} \bar{p}_+) (\bar{p}_+ \ell \times \ell^*) \right] ,$$

$$\delta_{00} C_{+0} = 21 \left[m^2 - \epsilon_+ \epsilon_- \right] h_- - 21 \left[m^2 - \epsilon_+ \epsilon_- \right] h_+ - 21 (\bar{p}_+ \bar{p}_+) g_- + 21 (\bar{p}_+ \bar{p}_+) g_+ + 21 \omega \epsilon_+ h_+ \\ - 21 \omega \epsilon_+ r_+ - 21 \omega \epsilon_+ r_- + 41 \left[m^2 (\bar{k} \bar{p}_+) - \omega \epsilon_+ (\bar{p}_+ \bar{p}_-) \right] (\bar{k} \ell^* \times \ell) \\ - 41 \omega \epsilon_+ (\bar{k} \bar{p}_+) (\bar{p}_- \ell \times \ell^*) - 4 \epsilon_+ \delta_0 (\bar{k} (\bar{k} \times \bar{p}_-) \times \bar{p}_+) ,$$

$$A_{+-} = 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) (\bar{k} \bar{p}_-) - 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_-) d + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) a \\ - 2 \omega \epsilon_+ (\bar{p}_- \bar{p}_-) d - 4 \omega \epsilon_- (\bar{p}_+ \bar{p}_+) a + 2 \omega \epsilon_+ (\bar{k} \bar{p}_+) (\bar{p}_- \bar{p}_-) - 2 \omega \epsilon_- (\bar{k} \bar{p}_-) (\bar{p}_+ \bar{p}_+) \\ + 2 \left[\omega^2 + 2a \right] (\bar{p}_+ \bar{p}_+) (\bar{p}_- \bar{p}_-) - 2 \left[2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} a + m^2 \omega^2 \right] (\bar{p}_+ \bar{p}_-) + 2 m^2 \omega^2 d ,$$

$$B_{+-} = 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) (\bar{k} \bar{p}_-) - 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) d + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_-) b \\ - 2 \omega \epsilon_- (\bar{p}_+ \bar{p}_+) d - 4 \omega \epsilon_+ (\bar{p}_- \bar{p}_-) b + 2 \omega \epsilon_- (\bar{k} \bar{p}_-) (\bar{p}_+ \bar{p}_+) - 2 \omega \epsilon_+ (\bar{k} \bar{p}_+) (\bar{p}_- \bar{p}_-) \\ + 2 \left[\omega^2 + 2a \right] (\bar{p}_- \bar{p}_-) (\bar{p}_+ \bar{p}_+) - 2 \left[2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} b + m^2 \omega^2 \right] (\bar{p}_+ \bar{p}_-) + 2 m^2 \omega^2 d ,$$

$$C_{+-} = 4 \omega^2 \epsilon_+ \epsilon_- d - 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) d - 2 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_-) d + 2 \omega \epsilon_- (\bar{p}_+ \bar{p}_+) d \\ + 2 \omega \epsilon_+ (\bar{p}_- \bar{p}_-) d + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{p}_+ \bar{p}_-) d - 4 (\bar{p}_+ \bar{p}_+) (\bar{p}_- \bar{p}_-) d + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) a \\ + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_-) b + 4 \omega \epsilon_- (\bar{p}_+ \bar{p}_+) a + 4 \omega \epsilon_+ (\bar{p}_- \bar{p}_-) b + 4 \left[m^2 - \epsilon_+ \epsilon_- \right] (\bar{k} \bar{p}_+) (\bar{k} \bar{p}_-) \\ - 4 \omega^2 \epsilon_+ \epsilon_- (\bar{p}_+ \bar{p}_-) ,$$

-8-

where $a = (\ell^+ p_-)(\ell p_-)$, $b = (\ell^+ p_+)(\ell p_+)$,
 $d = (\ell p_-)(\ell^+ p_+) + (\ell^+ p_-)(\ell p_+)$,
 $f = (\ell p_-)(\ell^+ p_+) - (\ell^+ p_-)(\ell p_+)$,
 $g_{\pm} = (\ell p_{\pm})(\ell^+ \vec{k} \times \vec{p}_{\pm}) - (\ell^+ p_{\pm})(\ell \vec{k} \times \vec{p}_{\pm})$,
 $h_{\pm} = (\ell p_{\pm})(\ell^+ \vec{k} \times \vec{p}_{\mp}) - (\ell^+ p_{\pm})(\ell \vec{k} \times \vec{p}_{\mp})$,
 $r_{\pm} = (\ell p_{\pm})(\ell^+ \vec{p}_{\mp} \times \vec{p}_{\pm}) - (\ell^+ p_{\pm})(\ell \vec{p}_{\mp} \times \vec{p}_{\pm})$,

δ_- refers to the spin direction of the negaton with respect to $\vec{p}_-$,

δ_+ refers to the spin direction of the positon with respect to $\vec{p}_+$,

δ_0 refers to the spin direction of the photon with respect to $\vec{k}$,

and use has been made of the relations

$$(kQ) = 0, \quad \ell_{\mu} = (\ell_1, \ell_2, \ell_3, 0). \quad (15)$$

V. EXPLICIT EVALUATION OF $d\sigma$ IN TERMS OF THE ANGLES INVOLVED IN THE PROBLEM

If one makes use of

$$\ell_2 = -i\delta_0 \ell, \quad k_{\mu} = (0, 0, k_3, \omega) \quad \ell^+ \ell = 1, \quad (16)$$

one gets relations such as

$$(\vec{k} \ell^+ \times \ell) = -i\delta_0 \frac{(\vec{k} \vec{A})}{\omega},$$

$$(\ell p_-)(\ell^+ p_+) + (\ell^+ p_-)(\ell p_+) = (\vec{p}_+ \vec{p}_-) - \frac{(\vec{k} \vec{p}_+)(\vec{k} \vec{p}_-)}{\omega^2}, \quad (17)$$

$$(\ell p_-)(\ell^+ p_+) - (\ell^+ p_-)(\ell p_+) = \frac{i\delta_0}{\omega} (\vec{k} \vec{p}_- \times \vec{p}_+)$$

etc.

Introducing the angles θ_+ , θ_- , and ϕ , where θ_+ and θ_- are the angles made by the outgoing positon and negaton respectively with the incoming photon, and ϕ is the angle between the $k - p_+$ and $k - p_-$ planes, one easily gets

-9-

$$\begin{aligned}
 (\ell p_-)(\ell^* p_+) + (\ell^* p_-)(\ell p_+) &= \epsilon_+ \epsilon_- \beta_+ \beta_- \sin \theta_+ \sin \theta_- \cos \phi, \\
 (\ell p_-)(\ell^* p_+) - (\ell^* p_-)(\ell p_+) &= i \epsilon_+ \epsilon_- \beta_+ \beta_- \sin \theta_+ \sin \theta_- \sin \phi, \\
 &\text{etc.},
 \end{aligned}
 \tag{18}$$

where β = velocity.

Now it is very simple to express the differential cross section in terms of these angles. As an example one easily finds

$$\begin{aligned}
 A_{+-} &= 2\omega \epsilon_+^2 \epsilon_-^2 \beta_+^2 \beta_-^2 + 2\omega \epsilon_+^2 \epsilon_-^2 \beta_+^2 \beta_-^2 \cos \theta_+ - 2\omega \epsilon_+^2 \epsilon_-^2 \beta_+^2 \beta_-^2 \cos \theta_- \\
 &+ \left\{ 2\epsilon_-^4 \epsilon_+^2 \beta_+^2 \beta_-^4 - 2\omega \epsilon_+^2 \epsilon_-^3 \beta_+^2 \beta_-^2 \right\} \sin^2 \theta_- - \omega \epsilon_+^2 \epsilon_-^2 \beta_+ \beta_- \cos \theta_+ \cos \theta_- \\
 &+ 2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} \omega \epsilon_-^2 \epsilon_+ \beta_+ \beta_-^2 \sin^2 \theta_- \cos \theta_+ \\
 &- 2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} \epsilon_-^3 \epsilon_+ \beta_-^3 \beta_+ \sin^2 \theta_- \cos \theta_- \cos \theta_+ \\
 &- 2\omega \epsilon_-^2 \epsilon_+^3 \beta_+ \beta_-^3 \sin \theta_+ \sin \theta_- \cos \phi - 2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} \epsilon_-^3 \epsilon_+ \beta_+ \beta_-^3 \sin \theta_+ \sin^3 \theta_- \cos \phi \\
 &- 2 \left\{ m^2 - \epsilon_+ \epsilon_- \right\} \omega \epsilon_+ \epsilon_-^2 \beta_+ \beta_-^2 \sin \theta_+ \sin \theta_- \cos \theta_- \cos \phi.
 \end{aligned}$$

The other coefficients can also easily be expressed in terms of θ_+ , θ_- , and ϕ .