UC Davis UC Davis Previously Published Works

Title Single-cell analysis: Advances and future perspectives.

Permalink https://escholarship.org/uc/item/16c7v6p5

Journal Bosnian journal of basic medical sciences, 16(4)

ISSN 1512-8601

Author Hodzic, Emir

Publication Date 2016-11-01

DOI 10.17305/bjbms.2016.1371

Peer reviewed

Single-cell analysis: Advances and future perspectives

Emir Hodzic*

Real-time PCR Research and Diagnostic Core Facility, School of Veterinary Medicine, University of California at Davis, California, United States of America

ABSTRACT

The last several years have seen rapid development of technologies and methods that permit a detailed analysis of the genome and transcriptome of a single cell. Recent evidence from studies of single cells reveals that each cell type has a distinct lineage and function. The lineage and stage of development of each cell determine how they respond to each other and the environment. Experimental approaches that utilize single-cell analysis are effective means to understand how cell networks work in concert to coordinate a response at the population level; recent progress in single-cell analysis is offering a glimpse at the future.

KEY WORDS: Single cell; analysis; DNA; RNA; genome DOI: http://dx.doi.org/10.17305/bjbms.2016.1371

Bosn J Basic Med Sci. 2016;16(4):313-314.

Introduction: In living tissues, there are a large number of cell types with the assumption that each cell type has a distinct lineage and function. However, recent evidence from studies of single cells reveals that this assumption is incorrect. Cells may be morphologically and genetically identical but are actually heterogeneous, made up of individual cells that differ dramatically. These differences can have important consequences for the health and function of the entire population. Single-cell analysis allows the study of cell-to-cell variation within a cell population (organ, tissue, and cell culture). In order to study diseases and drug development, in-depth analysis of stem cell differentiation, cancer, physiological functions in embryos and adults can only be accomplished with single-cell analysis [1-4]. Sample Preparation: To isolate a single cell from a heterogeneous population of cells in a manner that preserves biological integrity, there are several approaches available [5]. Besides several well-established methods, such as micromanipulation, laser-capture microdissection and fluorescence-activated cell sorting, there is an increasing number of novel techniques to isolate single cells with even greater accuracy and specificity [6,7].

Submitted: 05 May 2016/Accepted: 06 May 2016

© 2016 Emir Hodzic; licensee ABMSFBIH. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

In recent years, innovative technologies for single-cell genomics, transcriptomics, and proteomics, with a particular emphasis on quantification and multiplicity that can perform measurements on many single cells in a given experimental run, include microfluidics, optical tweezers, transcriptome *in vivo* analysis, and mass cytometry [4,8-10].

Applications: Advances in whole-genome and whole-transcriptome amplification have permitted the gene expression profiling and sequencing of the minute amounts of DNA and RNA present in a single cell, offering a window into the extent and nature of genomic and transcriptomic heterogeneity which occurs in both normal development and disease. Each cell in the body has a unique genomic structure, which allows the reconstruction of cell lineage trees with very high precision that can predict the existence of small population of steam cells. This information is important for cancer research for detection of rare tumor cells, preimplantation, and genetic diagnosis [11-13]. Single-cell approaches have been utilized for understanding the intricate cellular interplay involved in immune response that requires single-cell resolution, especially with rare antigen-specific T- or B-cells [14,15]. In addition, protein expression analysis is vital to understand the true metabolic or functional state of cells and the single-cell approach enables simultaneous analysis of more than 35 proteins of individual cells [16,17]. Currently, researchers are starting to combine single-cell genomics with single-cell proteomics to tackle important questions in fields including cancer, stem cell biology, neuroscience, developmental biology, and infectious disease. As more investigators explore heterogeneity in cell populations, knowledge of intricate biological

^{*}Corresponding author: Emir Hodzic, Real-time PCR Research and Diagnostic Core Facility, School of Veterinary Medicine, University of California at Davis, One Shields Avenue, Davis, CA 95616, USA. E-mail: ehodzic@ucdavis.edu

cellular networks will empower researchers to discover new ways to diagnose and treat disease.

Challenges: The advances of single-cell analysis over the past 5 years have happened at a lightning pace, and the potential for their use in various fields is high. However, the novelty of these single-cell techniques also implies various limitations. There is a lack of cross-disciplinary collaboration to more effectively utilize advantage of single-cell analysis. Another caveat is that algorithms for the analysis of single-cell data are even less mature than the experimental platforms, and effective interpretations of what are increasingly large datasets remain challenging, with techniques that vary across research groups [4,11,17].

DECLARATION OF INTERESTS

The author declares no conflict of interests.

REFERENCES

- Anselmetti D. Single cell analysis: Technologies and applications. Hanover, Germany: Wiley-VCH Verlag GmbH and Co. KGaA; 2009. p. 284.
- [2] Tang F, Lao K, Surani MA. Development and applications of single-cell transcriptome analysis. Nat Methods. 2011;8(4 Suppl):S6-11. http://dx.doi.org/10.1038/nmeth.1557.
- [3] Rubakhin SS, Romanova EV, Nemes P, Sweedler JV. Profiling metabolites and peptides in single cells. Nat Methods. 2011;8(4 Suppl):S20-9.

http://dx.doi.org/10.1038/nmeth.1549.

- [4] Heath JR, Ribas A, Mischel PS. Single-cell analysis tools for drug discovery and development. Nat Rev Drug Discov. 2016;15(3):204-16. http://dx.doi.org/10.1038/nrd.2015.16.
- [5] Navin N, Hicks J. Future medical applications of single-cell sequencing in cancer. Genome Med. 2011;3(5):31. http://dx.doi.org/10.1186/gm247.
- [6] Yilmaz S, Singh AK. Single cell genome sequencing. Curr Opin Biotechnol. 2012;23(3):437-43.

http://dx.doi.org/10.1016/j.copbio.2011.11.018.

[7] Saliba AE, Westermann AJ, Gorski SA, Vogel J. Single-cell RNA-seq: Advances and future challenges. Nucleic Acids Res. 2014;42(14):8845-60.

http://dx.doi.org/10.1093/nar/gku555.

[8] Yoshimoto N, Kida A, Jie X, Kurokawa M, Iijima M, Niimi T, et al. An automated system for high-throughput single cell-based breeding. Sci Rep. 2013;3:1191. http://dx.doi.org/10.1038/srep01191.

 [9] Lovatt D, Ruble BK, Lee J, Dueck H, Kim TK, Fisher S, et al. Transcriptome *in vivo* analysis (TIVA) of spatially defined single cells in live tissue. Nat Methods. 2014;11(2):190-6.

http://dx.doi.org/10.1038/nmeth.2804.
[10] Bendall SC, Simonds EF, Qiu P, Amir el-AD, Krutzik PO, Finck R, et al. Single-cell mass cytometry of differential immune and drug responses across a human hematopoietic continuum. Science. 2011;332(6030):687-96.

http://dx.doi.org/10.1126/science.1198704.

- [11] Macaulay IC, Voet T. Single cell genomics: Advances and future perspectives. PLoS Genet. 2014;10(1):1004126. http://dx.doi.org/10.1371/journal.pgen.1004126.
- [12] Navin N, Kendall J, Troge J, Andrews P, Rodgers L, McIndoo J, et al. Tumour evolution inferred by single-cell sequencing. Nature. 2011;472(7341):90-4. http://dx.doi.org/10.1038/nature09807.
- [13] Shapiro E, Biezuner T, Linnarsson S. Single-cell sequencing-based technologies will revolutionize whole-organism science. Nat Rev Genet. 2013;14(9):618-30.

http://dx.doi.org/10.1038/nrg3542.

[14] Shalek AK, Satija R, Adiconis X, Gertner RS, Gaublomme JT, Raychowdhury R, et al. Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. Nature. 2013;498(7453):236-40.

http://dx.doi.org/10.1038/nature12172.

- [15] Georgiou G, Ippolito GC, Beausang J, Busse CE, Wardemann H, Quake SR. The promise and challenge of high-throughput sequencing of the antibody repertoire. Nat Biotechnol. 2014;32(2):158-68. http://dx.doi.org/10.1038/nbt.2782.
- [16] Ståhlberg A, Thomsen C, Ruff D, Åman P. Quantitative PCR analysis of DNA, RNAs, and proteins in the same single cell. Clin Chem. 2012;58(12):1682-91. http://dx.doi.org/10.1373/clinchem.2012.191445.
- [17] Albayrak C, Jordi CA, Zechner C, Lin J, Bichsel CA, Khammash M, et al. Digital quantification of proteins and mRNA in single mammalian cells. Mol Cell. 2016;61(6):914-24. http://dx.doi.org/10.1016/j.molcel.2016.02.030.