

UC Berkeley

UC Berkeley Previously Published Works

Title

Diverse Bacterial Genes Modulate Plant Root Association by Beneficial Bacteria

Permalink

<https://escholarship.org/uc/item/16g8p0vw>

Journal

mBio, 11(6)

ISSN

2161-2129

Authors

do Amaral, Fernanda Plucani
Tuleski, Thalita Regina
Pankiewicz, Vania Carla Silva
et al.

Publication Date

2020-12-22

DOI

10.1128/mbio.03078-20

Peer reviewed



Diverse Bacterial Genes Modulate Plant Root Association by Beneficial Bacteria

Fernanda Plucani do Amaral,^a Thalita Regina Tuleski,^b Vania Carla Silva Pankievicz,^{b*} Ryan A. Melnyk,^{c*} Adam P. Arkin,^c Joel Griffitts,^d Michelle Zibetti Tadra-Sfeir,^b Emanuel Maltempi de Souza,^b Adam Deutschbauer,^c Rose Adele Monteiro,^b Gary Stacey^a

^aDivisions of Plant Science and Biochemistry, C. S. Bond Life Science Center, University of Missouri, Columbia, Missouri, USA

^bDepartment of Biochemistry and Molecular Biology, Universidade Federal do Parana, Curitiba, Parana, Brazil

^cEnvironmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA

^dDepartment of Microbiology and Molecular Biology, Brigham Young University, Provo, Utah, USA

Fernanda Plucani do Amaral, Thalita Regina Tuleski, and Vania Carla Silva Pankievicz contributed equally to this article. The author order was determined by contribution to writing the manuscript.

ABSTRACT The plant rhizosphere harbors a diverse population of microorganisms, including beneficial plant growth-promoting bacteria (PGPB), that colonize plant roots and enhance growth and productivity. In order to specifically define bacterial traits that contribute to this beneficial interaction, we used high-throughput transposon mutagenesis sequencing (TnSeq) in two model root-bacterium systems associated with *Setaria viridis*: *Azoarcus olearius* DQS4^T and *Herbaspirillum seropedicae* SmR1. This approach identified ~100 significant genes for each bacterium that appeared to confer a competitive advantage for root colonization. Most of the genes identified specifically in *A. olearius* encoded metabolism functions, whereas genes identified in *H. seropedicae* were motility related, suggesting that each strain requires unique functions for competitive root colonization. Genes were experimentally validated by site-directed mutagenesis, followed by inoculation of the mutated bacteria onto *S. viridis* roots individually, as well as in competition with the wild-type strain. The results identify key bacterial functions involved in iron uptake, polyhydroxybutyrate metabolism, and regulation of aromatic metabolism as important for root colonization. The hope is that by improving our understanding of the molecular mechanisms used by PGPB to colonize plants, we can increase the adoption of these bacteria in agriculture to improve the sustainability of modern cropping systems.

IMPORTANCE There is growing interest in the use of associative, plant growth-promoting bacteria (PGPB) as biofertilizers to serve as a sustainable alternative for agriculture application. While a variety of mechanisms have been proposed to explain bacterial plant growth promotion, the molecular details of this process remain unclear. The current research supports the idea that PGPB use in agriculture will be promoted by gaining more knowledge as to how these bacteria colonize plants, promote growth, and do so consistently. Specifically, the research seeks to identify those bacterial genes involved in the ability of two, PGPB strains, *Azoarcus olearius* and *Herbaspirillum seropedicae*, to colonize the roots of the C4 model grass *Setaria viridis*. Applying a transposon mutagenesis (TnSeq) approach, we assigned phenotypes and function to genes that affect bacterial competitiveness during root colonization. The results suggest that each bacterial strain requires unique functions for root colonization but also suggests that a few, critical functions are needed by both bacteria, pointing to some common mechanisms. The hope is that such information can be exploited to improve the use and performance of PGPB in agriculture.

KEYWORDS bacterium-root colonization, beneficial bacteria, gene functionality, transposon mutagenesis

Citation do Amaral FP, Tuleski TR, Pankievicz VCS, Melnyk RA, Arkin AP, Griffitts J, Tadra-Sfeir MZ, Maltempi de Souza E, Deutschbauer A, Monteiro RA, Stacey G. 2020. Diverse bacterial genes modulate plant root association by beneficial bacteria. *mBio* 11:e03078-20. <https://doi.org/10.1128/mBio.03078-20>.

Editor B. Gillian Turgeon, Cornell University

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to Gary Stacey, staceyg@missouri.edu.

* Present address: Vania Carla Silva Pankievicz, Department of Bacteriology, University of Wisconsin, Madison, Wisconsin, USA; Ryan A. Melnyk, Pivot Bio, Berkeley, California, USA.

This article is a direct contribution from Gary Stacey, a Fellow of the American Academy of Microbiology, who arranged for and secured reviews by Euan James, University of Dundee; Philip Poole, Oxford University; and Michael Sadowsky, University of Minnesota.

Received 30 October 2020

Accepted 9 November 2020

Published 15 December 2020

Plant health and development are influenced by a broad range of microorganisms inhabiting the root rhizosphere, including beneficial associative bacteria. These bacteria are usually minor components of the rhizosphere microbial community but have been shown to significantly enhance plant growth and yield (1, 2). This ability to enhance plant growth has been attributed to diverse mechanisms, including biological nitrogen fixation, production of phytohormones, enhancement of nutrient uptake (siderophore and phosphate solubilization), and biocontrol of pathogens and pests (3). Plant growth-promoting bacteria (PGPB) can colonize roots either on the root surface or as endophytes without eliciting a noticeable plant defense response (4, 5). Unlike some other bacterium-plant interactions that show strict host specificity, many PGPB can colonize a wide variety of plant species, including agriculturally important members of the *Poaceae* family, such as rice, maize, wheat, and a variety of bioenergy grasses (6, 7). PGPB strains have been described for several host plants, including *Setaria viridis*, a model C_4 plant, which is a close relative to a variety of bioenergy grasses. In a previous study, we demonstrated that *S. viridis* under lab conditions can obtain up to 100% of its nitrogen needs through biological nitrogen fixation mediated by diazotrophic PGPB (8). Among the strains used were the betaproteobacteria *Azoarcus olearius* and *Herbaspirillum seropedicae*, which exhibit strong growth-promoting ability (4, 9–12). *Azoarcus olearius* DQS4^T was originally isolated from oil-contaminated soil (13) and, based on its genome sequence, this strain shows high similarity with the well-studied *A. olearius* strain BH72 (14). Strain DQS4^T can colonize the roots of rice and *Setaria* to high levels and increase below- and above-ground biomass (9). *H. seropedicae* SmR1 is a well-studied endophytic bacterium that colonizes several plants, including maize, wheat, and *Setaria*. Recently, *in situ* metabolomic profiling of *S. viridis* roots colonized by SmR1 demonstrated that inoculation induced a wide variety of plant metabolic changes, including those affecting nitrogen and phytohormone levels (15). Thus, similar to many other, better-studied plant-microbe associations, PGPB inoculation appears to profoundly affect the metabolism of its host.

To identify specific PGPB genes that contribute to plant root colonization, we applied transposon mutagenesis sequencing (TnSeq) coupled with random barcoding (RB-TnSeq). This technique assesses gene functionality in mutant strains through the generation of reusable libraries of unique and mapped mutant insertions (16, 17). TnSeq is a powerful and sensitive approach to identify bacterial gene functions that play roles in bacterial fitness under researcher-defined growth conditions (16–18). Starting with information on the location and frequency of each mutation in the population, the change in mutational frequency after passage of the population through an environmental challenge (e.g., plant root colonization) leads to the identification of genes that are either essential, important, or detrimental to growth under that condition. For example, TnSeq of *Streptococcus pneumoniae* identified genes essential for bacterial basal growth, as well as genes involved in transcriptional regulation and carbohydrate transport (17). In the PGPB *Pseudomonas simiae*, TnSeq revealed genes involved in carbon metabolism and motility that could enhance or suppress colonization of *Arabidopsis* roots (19). In the present work, we applied standard TnSeq and a variant of TnSeq that uses random DNA barcodes to measure strain abundance (RB-TnSeq) to construct mutant libraries of strains DQS4^T and SmR1, respectively, and to then evaluate genetic contributions to fitness on the host plant *Setaria viridis*.

RESULTS

Overview of TnSeq analysis. Transposon sequencing is a high-throughput tool used to generate a large bacterial mutant population. Gene essentiality is defined as its importance in maintaining competitive fitness under a condition of interest (Fig. 1A).

For this study, two PGPB bacterial species were used, *Azoarcus olearius* DQS4^T and *Herbaspirillum seropedicae* SmR1, both diazotrophic and capable of promoting plant growth (8, 9, 11–13, 20). The mutagenized pool of each strain was inoculated onto *Setaria viridis* roots or a carbon augmented soil control with no plants. Root-associated




FIG 1 TnSeq representation and determination. (A) Insertion mutant library generated by a mariner transposon system from wild-type bacteria. A mutant pool of each bacterium (DQS4^T or SmR1) was inoculated onto *Setaria viridis*. Surviving mutant strains were recovered from the roots or soil without plants, at 10 days after inoculation, and the abundance of each insertion was quantified by TnSeq. (B) Categorization based on gene essentiality. Genes with low insertion counts within the root population generating negative fitness scores (<-1) were categorized as decreased root colonization (shown as square in both graphs). Genes with overrepresented insertion counts were given positive fitness scores (>1) and categorized as enhanced root colonization (shown as dots in both graphs). Plots show fitness scores obtained from the average values of root samples across biological replicates ($n=4$).

bacterial cells were recovered, transposon DNA was sequenced, and resulting reads were mapped to each bacterial genome. The relative incidence of a specific mutation (or lack thereof) was then determined and used to calculate fitness values. By high-throughput sequence analysis of insertion mutants, we covered about 3,692 genes of DQS4^T and 3,878 genes of SmR1 distributed throughout either strain's genome. Thus, regardless of method, very good coverage of each genome was achieved. In the case of DQS4^T transposon mutagenesis, 89.2% of 4,135 total genes were mapped to a gene harboring at least 1 insertion event. For the SmR1 library, 81.1% of the 4781 genes had identified mutations. We assume that genes without an insertion in the library likely identify those essential for bacterial growth in culture.

Our screening identified 89 and 130 gene mutations that significantly affected the ability of strains DQS4^T and SmR1 to colonize *S. viridis* roots, respectively. Given that our interest was to identify genes that exclusively affected root colonization, we excluded from further consideration any genes that also affected fitness under soil conditions (see Fig. S1a). The genes affecting root colonization were categorized based on the phenotypes as either (i) enhanced fitness value, mutations that increased root association (fitness score ≥ 1); or (ii) decreased fitness value, mutations that impaired root association (fitness score ≤ -1) (Fig. 1B; see also Fig. S1b in the supplemental material). For the complete list of gene mutations affecting fitness under each condition, see Table S2.

Unsurprisingly, many of the genes identified lacked a clear functional annotation, especially for *A. olearius* DQS4^T. Even though each strain showed a distinct pattern of mutations that affected fitness, a few common gene functions (e.g., those involved in chemotaxis and cell wall recycling) were identified impairing root colonization in both strains. The most common COG categories important for SmR1 root colonization were those involved in amino acid transport and metabolism, followed by energy production and conversion and coenzyme transport and metabolism. In the case of DQS4^T




FIG 2 Cluster of orthologous group (COG) categories of colonization genes. The distributions of genes significantly increasing or decreasing root colonization when mutated are shown. The color legend of dominant COG categories is shown separately for *Azoarcus olearius* DQS4⁺ and *Herbaspirillum seropedicae* SmR1.

the most common COGs were cell wall/membrane/envelope biogenesis and signal transduction, followed by translation, ribosomal structure and biogenesis, and inorganic ion transport and metabolism (Fig. 2). Although we cannot rule out that the different mutagenesis methods could contribute to the different fitness profiles, the overall mutational coverage of the genomes of both strains suggest that such effects are likely minor. Hence, the results seem to suggest that any given PGPB strain will have different major requirements for root colonization reflecting the specific metabolic needs of the bacterium.

Transposon mutations that benefit bacterial root association. Gene mutations that positively impacted the ability of bacterial strains to colonize roots were defined as those with a fitness score ≥ 1 . We identified 8 and 14 genes in this category in *A. olearius* and *H. seropedicae*, respectively (Table 1). This list included the *Azoarcus* gene predicted to encode a pyrroloquinoline quinone (PQQ)-dependent alcohol dehydrogenase (ADH; DQS_RS19730), which is involved in the oxidoreductase process of ADH. Based on sequence alignment with *Azoarcus olearius* BH72, the DQS_RS19730 gene encodes an ExaA5 protein that was previously reported to be induced upon exposure to ethanol as a carbon source (21). In our study, we found that *exaA5* plays an important role in root colonization. Disruption of this gene enhanced root colonization of *S. viridis* roots (Fig. 3A). However, the mutant was less competitive in root colonization than the wild type at 1 and 3 days after inoculation but seemed to recover to wild-type levels by 5 days after inoculation (d.a.i.) (Fig. 3B). An insertion mutation in the gene encoding a diguanylate cyclase (DQS_RS13665) significantly increased bacterial root colonization 3 and 5 days postinoculation (Fig. 3A) and showed the same trend in competition with the wild type (Fig. 3B). Diguanylate cyclases (DGCs) with a GGDEF active-site motif produce cyclic di-GMP (c-di-GMP) and play a major role in the transition between motile and sessile bacterial lifestyles (22–27). We also identified an iron regulator outer membrane protein, TonB (DQS_RS15430), presumably required for Fe³⁺

TABLE 1 Mutations that enhanced DQS4^T or SmR1 root fitness colonization of *S. viridis*

Strain ^a	Locus ID	Gene annotation	Fitness score	
DQS4 ^T	DQS_RS19730	PQQ-binding-like beta-propeller repeat protein ExaA5	1.36	
	DQS_RS15440	Peptidase	2.45	
	DQS_RS15145	Sigma-54-dependent Fis family transcriptional regulator	1.89	
	DQS_RS13665	Histidine kinase sensor domain-containing diguanylate cyclase	1.92	
	DQS_RS15430	TonB-dependent receptor	1.72	
	DQS_RS08695	Ribonuclease III	3.09	
	DQS_RS16645	DUF502 domain-containing protein	1.82	
	DQS_RS15445	TonB-dependent siderophore receptor	3.47	
	SmR1	HSERO_RS14975	Flagellar motor protein MotA	1.05
		HSERO_RS10140	Flagellar biosynthesis protein FlhB	1.06
HSERO_RS10305		Flagellar motor switch protein FliG	1.06	
HSERO_RS06285		Methyl-accepting chemotaxis protein	1.07	
HSERO_RS02815		HxIR family transcriptional regulator	1.20	
HSERO_RS10150		Flagellar biosynthesis regulator FlhF	1.24	
HSERO_RS10255		Flagellar biosynthesis protein FliQ	1.31	
HSERO_RS10310		Flagellar M-ring protein FliF	1.33	
HSERO_RS23885		XRE family transcriptional regulator	1.41	
HSERO_RS13885		Histidine kinase	1.73	
HSERO_RS14985		Transcriptional regulator	1.83	
HSERO_RS13890		LuxR family transcriptional regulator	1.85	
HSERO_RS20835		ABC transporter substrate-binding protein	1.93	
HSERO_RS08080		Poly[D(-)-3-hydroxyalkanoate] depolymerase	2.80	

^aDQS4^T, *Azoarcus olearius*; SmR1, *Herbaspirillum seropedicae*.

uptake (Fig. 3C). This protein was reported to interact with ExbA and ExbB, forming a complex that regulates iron acquisition (28).

In SmR1, 6 of the 14 gene mutations that increased root colonization are predicted to play a role in flagellar biosynthesis. FliF, FliG, FlhB, and FliQ are structural components of the flagellar basal body (see Fig. S2 in the supplemental material). MotA is complexed with MotB and functions as a proton channel for torque generation (29). Our testing showed that the absence of flagellar genes in the SmR1 strain conveyed an advantage for root colonization in *S. viridis* 10 days after inoculation. We also identified a poly-3-hydroxybutyrate (PHB)-related gene, a poly-3-hydroxyalkanoate depolymerase PhaZ1 (HSERO_RS08080), that enhanced colonization ability when mutated (Fig. 4). PhaZ1 is a PHB depolymerization enzyme required for the granule mobilization (30). Furthermore, mutations in the genes encoding transcriptional regulators and ABC transporters also resulted in increased root colonization by strain SmR1 (Table 1).

Insertions that impaired root association. Mutant strains corresponding to 81 DQS4^T and 114 SmR1 genes were identified as significantly reducing root colonization (i.e., fitness score ≤ -1) (Table 2; see also Table S2). Therefore, these gene functions are normally required for efficient colonization by wild-type bacteria.

Within this group of mutations, we identified two genes present in both DQS4^T and SmR1: *cheY* (DQS_RS02075 and HSERO_RS09745), which is related to chemotaxis, and *ampD* (DQS_RS17190 and HSERO_RS01880), which is involved in peptidoglycan degradation. Chemotaxis is directly involved in modulating the movement of the flagellum in response to attractants, helping bacterial orientation and active motion for survival toward favorable conditions (31–33). In addition, four other genes encoding transmembrane chemoreceptors, also known as methyl-accepting chemotaxis proteins (MCPs), were identified in *H. seropedicae*. The protein CheA (HSERO_RS15535) undergoes autophosphorylation induced by MCPs, and CheR (HSERO_RS10115) and CheW (HSERO_RS14950) acts to modulate the phosphorylation state of CheY (34) (see Fig. S3). Mutations in these genes impaired the capacity of *H. seropedicae* to colonize roots, consistent with a previous report (35).




FIG 3 Root associated bacterial cells recovered from *Setaria viridis* roots at 1, 3, and 5 d.a.i. (A) Root colonization after single inoculation with wild-type DQS4^T or with the mutant lines diguanylate cyclase-DGC (DQS_RS13665) or *ExaA5* (DQS_RS19730). The data are expressed as the log₁₀ CFU per g of root fresh weight (RFW). The graphs show an increase of root colonization by the mutant strains. (B) Coinoculation of *S. viridis* roots with a mixture of equal amounts of wild-type and mutant strains. (C) Map of Iron uptake cluster identified in DQS4^T by TnSeq as enhancing bacterial root attachment. (D) Root colonization of TonB mutant (DQS_RS15445) in CFU recovered from *S. viridis* roots after individual inoculation or after coinoculation with the wild type. Bars show mean averages ± the standard errors (SE) (*n*=20). Statistical significance, determined using a Student *t* test, is indicated by asterisks (*, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001).

Analysis of the DQS4^T TnSeq data identified a gene (DQS_RS09125) predicted to encode a two-component sensor histidine kinase containing a HAMP domain (histidine kinase, adenyl cyclases, methyl-binding proteins, and phosphatases) involved in signal conversion between the transmembrane-sensing and kinase activity control (36). One assumes that this histidine kinase is reacting to an unknown factor in the rhizosphere that normally enhances root association in wild-type cells. We also observed genes involved in cell wall formation, such as *murl* (HSERO_RS07755), a glutamate racemase responsible for the conversion of L-glutamate in D-glutamate, and *murA* (HSERO_RS20355), a UDP-*N*-acetylglucosamine 1-carboxyvinyltransferase responsible for converting UDP-*N*-acetylglucosamine to




FIG 4 *Setaria viridis* root colonization by mutants defective in PHB-related genes. Mutant strains (Δ *phaP1* involved in production or Δ *phaZ1* mobilization of PHB) were recovered from root samples. The assays were performed as colonizations 30 min, 1 day, and 5 days after inoculation. The strains were inoculated individually or coinoculated in the proportion of 1:1 with wild-type SmR1. The data are expressed as the log CFU per g of root fresh tissue. Bars indicate averages ± the SE. Statistical significance, determined using a Student *t* test, is indicated by asterisks (*, *P* ≤ 0.01).

TABLE 2 Partial list of mutations that impaired DQS4^T or SmR1 root fitness colonization of *S. viridis*^a

Strain	Locus ID	Fitness score	Description
DQS4 ^T	DQS_RS00855	−2.70	Hypothetical protein
	DQS_RS02275	−2.33	Diguanylate cyclase
	DQS_RS01300	−2.69	RND efflux transporter, permease protein
	DQS_RS18440	−2.89	Methyltransferase domain-containing protein
	DQS_RS20485	−1.97	Putative two-component system sensor protein
	DQS_RS07570	−1.85	Putative cobalt-zinc-cadmium resistance protein
SmR1	HSERO_RS01265	−2.93	Acetyl-CoA acetyltransferase
	HSERO_RS13965	−2.79	Single-stranded DNA exonuclease
	HSERO_RS17980	−2.78	GTP-binding protein
	HSERO_RS14955	−1.99	Chemotaxis protein
	HSERO_RS00080	−1.95	Histidine kinase
	HSERO_RS20750	−1.94	Cytochrome c oxidase subunit I

^aA complete list of mutations is available in Table S2 in the supplemental material. DQS4^T, *Azoarcus olearius*; SmR1, *Herbaspirillum seropedicae*.

UDP-*N*-acetylglucosamine enolpyruvate. A previous study of *H. seropedicae* reported the importance of the *mur* genes for maize root association and also showed that their expression was repressed in the presence of naringenin (37). In addition, a *N*-acetyl-anhydromuramyl-L-alanine amidase AmpD, likely involved in the degradation of peptidoglycan by hydrolysis of mucopeptides that can serve as signals for induction of β -lactamase (38–40), was identified as important for root colonization in both strains. We also identified a putative transcriptional regulator (DQS_RS12710) of catechol 2,3-dioxygenase, a flavonoid-related compound that might be involved in the degradation of aromatic compounds important for plant-bacterial interactions (41, 42). Among such compounds are flavonoids that can serve as chemoattractants for rhizobia in legumes (43, 44). Flavonoids can also mediate PGPB colonization, as reported for *H. seropedicae* colonization of *Arabidopsis* and the colonization of rice by *Serratia* spp. (45, 46). Interestingly, mutations in genes involved in PHB metabolism—including PhaP1 (HSERO_RS08150), a phasin family protein that affects PHB production, and PhbA1 (HSERO_RS01265), an acetyl-CoA acetyltransferase—decreased root colonization. The importance of PHB metabolism in bacteria during plant root colonization was reported previously in studies of *Setaria-Herbaspirillum* and legume-rhizobium symbiosis (12, 47).

Validation of candidate genes by insertional mutagenesis. To more definitively test the importance of individual genes identified in our screen, we selected candidate genes and created insertion mutations in 15 separate genes. These genes were selected to cover a diversity of putative functions representing operons containing multiple genes, as well as single genes, including those associated with both positive and negative fitness scores (Table 3). Using a gnotobiotic system, a competitive colonization assay was performed in which colonization of *S. viridis* roots by the individual mutants was measured, as well as their ability to compete for colonization when coinoculated (1:1) with the corresponding wild-type strain. Since root colonization is a dynamic process and might change over time, we evaluated competition at three different time points.

For *A. olearius* DQS4^T, we selected six genes where mutations reduced colonization, specifically a transcriptional regulator, *ybeZ*, ketoacyl ACP synthase (KAS), *peaF*, *pcm*, and *purU* (see the fitness score in Table 2).

Mutants defective in genes that resulted in reduced colonization in competition with the wild-type strain showed delayed colonization of *Setaria* roots. Mutation of the transcriptional regulator gene represents the most extreme case where colonization was completely blocked when inoculated singly onto *Setaria* roots (Fig. 5A and B; see also Fig. S4a and c). This gene encodes an AphS protein known to be involved in the phenol degradation pathway, where phenol is converted into catechol via catechol 2,3-dioxygenase (48). Analysis of *A. olearius* DQS4^T and BH72 genomes showed two

TABLE 3 Candidate genes selected for validation by gene knockout^a

Locus ID	Gene	Fitness	Description
DQS_RS04175	<i>ybeZ</i>	-3.26	Phosphate starvation-inducible protein
DQS_RS19965	<i>dqs_4056</i>	-4.48	Beta-ketoacyl synthase, N-terminal domain
DQS_RS06650	<i>peaF</i>	-4.67	FAD-dependent oxidoreductase
DQS_RS04520	<i>pcm</i>	-1.38	Protein-L-isoaspartate (D-aspartate) O-methyltransferase
DQS_RS19155	<i>purU</i>	-2.49	Formyltetrahydrofolate deformylase
DQS_RS12710	<i>aphS</i>	-3.72	Transcriptional regulator
HSERO_RS13265	<i>Hsero_2648</i>	-1.17	G3E family GTPase
HSERO_RS01870	<i>Hsero_0377</i>	-1.75	ABC transporter permease
HSERO_RS14960	<i>cheY</i>	-2.11	Fis family transcriptional regulator
HSERO_RS08535	<i>opuBC</i>	-1.71	Glycine/betaine ABC transporter substrate-binding protein
HSERO_RS17980	<i>typA</i>	-2.78	GTP-binding protein
HSERO_RS01880	<i>ampD</i>	-2.23	N-Acetyl-anhydromuranmyl-L-alanine amidase
HSERO_RS07970	<i>purD</i>	-3.06	Formyltetrahydrofolate deformylase
HSERO_RS03440	<i>exbD</i>	-2.43	Biopolymer transporter ExbD
HSERO_RS03445	<i>exbB</i>	-2.61	Biopolymer transport transmembrane protein

^aThe descriptions are based on eggNog annotation and UniProt. DQS4^T, *Azoarcus olearius*; SmR1, *Herbaspirillum seropedicae*.

operons potentially involved in the degradation of aromatic compounds located adjacent to the *aphS* gene (Fig. 6). However, consistent with the TnSeq results, when coinoculated with the wild-type DQS4^T, the mutant reduced colonization more modestly (see Fig. S4b and d). This result suggests the need for an unknown metabolite or signal




FIG 5 Candidate genes selected for validation during *S. viridis* root colonization. Root colonization by *Azoarcus olearius* DQS4^T or *Herbaspirillum seropedicae* SmR1 wild type and mutants defective in genes that decreased bacterial root colonization. (A) Root colonization by DQS4^T wild type (solid gray bars) and six mutant strains (white bars) when inoculated individually, recovered at 1 d.a.i. (B) Coinoculation of *S. viridis* roots with a mixture of equal amounts of wild type DQS4^T and mutant strains. (C) Root colonization of SmR1 wild type and nine mutant strains at 1 d.a.i. when inoculated individually. (D) Coinoculation with a mixture of equal amounts of wild-type SmR1 and mutant strains. The data are expressed in log of CFU per g of root fresh tissue. Bars indicate mean averages \pm the SE ($n=20$). Statistical significance, determined using a Student *t* test, is indicated by asterisks (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$). NR, not recovered; KAS, ketoacyl ACP synthase; TR, transcriptional regulator; ABC, ABC transporters.




FIG 6 Operon organization of aromatic metabolism in *Azoarcus olearius* DQS4^T. The transcriptional regulator *aphS* (DQS_RS12710) identified by TnSeq regulates genes related to phenol degradation and utilization. Genes and operons are indicated by arrows; use of the same color within a gray shade indicates genes that belong to an operon. DQS_RS12665 indicates the degradation protein meta in *A. olearius*.

that is produced by the wild type that can compensate for its loss in the mutant. As for the other mutants that impaired *A. olearius* DQS4^T colonization *ybeZ*, ketoacyl ACP synthase (*KAS*), *peaF*, *pcm*, and *purU*, all showed a significant reduction in colonization that was most pronounced 3 and 5 days after inoculation both when inoculated individually or in competition with the wild-type strain (see Fig. S4a and b).

Interestingly, *H. seropedicae* SmR1 genes related to PHB metabolism appeared to affect fitness by enhancing or decreasing root colonization as shown in Fig. 4. Among the nine selected genes where mutations impaired colonization by *H. seropedicae* (Table 2), six mutants—specifically the *opuBC*, *typA*, *ampD*, *purD*, *exbD*, and *exbB* mutants—showed a significant reduction in root colonization either when inoculated separately or coinoculated with the wild type (Fig. 5C and D; see also Fig. S5a and d).

DISCUSSION

Applying the transposon mutagenesis sequencing (TnSeq) approach to both *A. olearius* DQS4^T and *H. seropedicae* SmR1 revealed many genes required for these bacteria to competitively colonize *Setaria viridis* roots.

Our experiments identified 89 and 130 genes where mutations significantly affected the ability of *A. olearius* or *H. seropedicae*, respectively, to colonize *S. viridis* roots, including some genes previously reported to play a role in the plant-microbe associations. This result alone argues that root colonization is not a simple process but one that involves a variety of bacterial functions. General gene classes include those involved in cell wall biosynthesis, motility, chemotaxis and defense, and amino acid metabolism (19, 49–51).

Colonization assays with the 15 candidate genes selected for further confirmation by insertional mutagenesis showed a strong correlation with the results obtained by TnSeq, giving confidence that most if not all of the genes identified are likely important for root colonization by these PGPB strains.

Genes where mutations benefited root colonization can be assumed to normally play a role in suppressing colonization in wild-type cells. Among such genes in *A. olearius* DQS4^T is a homolog of the BH72 *exaA5* gene (azo3865), predicted to encode a pyrroloquinoline quinone-dependent alcohol dehydrogenase (DQS_RS19730) involved in methanol oxidation. Consistent with these findings, previous reports showed that mutation of ADH genes inhibited competitive colonization of rice roots by *A. olearius* BH72 (21, 52). Similarly, *Methylobacterium* spp. mutants defective in methanol oxidation were less competitive for colonization of *Medicago truncatula* roots when coinoculated with wild-type cells (53, 54). These data suggest that methanol metabolism is important for bacterial growth on the root surface and, perhaps more importantly, colonization is a very dynamic and likely heavily influenced by the overall microbial community.

We observed genes that increased fitness scores clustered within an operon presumably involved in iron uptake in *Azoarcus*. These genes are predicted to encode an outer membrane, ferric coprogen protein FhuE (DQS_RS15430), and a TonB-dependent siderophore receptor (DQS_RS15445), both of which were previously implicated in the

ability of this bacterium to colonize roots (9, 14). Bacterial iron uptake is complex, perhaps involving multiple bacterial processes, and can also be coopted by plant-encoded mechanisms. Mutation of these genes increased fitness values conveying a phenotypic advantage during root colonization, although less competitive than the wild type. Given the general role that iron availability plays in the ability of microorganisms to thrive and compete in virtually any environment, it is not surprising that iron uptake is also a crucial function for root colonization (17, 55). Recently, analysis of iron content in maize treated with the PGPB *A. brasilense* revealed a significant increase in total iron accumulation in seeds and higher yield (56), suggesting that PGPB can contribute to the iron metabolism of the host plant.

Previous studies demonstrated an important role for bacterial genes involved in motility for both endophytic and rhizosphere colonization of host roots (38, 57). Many genes involved in cell motility were among the common COG categories that appear to provide a fitness advantage for root colonization by SmR1, specifically mutations in genes related to flagellum assembly. We showed previously that an *H. seropedicae* mutant in the flagellar regulatory gene *fliA* was unable to endophytically colonize *S. viridis* roots, although this mutation did not affect rhizosphere colonization (12). *fliA* encodes the sigma factor σ_{28} RNA polymerase that mediates the transcription of genes involved in motility and flagellar synthesis (58). It is hard to imagine how the loss of motility *per se* could enhance root colonization. However, bacterial flagella can be recognized by specific receptors in plant cell membranes and activate a cascade of immune responses controlling bacterial infection (59–61). Transcriptome analysis of SmR1 attached to wheat roots showed that the flagellar gene cluster was downregulated, suggesting that the bacteria might switch to a twitching type of motility mediated by type IV pili (62).

Under certain environmental stresses or nutritional conditions, bacteria can use different sources of energy for survival, including mobilization of polymers such as polyhydroxyalkanoates (PHA). PHB is the PHA produced by bacteria. The PHB granules act as carbon storage that can be mobilized under different conditions. We found that disruption of PHA depolymerase, encoded by the *phaZ1* gene, enhanced bacterial colonization when inoculated individually or in competition with the wild type. According to Silveira Alves et al. (12), plant biomass was significantly reduced in *S. viridis* colonized by Δ *phaZ1* mutant despite colonizing roots to the same level as the wild-type strain. In contrast, we identified PhaP1 encoding a phasin involved in the PHB production (62–66), where deletion of Δ *phaP1* affected fitness negatively, reducing root colonization. Corroborating our findings, an increase in gene expression of the phasin genes was reported during wheat root colonization (62).

Many genes involved in peptidoglycan and lipopolysaccharide (LPS) biosynthesis were predicted to impair bacterial colonization of plant roots. For instance, mutation of LPS biosynthetic genes or the addition of exogenous *N*-acetylglucosamine was previously shown to impair *H. seropedicae*-maize root association (67). In studies of rice roots colonized by *Azoarcus* BH72, mutation of an endoglucanase (enzyme that cleaves cellulose) reduced root colonization, suggesting its importance for successful host cell invasion (68, 69). We demonstrated that a mutation in a transcriptional regulator (*aphS*) involved in aromatic compound degradation completely impaired root colonization during single inoculation. AphS was predicted to regulate genes important for phenol degradation in *A. olearius* BH72 (48).

Only two genes, *cheY* involved in chemotaxis and *ampD* involved in peptidoglycan cell wall recycling (39, 70), affected colonization of both strains. Our data suggest that bacterial chemotaxis provides a competitive advantage to wild-type cells during colonization of the plant root tissue. This system is well characterized in several motile bacterial species, such as *E. coli* and beneficial bacteria such as *Azospirillum brasilense*, *S. meliloti*, and others (31, 71–75).

In summary, our data indicate that, rather than a single or small subset of crucial functions, each strain uses differing functions for colonization, reflecting the unique

characteristics of each bacterium. Given that our study was focused on bacterial root colonization, mutations related to soil survival were not considered. However, for *Azoarcus* a deeper investigation of mutations that affected survival in soil could be useful in explaining the different lifestyle and adaptations of each bacterium to their environment, especially considering that *Azoarcus* sp. BH72, closely related to *A. olearius* DQS4^T, is a strict endophyte and has not been reported to survive without a host (76).

In summary, similar to most plant-microbe interactions, PGPB-plant interactions are complex and reflect the ability of the plant host and bacterial symbiont to profoundly influence the metabolism of the other. The fact that PGPB have broad host ranges and can enhance crop yield under field conditions has contributed to a continuing interest in using PGPB inoculants in agriculture. This study provides insights to better understand those gene functions involved in PGPB-host interaction and hopefully will contribute to the further development of PGPB inoculants for an efficient, sustainable agriculture.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *Azoarcus olearius* DQS4^T (Sm^r Nal^r) and *Herbaspirillum seropedicae* SmR1 (Sm^r) were used as bacterial model systems for this study. Both strains were grown in NFB-HPN (high phosphate and nitrogen) medium containing malate as carbon source (77, 78) supplemented with NH₄Cl (10 mmol liter⁻¹ [DQS4^T] and 20 mmol liter⁻¹ [SmR1]) according to the strain's requirements. The bacterial cultures were grown at 37°C for DQS4^T and 30°C for SmR1, with shaking at 130 rpm overnight.

***Setaria viridis* seed sterilization and growth conditions.** *Setaria viridis* A10.1 seeds were treated with 5 ml of sulfuric acid for 15 min to break dormancy and washed with running water for 1 min. Upon washing, the seeds were surface sterilized with 1% bleach plus 0.1% Tween 20 (vol/vol) for 3 min and then rinsed three times with sterile distilled water. Sterile seeds were sown onto plates containing one half Murashige-Skoog medium with 5% sucrose (wt/vol) and 1% Phytigel (wt/vol). Plates were placed in the dark for 24 h and then 2 days in the light at 30°C for germination. After germination, 20 seedlings of similar sizes were transferred to a pot containing sterilized Turface (Turface MVP) and vermiculite in a 3:1 proportion and grown for 5 days prior to inoculation. Four biological replicates were performed, including soil controls without plants. Plants were watered every day, and an additional 0.2% malate solution was added to the soil control as a carbon source to enhance bacterial survival.

Transposon mutant library construction. We applied two different approaches to generate mutant libraries for each strain, using engineered Tn5 transposon delivery vectors. (i) *A. olearius* DQS4^T was conjugated with *E. coli* harboring pJG714, which is described in (79). Around 130,000 transposants were generated by selection on kanamycin (50 μg ml⁻¹). The mutant library was harvested from plates and stored in 10% glycerol at -80°C for further TnSeq experiments. (ii) An *H. seropedicae* SmR1 library was constructed as previously described (18, 80) using a random barcode (RB-TnSeq) method (16). Cell aliquots of 1 ml (optical density at 600 nm [OD₆₀₀] of 1) were stored in 10% glycerol at -80°C until the TnSeq experiments were performed.

Mutant library competitive fitness assay on plant roots. A single aliquot of the DQS4^T mutant library was thawed and 50 μl was inoculated into 50 ml of modified Luria-Bertani (LB) medium containing 5 g liter⁻¹ tryptone, 5 g liter⁻¹ yeast extract, 3 g liter⁻¹ NaCl, and 0.5 g liter⁻¹ MgSO₄·7H₂O supplemented with kanamycin. For SmR1, glycerol stocks of 1-ml aliquots of the mutant library were inoculated into 50 ml of LB medium with kanamycin. The cultures were grown until reaching an OD₆₀₀ of 1.0, pelleted, and washed prior to dilution to 2 × 10⁹ cells/ml (DQS4^T) or 2 × 10⁸ cells/ml (SmR1) in modified Hoagland's solution without nitrogen source (81). Then, 50 ml of diluted DQS4^T or 10 ml of SmR1 culture was added directly onto the soil in each pot containing 20 plants or soil control pots. A plate was placed on the bottom of the pots in order to collect the flowthrough culture, and the plants were reinoculated after 15 min. The plants were kept in a growth chamber (Conviron) at 25°C with 16-h light/8-h dark cycle for 15 days. After 15 days, mutants were recovered from *S. viridis* roots by removing the 20 plants from the soil. Using a sterile surface, a 3:1 vermiculite potting mix allowed for the removal of the plants from the potting mix with few adhering particles. Roots were immersed into 100 ml of BRM medium incubated at 37°C for DQS4^T or NFB-HPN-malate incubated at 30°C for SmR1, both containing kanamycin (50 μg/ml), and then incubated for 6 h under 120 rpm agitation. After 6 h of agitation, 50-ml portions of the culture were filtered through sterile filter paper to eliminate any soil debris and transferred to a new flask with fresh medium (Km^r) incubated at 37°C or 30°C, according to each strain, followed by shaking at 120 rpm overnight. The cultures were then centrifuged for 2 min at 14,000 rpm. The supernatant was discarded; the pellets were immediately frozen in liquid nitrogen and then stored at -80°C prior to sequencing library preparation. Soil culture was filtered after 3 h of incubation to settle down soil particles to enable their separation from the culture.

TnSeq library preparation. For *Azoarcus olearius* DQS4^T, a bacterial culture at mid-log phase grown at 37°C was collected for DNA extraction. Cells were pelleted by centrifugation and immediately frozen at -80°C until DNA extraction. Genomic DNA was extracted, and transposon insertion junctions were selectively amplified, as described previously (79). Insertion junction libraries were multiplexed and sequenced using the Illumina HiSeq 2500 system. For *Herbaspirillum seropedicae* SmR1, libraries were sequenced on either the HiSeq 2000 or HiSeq 2500 system (Illumina) to map a greater fraction of the

mutant population. Genomic DNA was extracted, and barcodes were amplified as described previously (16, 18, 80).

Analysis of transposon insertion impact on bacterial fitness. For *A. olearius*, TnSeq reads were mapped to the reference genome, and genes received fitness values representing the relative abundance of insertions across conditions. For *H. seropedicae* SmR1, we calculated gene fitness scores based on the barcode abundance of the individual strains in the library. The fitness values represent the count in each sample relative to the time zero sample (for details, see reference 6). Differences in fitness values with a *P* value of ≤ 0.05 were considered statistically significant and categorized into two groups based on their fitness score: (i) enhanced fitness ≥ 1 or (ii) decreased fitness ≤ -1 .

Confirmation of gene functionality by site-directed mutagenesis. The suicide plasmid pJG194 carrying Km^r (82) was used to generate targeted bacterial gene insertion mutations in various candidate genes via single crossover. Amplicons (~500 bp) for internal gene fragments were cloned into plasmid pJG194 using HindIII and EcoRI restriction sites (see Table S1 in the supplemental material). Bacterial template gDNA was extracted by using a DNeasy Blood & Tissue (Qiagen) according to the protocol for Gram-negative bacteria. Amplification reactions were carried out using Phusion high-fidelity DNA polymerase (Thermo Fisher) under the following cycling conditions: 98°C for 1 min; 30 cycles 98°C for 20 s, 57°C for 30 s, and 72°C for 30 s; and a final extension of 72°C for 10 min, 4°C. PCR products were verified on a 1% electrophoresis agarose gel for the correct 500-bp size. The correct products were purified using Wizard SV gel and PCR cleanup system (Promega) and then digested with FastDigest HindIII and EcoRI for 20 min each. Samples were ligated with T4 ligase at 4°C overnight. Ligation products were transformed into *E. coli* EC100 competent cells (Epicentre, Madison, WI) by heat shock (10 min on ice, 45 s at 42°C, and 1 min on ice) and then added to 200 μ l of LB medium, followed by shaking at 37°C for 1 h prior plating onto LB+Km^r. The inserts of the generated plasmids were confirmed by PCR and DNA sequencing prior to transformation into *E. coli* ST18 (83) or S17.1 (84) for mating with the wild-type DQS4^T or SmR1 strain, respectively. Mutants were confirmed to have the desired gene disruptions by PCR analysis and DNA sequencing.

Competition assay in planta. To assess bacterial gene function in plant colonization, we performed a colonization assay using *Setaria viridis* seedlings grown hydroponically. Seeds were germinated on plates as described above and transferred to a glass tube containing 20 ml of sterile modified Hoagland's solution and 8 g of polypropylene beads. Seedlings were inoculated with 1 ml of bacterial culture (10^{-6} cells ml⁻¹) of the wild-type strain or mutant strain separately or in a 1:1 proportion. Quantification of colonization was performed by recovering the attached bacteria from the roots (*n* = 20) at 1, 3, and 5 d.a.i. Briefly, seedlings were carefully removed from the glass tubes and placed in a 2-ml tube. The roots were weighed, and then 1 ml of 0.9% NaCl was added to each tube. To recover the surface-attached bacteria, tubes were vortexed at high speed for 1 min, and the solution containing suspended bacteria was serially diluted. The dilutions were plated onto NFB-HPN malate medium containing appropriate antibiotics, including kanamycin for the targeted gene disruption mutants, and then incubated at 30°C for 24 h prior to colony counting and conversion to CFU g⁻¹ of fresh tissue.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

FIG S1, TIF file, 2.1 MB.

FIG S2, TIF file, 2.6 MB.

FIG S3, TIF file, 2.4 MB.

FIG S4, TIF file, 2.3 MB.

FIG S5, TIF file, 1.8 MB.

TABLE S1, DOCX file, 0.02 MB.

TABLE S2, XLSX file, 0.02 MB.

ACKNOWLEDGMENTS

This study was financially supported by the Department of Energy, Office of Biological and Environmental Research.

We thank M. Muller Santos from Federal University of Parana for providing PHB mutant strains. We acknowledge L. P. Alves and M. Cotta for their help with *Setaria* plants and Ha N. Duong for drawing Fig. S2.

We declare that we have no conflicts of interest.

REFERENCES

1. Pankievicz VCS, Irving TB, Maia LGS, Ané J-M. 2019. Are we there yet? The long walk towards the development of efficient symbiotic associations between nitrogen-fixing bacteria and non-leguminous crops. *BMC Biol* 17:99. <https://doi.org/10.1186/s12915-019-0710-0>.
2. Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith DL. 2018. Plant growth-promoting rhizobacteria: context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Front Plant Sci* 9:1473. <https://doi.org/10.3389/fpls.2018.01473>.
3. Olanrewaju OS, Glick BR, Babalola OO. 2017. Mechanisms of action of plant growth promoting bacteria. *World J Microbiol Biotechnol* 33:197. <https://doi.org/10.1007/s11274-017-2364-9>.

4. Monteiro RA, Balsanelli E, Wassem R, Marin AM, Brusamarello-Santos LCC, Schmidt MA, Tadra-Sfeir MZ, Pankiewicz VCS, Cruz LM, Chubatsu LS, Pedrosa FO, Souza EM. 2012. Herbaspirillum-plant interactions: microscopical, histological, and molecular aspects. *Plant Soil* 356:175–196. <https://doi.org/10.1007/s11104-012-1125-7>.
5. Hurek T, Reinhold-Hurek B, Van Montagu M, Kellenberger E. 1994. Root colonization and systemic spreading of *Azoarcus* sp. strain BH72 in grasses. *J Bacteriol* 176:1913–1923. <https://doi.org/10.1128/jb.176.7.1913-1923.1994>.
6. Compant S, Clement C, Sessitsch A. 2010. Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonization, mechanisms involved, and prospects for utilization. *Soil Biol Biochem* 42:669–678. <https://doi.org/10.1016/j.soilbio.2009.11.024>.
7. Afzal I, Shinwari ZK, Sikandar S, Shahzad S. 2019. Plant beneficial endophytic bacteria: mechanisms, diversity, host range and genetic determinants. *Microbiol Res* 221:36–49. <https://doi.org/10.1016/j.micres.2019.02.001>.
8. Pankiewicz VCS, do Amaral FP, Santos KFDN, Agtuca B, Xu Y, Schueller MJ, Arisi ACM, Steffens MBR, de Souza EM, Pedrosa FO, Stacey G, Ferrieri RA. 2015. Robust biological nitrogen fixation in a model grass-bacterial association. *Plant J* 81:907–919. <https://doi.org/10.1111/tj.12777>.
9. Faoro H, Rene Menegazzo R, Battistoni F, Gyaneshwar P, do Amaral FP, Taulé C, Rausch S, Gonçalves Galvão P, de Los Santos C, Mitra S, Heijo G, Sheu S-Y, Chen W-M, Mareque C, Zibetti Tadra-Sfeir M, Ivo Baldani J, Maluk M, Paula Guimarães A, Stacey G, de Souza EM, Pedrosa FO, Magalhães Cruz L, James EK. 2017. The oil-contaminated soil diazotroph *Azoarcus olearius* DQS-4^T is genetically and phenotypically similar to the model grass endophyte *Azoarcus* sp. BH72. *Environ Microbiol Rep* 9:223–238. <https://doi.org/10.1111/1758-2229.12502>.
10. Chubatsu LS, Monteiro RA, de Souza EM, de Oliveira MAS, Yates MG, Wassem R, Bonatto AC, Huerger LF, Steffens MBR, Rigo LU, Pedrosa FDO. 2012. Nitrogen fixation control in *Herbaspirillum seropedicae*. *Plant Soil* 356:197–207. <https://doi.org/10.1007/s11104-011-0819-6>.
11. Pedrosa FO, Monteiro RA, Wassem R, Cruz LM, Ayub RA, Colauto NB, Fernandez MA, Fungaro MHP, Grisard EC, Hungria M, Madeira HMF, Nodari RO, Osaku CA, Petzl-Erler ML, Terenzi H, Vieira LGE, Steffens MBR, Weiss VA, Pereira LFP, Almeida MIM, Alves LR, Marin A, Araujo LM, Balsanelli E, Baura VA, Chubatsu LS, Faoro H, Favetti A, Friedemann G, Glienke C, Karp S, Kava-Cordeiro V, Raittz RT, Ramos HJO, Ribeiro EMSF, Rigo LU, Rocha SN, Schwab S, Silva AG, Souza EM, Tadra-Sfeir MZ, Torres RA, Dabul ANG, Soares MAM, Gasques LS, Gimenes CCT, Valle JS, Ciferri RR, Correa LC, Murace NK, et al. 2011. Genome of *Herbaspirillum seropedicae* strain SmR1, a specialized diazotrophic endophyte of tropical grasses. *PLoS Genet* 7:e1002064. <https://doi.org/10.1371/journal.pgen.1002064>.
12. Silveira Alves LP, Plucani do Amaral F, Kim D, Todo Bom M, Piñero Gavidia M, Silvano Teixeira C, Holthman F, de Oliveira Pedrosa F, Maltempi de Souza E, Chubatsu LS, Müller-Santos M, Stacey G. 2019. Importance of poly-3-hydroxybutyrate (PHB) metabolism to the ability of *Herbaspirillum seropedicae* to promote plant growth. *Appl Environ Microbiol* 85:e02586-18. <https://doi.org/10.1128/AEM.02586-18>.
13. Chen M-H, Sheu S-Y, James EK, Young C-C, Chen W-M. 2013. *Azoarcus olearius* sp. nov., a nitrogen-fixing bacterium isolated from oil-contaminated soil. *Int J Syst Evol Microbiol* 63:3755–3761. <https://doi.org/10.1099/ijs.0.050609-0>.
14. Krause A, Ramakumar A, Bartels D, Battistoni F, Bekel T, Boch J, Böhm M, Friedrich F, Hurek T, Krause L, Linke B, McHardy AC, Sarkar A, Schneiker S, Syed AA, Thauer R, Vorhölter F-J, Weidner S, Pühler A, Reinhold-Hurek B, Kaiser O, Goesmann A. 2006. Complete genome of the mutualistic, N₂-fixing grass endophyte *Azoarcus* sp strain BH72. *Nat Biotechnol* 24:1385–1391. <https://doi.org/10.1038/nbt1243>.
15. Agtuca BJ, Stopka SA, Tuleski TR, do Amaral FP, Evans S, Liu Y, Xu D, Monteiro RA, Koppelaar DW, Paša-Tolić L, Anderton CR, Vertes A, Stacey G. 2020. *In situ* metabolomic analysis of *Setaria viridis* roots colonized by beneficial endophytic bacteria. *Mol Plant Microbe Interact* 33:272–283. <https://doi.org/10.1094/MPMI-06-19-0174-R>.
16. Wetmore KM, Price MN, Waters RJ, Lamson JS, He J, Hoover CA, Blow MJ, Bristow J, Butland G, Arkin AP, Deutschbauer A. 2015. Rapid quantification of mutant fitness in diverse bacteria by sequencing randomly bar-coded transposons. *mBio* 6:e00306-15. <https://doi.org/10.1128/mBio.00306-15>.
17. Van Opijnen T, Bodi KL, Camilli A. 2009. Tn-Seq: high-throughput parallel sequencing for fitness and genetic interaction studies in microorganisms. *Nat Methods* 6:767–U21. <https://doi.org/10.1038/nmeth.1377>.
18. Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV, Melynk RA, Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM, Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP, Deutschbauer AM. 2018. Mutant phenotypes for thousands of bacterial genes of unknown function. *Nature* 557:503–509. <https://doi.org/10.1038/s41586-018-0124-0>.
19. Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald M, Yoshikuni Y, Malmstrom RR, Deutschbauer AM, Dangl JL, Visel A. 2017. Genome-wide identification of bacterial plant colonization genes. *PLoS Biol* 15:e2002860. <https://doi.org/10.1371/journal.pbio.2002860>.
20. Reinhold-Hurek B, Hurek T. 2011. Living inside plants: bacterial endophytes. *Curr Opin Plant Biol* 14:435–443. <https://doi.org/10.1016/j.pbi.2011.04.004>.
21. Krause A, Bischoff B, Miché L, Battistoni F, Reinhold-Hurek B. 2011. Exploring the function of alcohol dehydrogenases during the endophytic life of *Azoarcus* sp. strain BH72. *Mol Plant Microbe Interact* 24:1325–1332. <https://doi.org/10.1094/MPMI-05-11-0139>.
22. Hengge R. 2009. Principles of c-di-GMP signaling in bacteria. *Nat Rev Microbiol* 7:263–273. <https://doi.org/10.1038/nrmicro2109>.
23. Hengge R, Galperin MY, Ghigo J-M, Gomelsky M, Green J, Hughes KT, Jenal U, Landini P. 2016. Systematic nomenclature for GGDEF and EAL domain-containing cyclic di-GMP turnover proteins of *Escherichia coli*. *J Bacteriol* 198:7–11. <https://doi.org/10.1128/JB.00424-15>.
24. Romling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial second messenger. *Microbiol Mol Biol Rev* 77:1–52. <https://doi.org/10.1128/MMBR.00043-12>.
25. Romling U. 2015. Small molecules with big effects: cyclic di-GMP-mediated stimulation of cellulose production by the amino acid L-arginine. *Sci Signal* 8:fs12. <https://doi.org/10.1126/scisignal.aac4734>.
26. Kim HK, Harshey RM. 2016. A diguanylate cyclase acts as a cell division inhibitor in a two-step response to reductive and envelope stresses. *mBio* 7:e00822-16. <https://doi.org/10.1128/mBio.00822-16>.
27. O'Neal L, Ryu M-H, Gomelsky M, Alexandre G. 2017. Optogenetic manipulation of cyclic di-GMP (c-di-GMP) levels reveals the role of c-di-GMP in regulating aerotaxis receptor activity in *Azospirillum brasilense*. *J Bacteriol* 199:e00020-17. <https://doi.org/10.1128/JB.00020-17>.
28. Noinaj N, Guillier M, Barnard TJ, Buchanan SK. 2010. TonB-dependent transporters: regulation, structure, and function. *Annu Rev Microbiol* 64:43–60. <https://doi.org/10.1146/annurev.micro.112408.134247>.
29. Blair DF. 2003. Flagellar movement driven by proton translocation. *FEBS Lett* 545:86–95. [https://doi.org/10.1016/S0014-5793\(03\)00397-1](https://doi.org/10.1016/S0014-5793(03)00397-1).
30. Babel W, Ackermann JU, Breuer U. 2001. Physiology, regulation, and limits of the synthesis of poly(3HB). *Adv Biochem Eng Biotechnol* 71:125–157. https://doi.org/10.1007/3-540-40021-4_4.
31. Scharf BE, Hynes MF, Alexandre GM. 2016. Chemotaxis signaling systems in model beneficial plant-bacterium associations. *Plant Mol Biol* 90:549–559. <https://doi.org/10.1007/s11103-016-0432-4>.
32. Wong-Ng J, Celani A, Vergassola M. 2018. Exploring the function of bacterial chemotaxis. *Curr Opin Microbiol* 45:16–21. <https://doi.org/10.1016/j.mib.2018.01.010>.
33. Clarke CR, Hayes BW, Runde BJ, Markel E, Swingle BM, Vinatzer BA. 2016. Comparative genomics of *Pseudomonas syringae* pathovar tomato reveals novel chemotaxis pathways associated with motility and plant pathogenicity. *PeerJ* 4:e2570. <https://doi.org/10.7717/peerj.2570>.
34. Li MS, Hazelbauer GL. 2004. Cellular stoichiometry of the components of the chemotaxis signaling complex. *J Bacteriol* 186:3687–3694. <https://doi.org/10.1128/JB.186.12.3687-3694.2004>.
35. Balsanelli E, Tadra-Sfeir MZ, Faoro H, Pankiewicz VC, de Baura VA, Pedrosa FO, de Souza EM, Dixon R, Monteiro RA. 2016. Molecular adaptations of *Herbaspirillum seropedicae* during colonization of the maize rhizosphere. *Environ Microbiol* 18:2343–2356. <https://doi.org/10.1111/1462-2920.12887>.
36. Khursigara CM, Wu X, Zhang P, Lefman J, Subramaniam S. 2008. Role of HAMP domains in chemotaxis signaling by bacterial chemoreceptors. *Proc Natl Acad Sci U S A* 105:16555–16560. <https://doi.org/10.1073/pnas.0806401105>.
37. Tadra-Sfeir MZ, Faoro H, Camilios-Neto D, Brusamarello-Santos L, Balsanelli E, Weiss V, Baura VA, Wassem R, Cruz LM, De Oliveira Pedrosa F, Souza EM, Monteiro RA. 2015. Genomewide transcriptional profiling of *Herbaspirillum seropedicae* SmR1 grown in the presence of naringenin. *Front Microbiol* 6:491.
38. Jacobs C, Huang LJ, Bartowsky E, Normark S, Park JT. 1994. Bacterial cell wall recycling provides cytosolic muropeptides as effectors for beta-lactamase induction. *EMBO J* 13:4684–4694. <https://doi.org/10.1002/j.1460-2075.1994.tb06792.x>.

39. Hölft J, Kopp U, Ursinus A, Wiedemann B. 1994. The negative regulator of beta-lactamase induction AmpD is a N-acetyl-anhydromuramyl-L-alanine amidase. *FEMS Microbiol Lett* 122:159–164. <https://doi.org/10.1111/j.1574-6968.1994.tb07159.x>.
40. Zeng X, Lin J. 2013. Beta-lactamase induction and cell wall metabolism in Gram-negative bacteria. *Front Microbiol* 4:128.
41. Marin AM, Souza EM, Pedrosa FO, Souza LM, Sassaki GL, Baura VA, Yates MG, Wasseem R, Monteiro RA. 2013. Naringenin degradation by the endophytic diazotroph *Herbaspirillum seropedicae* SmR1. *Microbiology (Reading)* 159:167–175. <https://doi.org/10.1099/mic.0.061135-0>.
42. Marin AM. 2016. Genetic and functional characterization of a novel meta-pathway for degradation of naringenin in *Herbaspirillum seropedicae* SmR1. *Environ Microbiol* 18:4653–4661. <https://doi.org/10.1111/1462-2920.13313>.
43. Oldroyd GE, Murray JD, Poole PS, Downie JA. 2011. The rules of engagement in the legume-rhizobial symbiosis. *Annu Rev Genet* 45:119–144. <https://doi.org/10.1146/annurev-genet-110410-132549>.
44. Wang Q, Liu JE, Zhu HY. 2018. Genetic and molecular mechanisms underlying symbiotic specificity in legume-rhizobium interactions. *Front Plant Sci* 9:313. <https://doi.org/10.3389/fpls.2018.00313>.
45. Gough C, Galera C, Vasse J, Webster G, Cocking EC, Dénarié J. 1997. Specific flavonoids promote intercellular root colonization of *Arabidopsis thaliana* by *Azorhizobium caulinodans* ORS571. *Mol Plant Microbe Interact* 10:560–570. <https://doi.org/10.1094/MPMI.1997.10.5.560>.
46. Balachandar D, Sandhiya GS, Sugitha TCK, Kumar K. 2006. Flavonoids and growth hormones influence endophytic colonization and *in planta* nitrogen fixation by a diazotrophic *Serratia* sp. in rice. *World J Microbiol Biotechnol* 22:707–712. <https://doi.org/10.1007/s11274-005-9094-0>.
47. Trainer MA, Charles TC. 2006. The role of PHB metabolism in the symbiosis of rhizobia with legumes. *Appl Microbiol Biotechnol* 71:377–386. <https://doi.org/10.1007/s00253-006-0354-1>.
48. Suvorova IA, Gelfand MS. 2019. Comparative genomic analysis of the regulation of aromatic metabolism in *Betaproteobacteria*. *Front Microbiol* 10:642. <https://doi.org/10.3389/fmicb.2019.00642>.
49. Moe LA. 2013. Amino acids in the rhizosphere: from plants to microbes. *Am J Bot* 100:1692–1705. <https://doi.org/10.3732/ajb.1300033>.
50. Levy A, Salas Gonzalez I, Mittelviehhaus M, Clingenpeel S, Herrera Paredes S, Miao J, Wang K, Devescovi G, Stillman K, Monteiro F, Rangel Alvarez B, Lundberg DS, Lu T-Y, Lebeis S, Jin Z, McDonald M, Klein AP, Feltcher ME, Rio TG, Grant SR, Doty SL, Ley RE, Zhao B, Venturi V, Pelletier DA, Vorholt JA, Tringe SG, Woyke T, Dangl JL. 2017. Genomic features of bacterial adaptation to plants. *Nat Genet* 50:138–150. <https://doi.org/10.1038/s41588-017-0012-9>.
51. Badri DV, Vivanco JM. 2009. Regulation and function of root exudates. *Plant Cell Environ* 32:666–681. <https://doi.org/10.1111/j.1365-3040.2009.01926.x>.
52. Krause A, Julich H, Mankar M, Reinhold-Hurek B. 2017. The regulatory network controlling ethanol-induced expression of alcohol dehydrogenase in the endophyte *Azoarcus* sp. strain BH72. *Mol Plant Microbe Interact* 30:778–785. <https://doi.org/10.1094/MPMI-01-17-0013-R>.
53. Sy A, Timmers ACJ, Knief C, Vorholt JA. 2005. Methylotrophic metabolism is advantageous for *Methylobacterium extorquens* during colonization of *Medicago truncatula* under competitive conditions. *Appl Environ Microbiol* 71:7245–7252. <https://doi.org/10.1128/AEM.71.11.7245-7252.2005>.
54. Dourado MN, Camargo Neves AA, Santos DS, Araújo WL. 2015. Biotechnological and agronomic potential of endophytic pink-pigmented methylotrophic *Methylobacterium* spp. *Biomed Res Int* 2015:909016. <https://doi.org/10.1155/2015/909016>.
55. Miethke M, Marahiel MA. 2007. Siderophore-based iron acquisition and pathogen control. *Microbiol Mol Biol Rev* 71:413–451. <https://doi.org/10.1128/MMBR.00012-07>.
56. Scott S, Housh A, Powell G, Anstaett J, Gerheart A, Benoit M, Wilder S, Schueller M, Ferrieri R. 2020. Crop yield, ferritin, and Fe(II) boosted by *Azospirillum brasilense* (HM053) in corn. *Agronomy* 10:394. <https://doi.org/10.3390/agronomy10030394>.
57. Buschart A, Sachs S, Chen X, Herglotz J, Krause A, Reinhold-Hurek B. 2012. Flagella mediate endophytic competence rather than act as MAMPs in rice-*Azoarcus* sp. strain BH72 interactions. *Mol Plant Microbe Interact* 25:191–199. <https://doi.org/10.1094/MPMI-05-11-0138>.
58. Liu XY, Matsumura P. 1995. An alternative sigma-factor controls transcription of flagellar class III operons in *Escherichia coli*: gene sequence, overproduction, purification, and characterization. *Gene* 164:81–84. [https://doi.org/10.1016/0378-1119\(95\)00480-T](https://doi.org/10.1016/0378-1119(95)00480-T).
59. Gust AA, Pruitt R, Nurnberger T. 2017. Sensing danger: key to activating plant immunity. *Trends Plant Sci* 22:779–791. <https://doi.org/10.1016/j.tplants.2017.07.005>.
60. Zhang J, Zhou JM. 2010. Plant immunity triggered by microbial molecular signatures. *Mol Plant* 3:783–793. <https://doi.org/10.1093/mp/ssp035>.
61. Wheatley RM, Ford BL, Li L, Aroney STN, Knights HE, Ledermann R, East AK, Ramachandran VK, Poole PS. 2020. Lifestyle adaptations of *Rhizobium* from rhizosphere to symbiosis. *Proc Natl Acad Sci U S A* 117:23823–23834. <https://doi.org/10.1073/pnas.2009094117>.
62. Pankievicz VCS, Camilios-Neto D, Bonato P, Balsanelli E, Tadra-Sfeir MZ, Faoro H, Chubatsu LS, Donatti L, Wajnberg G, Passetti F, Monteiro RA, Pedrosa FO, Souza EM. 2016. RNA-seq transcriptional profiling of *Herbaspirillum seropedicae* colonizing wheat (*Triticum aestivum*) roots. *Plant Mol Biol* 90:589–603. <https://doi.org/10.1007/s11103-016-0430-6>.
63. Alves LPS, Teixeira CS, Tirapelle EF, Donatti L, Tadra-Sfeir MZ, Steffens MBR, de Souza EM, de Oliveira Pedrosa F, Chubatsu LS, Müller-Santos M. 2016. Backup expression of the PhaP2 phasin compensates for phaP1 deletion in *Herbaspirillum seropedicae*, maintaining fitness and PHB accumulation. *Front Microbiol* 7:739. <https://doi.org/10.3389/fmicb.2016.00739>.
64. Batista MB, Teixeira CS, Sfeir MZT, Alves LPS, Valdameri G, Pedrosa F. d O, Sassaki GL, Steffens MBR, de Souza EM, Dixon R, Müller-Santos M. 2018. PHB biosynthesis counteracts redox stress in *Herbaspirillum seropedicae*. *Front Microbiol* 9:472. <https://doi.org/10.3389/fmicb.2018.00472>.
65. Mezzina MP, Pettinari MJ. 2016. Phasins, multifaceted polyhydroxyalkanoate granule-associated proteins. *Appl Environ Microbiol* 82:5060–5067. <https://doi.org/10.1128/AEM.01161-16>.
66. Tirapelle EF, Müller-Santos M, Tadra-Sfeir MZ, Kadowaki MAS, Steffens MBR, Monteiro RA, Souza EM, Pedrosa FO, Chubatsu LS. 2013. Identification of proteins associated with polyhydroxybutyrate granules from *Herbaspirillum seropedicae* SmR1: old partners. *PLoS One* 8:e75066. <https://doi.org/10.1371/journal.pone.0075066>.
67. Balsanelli E, Tuleski TR, de Baura VA, Yates MG, Chubatsu LS, de Oliveira Pedrosa F, de Souza EM, Monteiro RA. 2013. Maize root lectins mediate the interaction with *Herbaspirillum seropedicae* via N-acetylglucosamine residues of lipopolysaccharides. *PLoS One* 8:e77001. <https://doi.org/10.1371/journal.pone.0077001>.
68. Reinhold-Hurek B, Hurek T, Gillis M, Hoste B, Vancanneyt M, Kersters K, De Ley J. 1993. *Azoarcus* gen. nov., nitrogen-fixing proteobacteria associated with roots of Kallar grass (*Leptochloa fusca* [L.] Kunth), and description of two species, *Azoarcus indigenus* sp. nov. and *Azoarcus communism* sp. nov. *Int J Syst Bacteriol* 43:574–584. <https://doi.org/10.1099/00207713-43-3-574>.
69. Reinhold-Hurek B, Maes T, Gemmer S, Van Montagu M, Hurek T. 2006. An endoglucanase is involved in infection of rice roots by the not-cellulose-metabolizing endophyte *Azoarcus* sp. strain BH72. *Mol Plant Microbe Interact* 19:181–188. <https://doi.org/10.1094/MPMI-19-0181>.
70. Carrasco-Lopez C. 2011. Crystal structures of bacterial peptidoglycan amidase AmpD and an unprecedented activation mechanism. *J Biol Chem* 286:31714–31722. <https://doi.org/10.1074/jbc.M111.264366>.
71. Di Paolo D, Afanzar O, Armitage JP, Berry RM. 2016. Single-molecule imaging of electroporated dye-labelled CheY in live *Escherichia coli*. *Philos Trans R Soc B Biol Sci* 371:20150492. <https://doi.org/10.1098/rstb.2015.0492>.
72. Reinhold B, Hurek T, Niemann EG, Fendrik I. 1986. Close association of *Azospirillum* and diazotrophic rods with different root zones of Kallar grass. *Appl Environ Microbiol* 52:520–526. <https://doi.org/10.1128/AEM.52.3.520-526.1986>.
73. Hurek T, Reinhold-Hurek B, Turner GL, Bergersen FJ. 1994. Augmented rates of respiration and efficient nitrogen fixation at nanomolar concentrations of dissolved O₂ in hyperinduced *Azoarcus* sp. strain BH72. *J Bacteriol* 176:4726–4733. <https://doi.org/10.1128/jb.176.15.4726-4733.1994>.
74. Bible A, Russell MH, Alexandre G. 2012. The *Azospirillum brasilense* Che1 chemotaxis pathway controls swimming velocity, which affects transient cell-to-cell clumping. *J Bacteriol* 194:3343–3355. <https://doi.org/10.1128/JB.00310-12>.
75. Huang Z, Pan X, Xu N, Guo M. 2019. Bacterial chemotaxis coupling protein: structure, function and diversity. *Microbiol Res* 219:40–48. <https://doi.org/10.1016/j.micres.2018.11.001>.
76. Dixon R, Hartmann A. 2017. Novel insights into ecological distribution and plant growth promotion by nitrogen-fixing endophytes: how specialized are they? *Environ Microbiol Rep* 9:179–181. <https://doi.org/10.1111/1758-2229.12529>.
77. Machado HB, Funayama S, Rigo LU, Pedrosa FO. 1991. Excretion of ammonium by *Azospirillum brasilense* mutants resistant to ethylenediamine. *Can J Microbiol* 37:549–553. <https://doi.org/10.1139/m91-092>.

78. Klassen G, Pedrosa FO, Souza EM, Funayama S, Rigo LU. 1997. Effect of nitrogen compounds on nitrogenase activity in *Herbaspirillum seropedicae* SmR1. *Can J Microbiol* 43:887–891. <https://doi.org/10.1139/m97-129>.
79. diCenzo GC, Benedict AB, Fondi M, Walker GC, Finan TM, Mengoni A, Griffiths JS. 2018. Robustness encoded across essential and accessory replicons of the ecologically versatile bacterium *Sinorhizobium meliloti*. *PLoS Genet* 14:e1007357. <https://doi.org/10.1371/journal.pgen.1007357>.
80. Price MN, Zane GM, Kuehl JV, Melnyk RA, Wall JD, Deutschbauer AM, Arkin AP. 2018. Filling gaps in bacterial amino acid biosynthesis pathways with high-throughput genetics. *PLoS Genet* 14:e1007147. <https://doi.org/10.1371/journal.pgen.1007147>.
81. Hoagland D, Arnon D. 1950. The water-culture method for growing plants without soil. University of California Agricultural Experiment Station, Berkeley, CA.
82. Griffiths JS, Long SR. 2008. A symbiotic mutant of *Sinorhizobium meliloti* reveals a novel genetic pathway involving succinoglycan biosynthetic functions. *Mol Microbiol* 67:1292–1306. <https://doi.org/10.1111/j.1365-2958.2008.06123.x>.
83. Thoma S, Schobert M. 2009. An improved *Escherichia coli* donor strain for diparental mating. *FEMS Microbiol Lett* 294:127–132. <https://doi.org/10.1111/j.1574-6968.2009.01556.x>.
84. Simon R, Priefer U, Pühler A. 1983. A broad-host-range mobilization system for *in vivo* genetic engineering: transposon mutagenesis in Gram-negative bacteria. *Nat Biotechnol* 1:784–791. <https://doi.org/10.1038/nbt1183-784>.