UC Irvine

UC Irvine Previously Published Works

Title

Carbon-13 variation with depth in soils of Brazil and climate change during the Quaternary.

Permalink

<https://escholarship.org/uc/item/1783k514>

Journal

Oecologia, 106(3)

ISSN

0029-8549

Authors

Martinelli, IA
Pessenda, LCR
Espinoza, E
et al.

Publication Date

1996-05-01

DOI

10.1007/bf00334565

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

L.A. Martinelli · L.C.R. Pessenda · E. Espinoza
P.B. Camargo · E.C. Telles · C.C. Cerri · R.L. Victoria
R. Aravena · J. Richey · S. Trumbore

Carbon-13 variation with depth in soils of Brazil and climate change during the Quaternary

Received: 23 December 1994 / Accepted: 18 November 1995

Abstract Paleocological and geomorphological studies indicate that, during the middle Holocene, there was a predominance of drier conditions with grassy savannahs replacing forests across the South American continent. Modern savannahs are composed mainly of C4 plants and soils developed under this type of vegetation show enrichment in ^{13}C compared to soils under C3 vegetation cover. If soils contain stabilized organic matter formed in the middle Holocene, we hypothesize that former C4 vegetation would be evidenced by a large enrichment of ^{13}C in soil organic matter (SOM). We investigate this possibility examining the depth variation of carbon isotopic composition in 21 soil profiles collected by different researchers at 14 different sites in Brazil. Of these, profiles from only three sites showed a marked increase of ^{13}C with depth (9–10‰ enrichment in $\delta^{13}\text{C}$ difference between the surface soil and deepest depth); two sites showed intermediate enrichment (4–5‰), and nine sites showed a small enrichment of approximately 2.5‰. The majority of sites showing all-C3 derived SOM were in the Amazon region. Possible causes for the absence of a large ^{13}C enrichment with depth are: (1) dominance of C3 rather than C4 grasses in mid-Holocene savannahs, (2) soil profiles did not preserve organic matter derived from mid-Holocene plants, (3) the retreat of forest areas did not occur on a regional scale, but was a much more localized phenomenon.

Key words Amazon · Vegetation change · Stable carbon isotope · Radiocarbon · Soil organic matter

Introduction

Soil organic matter (SOM) is a complex mixture of compounds from different pools, ranging from very labile compounds with very fast cycling times to refractory components that accumulate over millenia (Trumbore 1993). These attributes are susceptible to climate change, which can cause alterations in organic matter inputs and rates of decomposition (Jenkinson et al. 1991). The cumulative response of SOM behavior to past and future changes is a significant factor in the global carbon cycle (Prentice and Fung 1990; Tans et al. 1990; Jenkinson et al. 1991).

Paleocological and geomorphological studies suggest the occurrence of severe climatic changes in the South American continent. It has been hypothesized that there were drier periods during the Pleistocene and Holocene than the present, when the tropical forest was replaced by savannah-like vegetation, with predominance of grasses (Van der Hammen 1974; Absy and Van der Hammen 1976; Absy 1980; Ab'Saber 1982; Bigarella and Andrade-Lima 1982; Leyden 1985; Markgraf 1989; Bush et al. 1990; Bush and Colinvaux 1990; Markgraf 1991; Absy et al. 1991). The maximum in the proportion of grass pollen was found in the middle Holocene from ca. 6,000 to 4,000 years before present (BP) for several places in South America (Absy 1980; Servant et al. 1989; Markgraf 1989; Absy et al. 1991; Ledru 1992, 1993; Servant et al. 1993), from 10,500 to 10,400 years BP for central Brazil (Servant et al. 1989, 1993; Ledru 1993), and ca. 13,000 years BP for eastern Amazonia (Absy et al. 1991).

If these grasses were of the C4 type, it may be possible to find a residual isotopic signal preserved in soil profiles. Depending on the degree of ^{13}C enrichment in SOM with depth, two basic processes could explain the observed trends

L.A. Martinelli (✉) · L.C.R. Pessenda · E. Espinoza
P.B. Camargo · E.C. Telles · C.C. Cerri · R.L. Victoria
Centro de Energia Nuclear na Agricultura, Av. Centenário 303,
CEP 13416-000, Piracicaba, SP, Brazil
Fax: 55-194-228339

R. Aravena
Waterloo Center for Groundwater Research,
University of Waterloo, Waterloo, Ontario N2L3G1, Canada

J. Richey
School of Oceanography, WB-10,
University of Washington, Seattle, WA 98195, USA

S. Trumbore
Department of Earth System Science,
University of California, Irvine, CA 92717, USA

1. If the ^{13}C enrichment with depth is small, the decomposition of organic matter, which favors ^{12}C , is most likely the cause of the trend.

2. If the ^{13}C enrichment with depth is large, it is a stronger indication that the signal is due to the previous existence of ^{13}C -enriched vegetation, probably C4 grasses.

We explore those possibilities through the analyses of the stable carbon isotope composition of 21 soil profiles collected at 14 sites in Brazil, encompassing different soil types and climatic conditions (Volkoff et al. 1982; Desjardins et al. 1991; Valencia 1993; Pessenda et al. 1995; Trumbore et al. 1995). Radiocarbon data for the organic matter from the same soil profiles provide an estimate of the minimum age of stabilized C4 organic matter.

Methods

Locations of soil profiles are shown in Fig. 1. A total of 21 soil profiles at 14 sites, encompassing different climate regimes and soil types were collected (Table 1). Of these sites 7 were located in

the Amazon basin, which is characterized by high temperature and heavy rainfall (Fig. 1). Three other profiles were also collected in areas of high average temperatures: Juacema, Nhecolândia and Salitre. The first one is located in the northeast region of Brazil, which is mainly characterized by low rainfall (Fig. 1). The Nhecolândia profile was collected in the Brazilian Pantanal, one of the largest floodplains in the world, and the Salitre profile was collected in the southernmost region of Brazil (Fig. 1). In areas of lower average temperatures, six profiles in four different places (São Roque, Piracicaba, Londrina and Tunas) were collected (Fig. 1). Soil types differed among places (Table 1). The most common soil types in Brazil, *Latossolo* and *Podzólico*, which are equivalent to ultisols and oxisols in the American classification system, were present at most sampling sites (Table 1). Soils classified as *Terra Roxa Estruturada* (equivalent to alfisols) were present at two sites, and finally *Cambissolo* and *Areia Quartzosa*, both equivalent to inceptisols, were present at one site each. In all sampling sites the vegetation was the primary forest characteristic of each region.

Details of soil sampling and soil characteristics can be found elsewhere (Cerri 1979; Volkoff et al. 1982; Soubiès and Chauvel 1985; Martins et al. 1991; Higa 1989; Rocha 1990; Valencia 1993; Pessenda et al. 1996; Trumbore et al. 1995). Analysis of carbon isotopic composition, expressed as $\delta^{13}\text{C}$, were not always carried out by the same authors that collected soil samples. (A list of those authors can be found in Table 1.) Radiocarbon ages, expressed as years before present (years BP), were available for profiles located at Piracicaba, Londrina, Altamira, Salitre, Nhecolândia and Para-

Fig. 1 Upper panel map of Brazil showing the geographical location of soil profiles. Stars indicate sites where the variation with depth of $\Delta^{13}\text{C}$ values of soil organic matter clearly indicates the existence of previous C4 vegetation. Circles indicate sites with no trends. Squares indicate sites with equivocal $\Delta^{13}\text{C}$ trends. Lower panel annual average temperature and average rainfall of sample sites

Table 1 Sites of soil sampling (*numbers in parentheses indicate the number of profiles collected at each site*) following by soil type according to the Brazilian classification system; soil texture; mean annual temperature, rainfall and climate type

Code – Site	Brazilian classification	Texture	Mean annual temperature	Mean annual rainfall	Climate	References
Man – Manaus (1)	Latossolo Amarelo	Clay	27° C	2100 mm	Trop. humid	a
Alt – Altamira (2)	Terra Roxa Estruturada	Clay	27° C	1650 mm	Trop. humid	b, c, d
Cpo-Capitão Poço (1)	Latossolo Podzolizado	Silty-clay	27° C	2500 mm	Trop. humid	e, c
Par – Paragominas (3)	Latossolo Amarelo	Clay	27° C	1750 mm	Trop. humid	f
Ori – Oriente Novo (1)	Latossolo Vermelho-Amarelo	Silty-clay	25° C	2200 mm	Trop. humid	g, c
Dom – S. Domingos (1)	Podzólico Vermelho-Amarelo	Silty-clay	25° C	2200 mm	Trop. humid	g, c
Ten – Terra Nova (1)	Cambissolo	Silty-clay	25° C	2300 mm	Trop. humid	h, c
Jua-Juacema (1)	Latossolo	Silty-clay	25° C	500–1000 mm	Trop. semi-arid	g, c
Pan – Nhecolândia (1)	Areia Quartzosa	Sandy	25° C	1280 mm	Trop. humid	i
Sal – Salitre (2)	Latossolo Vermelho-Amarelo	Clay	24° C	1700 mm	Subtropical	j
Sro – São Roque (1)	Podzólico	Sandy-clay	21° C	1200 mm	Subtropical	g, c
Pir – Piracicaba (2)	Latossolo Vermelho-Amarelo	Clay	20° C	1200 mm	Suptropical	d, g
Lon – Londrina (2)	Terra Roxa Estruturada	Clay	19° C	1250 mm	Suptropical	d, g
Tun – Tunas (1)	Latossolo Podzolizado	Silty-clay	18° C	1400 mm	Temperate	g, c

^a Volkoff et al. (1982) soil description and $\delta^{13}\text{C}$ analysis

^b Rocha (1990) soil description

^c Desjardins et al. (1991) $\delta^{13}\text{C}$ analysis

^d Valencia (1993) $\delta^{13}\text{C}$ analysis

^e Martins et al. (1986) soil description

^f Trumbore et al. (1995) soil description and $\delta^{13}\text{C}$ analysis

^g Cerri (1979) soil description

^h Higa (1989) soil description

ⁱ Victoria et al. (1995) soil description and $\delta^{13}\text{C}$ analysis

^j Pessenda et al. (1996) soil description and $\delta^{13}\text{C}$ analysis

gominas and these analyses were conducted by Valencia (1993) for the first three profiles, and by Pessenda et al. (1996), Victoria et al. (1995), and Trumbore et al. (1995) for the three last profiles, respectively.

Results and discussion

The contemporary vegetation cover of all sites was primary forest (C3 plant type), and the $\delta^{13}\text{C}$ of the surface soil organic matter varies from -28.5 to 26.0‰ . In order to make the comparison among different profiles easier, the depth variability of the carbon isotopic composition in each profile was expressed as the difference of the $\delta^{13}\text{C}$ of a soil depth in relation to the $\delta^{13}\text{C}$ value of the most superficial sampling depth. We call this difference $\Delta^{13}\text{C}$, defined as:

$$\Delta^{13}\text{C} = \delta^{13}\text{C}_{\text{depth}} - \delta^{13}\text{C}_{\text{surface}}$$

where $\delta^{13}\text{C}_{\text{surface}}$ is the value of the surface soil and $\delta^{13}\text{C}_{\text{depth}}$ is the value at a given depth. $\Delta^{13}\text{C}$ values were mostly positive, reflecting the fact that $\delta^{13}\text{C}$ values increased with depth in nearly all soil profiles.

Results were grouped according to the $\Delta^{13}\text{C}$ values. Figure 2A and B shows profiles with the smallest $\Delta^{13}\text{C}$ values. The profiles with the highest $\Delta^{13}\text{C}$ are shown in Fig. 2D and profiles with intermediate to small values of $\Delta^{13}\text{C}$ are grouped in Fig. 2C.

Most of the soil profiles with the smallest $\Delta^{13}\text{C}$ values (Fig. 2A and 2B) are from the Amazon Basin (Fig. 1). In these profiles the highest $\Delta^{13}\text{C}$ value was 3.5‰ , but most of the values were smaller than 2.5‰ , which indicates that the major cause of ^{13}C enrichment with depth was probably fractionation during decomposition of SOM. The only exception to this pattern was the depth variability observed in the Terra Nova profile, where the $\Delta^{13}\text{C}$

value reached almost 8.5‰ near the bottom (Fig. 2D). Profiles from Piracicaba and Londrina showed similar ^{13}C enrichment (Fig. 1). The profile Lon-1 at Londrina reached a $\Delta^{13}\text{C}$ value of 11.5‰ at the bottom (Fig. 2D). In such cases the most likely cause for this large ^{13}C enrichment with depth is the existence of prior C4 vegetation (Dzurec et al. 1985; Mondenesi et al. 1986; Schwartz et al. 1987; Volkoff and Cerri 1987; Martin et al. 1990; Desjardins et al. 1991; McPherson et al. 1993; Wang et al. 1993; Mariotti and Peterschmitt 1994; Victoria et al. 1995). Both profiles of Salitre, Nhecolândia, and Tunas had intermediate $\Delta^{13}\text{C}$ values varying approximately from 4.0 to 6.0‰ (Fig. 2C). The Nhecolândia profile came from the Brazilian Pantanal, a large floodplain, highly dynamic in terms of geomorphology (Adamoli 1982), with a landscape that alternates high land forested areas with low land areas of savannah. This dynamism is reflected in the $\Delta^{13}\text{C}$ values, with reach a maximum of almost 6‰ at 100 cm depth, suggesting the replacement of the forest by a C4 savannah, and a decrease to almost 0.5‰ at 170 cm depth, indicating the dominance of a C3 vegetation type again (Victoria et al. 1995). The profiles from Tunas and Salitre (Sal-1 and Sal-2) show $\Delta^{13}\text{C}$ increasing from 4.0 to 6.0‰ , within the upper 30 cm. Below, $\Delta^{13}\text{C}$ values were constant to the bottom of the profiles. According to Pessenda et al. (1995) the cause of the increase of $\Delta^{13}\text{C}$ with depth at these sites is fractionation of isotopes during decomposition and not a past vegetation change. However, palynological evidence of a past vegetation change in favor of grasses that may have been C4 (Ledru 1993), and the $\Delta^{13}\text{C}$ value of 4.0 – 6.0‰ , suggest that paleovegetation change cannot be ruled out as an explanation.

Table 2 summarizes the radiocarbon ages of soil profiles for those where such data were available. The ^{14}C

Fig. 2 Variation of $\Delta^{13}\text{C}$ values in relation to soil depth: **A**, **B** sites with smallest $\Delta^{13}\text{C}$ values; **C** sites with intermediate $\Delta^{13}\text{C}$ values, and **D** sites with highest $\Delta^{13}\text{C}$ values. Site abbreviations are given in Table 1

ages reported represent the average "age" of a carbon atom in SOM, and certainly represent a mixture of both older and younger material. For samples from deep in the soil, the radiocarbon age may be taken as a minimum age for stabilized organic matter. For instance, although the ^{14}C age of organic matter at 100 cm depth in Paragominas soils is about 14,000 years BP, it was estimated by Trumbore et al. (1995) that this average age represents a mixture of 10% modern carbon with 90% radiocarbon-free carbon (i.e., 90% with average age >40,000 years BP).

One of the most widely accepted climatic changes in the past is the drought that occurred in the middle Holocene (Servant et al. 1993). Pollen records from Salitre (Ledru 1993) and Southern Serra dos Carajás, near Altamira (Absy et al. 1991) show sharp increase in grass species. The Piracicaba, Londrina and Terra Nova profiles clearly show a sharp increase in $\Delta^{13}\text{C}$ (Fig. 2C). The deepest dated depths (170–180 cm) in the Piracicaba and Lond-

rina profiles reach ages of ca. 3600 years BP and ca. 9300 years BP. These soils were presumably subjected to paleoclimatic and paleovegetation changes associated with the mid-Holocene. In Salitre the deepest dated depth (190–200 cm) is almost 7000 years BP, which means that these profiles also experienced the middle Holocene drought. However, the increase of $\Delta^{13}\text{C}$ with depth was not as sharp as in the Piracicaba and Londrina profiles, making it difficult to establish the cause for this increase. No other profiles showed the effects of the middle Holocene dryness in their carbon isotopic composition. In addition, seven other profiles collected in several areas of the Amazon region down to 4 m depth show no significant variation in carbon stable isotopic composition (Sanaiotti, unpublished work). As modern savannah soils clearly show the presence of C_4 grasses in their SOM (Dzurec et al. 1985; Mondenesi et al. 1982; Schwartz et al. 1987; Volkoff and Cerri 1987; Martin et al. 1990; Desjardins et al. 1991; McPherson et al. 1993; Wang et al. 1993; Mariotti and Peterschmitt 1994), the apparent absence of C_4 signal in majority of the profiles is intriguing.

There are several hypothesis that can explain the lack of a C_4 signal.

Table 2 Radiocarbon ages (years BP) of soil organic matter along some soil profiles

Depth (cm)	Altamira ^a Alt-2	Paragominas ^b Par-1	Nhecolândia ^c	Salitre ^d Sal-2	Piracicaba ^a Pir-2	Londrina ^a Lon-2
0–10	modern	modern	modern	240	modern	modern
20–30		2400				
30–40			modern			
40–50	1440	4850			1230	820
70–80	2790		639		2680	1920
90–100	3640	14190		3700	3030	2390
110–120			1483			
120–130	4800				3260	5450
150–160				5550		
160–170			2002			
170–180	4390				3640	9340
190–200				6940		
290–300		22470				

^a Valencia (1993); ^b Trumbore et al. (1995); ^c Victoria et al. (1995); ^d Boulet et al. (1995)

1. The lack of signal would be consistent with dominance by C3 rather than C4 grasses. The same hypothesis was raised by Guillet et al. (1988), who found a similar situation in the tropical region of Colombia. However, judging by the modern savannahs of the world, this is unlikely.

2. The retreat of forest areas was not on a regional scale, but was a much more localized phenomenon.

3. The C4 plants were not present long enough to leave their isotopic imprint on stabilized soil organic matter.

The refuge theory for the high degree of biodiversity in tropical South America suggests that the high diversity resulted from differential species evolution in forest patches that were isolated when forests were replaced with grassy savannahs during the Pleistocene and Holocene, and remained isolated until a new climatic change provoked the return of an intact continuous forest (Haffer 1969; Vanzolini 1970; Prance 1973, 1982; Brown 1974). If the findings of this study are confirmed in further analysis, it would be a strong indication that extensive areas of grass never existed in the Amazon, especially during the Holocene. In view of refuge theory this would have strong implications for interpretations of species evolution of the Amazon rain forest.

Acknowledgements We would like to thank J. Hedges, S. Wilhelm, M. McClain, and Eric Davidson for comments on the manuscript. Financial support was provided by Conselho Nacional de Pesquisa e Desenvolvimento and Fundação de Amparo à Pesquisa do Estado de São Paulo.

References

- Ab'Saber AN (1982) The paleoclimate and paleoecology of Brazilian Amazon. In: Prance GT (ed) Biological diversification in the tropics. Columbia University Press, New York, pp 41–59
- Absy ML (1980) Dados sobre as mudanças do clima e da vegetação da Amazônia durante o Quaternário. *Acta Amazonica* 10:929–932
- Absy ML, Van der Hammen T (1976) Some paleo-ecological data from Rondônia, southern part of Amazonian Basin. *Acta Amazonica* 6:293–299
- Absy ML, Cleef A, Fournier M, Martin L, Servant M, Sifeddine A, Ferreira da Silva M, Soubies F, Suguio K, Turcq B, Van der Hammen T (1991) Mise en évidence de quatre phases d'ouverture de la forêt dense dans le sud-est de l'Amazonie au cours des 60000 dernières années. Première comparaison avec d'autres régions tropicales. *CR Acad Sci Paris Sér II* 312:673–678
- Adámoli JM (1982) O Pantanal e suas relações fitogeográficas com os cerrados. Discussão sobre o conceito "Complexo Pantanal". In: Proceedings of Congresso Brasileiro de Botânica. Sociedade Brasileira de Botânica, Teresina, Brazil, pp 109–119
- Bigarella JJ, Andrade-Lima D (1982) Paleoenviromental changes in Brazil. In: Prance GT (ed) Biological diversification in the tropics. Columbia University Press, New York, pp 27–40
- Boulet R, Pessenda LCR, Telles ECC, Melfi AJ (1995) Une évaluation de la vitesse de l'accumulation superficielle de matière par la faune du sol à partir de la datation des charbons et de l'humine du sol. Exemple des latosols des versants du lac Campestre, Salitre, Minas Gerais, Brésil. *CR Acad Sci Paris Ser IIa* 320:287–294
- Brown KS Jr (1974) Quaternary refugia in tropical America: evidence for race formation in *Heliconius* butterflies. *Proc R Soc Lond B* 187:369–378
- Bush M, Colinvaux PA (1990) A pollen record of a complete glacial cycle from lowland Panama. *J Veg Sci* 1:105–118
- Bush MB, Colinvaux PA, Wiemann C, Piperno DR, Liu KB (1990) Late Pleistocene temperature depression and vegetation change in Ecuadorian Amazonia. *Quat Res* 34:330–345
- Cerri C (1979) Alteração e pedogênese em áreas graníticas situadas sob diferentes condições bioclimáticas encontradas no Brasil. Master Thesis, University of São Paulo
- Desjardins Th, Volkoff B, Andreux F, Cerri C (1991) Distribution du carbone total et de l'isotope ¹³C dans des sols ferrallitiques du Brésil. *Sci Sol* 29:175–187
- Dzurec RS, Boutton TW, Caldwell MM, Smith BN (1985) Carbon isotope ratios of soil organic matter and their use in assessing community composition changes in Curlew Valley, Utah. *Oecologia* 66:17–24
- Guillet B, Faivre P, Mariotti A, Khobzi J (1988) The ¹⁴C dates and ¹³C/¹²C ratios of soil organic matter as a means of studying the past vegetation in intertropical regions: examples from Colombia (South America). *Palaeogeogr Palaeoclim Palaeoecol* 65:51–58
- Haffer J (1969) Speciation in Amazonian forest birds. *Science* 165:131–137
- Higa NT (1989) Caracterização das fases sólida e líquida de solos e arenas derivados do granito sob floresta tropical Amazônica (Terra Nova do Norte – MT). Ph D Thesis, University of São Paulo
- Jenkinson DS, Adams DE, Wild A (1991) Model estimates of CO₂ emissions from soil in response to global warming. *Nature* 351:304–306

- Ledru MP (1992) Modifications de la végétation du Brésil Central entre la dernière époque glaciaire et l'interglaciaire actuel. *CR Acad Sci Paris Sér II* 314:117–123
- Ledru MP (1993) Late Quaternary environmental and climatic changes in central Brazil. *Quat Res* 39:90–98
- Leyden BW (1985) Late Quaternary aridity and holocene moisture fluctuations in the Lake Valencia basin, Venezuela. *Ecology* 66:1279–1295
- Mariotti A, Petershmitt E (1994) Forest savanna ecotone dynamics in India as revealed by carbon isotope ratios of soil organic matter. *Oecologia* 97:475–480
- Markgraf V (1989) Paleoclimates in Central and South America since 18,000 BP based on pollen and lake-level records. *Quat Sci Rev* 8:1–24
- Markgraf V (1991) Younger Dryas in southern South America? *Boreas* 20:63–69
- Martin A, Mariotti A, Balesdent J, Lavelle P, Vuattoux V (1990) Estimate of organic matter turnover rate in savanna soil by ^{13}C natural abundance measurements. *Soil Biol Biochem* 22:517–523
- Martins PFS, Cerri CC, Volkoff B, Andreux F, Chauvel A (1991) Consequences of clearing and tillage on the soil of a natural Amazonian ecosystem. *For Ecol Manage* 38:273–282
- McPherson GR, Boutton TW, Midwood AJ (1993) Stable carbon isotope analysis of soil organic matter illustrates vegetation change at the grassland/woodland boundary in southeastern Arizona, USA. *Oecologia* 93:95–101
- Mondenesi MC, Matsui E, Volkoff B (1986) Relação $^{13}\text{C}/^{12}\text{C}$ nos horizontes húmiferos superficiais e nos horizontes escuros profundos dos solos de campo e mata da região de Campos do Jordão, São Paulo, Brasil. In: Regional colloquium on soil organic matter studies. Centro de Energia Nuclear na Agricultura, Piracicaba, Brazil, pp 155–160
- Pessenda LCR, Aravena R, Melfi AJ, Telles ECC, Boulet R, Valencia EPE, Tomazello M (1995) The use of carbon isotopes (^{13}C , ^{14}C) in soil to evaluate vegetation changes during the Holocene in Central Brazil. *Radiocarbon* 38
- Prance GT (1973) Phytogeographic support for the theory of Pleistocene forest refuges in the Amazon Basin, based on evidence from distribution patterns in Caryocaraceae, Chrysolalanaceae, Dichapetalaceae and Lecythidaceae. *Acta Amazonica* 3:5–28
- Prance GT (1982) Forest refuges: evidence from woody angiosperms. In: Prance GT (ed) Biological diversification in the tropics. Columbia University Press, New York, pp 137–158
- Prentice KC, Fung IY (1990) The sensitivity of terrestrial carbon storage to climate change. *Nature* 346:48–51
- Rocha GC (1990) Características e dinâmica de coberturas pedológicas sobre rochas básicas nas regiões norte e sul do Brasil. Ph D Thesis, University of São Paulo
- Schwartz D, Mariotti A, Lanfranchi R, Guillet B (1987) $^{13}\text{C}/^{12}\text{C}$ ratios of soil organic matter as indicator of vegetation changes in the Congo. *Geoderma* 39:97–103
- Servant M, Fournier M, Soubiès F, Suguio K, Turcq B (1989) Sécheresse holocène au Brésil (18–20° latitude Sud). Implications paléométéorologiques. *CR Acad Sci Paris Sér II* 309:153–156
- Servant M, Maley J, Turcq B, Absy ML, Brenac P, Fournier M, Ledru MP (1993) Tropical forest changes during the Late Quaternary in African and South American lowlands. *Paleogeogr Paleoclimatol Palaeoecol* 7:1–16
- Tans PT, Fung IY, Takahashi T (1990) Observational constraints on the global atmospheric CO_2 budget. *Science* 247:1431–1438
- Trumbore S (1993) Comparison of carbon dynamics in tropical and temperate soils using radiocarbon measurements. *Global Biogeochem Cycles* 7:275–290
- Trumbore SE, Davidson EA, Camargo PB, Nepstad DC, Martinelli LA (1995) Below-ground cycling of carbon in forests and pastures of eastern Amazonia. *Global Biogeochem Cycles* 9:515–528
- Valencia EPE (1993) Datação por ^{14}C e razão $^{13}\text{C}/^{12}\text{C}$ de solos sob climas tropical e subtropical do Brasil. Master Thesis, University of São Paulo
- Van der Hammen T (1974) The Pleistocene changes of vegetation and climate in tropical South America. *J Biogeogr* 1:3–26
- Vanzolini PE (1970) Zoologia sistemática geografia e a origem das espécies. *Inst Geogr São Paulo* 3:1–56
- Victoria RL, Fernandez FA, Martinelli LA, Piccolo MC, Camargo PB, Trumbore S (1995) Past vegetation changes in the Brazilian Pantanal arboreal-grassy savannah ecotone by using carbon isotopes in the soil organic matter. *Global Change Biol* 1:165–171
- Volkoff B, Cerri CC (1987) Carbon isotopic fractionation in subtropical Brazilian grassland soils. Comparison with tropical forest soil. *Plant Soil* 102:27–31
- Volkoff B, Matsui E, Cerri CC (1982) Discriminação isotópica do carbono nos húmicos de latossolos e podzol na região amazônica do Brasil. In: Colóquio regional sobre matéria orgânica do solo. Piracicaba, pp 147–153
- Wang Y, Cerling TE, Effand WR (1993) Stable isotope ratios of soil carbonate and soil organic matter as indicators of forest invasion of prairie near Ames, Iowa. *Oecologia* 95:365–369