

UC Davis

UC Davis Previously Published Works

Title

Contribution of the neighborhood environment and obesity to breast cancer survival: the California Breast Cancer Survivorship Consortium.

Permalink

<https://escholarship.org/uc/item/1833d62c>

Journal

Cancer Epidemiology, Biomarkers and Prevention, 24(8)

Authors

Cheng, Iona
Shariff-Marco, Salma
Koo, Jocelyn
et al.

Publication Date

2015-08-01

DOI

10.1158/1055-9965.EPI-15-0055

Peer reviewed



Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2015 August ; 24(8): 1282–1290. doi:
10.1158/1055-9965.EPI-15-0055.

Contribution of the Neighborhood Environment and Obesity to Breast Cancer Survival: The California Breast Cancer Survivorship Consortium

Iona Cheng^{1,2}, Salma Shariff-Marco^{1,2,3}, Jocelyn Koo¹, Kristine R. Monroe⁴, Juan Yang¹, Esther M. John^{1,2,3}, Allison W. Kurian^{2,3}, Marilyn L. Kwan⁵, Brian E. Henderson⁴, Leslie Bernstein⁶, Yani Lu⁶, Richard Sposto⁷, Cheryl Vigen⁴, Anna H. Wu⁴, Scarlett Lin Gomez^{1,2,3}, and Theresa H.M. Keegan^{1,2,3}

¹Cancer Prevention Institute of California, Fremont, CA

²Stanford Cancer Institute, Stanford, CA

³Stanford University School of Medicine, Stanford, CA

⁴University of Southern California, Keck School of Medicine, Los Angeles, CA

⁵Division of Research, Kaiser Permanente Northern California, Oakland, CA

⁶City of Hope, Duarte, CA

⁷University of Southern California, Children's Hospital, Los Angeles, CA

Abstract

Little is known about neighborhood attributes that may influence opportunities for healthy eating and physical activity in relation to breast cancer mortality. We used data from the California Breast Cancer Survivorship Consortium and the California Neighborhoods Data System to examine the neighborhood environment, body mass index, and mortality after breast cancer. We studied 8,995 African American, Asian American, Latina, and non-Latina White women with breast cancer. Residential addresses were linked to the CNDS to characterize neighborhoods. We used multinomial logistic regression to evaluate the associations between neighborhood factors and obesity, and Cox proportional hazards regression to examine associations between neighborhood factors and mortality. For Latinas, obesity was associated with more neighborhood crowding (Quartile 4 (Q4) vs. Q1: Odds Ratio (OR)=3.24; 95% Confidence Interval (CI): 1.50-7.00); breast cancer-specific mortality was inversely associated with neighborhood businesses (Q4 vs. Q1: Hazard Ratio (HR)=0.46; 95% CI: 0.25-0.85) and positively associated with multi-family housing (Q3 vs. Q1: HR=1.98; 95% CI: 1.20-3.26). For non-Latina Whites, lower neighborhood socioeconomic status (SES) was associated with obesity (Quintile 1 (Q1) vs. Q5: OR=2.52; 95% CI: 1.31-4.84), breast cancer-specific (Q1 vs. Q5: HR=2.75; 95% CI: 1.47-5.12), and all-cause (Q1 vs. Q5: HR=1.75; 95% CI: 1.17-2.62) mortality. For Asian Americans, no associations were seen. For African Americans, lower neighborhood SES was

Corresponding author: Iona Cheng, Cancer Prevention Institute of California, 2201 Walnut Ave., Suite 300, Fremont, CA 94538; phone: 510-608-5055; fax: 510-608-5085; iona.cheng@cpic.org.

Conflicts of Interest: None

associated with lower mortality in a nonlinear fashion. Attributes of the neighborhood environment were associated with obesity and mortality following breast cancer diagnosis, but these associations differed across racial/ethnic groups.

Keywords

California Breast Cancer Survivorship Consortium; Neighborhood Environment; Body Mass Index; Survival; Mortality

Introduction

The obesity epidemic in the United States is a serious health priority for cancer care as an increasing number of cancer patients are obese at diagnosis, and numerous studies among Whites have demonstrated a higher mortality among obese, compared to normal weight, breast cancer patients (1, 2). In a meta-analysis of over 213,000 women with breast cancer, those who were obese (body mass index (BMI) >30 kg/m²) or overweight (BMI 25- <30 kg/m²) were at increased risk of all-cause mortality, regardless of when BMI was ascertained (i.e. before or after diagnosis) (2). Within our racially/ethnically diverse California Breast Cancer Survivorship Consortium (CBCSC), we have demonstrated increased risks of all-cause and breast cancer-specific mortality among morbidly obese (BMI > 40 kg/m²) non-Latina Whites and Latinas in comparison to normal weight women (1).

Interest in the relation between the neighborhood environment—social and man-made (“built”) physical attributes of an individual's surroundings (3, 4)—and levels of obesity is growing, as these attributes provide opportunities and/or barriers for healthy eating and physical activity, and may influence health outcomes. By using data on the neighborhood environment from the California Neighborhoods Data System (CNDS) (3) and building on our prior work in the CBCSC (1), we investigated the associations of the neighborhood environment with pre-diagnostic BMI in cross-sectional analyses and breast cancer-specific and all-cause mortality in prospective analyses among a racial/ethnically diverse cohort of breast cancer cases.

Materials and Methods

Study Participants

The CBCSC is comprised of six California-based epidemiologic studies of breast cancer etiology/prognosis (5). For this analysis, five studies contributed data, including three case-control studies: the Asian American Breast Cancer Study (AABCS) (6), Women's Contraceptive and Reproductive Experiences study (CARE) (7), and San Francisco Bay Area Breast Cancer Study (SFBCS) (8, 9); and two cohort studies: the California Teachers Study (CTS) (10) and Multiethnic Cohort (MEC) (11). Each study collected cases' data on reproductive, lifestyle, sociodemographic, and other breast cancer risk or prognostic factors, which were harmonized according to common definitions (5). Pre-diagnosis BMI was ascertained closest to the date of breast cancer diagnosis in order to best coincide with the

characterization of the neighborhood environment at the time of diagnosis. Clinicopathologic and treatment factors were obtained from the California Cancer Registry (5). Institutional Review Board approval was received from all participating institutions and from the California Protection for Human Subjects state institutional review board.

We excluded study participants with prior cancer diagnoses ($n=779$), in situ histology ($n=22$), follow-up time < 30 days ($n=19$), incomplete address ($n=240$), and those who were underweight ($BMI < 18.5 \text{ kg/m}^2$; $n=183$) or were missing BMI ($n=283$), leaving 8,995 breast cancer cases for analysis. Vital status and cause of death were ascertained from the California Cancer Registry as of December 31, 2010. Over a median follow-up time of 10.3 years, 1,284 women died of breast cancer among 2,426 total deaths.

California Neighborhoods Data System

Residential addresses at the time of breast cancer diagnosis were geocoded to latitude and longitude coordinates and linked census and business data of the California Neighborhoods Data Systems (3). Addresses were assigned to 1990 Census block groups (diagnoses 1994-1995) and 2000 Census block groups (diagnoses 1996-2007) to ascertain neighborhood levels of SES (created by principal component analysis of census and American Community Survey data on education, housing, employment, occupation, income, and poverty (12, 13)); population density; urbanicity, commute patterns; household crowding (i.e. housing with >1 occupant per room); proportion of multi-family housing units (i.e. housing structures with 2 or more units, apartment complexes); and were categorized into levels according to the state distribution (Supplemental Tables 1 and 2). Geocodes were also linked to business data to quantify neighborhood attributes of the retail/restaurant food environment; parks; recreational facilities; street connectivity(14) (i.e., gamma index, defined as the ratio of actual number of street segments to maximum possible number of intersections and expressed as the percentage of connectivity); and total businesses within a one mile pedestrian network distance of participant's residence, reflecting a reasonable distance to walk to a destination. Specifically, information on number of businesses was based on business listings derived from Walls & Associates' National Establishment Time-Series Database from 1990-2008 (15). Traffic density using previously described methods (16) was based on traffic counts from the California Department of Transportation (2004) (17) that were within a residential buffer area of a 500 meter radius based on the assumption that traffic close to a subject's residence influences walking/physical activity behaviors. These neighborhood business and traffic-related attributes were categorized according to the study participant distribution (Supplemental Tables 1 and 2). Study methods of these neighborhood data have been described previously (3, 18, 19). The Census block group (an area of approximately 1,500 residents) was considered our neighborhood unit.

Statistical Analysis

For cross-sectional analysis of the relationship between neighborhood factors and pre-diagnostic BMI, multivariate multinomial regression was conducted to estimate odds ratios (OR) of being overweight ($BMI=25-29.9$) or obese ($BMI= 30$) versus normal weight ($BMI=18.5-24.9$). All multinomial models were stratified on stage and study, and included all neighborhood variables and adjusted for variables listed in Table 1, which showed

significant associations with BMI in unadjusted models. For prospective mortality analyses, multivariable Cox proportional hazard regressions were conducted to estimate hazard ratios (HR) of breast cancer-specific and all-cause mortality. All Cox models included all neighborhood factors and were stratified on stage and study, and adjusted for variables listed in Tables 2 and 3, which showed significant univariate associations with BMI and/or breast cancer-specific and overall mortality, respectively. All models were adjusted for clustering within block groups by applying the sandwich estimator of the covariance structure, which has been shown to account for intracluster dependence and has yielded robust standard error estimates even under model misspecification (20). Multicollinearity in our models was assessed by examining variation inflation factors (VIF). All models met our criteria of non-multicollinearity with $VIF < 10$. All P values presented are two-sided. A P value threshold < 0.05 was used to determine statistical significance and no correction was applied for multiple hypothesis testing. Analyses were conducted using SAS (version 9.3, Cary, NC).

Results

Of the 8,995 breast cancer cases in the CBCSC, 47% were non-Latina White, 20% Latina, 19% African American, and 14% Asian American (Supplemental Table 3). The majority had Stage I (49%) or II (40%), 55% had estrogen receptor (ER+) or progesterone (PR+) positive tumors, 56% had breast conserving surgery, 40% received chemotherapy, and 51% received radiation treatment (Supplemental Table 4). Approximately 27% lived in low SES neighborhoods, 60% lived in suburban neighborhoods, and 21% lived in neighborhoods with >3 parks (Supplemental Table 1).

Overall, living in low versus high SES neighborhoods was associated with higher odds of being overweight (p trend < 0.01) or obese (p trend = 0.02) (Table 1). Significant SES-BMI associations were seen only among non-Latina Whites, although similar patterns were observed in African Americans. Among all breast cancer cases, living in high versus low household crowding (housing with >1 occupant per room) was associated with an increased odds of obesity (p trend=0.02). Latinas demonstrated the strongest association between obesity and household crowding (p trend < 0.01), with those living in neighborhoods in the highest versus lowest quartile of household crowding having a 3-fold higher odds of obesity (95% CI: 1.50-7.00). In addition, Latinas living in neighborhoods at the highest versus lowest quartile of street connectivity had an increased odds of obesity (OR=1.77; 95% CI: 1.06-2.95). For non-Latina Whites, living in neighborhoods with a higher proportion of multi-family housing units was associated with a lower odds of being overweight (Q4 vs. Q1 OR=0.72; 95% CI: 0.54-0.95; p trend < 0.01). Living in streets with high connectivity versus low connectivity was associated with a significant increased odds of obesity (p trend=0.02) in African Americans but there were no other significant BMI-neighborhood associations. No BMI-neighborhood associations were observed among Asian Americans.

Among all breast cancer cases, pre-diagnostic BMI was not associated with breast cancer-specific mortality (Table 2) and was marginally associated with all-cause mortality (p trend=0.05) (Table 3). For Latinas, those who were morbidly obese (BMI > 40 kg/m²) were at increased risks of breast-cancer specific (HR=2.13; 95% CI: 1.10-4.15) and all-cause (HR=2.15; 95% CI: 1.31-3.53) mortality versus normal weight women. Neighborhood-

mortality associations were most notable among Latinas. Latinas living in neighborhoods with a high versus low proportion of multi-family housing units were at increased risks of breast cancer-specific and all-cause mortality. Latinas living in neighborhoods with a high versus low number of businesses had a lower risk of breast cancer-specific mortality (HR=0.46; 95% CI: 0.25-0.85), while those living in neighborhoods with >1 park were at greater risk of breast cancer-specific mortality versus those living in neighborhoods with no parks (p trend=0.03).

Neighborhood SES was associated with mortality among non-Latina Whites and African Americans, but in opposite directions (Tables 2 and 3). Non-Latina Whites living in low versus high SES neighborhoods were at increased risk of breast cancer-specific (Q1 vs. Q5: HR=2.75; 95% CI: 1.47-5.12; p trend< 0.01) and all-cause (Q1 vs. Q5: HR=1.75; 95% CI: 1.17-2.62; p trend=0.01) mortality. Conversely, African Americans living in SES neighborhoods (Q1 to Q4) had decreased risks of breast cancer-specific and all-cause mortality versus those living in the highest SES (Q5) neighborhood, but these relationships were not linear. Because of the differing proportions of non-Latina Whites and African Americans in the higher SES groups (Q4 & Q5=70.2% and 24.5%, respectively), we examined SES and mortality associations using race/ethnicity specific cut-points and found similar mortality associations between the lowest vs. highest levels of SES in comparison to using the state-wide cut-points (data not shown). For Asian Americans, no neighborhood-mortality associations were observed.

Discussion

Our central aim of this large consortium study was to examine breast cancer mortality in relation to obesity and specific attributes of the neighborhood environment potentially related to obesity across diverse racial/ethnic groups. In cross-sectional analysis, we identified that greater household crowding and more street connectivity (among Latinas), and low neighborhood SES and less multi-family housing (among non-Latina Whites) were important risk factors for obesity. In addition, low neighborhood SES (among non-Latina Whites) and high multi-family housing neighborhoods (among Latinas) were associated with higher mortality in a prospective analysis; and lower neighborhood SES (among African Americans) and greater number of businesses (among Latinas) were associated with lower mortality. To our knowledge, this is one of the first studies to evaluate a comprehensive suite of neighborhood attributes and their associations with breast cancer mortality across multiple racial/ethnic groups.

In a previous pooled analysis (18) of 4,345 breast cancer cases from the San Francisco Bay Area that included SFBCS participants (21, 22), lower neighborhood SES was associated with higher overall mortality. Our findings confirm the inverse association between SES and mortality reported by Keegan et al. (18) and others (23-28) that have largely focused on Whites and examined SES alone and no other neighborhood attributes. Furthermore, we identified heterogeneous effects by race/ethnicity for the associations of neighborhood SES with overall mortality (p interaction<0.01) as evidenced by the higher risk of mortality with increasing SES for non-Latina Whites and the lack of clear associations in other racial/ethnic groups. In addition, we did not observe an association between the number of neighborhood

using Census boundaries does allow us to efficiently examine a number of social and built environment factors across a large number of geographic units that would have been costly to obtain through other sources (e.g., self-report, neighborhood audits); moreover, it is plausible that the attributes of census boundaries may highly correlate with perceived neighborhoods (51). In addition, we were unable to account for neighborhood disorder, safety, and deterioration (52), factors that could influence the associations that we observed (e.g., higher odds of obesity among Latinas and African Americans residing in neighborhoods with more connected streets). We tested *a priori* selected neighborhood factors and because no validated cumulative index of street connectivity exists for California, we were unable to examine such an index, which that may better capture physical activity environments. Lastly, we did not adjust for multiple testing and recognize that some of our findings may be due to chance. Future research should incorporate these elements when evaluating factors underlying the neighborhood associations with obesity and mortality. Such insight is important for identifying interventions to improve survival outcomes for breast cancer patients across all racial/ethnic populations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial Support: This work was supported by the California Breast Cancer Research Program (CBCRP) (Grants 16ZB-8001 A.H. Wu, Authors: Spoto, Vigen; 16ZB-8002 S.L. Gomez, T.H. Keegan, S. Shariff-Marco J. Koo, J. Yang, A. W. Kurian, E. M. John; 16ZB-8003 L. Bernstein, Y. Lu; 16ZB-8004 M.L. Kwan; 16ZB-8005 K.R. Monroe, I. Cheng, B.E. Henderson). The Asian American Breast Cancer Study was supported by CBCRP grants 1RB-0287, 3PB-0120, and 5PB-0018 to A. H. Wu. The San Francisco Bay Area Breast Cancer Study was supported by National Cancer Institute grants R01 CA063446 and R01 CA077305; by the U.S. Department of Defense (DOD) grant DAMD17-96-1-6071; and by the CBCRP grants 4JB-1106 and 7PB-0068 to E. M. John. The Women's CARE Study was funded by the National Institute of Child Health and Human Development (NICHD), through a contract with USC (N01-HD-3-3175), and the California Teachers Study was funded by the California Breast Cancer Act of 1993; National Cancer Institute grants (R01 CA77398 and K05 CA136967) to L. Bernstein; and the California Breast Cancer Research Fund (contract 97-10500) to L. Bernstein. The Multiethnic Cohort Study was supported by National Cancer Institute grants R01 CA54281, R37CA54281, and UM1 CA164973 to B.E. Henderson, L.N. Kolonel, L. Le Marchand, L.R. Wilkens. The Life After Cancer Epidemiology Study is supported by National Cancer Institute grant R01 CA129059 to B. J. Caan. The collection of cancer incidence data used in this study was supported by the California Department of Health Services as part of the statewide cancer reporting program mandated by California Health and Safety Code Section 103885; the National Cancer Institute's Surveillance, Epidemiology, and End Results Program under contract HHSN261201000140C awarded to the Cancer Prevention Institute of California, contract HHSN26120100035C awarded to the University of Southern California, and contract HHSN26120100034C awarded to the Public Health Institute; and the Centers for Disease Control and Prevention's National Program of Cancer Registries, under agreement #1U58 DP000807-01 awarded to the Public Health Institute. The ideas and opinions expressed herein are those of the authors, and endorsement by the State of California, the California Department of Health Services, the National Cancer Institute, or the Centers for Disease Control and Prevention or their contractors and subcontractors is not intended nor should be inferred.

References

1. Kwan ML, John EM, Caan BJ, Lee VS, Bernstein L, Cheng I, et al. Obesity and mortality after breast cancer by race/ethnicity: The California Breast Cancer Survivorship Consortium. *Am J Epidemiol.* 2014; 179:95–111. [PubMed: 24107615]
2. Chan DS, Vieira AR, Aune D, Bandera EV, Greenwood DC, McTiernan A, et al. Body mass index and survival in women with breast cancer-systematic literature review and meta-analysis of 82 follow-up studies. *Ann Oncol.* 2014; 25:1901–14. [PubMed: 24769692]

3. Gomez SL, Glaser SL, McClure LA, Shema SJ, Kealey M, Keegan TH, et al. The California Neighborhoods Data System: a new resource for examining the impact of neighborhood characteristics on cancer incidence and outcomes in populations. *Cancer Causes Control*. 2011; 22:631–47. [PubMed: 21318584]
4. Brownson RC, Hoehner CM, Day K, Forsyth A, Sallis JF. Measuring the built environment for physical activity: state of the science. *Am J Prev Med*. 2009; 36:S99–123 e12. [PubMed: 19285216]
5. Wu AH, Gomez SL, Vigen C, Kwan ML, Keegan TH, Lu Y, et al. The California Breast Cancer Survivorship Consortium (CBCSC): prognostic factors associated with racial/ethnic differences in breast cancer survival. *Cancer Causes Control*. 2013; 24:1821–36. [PubMed: 23864487]
6. Wu AH, Wan P, Hankin J, Tseng CC, Yu MC, Pike MC. Adolescent and adult soy intake and risk of breast cancer in Asian-Americans. *Carcinogenesis*. 2002; 23:1491–6. [PubMed: 12189192]
7. Marchbanks PA, McDonald JA, Wilson HG, Burnett NM, Daling JR, Bernstein L, et al. The NICHD Women's Contraceptive and Reproductive Experiences Study: methods and operational results. *Ann Epidemiol*. 2002; 12:213–21. [PubMed: 11988408]
8. John EM, Horn-Ross PL, Koo J. Lifetime physical activity and breast cancer risk in a multiethnic population: the San Francisco Bay area breast cancer study. *Cancer Epidemiol Biomarkers Prev*. 2003; 12:1143–52. [PubMed: 14652273]
9. John EM, Phipps AI, Davis A, Koo J. Migration history, acculturation, and breast cancer risk in Hispanic women. *Cancer Epidemiol Biomarkers Prev*. 2005; 14:2905–13. [PubMed: 16365008]
10. Bernstein L, Allen M, Anton-Culver H, Deapen D, Horn-Ross PL, Peel D, et al. High breast cancer incidence rates among California teachers: results from the California Teachers Study (United States). *Cancer Causes Control*. 2002; 13:625–35. [PubMed: 12296510]
11. Kolonel LN, Henderson BE, Hankin JH, Nomura AM, Wilkens LR, Pike MC, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *Am J Epidemiol*. 2000; 151:346–57. [PubMed: 10695593]
12. Yost K, Perkins C, Cohen R, Morris C, Wright W. Socioeconomic status and breast cancer incidence in California for different race/ethnic groups. *Cancer Causes Control*. 2001; 12:703–11. [PubMed: 11562110]
13. Yang, JSC.; Harrati, A.; Clarke, C.; Keegan, THM.; Gomez, SL. Developing an area-based socioeconomic measure from American Community Survey data. Fremont, California: Prevention Institute of California; 2014.
14. Berrigan D, Pickle LW, Dill J. Associations between street connectivity and active transportation. *Int J Health Geogr*. 2010; 9:20. [PubMed: 20412597]
15. National Establishment Time-Series (NETS) Database. 2009. Oakland, CA: 2008.
16. Gunier RB, Hertz A, Von Behren J, Reynolds P. Traffic density in California: socioeconomic and ethnic differences among potentially exposed children. *J Expo Anal Environ Epidemiol*. 2003; 13:240–6. [PubMed: 12743618]
17. California Department of Transportation. Highway Performance and Monitoring System. 2004
18. Keegan TH, Shariff-Marco S, Sangaramoorthy M, Koo J, Hertz A, Schupp CW, et al. Neighborhood influences on recreational physical activity and survival after breast cancer. *Cancer Causes Control*. 2014; 25:1295–308. [PubMed: 25088804]
19. Keegan TH, Hurley S, Goldberg D, Nelson DO, Reynolds P, Bernstein L, et al. The association between neighborhood characteristics and body size and physical activity in the California teachers study cohort. *Am J Public Health*. 2012; 102:689–97. [PubMed: 21852626]
20. Lin DYWL. The robust inference for the Cox Proportional Hazards Mode. *Journal of the American Statistical Association*. 1989; 84:1074–8.
21. John EM, Hopper JL, Beck JC, Knight JA, Neuhausen SL, Senie RT, et al. The Breast Cancer Family Registry: an infrastructure for cooperative multinational, interdisciplinary and translational studies of the genetic epidemiology of breast cancer. *Breast Cancer Res*. 2004; 6:R375–89. [PubMed: 15217505]
22. John EM, Miron A, Gong G, Phipps AI, Felberg A, Li FP, et al. Prevalence of pathogenic BRCA1 mutation carriers in 5 US racial/ethnic groups. *JAMA*. 2007; 298:2869–76. [PubMed: 18159056]

23. Sprague BL, Trentham-Dietz A, Gangnon RE, Ramchandani R, Hampton JM, Robert SA, et al. Socioeconomic status and survival after an invasive breast cancer diagnosis. *Cancer*. 2011; 117:1542–51. [PubMed: 21425155]
24. O'Malley CD, Le GM, Glaser SL, Shema SJ, West DW. Socioeconomic status and breast carcinoma survival in four racial/ethnic groups: a population-based study. *Cancer*. 2003; 97:1303–11. [PubMed: 12599239]
25. Byers TE, Wolf HJ, Bauer KR, Bolick-Aldrich S, Chen VW, Finch JL, et al. The impact of socioeconomic status on survival after cancer in the United States : findings from the National Program of Cancer Registries Patterns of Care Study. *Cancer*. 2008; 113:582–91. [PubMed: 18613122]
26. Du XL, Fang S, Meyer TE. Impact of treatment and socioeconomic status on racial disparities in survival among older women with breast cancer. *Am J Clin Oncol*. 2008; 31:125–32. [PubMed: 18391595]
27. Shariff-Marco S, Yang J, John EM, Sangaramoorthy M, Hertz A, Koo J, et al. Impact of neighborhood and individual socioeconomic status on survival after breast cancer varies by race/ethnicity: the Neighborhood and Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev*. 2014; 23:793–811. [PubMed: 24618999]
28. Bassett MT, Krieger N. Social class and black-white differences in breast cancer survival. *Am J Public Health*. 1986; 76:1400–3. [PubMed: 3777285]
29. Li F, Fisher KJ, Brownson RC, Bosworth M. Multilevel modelling of built environment characteristics related to neighbourhood walking activity in older adults. *J Epidemiol Community Health*. 2005; 59:558–64. [PubMed: 15965138]
30. McCormack GR, Giles-Corti B, Bulsara M. The relationship between destination proximity, destination mix and physical activity behaviors. *Prev Med*. 2008; 46:33–40. [PubMed: 17481721]
31. Kim J, Choi WJ, Jeong SH. The effects of physical activity on breast cancer survivors after diagnosis. *J Cancer Prev*. 2013; 18:193–200. [PubMed: 25337546]
32. Sugiyama T, Francis J, Middleton NJ, Owen N, Giles-Corti B. Associations between recreational walking and attractiveness, size, and proximity of neighborhood open spaces. *Am J Public Health*. 2010; 100:1752–7. [PubMed: 20634455]
33. Gomez SL, Shariff-Marco S, De Rouen M, Keegan THM, Yen IH, Mujahid M, Satariano WA, Glaser SL. The Impact of Neighborhood Social and Built Environment Factors across the Cancer Continuum: Current Research, Methodologic Considerations, and Future Directions. *Cancer*. 2015 in press.
34. Banegas MP, Tao L, Altekruze S, Anderson WF, John EM, Clarke CA, et al. Heterogeneity of breast cancer subtypes and survival among Hispanic women with invasive breast cancer in California. *Breast Cancer Res Treat*. 2014; 144:625–34. [PubMed: 24658879]
35. Gomez SL, Clarke CA, Shema SJ, Chang ET, Keegan THM, Glaser SL. Disparities in Breast Cancer Survival Among Asian Women by Ethnicity and Immigrant Status: A Population-Based Study. *American Journal of Public Health*. 2010; 100:861–9. [PubMed: 20299648]
36. Keegan T, Quach T, Shema S, Glaser S, Gomez S. The influence of nativity and neighborhoods on breast cancer stage at diagnosis and survival among California Hispanic women. *BMC Cancer*. 2010; 10:603. [PubMed: 21050464]
37. Patel MI, Schupp CW, Gomez SL, Chang ET, Wakelee HA. How Do Social Factors Explain Outcomes in Non-Small-Cell Lung Cancer Among Hispanics in California? Explaining the Hispanic Paradox. *J Clin Oncol*. 2013
38. Schupp CW, Press DJ, Gomez SL. Immigration factors and prostate cancer survival among Hispanic men in California: Does neighborhood matter? *Cancer*. 2014
39. Lim JW, Ashing-Giwa KT. Examining the effect of minority status and neighborhood characteristics on cervical cancer survival outcomes. *Gynecol Oncol*. 2011; 121:87–93. [PubMed: 21183210]
40. Eschbach K, Ostir GV, Patel KV, Markides KS, Goodwin JS. Neighborhood context and mortality among older Mexican Americans: is there a barrio advantage? *Am J Public Health*. 2004; 94:1807–12. [PubMed: 15451754]

41. Russell E, Kramer MR, Cooper HL, Thompson WW, Arriola KR. Residential racial composition, spatial access to care, and breast cancer mortality among women in Georgia. *J Urban Health*. 2011; 88:1117–29. [PubMed: 21847712]
42. Russell EF, Kramer MR, Cooper HL, Gabram-Mendola S, Senior-Crosby D, Jacob Arriola KR. Metropolitan area racial residential segregation, neighborhood racial composition, and breast cancer mortality. *Cancer Causes Control*. 2012; 23:1519–27. [PubMed: 22825071]
43. Warner ET, Gomez SL. Impact of neighborhood racial composition and metropolitan residential segregation on disparities in breast cancer stage at diagnosis and survival between black and white women in California. *J Community Health*. 2010; 35:398–408. [PubMed: 20358266]
44. Lochner KA, Kawachi I, Brennan RT, Buka SL. Social capital and neighborhood mortality rates in Chicago. *Soc Sci Med*. 2003; 56:1797–805. [PubMed: 12639596]
45. Keegan TH, Quach T, Shema S, Glaser SL, Gomez SL. The influence of nativity and neighborhoods on breast cancer stage at diagnosis and survival among California Hispanic women. *BMC Cancer*. 2010; 10:603. [PubMed: 21050464]
46. Gomez SL, Clarke CA, Shema SJ, Chang ET, Keegan TH, Glaser SL. Disparities in breast cancer survival among Asian women by ethnicity and immigrant status: a population-based study. *Am J Public Health*. 2010; 100:861–9. [PubMed: 20299648]
47. Martinez-Tyson D, Pathak EB, Soler-Vila H, Flores AM. Looking under the Hispanic umbrella: cancer mortality among Cubans, Mexicans, Puerto Ricans and other Hispanics in Florida. *J Immigr Minor Health*. 2009; 11:249–57. [PubMed: 18506623]
48. Gomez SL, Glaser SL, Horn-Ross PL, Cheng I, Quach T, Clarke CA, et al. Cancer research in Asian American, Native Hawaiian, and Pacific Islander populations: accelerating cancer knowledge by acknowledging and leveraging heterogeneity. *Cancer Epidemiol Biomarkers Prev*. 2014; 23:2202–5. [PubMed: 25368394]
49. Gomez SL, Quach T, Horn-Ross PL, Pham JT, Cockburn M, Chang ET, et al. Hidden breast cancer disparities in Asian women: disaggregating incidence rates by ethnicity and migrant status. *Am J Public Health*. 2010; 100(Suppl 1):S125–31. [PubMed: 20147696]
50. Yen IH, Scherzer T, Cubbin C, Gonzalez A, Winkleby MA. Women's perceptions of neighborhood resources and hazards related to diet, physical activity, and smoking: focus group results from economically distinct neighborhoods in a mid-sized U.S. city. *Am J Health Promot*. 2007; 22:98–106. [PubMed: 18019886]
51. Diez Roux AV. Neighborhoods and health: where are we and where do we go from here? *Rev Epidemiol Sante Publique*. 2007; 55:13–21. [PubMed: 17320330]
52. Fish JS, Ettner S, Ang A, Brown AF. Association of perceived neighborhood safety with [corrected] body mass index. *Am J Public Health*. 2010; 100:2296–303. [PubMed: 20864717]

Table 1
Association between pre-diagnosis BMI and the neighborhood environment, California Breast Cancer Survivorship Consortium

	All n=8995			African Americans n=1719			Asian Americans n=1234			
	Overweight vs. Normal Weight		Obese vs. Normal Weight	Overweight vs. Normal Weight		Obese vs. Normal Weight	Overweight vs. Normal Weight		Obese vs. Normal Weight	
	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI
Socioeconomic status ^{b,c}										
Q5-high	1.00		1.00		1.00		1.00		1.00	
Q4	1.34	(1.15-1.55)	1.23	(1.01-1.49)	1.33	(0.78-2.28)	1.25	(0.69-2.27)	0.84	(0.53-1.36)
Q3	1.51	(1.25-1.82)	1.36	(1.08-1.71)	1.53	(0.84-2.79)	1.51	(0.80-2.85)	0.94	(0.51-1.73)
Q2	1.44	(1.15-1.81)	1.35	(1.03-1.78)	1.76	(0.93-3.34)	1.66	(0.84-3.28)	0.95	(0.46-1.97)
Q1-low	1.72	(1.28-2.30)	1.43	(1.01-2.02)	1.93	(0.93-4.02)	1.66	(0.76-3.64)	1.26	(0.52-3.04)
p trend	<0.01		0.02		0.06		0.16		0.68	
Population density ^c										
Q1-low	1.00		1.00		1.00		1.00		1.00	
Q2	1.09	(0.91-1.30)	1.04	(0.83-1.30)	1.51	(0.77-2.97)	1.04	(0.52-2.08)	1.32	(0.71-2.44)
Q3	1.07	(0.88-1.31)	1.08	(0.84-1.39)	1.53	(0.78-2.98)	0.95	(0.47-1.92)	1.48	(0.77-2.85)
Q4-high	1.09	(0.84-1.40)	1.14	(0.83-1.55)	1.44	(0.70-2.96)	0.83	(0.38-1.79)	1.05	(0.48-2.28)
p trend	0.53		0.23		0.22		0.84		0.83	
Urbanicity ^c										
metropolitan suburban	1.00		1.00		1.00		1.00		1.00	
metropolitan urban	0.86	(0.70-1.07)	0.88	(0.69-1.12)	1.02	(0.68-1.53)	1.05	(0.69-1.61)	0.82	(0.46-1.47)
city	1.00	(0.85-1.17)	1.08	(0.88-1.33)	1.23	(0.59-2.56)	1.26	(0.59-2.73)	0.80	(0.30-2.15)
town	1.28	(0.85-1.92)	1.16	(0.65-2.05)	--	--	0.68	(0.19-2.44)	--	--
rural	0.82	(0.59-1.14)	1.08	(0.70-1.67)	0.18	(0.01-2.84)	--	--	--	--
% Foreign Born ^c										
Q1-low %	1.00		1.00		1.00		1.00		1.00	
Q2	1.07	(0.92-1.24)	1.09	(0.90-1.31)	0.78	(0.53-1.16)	0.95	(0.63-1.45)	1.13	(0.52-2.44)
Q3	1.06	(0.89-1.25)	1.05	(0.84-1.30)	0.89	(0.58-1.38)	0.95	(0.59-1.54)	1.90	(0.92-3.93)
Q4-high %	1.06	(0.86-1.32)	1.05	(0.81-1.37)	0.81	(0.48-1.36)	1.33	(0.78-2.26)	1.60	(0.75-3.39)
p trend	0.69		0.89		0.78		0.26		0.27	
% Commuting by public transportation/walk/bike ^c										
Q1-low %	1.00		1.00		1.00		1.00		1.00	
										0.98

	All n=8995						African Americans n=1719						Asian Americans n=1234					
	Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight			
	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI		
	0.97	(0.80-1.16)	1.10	(0.89-1.38)	1.22	(0.72-2.05)	1.71	(0.97-3.01)	0.67	(0.37-1.22)	0.74	(0.30-1.82)						
Q4-high %																		
p trend		0.32		0.69		0.89		0.07		0.33		0.14						
Household crowding ^{c,d}																		
Q1-low	1.00		1.00		1.00		1.00		1.00		1.00		1.00		1.00			
Q2	1.10	(0.95-1.28)	1.20	(0.98-1.46)	0.72	(0.43-1.21)	0.65	(0.37-1.16)	1.49	(0.87-2.53)	2.98	(0.86-10.35)						
Q3	1.06	(0.88-1.29)	1.35	(1.06-1.72)	0.61	(0.35-1.07)	0.72	(0.39-1.32)	1.30	(0.70-2.38)	1.98	(0.51-7.66)						
Q4-high	1.24	(0.94-1.63)	1.54	(1.10-2.16)	0.57	(0.28-1.17)	0.63	(0.29-1.37)	2.17	(0.98-4.80)	3.67	(0.73-18.42)						
p trend		0.41		0.02		0.07		0.22		0.12		0.16						
% Multi-family housing units ^{c,e}																		
Q1-low %	1.00		1.00		1.00		1.00		1.00		1.00		1.00		1.00			
Q2	0.95	(0.82-1.10)	0.87	(0.72-1.05)	1.10	(0.69-1.73)	1.02	(0.63-1.66)	0.97	(0.62-1.51)	0.88	(0.41-1.90)						
Q3	0.80	(0.68-0.95)	0.89	(0.73-1.08)	0.89	(0.56-1.42)	1.02	(0.63-1.66)	0.66	(0.41-1.07)	1.14	(0.52-2.50)						
Q4-high %	0.89	(0.74-1.07)	0.92	(0.74-1.15)	1.29	(0.78-2.11)	1.35	(0.80-2.26)	0.91	(0.52-1.60)	0.83	(0.31-2.21)						
p trend		0.04		0.39		0.64		0.44		0.21		0.47						
Street connectivity: Gamma ^{f,h}																		
Q1-low connectivity	1.00		1.00		1.00		1.00		1.00		1.00		1.00		1.00			
Q2	1.00	(0.86-1.16)	1.04	(0.86-1.27)	0.87	(0.52-1.45)	0.87	(0.49-1.53)	1.03	(0.65-1.62)	1.14	(0.56-2.29)						
Q3	0.90	(0.77-1.07)	1.11	(0.90-1.37)	0.99	(0.59-1.67)	1.57	(0.88-2.80)	1.03	(0.64-1.65)	0.68	(0.31-1.51)						
Q4-high connectivity	0.98	(0.81-1.19)	1.19	(0.94-1.50)	1.04	(0.61-1.77)	1.59	(0.88-2.87)	1.36	(0.79-2.33)	0.99	(0.37-2.63)						
p trend		0.66		0.28		0.50		0.02		0.17		0.68						
Number of businesses ^h																		
Q1-low	1.00		1.00		1.00		1.00		1.00		1.00		1.00		1.00			
Q2	0.94	(0.79-1.12)	0.97	(0.78-1.22)	0.74	(0.43-1.29)	0.97	(0.50-1.89)	0.93	(0.53-1.65)	0.79	(0.30-2.10)						
Q3	1.01	(0.83-1.24)	1.07	(0.83-1.38)	0.87	(0.48-1.55)	1.08	(0.54-2.16)	0.95	(0.51-1.78)	0.99	(0.33-2.92)						
Q4-high	0.86	(0.68-1.08)	0.74	(0.56-0.99)	0.66	(0.35-1.27)	0.68	(0.32-1.45)	1.01	(0.50-2.04)	0.61	(0.17-2.19)						
p trend		0.55		0.07		0.37		0.13		0.91		0.31						

	All n=8995				African Americans n=1719				Asian Americans n=1234			
	Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight	
	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI
Restaurant environment index ^{g,h}	1.00		1.00		1.00		1.00		1.00		1.00	
only non-fast food restaurants												
<median	1.11	(0.94-1.32)	1.05	(0.85-1.29)	1.13	(0.67-1.89)	0.96	(0.55-1.67)	0.85	(0.52-1.42)	0.95	(0.36-2.51)
>median	1.07	(0.92-1.25)	1.07	(0.88-1.30)	1.27	(0.79-2.06)	0.97	(0.58-1.63)	0.90	(0.55-1.46)	1.38	(0.58-3.28)
No Business	0.97	(0.78-1.21)	0.99	(0.74-1.32)	1.45	(0.60-3.49)	1.26	(0.51-3.14)	0.77	(0.34-1.72)	0.63	(0.13-3.02)
p trend		0.42		0.15		0.28		0.92		0.55		0.10
Number of parks ^h	1.00		1.00		1.00		1.00		1.00		1.00	
0												
1	0.86	(0.75-0.99)	0.97	(0.82-1.15)	1.00	(0.69-1.47)	0.89	(0.60-1.33)	0.84	(0.57-1.24)	1.67	(0.76-3.67)
2	0.98	(0.84-1.14)	0.99	(0.82-1.19)	1.19	(0.79-1.80)	0.85	(0.55-1.32)	0.91	(0.59-1.40)	2.17	(0.91-5.19)
3	0.97	(0.83-1.15)	1.08	(0.89-1.31)	1.18	(0.77-1.81)	1.10	(0.70-1.73)	0.70	(0.42-1.17)	1.44	(0.55-3.80)
p trend		0.76		0.32		0.31		0.65		0.21		0.27
Socioeconomic status ^{b,c}	1.00		1.00		1.00		1.00		1.00		1.00	
Q5-high												
Q4	1.44	(0.97-2.14)	1.02	(0.65-1.62)	1.44	(1.18-1.77)	1.43	(1.09-1.88)				
Q3	1.17	(0.73-1.86)	0.74	(0.44-1.25)	1.77	(1.36-2.31)	1.73	(1.23-2.44)				
Q2	1.16	(0.66-2.04)	0.97	(0.54-1.76)	1.49	(1.05-2.13)	1.69	(1.08-2.65)				
Q1-low	1.21	(0.60-2.45)	0.78	(0.37-1.65)	2.50	(1.49-4.18)	2.52	(1.31-4.84)				
p trend		0.96		0.69		<0.01		<0.01				
Population density ^c	1.00		1.00		1.00		1.00		1.00		1.00	
Q1-low												
Q2	1.17	(0.70-1.98)	0.87	(0.50-1.51)	1.01	(0.82-1.26)	1.19	(0.88-1.62)				
Q3	1.28	(0.75-2.17)	0.89	(0.51-1.56)	0.90	(0.69-1.17)	1.25	(0.88-1.79)				
Q4-high	1.25	(0.67-2.35)	0.92	(0.47-1.79)	1.06	(0.69-1.62)	1.59	(0.94-2.67)				
p trend		0.55		0.97		0.87		0.06				
Urbanicity ^c	1.00		1.00		1.00		1.00		1.00		1.00	
metropolitan suburban												
metropolitan urban	0.92	(0.57-1.47)	1.01	(0.62-1.65)	0.86	(0.56-1.32)	0.79	(0.48-1.30)				
city	1.03	(0.64-1.66)	1.25	(0.74-2.10)	1.00	(0.82-1.22)	0.99	(0.76-1.28)				

	All n=8995						African Americans n=1719						Asian Americans n=1234					
	Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight			
	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI	OR ^d	95% CI		
	--	(0.18-2.57)	1.69	(0.56-5.13)	1.16	(0.37-3.67)	1.27	(0.82-1.98)	1.11	(0.59-2.07)								
town	0.68		1.00		1.00		1.00		0.96									
rural	1.00		1.00		1.00		1.00		1.00									
% Foreign Born ^c	1.11	(0.67-1.86)	1.88	(1.06-3.34)	1.46	(0.78-2.75)	1.24	(1.03-1.50)	0.95	(0.74-1.23)								
Q1-low %	0.85	(0.49-1.47)	1.46	(0.78-2.75)	1.14	(0.90-1.44)	1.14	(0.90-1.44)	0.97	(0.70-1.35)								
Q2	0.89	(0.48-1.66)	1.10	(0.55-2.19)	1.18	(0.82-1.69)	1.18	(0.82-1.69)	1.00	(0.63-1.57)								
Q3	0.76		0.30		0.30		0.26		0.90									
Q4-high %	1.00		1.00		1.00		1.00		1.00									
p trend	0.82		0.42		0.42		0.37		0.99									
% Commuting by public transportation/walk/bike ^c	1.00		1.00		1.00		1.00		1.00									
Q1-low %	1.19	(0.75-1.89)	0.83	(0.52-1.34)	0.83	(0.52-1.34)	1.04	(0.86-1.25)	1.22	(0.95-1.56)								
Q2	0.88	(0.55-1.41)	0.74	(0.46-1.20)	0.91	(0.74-1.13)	0.91	(0.74-1.13)	0.98	(0.74-1.28)								
Q3	1.11	(0.66-1.86)	0.77	(0.45-1.32)	0.89	(0.68-1.17)	0.89	(0.68-1.17)	1.12	(0.80-1.58)								
Q4-high %	0.82		0.42		0.42		0.37		0.99									
p trend	1.00		1.00		1.00		1.00		1.00									
Household crowding ^{c,d}	1.81	(1.12-2.92)	1.97	(1.13-3.42)	1.97	(1.13-3.42)	1.02	(0.84-1.23)	1.14	(0.88-1.48)								
Q1-low	1.66	(0.96-2.87)	1.93	(1.02-3.65)	1.04	(0.79-1.37)	1.04	(0.79-1.37)	1.45	(1.02-2.05)								
Q2	2.16	(1.08-4.31)	3.24	(1.50-7.00)	1.07	(0.68-1.67)	1.07	(0.68-1.67)	1.09	(0.61-1.95)								
Q3	0.06		<0.01		<0.01		0.70		0.39									
Q4-high	1.00		1.00		1.00		1.00		1.00									
p trend	1.16	(0.80-1.69)	1.01	(0.67-1.52)	0.88	(0.72-1.08)	0.88	(0.72-1.08)	0.86	(0.66-1.12)								
% Multi-family housing units ^{c,e}	1.14	(0.75-1.71)	1.32	(0.86-2.04)	1.32	(0.86-2.04)	0.79	(0.63-0.99)	0.68	(0.50-0.91)								
Q1-low %	1.14	(0.72-1.79)	1.16	(0.71-1.89)	1.16	(0.71-1.89)	0.72	(0.54-0.95)	0.77	(0.54-1.10)								
Q2	0.38		0.21		0.21		<0.01		0.03									
Q3	1.00		1.00		1.00		1.00		1.00									
Q4-high %	1.07	(0.73-1.58)	1.30	(0.84-2.00)	0.99	(0.81-1.20)	0.99	(0.81-1.20)	0.97	(0.74-1.27)								
p trend	1.00		1.00		1.00		1.00		1.00									
Street connectivity: Gamma/h	1.07	(0.73-1.58)	1.30	(0.84-2.00)	0.99	(0.81-1.20)	0.99	(0.81-1.20)	0.97	(0.74-1.27)								
Q1-low connectivity	1.00		1.00		1.00		1.00		1.00									
Q2	1.07	(0.73-1.58)	1.30	(0.84-2.00)	0.99	(0.81-1.20)	0.99	(0.81-1.20)	0.97	(0.74-1.27)								

	All n=8995				African Americans n=1719				Asian Americans n=1234			
	Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight		Overweight vs. Normal Weight		Obese vs. Normal Weight	
	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI	OR ^a	95% CI
Q3	1.05	(0.70-1.59)	1.33	(0.84-2.11)	0.82	(0.65-1.03)	1.06	(0.79-1.44)				
Q4-high connectivity	1.17	(0.72-1.89)	1.77	(1.06-2.95)	0.78	(0.59-1.04)	0.86	(0.60-1.25)				
p trend	0.38		0.03		0.03		0.17					
Number of businesses ^h	1.00		1.00		1.00		1.00					
Q1-low	0.95	(0.61-1.49)	1.08	(0.66-1.76)	1.01	(0.80-1.28)	0.90	(0.66-1.24)				
Q2	0.88	(0.54-1.45)	0.97	(0.56-1.69)	1.12	(0.84-1.49)	1.10	(0.76-1.58)				
Q3	0.69	(0.38-1.25)	0.79	(0.42-1.49)	0.94	(0.67-1.32)	0.61	(0.39-0.96)				
Q4-high	0.51		0.31		0.90		0.44					
p trend	0.57		0.20		0.81		0.59					
only non-fast food restaurants	1.00		1.00		1.00		1.00					
Restaurant environment index ^{g/h}	1.71	(1.12-2.60)	1.21	(0.77-1.89)	1.07	(0.84-1.37)	1.17	(0.85-1.60)				
<median	1.28	(0.85-1.91)	1.24	(0.81-1.91)	1.02	(0.82-1.26)	1.01	(0.76-1.33)				
>median	1.43	(0.71-2.89)	1.02	(0.48-2.18)	0.88	(0.68-1.14)	1.06	(0.73-1.54)				
No Business	0.82		0.60		0.26		0.78					
p trend	0.82		0.60		0.26		0.78					
Number of parks ^h	1.00		1.00		1.00		1.00					
0	0.98	(0.67-1.42)	1.05	(0.70-1.56)	0.81	(0.66-0.99)	0.98	(0.75-1.27)				
1	1.16	(0.77-1.73)	1.01	(0.65-1.55)	0.89	(0.71-1.12)	1.06	(0.80-1.42)				
2	0.84	(0.55-1.27)	1.07	(0.69-1.65)	1.12	(0.88-1.41)	1.00	(0.73-1.37)				
3	0.82		0.60		0.26		0.78					
p trend	0.82		0.60		0.26		0.78					

^a Stratified by stage (AJCC) and study (AABCS, CARE, CTS, MEC, SFBCS). Adjusted for age, log (age), year of diagnosis, block group clustering, education, number of births, smoking status, alcohol consumption, hypertension, diabetes. Analysis for all groups combined also adjusted for race/ethnicity

^b Based on SES composite index of seven indicator variables for Census block groups (Liu education index, proportion blue collar job, proportion older than age 16 in the workforce without a job, median household income, percent below 200% of federal poverty line, median rent, median house value)

^c U.S. census data; categories based on CA state-wide distribution

^d Percent occupied housing with 1 occupant per room

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Percent of housing structures with ≥ 2 units
 f_j Ratio of actual number of street segments to maximum possible number of intersections
 g_j Ratio of the number of fast food restaurants to non-fast food restaurants
 h_j Business or traffic data; categories based on study participant distribution

Table 2
Association between pre-diagnosis BMI, the neighborhood environment, and breast cancer-specific mortality, California Breast Cancer Survivorship Consortium

	All n=8995		African Americans n=1719		Asian Americans n=1234		Latinas n=1754		Non-Latina Whites n=4234	
	Deaths (n)	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	
BMI (m/kg ²)										
Normal weight	548	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Overweight	393	1.04 (0.90-1.19)	0.79 (0.60-1.05)	1.44 (0.93-2.22)	0.99 (0.68-1.44)	1.09 (0.86-1.37)	1.00	1.09 (0.68-1.44)	1.09 (0.86-1.37)	
Obese	213	1.08 (0.90-1.29)	0.83 (0.60-1.14)	1.96 (0.96-3.98)	1.06 (0.69-1.64)	1.21 (0.87-1.69)	1.00	1.06 (0.69-1.64)	1.21 (0.87-1.69)	
Severely obese	75	1.06 (0.81-1.37)	0.87 (0.56-1.37)	--	0.88 (0.45-1.71)	1.37 (0.86-2.19)	1.00	0.88 (0.45-1.71)	1.37 (0.86-2.19)	
Morbidly obese	55	1.22 (0.89-1.68)	1.00 (0.61-1.64)	--	2.13 (1.10-4.15)	0.94 (0.46-1.92)	1.00	2.13 (1.10-4.15)	0.94 (0.46-1.92)	
p trend		0.21	0.66	0.15	0.23	0.24	0.15	0.23	0.24	
Socioeconomic status ^{b,c}										
Q5-high	308	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
Q4	267	0.95 (0.79-1.14)	0.59 (0.37-0.93)	0.66 (0.36-1.22)	0.79 (0.48-1.30)	1.19 (0.91-1.56)	1.00	0.79 (0.48-1.30)	1.19 (0.91-1.56)	
Q3	247	1.00 (0.81-1.24)	0.69 (0.43-1.11)	0.85 (0.38-1.86)	1.11 (0.64-1.93)	1.19 (0.85-1.68)	1.00	1.11 (0.64-1.93)	1.19 (0.85-1.68)	
Q2	251	1.14 (0.89-1.46)	0.73 (0.45-1.20)	0.93 (0.38-2.27)	0.76 (0.40-1.42)	1.73 (1.12-2.67)	1.00	0.76 (0.40-1.42)	1.73 (1.12-2.67)	
Q1-low	210	1.19 (0.87-1.62)	0.65 (0.37-1.13)	1.20 (0.41-3.53)	1.16 (0.54-2.52)	2.75 (1.47-5.12)	1.00	1.16 (0.54-2.52)	2.75 (1.47-5.12)	
p trend		0.10	0.91	0.90	0.96	< 0.01	0.90	0.96	< 0.01	
Household crowding ^{c,d}										
Q1-low	272	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
Q2	269	0.89 (0.74-1.07)	0.59 (0.38-0.92)	0.72 (0.36-1.43)	0.86 (0.45-1.65)	0.96 (0.75-1.24)	1.00	0.86 (0.45-1.65)	0.96 (0.75-1.24)	
Q3	345	0.99 (0.81-1.22)	0.87 (0.57-1.32)	1.10 (0.54-2.26)	1.17 (0.62-2.21)	0.76 (0.55-1.05)	1.00	1.17 (0.62-2.21)	0.76 (0.55-1.05)	
Q4-high	397	0.90 (0.70-1.17)	0.82 (0.50-1.34)	0.62 (0.25-1.54)	0.93 (0.45-1.92)	0.73 (0.45-1.18)	1.00	0.93 (0.45-1.92)	0.73 (0.45-1.18)	
p trend		0.67	0.90	0.82	0.96	0.10	0.90	0.96	0.10	
% Multi-family housing units ^{c,e}										
Q1-low %	274	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
Q2	310	1.19 (1.00-1.42)	1.10 (0.73-1.64)	1.08 (0.61-1.90)	1.91 (1.17-3.10)	1.00 (0.77-1.30)	1.00	1.91 (1.17-3.10)	1.00 (0.77-1.30)	
Q3	343	1.13 (0.94-1.37)	1.27 (0.85-1.89)	0.69 (0.36-1.33)	1.98 (1.20-3.26)	0.79 (0.58-1.07)	1.00	1.98 (1.20-3.26)	0.79 (0.58-1.07)	
Q4-high %	356	1.08 (0.89-1.32)	1.07 (0.71-1.63)	1.34 (0.69-2.58)	1.67 (0.96-2.88)	0.90 (0.64-1.26)	1.00	1.67 (0.96-2.88)	0.90 (0.64-1.26)	
p trend		0.87	0.98	0.88	0.14	0.16	0.88	0.14	0.16	
Street connectivity: Gamma ^{f,g}										
Q1-low %	276	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	

	Deaths (n)	All n=8995		African Americans n=1719		Asian Americans n=1234		Latinas n=1754		Non-Latina Whites n=4234	
		HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI
Q2	284	0.87	(0.73-1.05)	0.89	(0.56-1.41)	1.36	(0.76-2.45)	0.82	(0.50-1.35)	0.77	(0.59-0.99)
Q3	339	0.91	(0.75-1.10)	0.86	(0.54-1.35)	1.75	(0.92-3.35)	1.01	(0.61-1.65)	0.76	(0.56-1.03)
Q4-high %	385	0.95	(0.77-1.17)	0.87	(0.55-1.39)	1.30	(0.65-2.61)	1.53	(0.88-2.66)	0.76	(0.54-1.08)
p trend			0.77		0.74		0.48		0.05		0.12
Number of businesses ^g	277	1.00		1.00		1.00		1.00		1.00	
Q2	323	1.01	(0.84-1.21)	0.95	(0.61-1.49)	1.27	(0.68-2.38)	0.68	(0.42-1.12)	1.04	(0.80-1.36)
Q3	369	1.07	(0.88-1.30)	1.28	(0.81-2.01)	0.91	(0.47-1.76)	0.55	(0.32-0.96)	1.19	(0.87-1.63)
Q4-high	314	0.97	(0.77-1.22)	1.18	(0.72-1.95)	0.54	(0.25-1.16)	0.46	(0.25-0.85)	1.09	(0.73-1.61)
p trend			0.82		0.27		0.10		0.04		0.39
Number of parks ^g	337	1.00		1.00		1.00		1.00		1.00	
1	398	1.01	(0.86-1.18)	1.10	(0.81-1.50)	0.79	(0.49-1.28)	1.66	(1.01-2.73)	1.01	(0.78-1.30)
2	284	1.07	(0.90-1.28)	1.15	(0.82-1.60)	0.96	(0.56-1.64)	1.75	(1.05-2.90)	0.99	(0.73-1.36)
3	264	0.97	(0.80-1.16)	0.98	(0.69-1.40)	0.70	(0.36-1.37)	2.02	(1.19-3.43)	0.92	(0.67-1.26)
p trend			0.90		0.87		0.44		0.03		0.60

^a Stratified by age (AJCC) and study (AABCS, CARE, CTS, MEC, SFBACS). Adjusted for age, log (age), year of diagnosis, histology, grade, ER/PR status, nodal involvement, tumor size, second primary tumor, multiple primary tumor, days from diagnosis of index tumor to secondary primary diagnosis, days from diagnosis of index tumor to multiple primary tumor, surgery type, chemotherapy, radiation, clustering by block group, education, parity, smoking, alcohol consumption, hypertension, diabetes. Analysis for all groups combined also adjusted for race/ethnicity

^b Based on SES composite index of seven indicator variables for Census block groups (Lin education index, proportion blue collar job, proportion older than age 16 in the workforce without a job, median household income, percent below 200% of federal poverty line, median rent, median house value).

^c U.S. census data; categories based on CA state-wide distribution

^d Percent occupied housing with 1 occupant per room

^e Percent of housing structures with 2 units

^f Ratio of actual number of street segments to maximum possible number of intersections

^g Business or traffic data; categories based on study participant distribution

Table 3
Association between pre-diagnosis BMI, the neighborhood environment, and all cause mortality, California Breast Cancer Survivorship Consortium

	All n=8995		African Americans n=1719		Asian Americans n=1234		Latinas n=1754		Non-Latina Whites n=4234	
	Deaths (n)	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	HR ^a 95% CI	
BMI (m/kg ²)										
Normal weight	1008	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Overweight	753	1.01 (0.91-1.12)	0.87 (0.7-1.07)	1.23 (0.88-1.71)	1.00 (0.75-1.33)	1.01 (0.87-1.17)				
Obese	419	1.07 (0.94-1.22)	0.81 (0.64-1.04)	1.45 (0.84-2.48)	1.20 (0.88-1.65)	1.18 (0.96-1.46)				
Severely obese	150	1.12 (0.93-1.35)	0.86 (0.62-1.2)	--	1.33 (0.86-2.06)	1.41 (1.02-1.94)				
Morbidly obese	96	1.24 (0.98-1.57)	1.04 (0.72-1.5)	--	2.15 (1.31-3.53)	1.06 (0.64-1.76)				
p trend		0.05	0.42	0.42	<0.01	0.05				
Socioeconomic status ^{b,c}										
Q5-high	584	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Q4	501	0.95 (0.83-1.08)	0.54 (0.38-0.77)	0.87 (0.53-1.43)	0.90 (0.62-1.31)	1.05 (0.88-1.26)				
Q3	490	1.08 (0.93-1.26)	0.72 (0.50-1.03)	1.16 (0.63-2.14)	1.30 (0.86-1.96)	1.13 (0.91-1.40)				
Q2	472	1.12 (0.94-1.34)	0.66 (0.45-0.95)	0.87 (0.42-1.78)	0.92 (0.57-1.48)	1.42 (1.07-1.87)				
Q1-low	376	1.11 (0.89-1.38)	0.58 (0.38-0.89)	1.11 (0.48-2.58)	1.07 (0.61-1.90)	1.75 (1.17-2.62)				
p trend		0.16	0.27	0.78	0.71	0.01				
Household crowding ^{c,d}										
Q1-low	516	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Q2	535	1.03 (0.90-1.17)	0.82 (0.58-1.15)	1.20 (0.68-2.09)	0.98 (0.61-1.57)	1.07 (0.91-1.27)				
Q3	646	1.11 (0.96-1.29)	1.09 (0.78-1.51)	1.45 (0.80-2.63)	0.98 (0.61-1.58)	1.03 (0.84-1.27)				
Q4-high	726	1.04 (0.87-1.25)	1.03 (0.71-1.49)	1.12 (0.55-2.30)	0.96 (0.57-1.62)	0.89 (0.66-1.21)				
p trend		0.41	0.52	0.39	0.78	0.86				
% Multi-family housing units ^{c,e}										
Q1-low %	490	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Q2	573	1.12 (0.98-1.27)	1.00 (0.74-1.35)	1.16 (0.74-1.81)	1.62 (1.15-2.28)	1.04 (0.87-1.25)				
Q3	650	1.09 (0.95-1.25)	1.22 (0.91-1.65)	0.87 (0.53-1.42)	1.47 (1.03-2.09)	0.90 (0.74-1.11)				
Q4-high %	710	1.12 (0.97-1.3)	1.15 (0.85-1.56)	1.12 (0.66-1.91)	1.39 (0.94-2.04)	1.02 (0.82-1.27)				
p trend		0.25	0.28	0.92	0.28	0.62				
Street connectivity: Gamma ^{f,g}										
Q1-low %	505	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

		All n=8995			African Americans n=1719		Asian Americans n=1234		Latinas n=1754		Non-Latina Whites n=4234	
		Deaths (n)	HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI	HR ^a	95% CI
		587	0.98	(0.87-1.12)	1.06	(0.75-1.52)	1.32	(0.84-2.09)	1.08	(0.76-1.56)	0.92	(0.78-1.09)
	Q2	610	0.93	(0.81-1.06)	1.05	(0.74-1.49)	1.45	(0.88-2.39)	0.95	(0.66-1.37)	0.80	(0.65-0.97)
	Q3	723	1.03	(0.89-1.19)	1.08	(0.76-1.53)	1.18	(0.68-2.03)	1.42	(0.95-2.14)	0.91	(0.73-1.14)
	Q4-high %											
	p trend			0.86		0.70		0.41		0.13		0.21
	Q1-low	519	1.00		1.00		1.00		1.00		1.00	
	Q2	614	1.01	(0.89-1.15)	0.91	(0.65-1.28)	1.09	(0.66-1.79)	0.84	(0.59-1.21)	1.06	(0.88-1.26)
	Q3	679	1.02	(0.89-1.18)	1.00	(0.71-1.41)	0.98	(0.58-1.66)	0.83	(0.56-1.24)	1.12	(0.91-1.37)
	Q4-high	611	0.93	(0.79-1.09)	1.00	(0.69-1.46)	0.61	(0.34-1.1)	0.70	(0.44-1.09)	0.99	(0.77-1.27)
	p trend			0.38		0.67		0.05		0.20		0.73
	Number of parks ^g	617	1.00		1.00		1.00		1.00		1.00	
	1	734	1.00	(0.9-1.13)	1.07	(0.84-1.35)	0.74	(0.51-1.08)	1.27	(0.90-1.80)	1.02	(0.86-1.21)
	2	537	1.05	(0.92-1.19)	1.11	(0.86-1.44)	0.87	(0.58-1.32)	1.30	(0.91-1.85)	1.05	(0.86-1.28)
	3	535	1.03	(0.90-1.17)	1.19	(0.91-1.55)	0.76	(0.46-1.25)	1.26	(0.86-1.85)	0.99	(0.81-1.22)
	p trend			0.55		0.27		0.31		0.33		0.95

^a Stratified by age (AJCC) and study (AABCS,CARE,CTS, MEC, SFBACS). Adjusted for age, log (age), year of diagnosis, histology, grade, ER/PR status, nodal involvement, tumor size, second primary tumor, multiple primary tumor, days from diagnosis of index tumor to secondary primary diagnosis, days from diagnosis of index tumor to multiple primary tumor, surgery type, chemotherapy, radiation, clustering by block group, education, parity, smoking, alcohol consumption, hypertension, diabetes. Analysis for all groups combined also adjusted for race/ethnicity

^b Based on SES composite index of seven indicator variables for Census block groups (Lin education index, proportion blue collar job, proportion older than age 16 in the workforce without a job, median household income, percent below 200% of federal poverty line, median rent, median house value).

^c U.S. census data; categories based on CA state-wide distribution

^d Percent occupied housing with 1 occupant per room

^e Percent of housing structures with 2 units

^f Ratio of actual number of street segments to maximum possible number of intersections

^g Business or traffic data; categories based on study participant distribution