

Lawrence Berkeley National Laboratory

LBL Publications

Title

Near-Complete Genome Sequence of *Thalassospira* sp. Strain KO164 Isolated from a Lignin-Enriched Marine Sediment Microcosm

Permalink

<https://escholarship.org/uc/item/18v224kx>

Journal

Microbiology Resource Announcements, 4(6)

ISSN

2576-098X

Authors

Woo, Hannah L

O'Dell, Kaela B

Utturkar, Sagar

et al.

Publication Date


2016-12-29

DOI

10.1128/genomea.01297-16

Peer reviewed

Near-Complete Genome Sequence of *Thalassospira* sp. Strain KO164 Isolated from a Lignin-Enriched Marine Sediment Microcosm

Hannah L. Woo,^a Kaela B. O'Dell,^b Sagar Utturkar,^c Kathryn R. McBride,^b Marcel Huntemann,^d Alicia Clum,^d Manoj Pillay,^d Krishnaveni Palaniappan,^d Neha Varghese,^d Natalia Mikhailova,^d Dimitrios Stamatis,^d T. B. K. Reddy,^d Chew Yee Ngan,^d Chris Daum,^d Nicole Shapiro,^d Victor Markowitz,^d Natalia Ivanova,^d Nikos Kyrpides,^d Tanja Woyke,^d Steven D. Brown,^{c,e}  Terry C. Hazen^{a,b,c,e,f}

Department of Civil and Environmental Engineering, University of Tennessee, Knoxville, Tennessee, USA^a; Department of Microbiology, University of Tennessee, Knoxville, Tennessee, USA^b; Graduate School of Genome Science and Technology, University of Tennessee, Knoxville, Tennessee, USA^c; DOE Joint Genome Institute, Walnut Creek, California, USA^d; Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^e; Department of Earth and Planetary Sciences, University of Tennessee, Knoxville, Tennessee, USA^f

***Thalassospira* sp. strain KO164 was isolated from eastern Mediterranean seawater and sediment laboratory microcosms enriched on insoluble organosolv lignin under oxic conditions. The near-complete genome sequence presented here will facilitate analyses into this deep-ocean bacterium's ability to degrade recalcitrant organics such as lignin.**

Received 3 October 2016 Accepted 7 October 2016 Published 23 November 2016

Citation Woo HL, O'Dell KB, Utturkar S, McBride KR, Huntemann M, Clum A, Pillay M, Palaniappan K, Varghese N, Mikhailova N, Stamatis D, Reddy TBK, Ngan CY, Daum C, Shapiro N, Markowitz V, Ivanova N, Kyrpides N, Woyke T, Brown SD, Hazen TC. 2016. Near-complete genome sequence of *Thalassospira* sp. strain KO164 isolated from a lignin-enriched marine sediment microcosm. *Genome Announc* 4(6):e01297-16. doi:10.1128/genomeA.01297-16.

Copyright © 2016 Woo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Terry C. Hazen, tchazen@utk.edu.

Lignin derived from the plant cell wall is difficult for microbes to degrade due to its complex, aromatic, and heterogeneous structure. The physiochemical conditions and microbial controls on lignin degradation is relatively unknown. This knowledge gap is especially ill studied in marine environments, where lignin can enter as allochthonous organic matter through rivers (1). Few microorganisms other than white-rot fungi have been found to degrade lignin (2). In deep marine environments, where extreme oligotrophy and salinity can limit fungal growth, bacterial microorganisms may be primarily responsible for any lignin degradation. A better understanding of marine lignin degradation by microbes may discover potentially novel and stress-tolerant biocatalysts for lignin valorization application in industry.

Seawater and sediment were sampled from the Nile Deep Sea Fan, an oxic and hypersaline deep-ocean environment at 700-m depth (3). The sample was then enriched with 0.05% organosolv lignin for several weeks and then serially diluted and plated onto minimal media (ONR7a–DSMZ medium 950) agar containing 0.05% weight/volume alkali lignin. A colony appeared on the agar within two weeks of being incubated at 14°C that was later identified as belonging to the genus *Thalassospira* by 16S rRNA gene PCR and Sanger sequencing.

Genomic DNA was isolated from KO164 using a MoBio Ultra-Clean extraction kit, and sequencing was performed at the Joint Genome Institute (JGI, Walnut Creek, CA, USA). A Pacific Biosciences standard template preparation protocol was used for creating >10-kb libraries. Genome sequencing was performed using a PacBio RS II instrument as described previously, using C4 chemistry and polymerase version P6 (4, 5). The sequencing run generated 166,402 filtered subreads totaling 718.9 Mb. All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov>. The raw reads were assembled using HGAP version: 2.3.0 (6). The final assembly con-

tained two contigs, with a total genome size of 4.81 Mb. The input read coverage was 87.3×. Gene identification and annotation were performed at Oak Ridge National Laboratory using methods described previously (7).

The genome contains 4,472 candidate protein-coding genes and includes putative oxidoreductases, aromatic-ring opening dioxygenases, and peroxidases which may present targets for follow-up studies. Based on a BLASTn search of its four 16S rRNA gene copies, KO164 was most similar to *Thalassospira xiamenensis*, a strain isolated from oil-contaminated coastal water (8) and *T. permensis*, a halotolerant naphthalene-degrading strain from soil (9). In Biolog growth assays, KO164 was halotolerant up to 8% NaCl and grew optimally at pH 9.5. Among 200 different carbon sources tested, KO164 grew best in L-glutamate, acetic acid, L-pyroglutamate, malate, L-ornithine, glycyl-L-proline, itaconic acid, glycyl-L-glutamate, N-acetyl-L-glutamate, and L-glutamine. KO164 possesses the pathways to produce L-glutamate from alpha-ketoglutarate and ammonium but benefits from the exogenous addition of glutamate or glutamine. This genome sequence will facilitate studies in the breakdown of lignin and broader carbon and nitrogen cycling analyses of marine microbiomes. In addition, insights into growth in high-pH environments will be enabled by comparative genomics studies with other alkaliphiles such as *Caldalkalibacillus thermarum* strain TA2.A1 (10) and *Clostridium paradoxum* strain JW-YL-7 (11).

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [MAJC00000000](https://www.ncbi.nlm.nih.gov/nuclink/MAJC00000000).

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under grant no. DGE-1452154. This work was also supported by the Office of Biological

and Environmental Research in the DOE Office of Science through the BES, a DOE Bioenergy Research Center. Oak Ridge National Laboratory is managed by UT-Battelle, LLC, Oak Ridge, Tennessee, USA, for the DOE under contract DE-AC05-00OR22725. The work conducted by the DOE Joint Genome Institute is supported by the DOE Office of Science under contract no. DE-AC02-05CH11231.

FUNDING INFORMATION

This work, including the efforts of Hannah Woo, was funded by NSF Graduate Research Fellowship Program (DGE-1452154). This work, including the efforts of Marcel Huntemann, Alicia Clum, Manoj Pillay, Krishna Palaniappan, Neha Varghese, Natalia Mikhailova, Dimitrios Stamatis, T.B.K. Reddy, Chew Ngan, Chris Daum, Nicole Shapiro, Victor Markowitz, Natalia Ivanova, Nikos C. Kyrpides, and Tanja Woyke, was funded by DOE Joint Genome Institute (DE-AC02-05CH11231). This work, including the efforts of Kaela B. O'Dell, Sagar Utturkar, Kathryn R. McBride, Steven D. Brown, and Terry C. Hazen, was funded by DOE | LDRD | Oak Ridge National Laboratory (ORNL) (DE-AC05-00OR22725).

REFERENCES

- Hedges JI, Keil RG, Benner R. 1997. What happens to terrestrial organic matter in the ocean? *Org Geochem* 27:195–212. [http://dx.doi.org/10.1016/S0146-6380\(97\)00066-1](http://dx.doi.org/10.1016/S0146-6380(97)00066-1).
- Kirk TK, Farrell RL. 1987. Enzymatic “combustion”: the microbial degradation of lignin. *Annu Rev Microbiol* 41:465–501. <http://dx.doi.org/10.1146/annurev.mi.41.100187.002341>.
- Techtmann SM, Fortney JL, Ayers KA, Joyner DC, Linley TD, Pfiffner SM, Hazen TC. 2015. The unique chemistry of eastern Mediterranean water masses selects for distinct microbial communities by depth. *PLoS One* 10:e0120605. <http://dx.doi.org/10.1371/journal.pone.0120605>.
- Brown SD, Nagaraju S, Utturkar S, De Tissera S, Segovia S, Mitchell W, Land ML, Dassanayake A, Köpke M. 2014. Comparison of single-molecule sequencing and hybrid approaches for finishing the genome of *Clostridium autoethanogenum* and analysis of CRISPR systems in industrial relevant clostridia. *Biotechnol Biofuels* 7:40. <http://dx.doi.org/10.1186/1754-6834-7-40>.
- Utturkar SM, Klingeman DM, Bruno-Barcena JM, Chinn MS, Grunden AM, Köpke M, Brown SD. 2015. Sequence data for *Clostridium autoethanogenum* using three generations of sequencing technologies. *Sci Data* 2:150014. <http://dx.doi.org/10.1038/sdata.2015.14>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
- Woo HL, Utturkar S, Klingeman D, Simmons BA, DeAngelis KM, Brown SD, Hazen TC. 2014. Draft genome sequence of the lignin-degrading *Burkholderia* sp. strain LIG30, isolated from wet tropical forest soil. *Genome Announc* 2(3):e00637-14. <http://dx.doi.org/10.1128/genomeA.00637-14>.
- Lai Q, Shao Z. 2012. Genome sequence of *Thalassospira xiamenensis* type strain M-5. *J Bacteriol* 194:6957. <http://dx.doi.org/10.1128/JB.01904-12>.
- Plotnikova EG, Anan'ina LN, Krausova VI, Ariskina EV, Prisyazhnaya NV, Lebedev AT, Demakov VA, Evtushenko LI. 2011. *Thalassospira permensis* sp. nov., a new terrestrial halotolerant bacterium isolated from a naphthalene-utilizing microbial consortium. *Mikrobiologiya* 80:691–699. <http://dx.doi.org/10.1134/S0026261711050109>.
- Kalamorz F, Keis S, McMillan DG, Olsson K, Stanton J-A, Stockwell P, Black MA, Klingeman DM, Land ML, Han CS, Martin SL, Becher SA, Peddie CJ, Morgan HW, Matthies D, Preiss L, Meier T, Brown SD, Cook GM. 2011. Draft genome sequence of the thermoalkaliphilic *Caldalkalibacillus thermarum* strain TA2.A1. *J Bacteriol* 193:4290–4291. <http://dx.doi.org/10.1128/JB.05035-11>.
- Lancaster WA, Utturkar SM, Poole FL, Klingeman DM, Elias DA, Adams MW, Brown SD. 2016. Near-complete genome sequence of *Clostridium paradoxum* strain JW-YL-7. *Genome Announc* 4(3):e00229-16. <http://dx.doi.org/10.1128/genomeA.00229-16>.