

UC Davis

UC Davis Previously Published Works

Title

Correction: The persimmon genome reveals clues to the evolution of a lineage-specific sex determination system in plants

Permalink

<https://escholarship.org/uc/item/19515818>

Journal

PLOS Genetics, 16(5)

ISSN

1553-7390

Authors

Akagi, Takashi
Shirasawa, Kenta
Nagasaki, Hideki
[et al.](#)

Publication Date

2020

DOI

10.1371/journal.pgen.1008845

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

CORRECTION

Correction: The persimmon genome reveals clues to the evolution of a lineage-specific sex determination system in plants

Takashi Akagi, Kenta Shirasawa, Hideki Nagasaki, Hideki Hirakawa, Ryutaro Tao, Luca Comai, Isabelle M. Henry

The images for Figs 5 and 6 are incorrectly switched. The image that appears as Fig 5 should be Fig 6, and the image that appears as Fig 6 should be Fig 5. The figure captions appear in the correct order. Please see the correct Figs 5 and 6 here.



OPEN ACCESS

Citation: Akagi T, Shirasawa K, Nagasaki H, Hirakawa H, Tao R, Comai L, et al. (2020) Correction: The persimmon genome reveals clues to the evolution of a lineage-specific sex determination system in plants. PLoS Genet 16(5): e1008845. <https://doi.org/10.1371/journal.pgen.1008845>

Published: May 26, 2020

Copyright: © 2020 Akagi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 5. Functional differentiation between *MeGI* and *SiMeGI*. **a-h**, *N. tabacum* transgenic lines expressing either of *MeGI* or *SiMeGI* under the control of the 35S promoter. The lines expressing *MeGI* (**a-c**) showed rudimental anthers (**a**) which did not produce functional pollen grains (**b**), and severe dwarfism with chlorophyll starvation and narrow leaves (**c**, see S7 Fig for the detail). The lines expressing *SiMeGI* (**d-f**) developed regular anthers (**d**) which produced fertile pollen (**e**), and showed moderate dwarfism (**f**). pis: pistil, ra: rudimental anthers, an: anthers. **g-h**, Both *MeGI*- and *SiMeGI*-overexpressing lines were phenotypically different from the control plants transformed with empty vectors (cont), but the *MeGI*-expressing lines exhibited more severe departure from the WT controls for specific traits, such as leaves width (see S9 Fig). Bars indicate 5mm for **a** and **d**, 50mm for **c**, **f**, **g**, and **h**. **i-j**, expression patterns of *MeGI*, *SiMeGI*, and *PI*, with *actin* as a positive control, in the transgenic lines transformed with CaMV35S-*MeGI* (**i**) and CaMV35S-*SiMeGI* (**j**). **k**, DNA motifs identified as preferentially bound to following transcription factors all nested within the *MeGI*/*SiMeGI* clade: *MeGI* [25], *SiMeGI* (our experiments), and three Arabidopsis HD-ZIP1 genes [29], using DAP-Seq analyses (see Methods). **l-n**, expression patterns of *MeGI* and *SiMeGI* in buds and flower primordia were highly correlated (Pearson's $r > 0.7$). Expression levels in female (**l**) and male (**m**) are expressed as RPKM values. **n**, Developmental stages.

<https://doi.org/10.1371/journal.pgen.1008845.g001>




Fig 6. Genomic context of the Y-chromosomal region surrounding OGI. a, Read coverage from male (blue) and female (pink) samples and male/female coverage ratio across the scaffolds covering the male-specific region of the Y-chromosome. For both the male and female reads, expected coverage a single-copy sites is approximately 20 (grey lines across). This male-specific region was assembled via anchoring of the scaffolds with BAC sequences. Approximately 1.3Mb region was covered by Y-allelic scaffolds. More than 400kb of long repetitive sequences (dotted lines), flank OGI. Outer regions of these hyper repetitive sequences contain male-specific sequences (blue bands in M/F rate) and pseudo autosomal region (PAR)-like sequences (orange lines), where M/F rate was less than 70%, and the percentage of repetitive sequences was much lower. b, The silent divergence rate (dS) between X and Y alleles of the genes located in the PAR-like sequences (orange circles) decreases with distance to OGI. Still, for most of these genes, the dS value between the X and Y alleles was larger than the average interspecific dS between the X alleles of *D. lotus* and *D. mespiliformis* (green square and dotted line), *D. lotus* and *D. virginiana* (blue square and dotted line), and *D. lotus* and *D. kaki* (red square dotted line). These results suggest that, in these PAR-like sequences, recombination between the X and Y alleles was suppressed before the divergence of *Diospyros* species, or at least predates the divergence between *D. lotus* and *D. kaki*. dS values for genes located in the regions closest to OGI are comparable to dS values between OGI and MeGI (gray circle, dS = 0.205), which suggest that little or no recombination occurred between these sequences after the establishment of OGI. In comparison, dS values between the X and Y alleles of genes located in the recombining region of chromosomes 15 are much lower (white circles on the right).

<https://doi.org/10.1371/journal.pgen.1008845.g002>

Reference

1. Akagi T, Shirasawa K, Nagasaki H, Hirakawa H, Tao R, Comai L, et al. (2020) The persimmon genome reveals clues to the evolution of a lineage-specific sex determination system in plants. *PLoS Genet* 16 (2): e1008566. <https://doi.org/10.1371/journal.pgen.1008566> PMID: 32069274