

UC Merced

UC Merced Previously Published Works

Title

Draft Genome Sequence of Syntrophorhabdus aromaticivorans Strain UI, a Mesophilic Aromatic Compound-Degrading Syntroph

Permalink

<https://escholarship.org/uc/item/19h5q7j7>

Journal

Microbiology Resource Announcements, 2(1)

ISSN

2576-098X

Authors

Nobu, Masaru K
Narihiro, Takashi
Tamaki, Hideyuki
[et al.](#)

Publication Date

2014-02-27

DOI

10.1128/genomea.01064-13

Peer reviewed

Draft Genome Sequence of *Syntrophorhabdus aromaticivorans* Strain UI, a Mesophilic Aromatic Compound-Degrading Syntroph

Masaru K. Nobu,^a Takashi Narihiro,^{a,b} Hideyuki Tamaki,^b Yan-Ling Qiu,^c Yuji Sekiguchi,^d Tanja Woyke,^e Lynne Goodwin,^f Karen W. Davenport,^f Yoichi Kamagata,^{b,g} Wen-Tso Liu^a

Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA^a; Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki, Japan^b; Key Laboratory of Biofuels, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, Shandong Province, People's Republic of China^c; Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki, Japan^d; DOE Joint Genome Institute, Walnut Creek, California, USA^e; Los Alamos National Laboratory, Los Alamos, New Mexico, USA^f; Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Sapporo, Hokkaido, Japan^g

***Syntrophorhabdus aromaticivorans* strain UI is a mesophilic bacterium capable of degrading aromatic substrates in syntrophic cooperation with a partner methanogen. The draft genome sequence is 3.7 Mb, with a G+C content of 52.0%.**

Received 8 November 2013 Accepted 21 January 2014 Published 6 February 2014

Citation Nobu MK, Narihiro T, Tamaki H, Qiu Y-L, Sekiguchi Y, Woyke T, Goodwin L, Davenport KW, Kamagata Y, Liu W-T. 2014. Draft genome sequence of *Syntrophorhabdus aromaticivorans* strain UI, a mesophilic aromatic compound-degrading syntroph. *Genome Announc.* 2(1):e01064-13. doi:10.1128/genomeA.01064-13.

Copyright © 2014 Nobu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Wen-Tso Liu, wtliu@illinois.edu.

Syntrophorhabdus aromaticivorans strain UI is one of the few isolated bacteria capable of anaerobic degradation of aromatic compounds in syntrophic association with a methanogen (1). *S. aromaticivorans* UI was isolated from a mesophilic bioreactor treating aromatic compound-containing wastewater as a representative of the family *Syntrophorhabdaceae* of the class *Deltaproteobacteria*. Other isolates include members of the genus *Syntrophus* (*S. aciditrophicus* strain SB, *S. buswellii* strain DM-2, and *S. gentianae* strain HQGö1) of the class *Deltaproteobacteria* and members of the genera *Pelotomaculum* (*P. isophthalicum* strain JI and *P. terephthalicum* strain JT) and *Sporotomaculum* (*S. hydroxybenzoicum* strain BT and *S. syntrophicum* strain FB) of the class *Clostridia* (2–7). Of these aromatic compound-degrading syntrophs, strain UI is uniquely capable of degrading phenol (1). It is also known to syntrophically degrade *p*-cresol, 4-hydroxybenzoate, isophthalate, and benzoate with a hydrogenotrophic methanogen (1), suggesting that *Syntrophorhabdaceae* members likely play an important role in anaerobic treatment of industrial wastewater-containing aromatic compounds.

The draft genome of *S. aromaticivorans* strain UI was generated from a coculture of strain UI and *Methanospirillum hungatei* strain JF-1 at the DOE Joint Genome Institute (JGI) with a combined approach using Illumina (8) and 454 (9) technologies. We constructed and sequenced an Illumina GAI shotgun library (69,270,135 reads) totaling 5,264.5 Mb and a paired-end 454 library (322,727 reads, an average insert size of 8 kb) totaling 64 Mb of 454 data. Assemblies were performed using a combination of Newbler version 2.3-PreRelease-6/30/2009, Velvet version 1.0.13 (10), and parallel Phrap version SPS-4.24 (High Performance Software, LLC). The software Consed (11–13) was used in the following finishing process. Illumina data were used to correct potential base errors and increase consensus quality using the software Polisher developed at JGI (A. Lapidus, unpublished data). Possible misassemblies were corrected using gapResolution (C. Han, unpublished data) or DupFinisher (14) or se-

quencing cloned bridging PCR fragments with subcloning. Gaps between contigs were closed by editing in Consed, by PCR, and by bubble PCR (J.-F. Cheng, unpublished data) primer walks. A total of 308 additional reactions were necessary to close gaps and to raise the quality of the finished sequence. *M. hungatei* strain JF-1 sequences were removed using scaffold read coverage and BLAST. The estimated genome size is 3.7 Mb, and the final assembly is based on 30.3 Mb of 454 draft data and 5,235.1 Mb of Illumina draft data, which provide an average 8.2× coverage and 1,414.9× coverage of the genome, respectively.

The strain UI draft genome is composed of three linear scaffolds and has a G+C content of 52.0%. This genome contains 3,691 genes, 3,632 protein-coding and 59 RNA genes. The Integrated Microbial Genomes (IMG) pipeline (15) annotated 71.82% of the protein-coding genes with function prediction. The 59 RNA genes consisted of 3 rRNA genes (5S, 16S, and 23S), 51 tRNA genes, and 5 other RNA genes. We are currently exploring this genome to identify novel syntrophic aromatic degradation genes and pathways for further comparison with benzoate-degrading syntrophs.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [AJUN00000000](https://www.ncbi.nlm.nih.gov/nuccore/AJUN00000000). The version described in this paper is the first version, [AJUN01000000](https://www.ncbi.nlm.nih.gov/nuccore/AJUN01000000).

ACKNOWLEDGMENT

The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231.

REFERENCES

1. Qiu YL, Hanada S, Ohashi A, Harada H, Kamagata Y, Sekiguchi Y. 2008. *Syntrophorhabdus aromaticivorans* gen. nov., sp. nov., the first cultured anaerobe capable of degrading phenol to acetate in obligate syntrophic associations with a hydrogenotrophic methanogen. *Appl. Environ. Microbiol.* 74:2051–2058. [http://dx.doi.org/10.1128/AEM.02378-07](https://doi.org/10.1128/AEM.02378-07).

2. Mountfort DO, Brulla WJ, Krumholz LR, Bryant MP. 1984. *Syntrophus buswellii* gen. nov., sp. nov.: a benzoate catabolizer from methanogenic ecosystems. *Int. J. Syst. Bacteriol.* 34:216–217. <http://dx.doi.org/10.1099/00207713-34-2-216>.
3. Wallrabenstein C, Gorny N, Springer N, Ludwig W, Schink B. 1995. Pure culture of *Syntrophus buswellii*, definition of its phylogenetic status, and description of *Syntrophus gentianae* sp. nov. *Syst. Appl. Microbiol.* 18:62–66. [http://dx.doi.org/10.1016/S0723-2020\(11\)80449-2](http://dx.doi.org/10.1016/S0723-2020(11)80449-2).
4. Jackson BE, Bhupathiraju VK, Tanner RS, Woese CR, McInerney MJ. 1999. *Syntrophus aciditrophicus* sp. nov., a new anaerobic bacterium that degrades fatty acids and benzoate in syntrophic association with hydrogen-using microorganisms. *Arch. Microbiol.* 171:107–114. <http://dx.doi.org/10.1007/s002030050685>.
5. Brauman A, Müller JA, Garcia JL, Brune A, Schink B. 1998. Fermentative degradation of 3-hydroxybenzoate in pure culture by a novel strictly anaerobic bacterium, *Sporotomaculum hydroxybenzoicum* gen. nov., sp. nov. *Int. J. Syst. Bacteriol.* 48(Pt 1):215–221. <http://dx.doi.org/10.1099/00207713-48-1-215>.
6. Qiu YL, Sekiguchi Y, Imachi H, Kamagata Y, Tseng IC, Cheng SS, Ohashi A, Harada H. 2003. *Sporotomaculum syntrophicum* sp. nov., a novel anaerobic, syntrophic benzoate-degrading bacterium isolated from methanogenic sludge treating wastewater from terephthalate manufacturing. *Arch. Microbiol.* 179:457. <http://link.springer.com/article/10.1007/s00203-003-0557-0>.
7. Qiu YL, Sekiguchi Y, Hanada S, Imachi H, Tseng IC, Cheng SS, Ohashi A, Harada H, Kamagata Y. 2006. *Pelotomaculum terephthalicum* sp. nov. and *Pelotomaculum isophthalicum* sp. nov.: two anaerobic bacteria that degrade phthalate isomers in syntrophic association with hydrogenotrophic methanogens. *Arch. Microbiol.* 185:172–182. <http://dx.doi.org/10.1007/s00203-005-0081-5>.
8. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
9. Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen ZT, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer MLI, Jarvie TP, Jirage KB, Kim JB, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT, Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu PG, Begley RF, Rothberg JM. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437:376–380. <http://dx.doi.org/10.1038/nature03959>.
10. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
11. Ewing B, Green P. 1998. Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res.* 8:186–194.
12. Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* 8:175–185. <http://dx.doi.org/10.1101/gr.8.3.175>.
13. Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* 8:195–202. <http://dx.doi.org/10.1101/gr.8.3.195>.
14. Han C, Chain P. 2006. Finishing repeat regions automatically with Dup-Finisher, p 141–146. *In* Arabnia HR, Valafar H (ed), *Proceedings of the 2006 International Conference on Bioinformatics & Computational Biology*. CSREA Press, Las Vegas, NV.
15. Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Grechkin Y, Ratner A, Anderson I, Lykidis A, Mavromatis K, Ivanova NN, Kyrpides NC. 2010. The integrated microbial genomes system: an expanding comparative analysis resource. *Nucleic Acids Res.* 38:D382–D390. <http://dx.doi.org/10.1093/nar/gkp887>.