

UCSF

UC San Francisco Previously Published Works

Title

The role of acalabrutinib in adults with chronic lymphocytic leukemia

Permalink

<https://escholarship.org/uc/item/19p99299>

Authors

Fakhri, Bitu
Andreadis, Charalambos

Publication Date

2021

DOI

10.1177/2040620721990553

Peer reviewed

The role of acalabrutinib in adults with chronic lymphocytic leukemia

Bitá Fakhri^{ID} and Charalambos Andreadis

Ther Adv Hematol

2021, Vol. 12: 1–8

DOI: 10.1177/
2040620721990553

© The Author(s), 2021.
Article reuse guidelines:
[sagepub.com/journals-
permissions](https://sagepub.com/journals-permissions)

Abstract: The treatment landscape of chronic lymphocytic leukemia (CLL) has significantly changed in the past decade. This paradigm shift is due to the introduction of novel agents to the field. The two major classes of drugs that have contributed to this dramatic evolution include the Bruton tyrosine kinase (BTK) inhibitors and BCL2 inhibitors. Ibrutinib was the first-in-class drug which was initially approved by the US Food and Drug Administration (FDA) for the treatment of patients with relapsed/refractory and later for patients with treatment-naïve CLL. Despite encouraging efficacy outcomes, its use has been associated with cardiovascular and gastrointestinal toxicities likely due to off-target inhibition of ITK, TEC and EGFR family kinases. The next generation of BTK inhibitors was developed to be more selective with less off-target inhibition with the prospect to improve tolerability without compromising efficacy. Acalabrutinib, a selective covalent BTK inhibitor, is a second generation BTK inhibitor. The focus of this review is on two major phase III trials that resulted in the FDA approval of acalabrutinib in 2019. The ELEVATE TN trial investigated acalabrutinib with or without obintuzumab *versus* chlorambucil–obintuzumab in older and frail patients with previously untreated CLL. The ASCEND trial explored acalabrutinib *versus* chemoimmunotherapy in patients with relapsed/refractory CLL. Both trials demonstrated superiority of the acalabrutinib-containing arms in terms of both efficacy and tolerability. Unfortunately, the availability of new generation BTK inhibitors has not resulted in mitigating the financial toxicities associated with these potentially life-long treatments.

Keywords: chronic lymphocytic leukemia, small lymphocytic lymphoma, bruton tyrosine kinase inhibitors, BTK inhibitors, acalabrutinib, ibrutinib

Received: 29 July 2020; revised manuscript accepted: 6 January 2021.

Introduction

The treatment landscape of chronic lymphocytic leukemia (CLL) has drastically changed in the past 5 years. Inhibition of the B-cell receptor pathway, particularly of Bruton tyrosine kinase (BTK), is a major therapeutic target in B-cell malignancies, including CLL. The introduction of BTK inhibitors has revolutionized the field of CLL. The first-in-class BTK inhibitor, ibrutinib, was approved for the treatment of patients with relapsed/refractory CLL by the US Food and Drug Administration (FDA) in February 2014. The initial approval was based on the phase III RESONATE trial, which compared single agent ibrutinib to ofatumumab in high-risk patients with relapsed/refractory CLL.¹ Long-term follow-up of

the RESONATE trial showed that at a median follow-up of 65.3 months, the median progression-free survival (PFS) of patients in the ibrutinib arm was 44.1 months *versus* 8.1 months in the ofatumumab arm [hazard ratio (HR) 0.148; 95% confidence interval (CI) 0.113–0.196]. This superiority in PFS was observed consistently across all baseline disease and patient characteristics, including patients with high-risk features.^{2,3} Two years later, in March 2016, the FDA approved ibrutinib as a first-line treatment for patients with untreated CLL. The approval was based on the results of the RESONATE-2 trial, which investigated the use of ibrutinib *versus* chlorambucil in 269 treatment-naïve patients with CLL aged 65 years or older.⁴ The long-term

Correspondence to:

Bitá Fakhri
Department of Medicine,
Division of Hematology/
Blood and Marrow
Transplantation, University
of California, 400
Parnassus Avenue, San
Francisco, CA, USA
Bitá.Fakhri@ucsf.edu

Charalambos Andreadis
Department of Medicine,
Division of Hematology/
Blood and Marrow
Transplantation, University
of California, San
Francisco, CA, USA

follow-up of the RESONATE-2 trial indicated a sustained PFS and overall survival (OS) benefit with ibrutinib monotherapy *versus* chlorambucil in patients with previously untreated CLL including those with high-risk features. At a median follow-up of 60 months, the PFS estimates were 70% *versus* 12% (HR 0.146; 95% CI 0.098–0.218), and the OS estimates were 83% *versus* 68% (HR 0.450; 95% CI 0.266–0.761) – both favoring the ibrutinib arm.^{5,6}

Given the promising long-term outcomes of both RESONATE and RESONATE-2 trials, the next rational step was to compare the ibrutinib-based treatment combinations to traditionally used chemoimmunotherapy regimens in the frontline setting. As such, the Alliance group designed a phase III multicenter study randomly assigning 547 patients older than 65 years of age with treatment-naïve CLL to chemoimmunotherapy with bendamustine–rituximab *versus* single agent ibrutinib *versus* ibrutinib and rituximab combination. At a median follow-up of 38 months, the 2-year PFS was 74% with bendamustine–rituximab (BR) *versus* 87% with single agent ibrutinib (HR 0.39; 95% CI 0.26–0.58; $p < 0.001$) *versus* 88% in the ibrutinib plus rituximab arm (HR 0.38; 95% CI 0.25–0.59; $p < 0.001$). There was no significant difference between the ibrutinib arms with regard to PFS.⁷ In a similar fashion, the ECOG-ACRIN group randomly assigned 529 patients younger than 70 years of age with untreated CLL to chemoimmunotherapy with fludarabine, cyclophosphamide, rituximab (FCR) *versus* ibrutinib plus rituximab. At a median follow-up of 33.6 months, the results of the analysis for both PFS and OS endpoints favored ibrutinib plus rituximab over chemoimmunotherapy in patient with *IGHV* unmutated status. The 3-year PFS rate was 89.4% *versus* 72.9% (HR 0.35; 95% CI 0.22–0.56; $p < 0.001$) favoring the ibrutinib-containing arm. It is important to note that the difference in PFS at 3 years among patients with *IGHV* mutated CLL was not significant (87.7% in the ibrutinib–rituximab group and 88.0% in the chemoimmunotherapy group; HR 0.44; 95% CI 0.14–1.36).⁸ Parallel to the afore-mentioned intergroup studies, the iLLUMINATE trial randomly assigned patients with treatment-naïve CLL and older than 65 years of age or younger than 65 years with comorbidities to receive either obinutuzumab and chlorambucil *versus* obinutuzumab and ibrutinib. After a median follow-up of

31 months, the estimated 30-month PFS was 79% in the ibrutinib plus obinutuzumab group and 31% in the chlorambucil plus obinutuzumab group.⁹ These studies have provided support for adopting targeted agents in select patient populations with treatment-naïve CLL.

Despite promising results associated with ibrutinib, its use has been limited by its toxicity profile. In the largest real-world analysis, with a median follow-up of 17 months, an estimated 41% of patients discontinued ibrutinib mainly due to unfavorable side effects. The most common toxicities include cardiovascular (atrial fibrillation, hypertension), diarrhea, skin rash and arthritis.¹⁰ The availability of new generations of the same class, with more selective target inhibition, can improve on the side effect profile of ibrutinib. In this review, we elaborate on the landmark phase III trials that established a place for acalabrutinib in the treatment of patients with CLL and focus on unmet needs and future directions.

Acalabrutinib mechanism of action and pharmacodynamics

BTK is a proximal component of the B-cell receptor signaling pathway and a therapeutic target for a wide range of B-cell malignancies including CLL. Acalabrutinib forms a covalent bond with Cys481 in the ATP-binding pocket which results in irreversible BTK inhibition. Unlike ibrutinib, acalabrutinib is more selective and does not inhibit other kinases, including EGFR, TEC, or ITK.^{11,12} Inhibition of both on-target and off-target kinases is known to contribute to ibrutinib-related adverse events such as an increased risk of atrial fibrillation, bleeding (through TEK, ITK) and diarrhea (through EGFR).^{13–15} Toxicity is the most common reason for discontinuation of ibrutinib.¹⁰ For patients who discontinue ibrutinib due to intolerance, treatment with acalabrutinib is commonly well tolerated.¹⁶

In terms of dosing schedule, the phase I study of acalabrutinib in B-cell malignancies demonstrated that all tested doses (100–400 mg daily and 100 mg twice daily) achieved superb BTK inhibition 4 h after dosing, but the 100 mg twice daily dosing schedule showed optimal inhibition 24 h after dosing.¹¹ However, BTK occupancy measured in the phase I studies of ibrutinib and acalabrutinib was performed in peripheral blood

mononuclear cells rather than in tissue compartments.^{11,17} Given the fact that most B-cell receptor activation and ensuing cell proliferation occur within lymph nodes, it is imperative to study the relationship between different dosing schedules and maximal BTK occupancy in lymph nodes. To pursue this goal, Sun *et al.* performed a thorough analysis of BTK occupancy by acalabrutinib at two different dosing schedules and the subsequent biological consequences in different tissue compartments. Their randomized phase II study investigated the safety, efficacy, and pharmacodynamics of acalabrutinib at 100 mg twice daily or 200 mg once daily in 48 patients with treatment-naïve or relapsed/refractory CLL. Twice daily dosing achieved higher BTK occupancy and more potent pathway inhibition in lymph nodes compared with once daily dosing. In the twice daily cohort, acalabrutinib was well tolerated. The overall response rate (ORR) in the 100 mg twice daily cohort was 96% (95% CI 78.9–99.9) as compared to 79% (95% CI 58–93) in the 200 mg once daily cohort. The 2-year PFS rate was 92% (95% CI 70–98) in the twice daily group *versus* 87% (95% CI 57–97) in the once daily group.¹⁸ The study was not powered to investigate the impact of BTK occupancy on long-term clinical outcomes. Based on the early phase findings, the phase III clinical trials of acalabrutinib have all adopted the 100 mg twice daily dosing schedule.

Acalabrutinib in treatment of patients with treatment-naïve CLL

ELEVATE TN is a phase III, multicenter, international study in patients with treatment-naïve CLL comparing the efficacy of acalabrutinib with or without obinutuzumab against chlorambucil with obinutuzumab. The trial was designed to include older and frail patient populations. The eligibility criteria included patients with untreated CLL aged 65 years or older, or younger than 65 years with creatinine clearance of 30–69 mL/min or a cumulative illness rating scale for geriatrics score greater than 6. Additional criteria included an Eastern Cooperative Oncology Group (ECOG) performance status score of 0–2 and adequate liver, kidney and hematological function. Patients with significant cardiovascular disease were excluded. Of 535 patients, 179 patients were randomly assigned to receive acalabrutinib plus obinutuzumab, 179 were assigned to single

agent acalabrutinib, and 177 patients were assigned to obinutuzumab and oral chlorambucil in 28-day cycles. To reduce infusion-related reactions, acalabrutinib was administered for one cycle before obinutuzumab administration. Oral acalabrutinib was administered at 100 mg twice a day until disease progression or unacceptable toxicity. Oral chlorambucil was given (0.5 mg/kg) on days 1 and 15 of each cycle for a total of six cycles. The primary endpoint was PFS. Crossover to acalabrutinib was allowed in patients who progressed on obinutuzumab–chlorambucil.¹⁹

At a median follow-up of 28.3 months, median PFS was longer in acalabrutinib-containing arms compared with obinutuzumab–chlorambucil. Estimated 2-year PFS was 93% with acalabrutinib–obinutuzumab (95% CI 87–96%), 87% with acalabrutinib monotherapy (95% CI 81–92%), and 47% with obinutuzumab–chlorambucil (95% CI 39–55%). The most common grade 3 or higher adverse event was neutropenia among all three arms. Grade 3 or higher infections occurred in 21%, 14% and 8% of patients in acalabrutinib–obinutuzumab, acalabrutinib monotherapy, and obinutuzumab–chlorambucil cohorts, respectively. Importantly, grade 3 or higher atrial fibrillation and hypertension were 0.6% and 2.8% in the acalabrutinib plus obinutuzumab arm, 0 and 2.2% in the acalabrutinib monotherapy arm, and 0 and 3% in the obinutuzumab–chlorambucil arm. In terms of difference between the PFS in acalabrutinib-containing arms, the study was not powered to draw statistically significant conclusions between those two arms.^{19,20} Based on this study, the FDA granted approval to acalabrutinib – either as a single agent or in combination with obinutuzumab – for patients with untreated CLL.

Acalabrutinib in treatment of patients with relapsed/refractory CLL

Ascend is a phase III, multicenter, international study investigating acalabrutinib in patients with relapsed/refractory CLL. Eligible patients older than 18 years were randomly assigned and stratified by del(17p) status, ECOG performance status score, and number of prior lines of therapy. Patients received acalabrutinib monotherapy or investigator's choice (idelalisib plus rituximab or bendamustine plus rituximab). The primary endpoint was PFS. Key secondary endpoints included ORR, OS, and safety. Of 310 patients, 155 were

randomly assigned to acalabrutinib monotherapy and 155 patients were randomly assigned to investigator's choice (idelalisib–rituximab, $n=119$; bendamustine–rituximab, $n=36$). Patients had received a median of two prior therapies (range 1–10). At a median follow-up of 16.1 months (range 0.03–22.4 months), median PFS was significantly longer with acalabrutinib monotherapy compared with investigator's choice [not reached *versus* 16.5 months (95% CI 14–17 months); HR 0.31 (95% CI 0.20–0.49); $p < 0.0001$]. Estimated one-year PFS was 88% (95% CI 81–92) for acalabrutinib and 68% (95% CI 59–75) for investigator's choice. The PFS superiority in the acalabrutinib arm was consistent among high-risk subgroups including *IGHV* unmutated status, *TP53* mutation and the presence of del(17p). Serious adverse events occurred in 29%, 56% and 26% of patients treated with acalabrutinib monotherapy, idelalisib–rituximab, and bendamustine–rituximab, respectively. Grade 3 or higher diarrhea was reported in 1%, 24% and 0% of patients treated with acalabrutinib monotherapy, idelalisib–rituximab, and bendamustine–rituximab, respectively. The difference in adverse events profile could cautiously reflect the tolerability of BTK inhibitors in general as compared to PI3 kinase inhibitors.²¹ This trial, alongside with the ELEVATE TN trial, resulted in the approval of acalabrutinib in patients with treatment-naïve or relapsed/refractory CLL.

In addition to the ASCEND trial, acalabrutinib is currently being compared in a head-to-head fashion to ibrutinib in patients with relapsed/refractory CLL (NCT02477696). The accrual target is 533 participants, which started in October 2015 and is anticipated to complete in March 2021. The primary endpoint of this study is PFS, and key secondary endpoints include the incidence of grade 3 and higher treatment-emergent infections, Richter's syndrome, atrial fibrillation and overall survival between the two arms. A summary of the landmark phase III trials of BTK inhibitors for patients with treatment-naïve and relapsed/refractory CLL is provided in Table 1.

Discussion and future directions

The ELEVATE TN trial in untreated patients with CLL and the ASCEND trial in patients with relapsed/refractory CLL indicate that acalabrutinib is an effective treatment option with an acceptable side effect profile. In the ELEVATE

TN trial,²⁰ after a median follow-up of 28.3 months, the 2-year PFS was 90% *versus* 47% favoring the acalabrutinib-containing arms as opposed to the previously acceptable regimen of obinutuzumab and chlorambucil in the older and frail patient population.²² Similarly the ASCEND trial at a median follow-up of 16 months showed a one-year PFS of 88% *versus* 68% favoring the acalabrutinib arm as opposed to the investigator's choice of rituximab plus idelalisib/bendamustine. Both trials met their primary endpoint of efficacy. In addition, they proved that acalabrutinib is indeed associated with a favorable side effect profile due to less off-target kinase inhibition. In the ELEVATE TN trial, the rate of grade ≥ 3 atrial fibrillation and hypertension was roughly 0.5% and 2.5% in patients randomly assigned to the acalabrutinib arms.²⁰ In the ASCEND trial the rate of grade ≥ 3 diarrhea in patients receiving acalabrutinib was 1% – corroborating acalabrutinib less off-target inhibition of the EGFR kinase family.²¹ The final analysis of the RESONATE trial, showed that all-grade (grade ≥ 3) hypertension and atrial fibrillation occurred in 21% (9%) and 12% (6%) of patients, respectively.³ It is critical to acknowledge that indirect cross-trial comparisons are fraught with bias, and one should reserve judgment after the phase III non-inferiority trial of direct comparison between ibrutinib and acalabrutinib completes accrual (NCT02477696). In a real-world analysis of 69 patients with follicular lymphoma or relapsed/refractory CLL who were started on acalabrutinib outside a clinical trial, prior intolerance to ibrutinib was the reason for starting acalabrutinib in the majority of patients; the most common adverse events resulting in discontinuation of ibrutinib included rash (22%), atrial fibrillation/flutter (17%), arthralgia (17%), bleeding (13%), infection (13%) and fatigue (13%). The most common toxicities associated with acalabrutinib included fatigue (13%), infection (13%), diarrhea/colitis (10%), nausea/vomiting (9%), headache in five (7%), rash (7%), bleeding (3%) and atrial fibrillation/flutter (1%).²³ Despite an improved side effect profile in both original trials and in the real-world settings, the twice daily dosing schedule with acalabrutinib as compared to once daily dosing with ibrutinib can inconvenience some patients.

In addition to CLL, both in upfront and relapsed/refractory settings, acalabrutinib has been approved in patients with relapsed/refractory mantle cell lymphoma based on a phase II trial.²⁴ In this trial,

Table 1. Summary of phase III trials of BTK inhibitors for patients with treatment-naïve and relapsed/refractory CLL.

Treatment setting	Study	n	Arms of the study	Median follow-up	PFS	Ggrade ≥3 non-hematological AEs in BTKi arm
Previously untreated	RESONATE-2 ⁶	269	Ibrutinib (n = 136) versus chlorambucil (n = 133)	60 months	5-year PFS: 70% versus 12%	HTN (8%), Afib (5%), diarrhea (4%),
	Alliance ⁷	547	Ibrutinib (n = 182) versus Ibrutinib + rituximab (n = 182) versus bendamustine + rituximab (n = 183)	38 months	2-year PFS 87% versus 88% versus 74%	HTN (29%), Afib (9%), bleeding (2%)
	ECOG-ACRIN ⁸	529	Ibrutinib + rituximab (n = 354) versus FCR (n = 175)	33.6 months	3-year PFS: 89% versus 73%	HTN (19%), Afib (3%), diarrhea (4%)
	iLLUMINATE ⁹	229	Ibrutinib + obinutuzumab (n = 113) versus chlorambucil + Obinutuzumab (n = 119)	31 months	30-month PFS 79% versus 31%	HTN (4%) Afib (5%) diarrhea (3%)
	ELEVATE TN ²⁰	535	Acalabrutinib (n = 179) versus acalabrutinib + obinutuzumab (n = 179) versus chlorambucil + obinutuzumab (n = 177)	28.3 months	2-year PFS 87% versus 93% versus 47%	Headache (1%), HTN (3%), Afib (0.6%), bleeding (2%), diarrhea (4.5%)
Relapsed/refractory	RESONATE ³	391	Ibrutinib (n = 195) versus ofatumumab (n = 196)	65.3 months	5-year PFS 40% versus 3%	HTN (9%), Afib (6%), major hemorrhage (10%) diarrhea (7%)
	ASCEND ²¹	310	Acalabrutinib (155) versus idelalisib-rituximab (119) versus bendamustine-rituximab (36)	16.1 months	1-year PFS 88% versus 68%	Headache (1%), HTN (2%), Afib (2%), bleeding (1%), diarrhea (1%)
	NCT02477696	Accrual Target: 533	Acalabrutinib versus ibrutinib	N/A	<ul style="list-style-type: none"> Primary endpoint: PFS Secondary endpoints: incidence of treatment-emergent grade ≥3 infections, Richter's syndrome, atrial fibrillation, and OS 	Currently enrolling
	ALPINE NCT03734016	Accrual target: 600	Zanubrutinib versus ibrutinib	N/A	<ul style="list-style-type: none"> Primary endpoint: ORR Secondary endpoint: PFS, OS, DOR 	Currently enrolling

Afib, atrial fibrillation; BTK, Bruton tyrosine kinase; CLL, chronic lymphocytic leukemia; DOR, duration of response; FCR, fludarabine, cyclophosphamide, rituximab; HTN, hypertension; OS, overall survival; ORR, overall response rate; PFS, progression-free survival.

acalabrutinib at the recommended dose of 100 mg twice daily was proved to be well tolerated as evidenced by only 1.6% of patients requiring dose reductions and only 6.5% of patients discontinuing acalabrutinib due to adverse events. Of note, atrial fibrillation was not observed in any patient. The most common side effects were headaches (36%) and diarrhea (38%), both of which were typically grades 1–2 and self-limited.^{24,25}

It is critical to note that despite the FDA approval of acalabrutinib for both upfront and relapsed/refractory settings in patients with CLL, there are currently no data on acalabrutinib in young and fit patients. As well as the currently FDA approved ibrutinib and acalabrutinib in the treatment of patients with CLL, other BTK inhibitors are under investigation in multiple national and international trials. To name a few, zanubrutinib – another covalent BTK inhibitor, has shown high overall response rates in patients with relapsed/refractory CLL.²⁶ Moreover, LOXO-305 is a non-covalent BTK inhibitor with activity against the most common acquired BTK inhibitor resistance mutations in CLL (Cys485 and phospholipase C gamma-2).²⁷ The results of phase I of LOXO-305 in patients with relapsed/refractory CLL was presented at the 2020 Annual ASH Meeting, with outstanding response rates in patients with heavily pre-treated CLL.²⁸

Before addressing the future directions, it is necessary to talk about the German CLL14 trial. In this trial, 432 patients with untreated CLL were randomly assigned to BCL2 inhibitor venetoclax plus obinutuzumab for one year *versus* chlorambucil plus obinutuzumab. At a median follow-up of 28 months, the 2-year PFS was significantly higher in the venetoclax plus obinutuzumab group than in the chlorambucil plus obinutuzumab group: 88% as compared with 64% (HR 0.35; 95% CI 0.23–0.53). The study established time-limited venetoclax and obinutuzumab combination as another standard treatment option for patients with untreated CLL.²⁹ The recently published longer follow-up showed that at a median follow-up of about 40 months, the median PFS was not reached in the venetoclax plus obinutuzumab group *versus* 36 months in the chlorambucil plus obinutuzumab group.³⁰

Considering the growing armamentarium of treatment options available for patients with treatment-naïve CLL, the next reasonable step is

to study the combination of BTK inhibitors and BCL2 inhibitors. In doing so, the design of the trials should focus on time-limited approaches with the potential use of minimal residual disease based on next generation sequencing to guide the treatment duration in select populations of patients. The phase II CAPTIVATE trial combined ibrutinib and venetoclax in previously untreated high-risk and older patients with CLL. Patients received ibrutinib monotherapy for three cycles, followed by the addition of venetoclax for a total of 24 cycles. Overall, 92% of the patients had negative prognostic features – unmutated *IGHV*, *TP53* aberration, or chromosome 11q deletion. After 12 cycles of combined treatment, 88% of the patients had complete remission or complete remission with incomplete count recovery, and 61% had remission with undetectable minimal residual disease by flow cytometry in bone marrow.³¹ To evaluate further the efficacy of a combination of BTK inhibitors and BCL2 inhibitors, both the Alliance and ECOG cooperative groups are currently conducting two important phase III trials in patients with untreated CLL. Both trials are comparing ibrutinib–obinutuzumab *versus* ibrutinib–venetoclax–obinutuzumab. The Alliance trial is including patients below the age of 70 years (NCT03701282), whereas the ECOG group is targeting patients 70 years of age and older (NCT03737981). As demonstrated in both ELEVATE TN and ASCEND trials, the efficacy of acalabrutinib as a single agent with or without CD20 monoclonal antibodies is comparable to ibrutinib. Whether the efficacy of acalabrutinib in combination regimens with venetoclax exceeds that of ibrutinib is yet to be elucidated. In an effort to answer this question and parallel to the above-mentioned studies by the Alliance and ECOG cooperative groups, the ACE-CL-311 phase III study is designed to compare the efficacy and safety of acalabrutinib in combination with venetoclax with and without obinutuzumab compared to investigator's choice of chemoimmunotherapy in subjects with previously untreated CLL without chromosome 17 deletion or *TP53* mutation. The primary endpoint of the study is PFS and the accrual target is 780 participants. The study opened to accrual in February 2019 and is expected to complete in April 2024 (NCT03836261). While the field of CLL has made great progress by examining different BTK inhibitors and their combination with BCL2 inhibitors, it is imperative to expand our search for new

molecular targets to benefit further patients who will eventually progress on both classes of drugs. In addition, the future trials should investigate the suitability of acalabrutinib and other BTK inhibitors in the treatment of Richter's syndrome.³²

In summary, the future outlook for patients with CLL has never been brighter. With the development of new targeted agents, the field has entered a new era. Physicians and patients are not choosing between the lesser of two evils anymore. Oncologists have the luxury of tailoring treatment strategies to individual patients' needs and circumstances. As we celebrate the recent advances, it is critical to bear in mind that opportunity translates into responsibility. Unfortunately, illness is universal but access to care is not. The countries with access to targeted agents have a moral and ethical responsibility to facilitate access to these agents to underserved communities through a variety of national and global health initiatives until every patient with CLL can enjoy the benefits of these new treatments.

Conflict of interest

The authors declare that there is no conflict of interest.

Funding


The authors received no financial support for the research, authorship, and/or publication of this article.

Ethics and consent

The authors confirm that ethical committee approval was sought where necessary and is acknowledged within the text of the submitted manuscript.

As the manuscript is a review of literature and no patient information has been included in the text or analysis, the guidelines on patient consent do not apply to this manuscript.

ORCID iD

Bita Fakhri  <https://orcid.org/0000-0002-2124-7243>

References

1. Byrd JC, Brown JR, O'Brien S, *et al.* Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med* 2014; 371: 213–223.
2. Byrd JC, Hillmen P, O'Brien S, *et al.* Long-term follow-up of the RESONATE phase 3 trial of ibrutinib vs ofatumumab. *Blood* 2019; 133: 2031–2042.
3. Munir T, Brown JR, O'Brien S, *et al.* Final analysis from RESONATE: up to six years of follow-up on ibrutinib in patients with previously treated chronic lymphocytic leukemia or small lymphocytic lymphoma. *Am J Hematol* 2019; 94: 1353–1363.
4. Burger JA, Tedeschi A, Barr PM, *et al.* Ibrutinib as initial therapy for patients with chronic lymphocytic leukemia. *N Engl J Med* 2015; 373: 2425–2437.
5. Barr PM, Robak T, Owen C, *et al.* Sustained efficacy and detailed clinical follow-up of first-line ibrutinib treatment in older patients with chronic lymphocytic leukemia: extended phase 3 results from RESONATE-2. *Haematologica* 2018; 103: 1502–1510.
6. Burger JA, Barr PM, Robak T, *et al.* Long-term efficacy and safety of first-line ibrutinib treatment for patients with CLL/SLL: 5 years of follow-up from the phase 3 RESONATE-2 study. *Leukemia*. Epub ahead of print 18 October 2019. DOI: 10.1038/s41375-019-0602-x
7. Woyach JA, Ruppert AS, Heerema NA, *et al.* Ibrutinib regimens versus chemoimmunotherapy in older patients with untreated CLL. *N Engl J Med* 2018; 379: 2517–2528.
8. Shanafelt TD, Wang XV, Kay NE, *et al.* Ibrutinib–rituximab or chemoimmunotherapy for chronic lymphocytic leukemia. *N Engl J Med* 2019; 381: 432–443.
9. Moreno C, Greil R, Demirkan F, *et al.* Ibrutinib plus obinutuzumab versus chlorambucil plus obinutuzumab in first-line treatment of chronic lymphocytic leukaemia (iLLUMINATE): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2019; 20: 43–56.
10. Mato AR, Nabhan C, Thompson MC, *et al.* Toxicities and outcomes of 616 ibrutinib-treated patients in the United States: a real-world analysis. *Haematologica* 2018; 103: 874–879.
11. Byrd JC, Harrington B, O'Brien S, *et al.* Acalabrutinib (ACP-196) in relapsed chronic lymphocytic leukemia. *N Engl J Med* 2016; 374: 323–332.
12. Herman SEM, Montraveta A, Niemann CU, *et al.* The Bruton Tyrosine Kinase (BTK)

- inhibitor acalabrutinib demonstrates potent on-target effects and efficacy in two mouse models of chronic lymphocytic leukemia. *Clin Cancer Res* 2017; 23: 2831–2841.
13. Brown JR. How I treat CLL patients with ibrutinib. *Blood* 2018; 131: 379–386.
 14. Lipsky AH, Farooqui MZ, Tian X, *et al.* Incidence and risk factors of bleeding-related adverse events in patients with chronic lymphocytic leukemia treated with ibrutinib. *Haematologica* 2015; 100: 1571.
 15. Wang A, Yan XE, Wu H, *et al.* Ibrutinib targets mutant-EGFR kinase with a distinct binding conformation. *Oncotarget* 2016; 7: 69760–69769.
 16. Awan FT, Schuh A, Brown JR, *et al.* Acalabrutinib monotherapy in patients with chronic lymphocytic leukemia who are intolerant to ibrutinib. *Blood Adv* 2019; 3: 1553–1562.
 17. Advani RH, Buggy JJ, Sharman JP, *et al.* Bruton tyrosine kinase inhibitor ibrutinib (PCI-32765) has significant activity in patients with relapsed/refractory B-cell malignancies. *J Clin Oncol* 2013; 31: 88–94.
 18. Sun C, Nierman P, Kendall EK, *et al.* Clinical and biological implications of target occupancy in CLL treated with the BTK inhibitor acalabrutinib. *Blood* 2020; 136: 93–105.
 19. Sharman JP, Banerji V, Fogliatto LM, *et al.* ELEVATE TN: phase 3 study of acalabrutinib combined with Obinutuzumab (O) or alone vs o plus Chlorambucil (Clb) in Patients (Pts) with treatment-naive Chronic Lymphocytic Leukemia (CLL). *Blood* 2019; 134 (Suppl. 1): 31.
 20. Sharman JP, Egyed M, Jurczak W, *et al.* Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naive chronic lymphocytic leukaemia (ELEVATE TN): a randomised, controlled, phase 3 trial. *Lancet* 2020; 395: 1278–1291.
 21. Ghia P, Pluta A, Wach M, *et al.* ASCEND: phase III, randomized trial of acalabrutinib versus idelalisib plus rituximab or bendamustine plus rituximab in relapsed or refractory chronic lymphocytic leukemia. *J Clin Oncol* 2020; 28: 2849–2861.
 22. Goede V, Fischer K, Busch R, *et al.* Obinutuzumab plus chlorambucil in patients with CLL and coexisting conditions. *N Engl J Med* 2014; 370: 1101–1110.
 23. Yazdy MS, Mato AR, Roeker LE, *et al.* Toxicities and outcomes of acalabrutinib-treated patients with chronic lymphocytic leukemia: a retrospective analysis of real world patients. *Blood* 2019; 134 (Suppl. 1): 4311.
 24. Wang M, Rule S, Zinzani PL, *et al.* Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): a single-arm, multicentre, phase 2 trial. *Lancet* 2018; 391: 659–667.
 25. Wang M, Rule S, Zinzani PL, *et al.* Long-term follow-up of acalabrutinib monotherapy in patients with relapsed/refractory mantle cell lymphoma. *Blood* 2018; 132 (Suppl. 1): 2876.
 26. Hillmen P, Brown JR, Byrd JC, *et al.* ALPINE: phase III zanubrutinib (BGB-3111) versus ibrutinib in patients with relapsed/refractory (R/R) chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL). *J Clin Oncol* 2019; 37 (Suppl. 15): TPS7572.
 27. Woyach JA, Ruppert AS, Guinn D, *et al.* BTK(C481S)-mediated resistance to ibrutinib in chronic lymphocytic leukemia. *J Clin Oncol* 2017; 35: 1437–1443.
 28. Mato AR, Flinn IW, Pagel JM, *et al.* Results from a first-in-human, proof-of-concept phase 1 trial in pretreated B-cell malignancies for Loxo-305, a next-generation, highly selective, non-covalent BTK inhibitor. *Blood* 2019; 134 (Suppl. 1): 501.
 29. Fischer K, Al-Sawaf O, Bahlo J, *et al.* Venetoclax and obinutuzumab in patients with CLL and coexisting conditions. *N Engl J Med* 2019; 380: 2225–2236.
 30. Al-Sawaf O, Zhang C, Tandon M, *et al.* Venetoclax plus obinutuzumab versus chlorambucil plus obinutuzumab for previously untreated chronic lymphocytic leukaemia (CLL14): follow-up results from a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 2020; 21: 1188–1200.
 31. Jain N, Keating M, Thompson P, *et al.* Ibrutinib and venetoclax for first-line treatment of CLL. *N Engl J Med* 2019; 380: 2095–2103.
 32. Hillmen P, Schuh A, Eyre TA, *et al.* Acalabrutinib monotherapy in patients with richter transformation from the phase 1/2 ACE-CL-001 clinical study. *Blood* 2016; 128: 60.