

UC Irvine

UC Irvine Electronic Theses and Dissertations

Title

Refining Multivariate Value Set Bounds

Permalink

<https://escholarship.org/uc/item/1c11x6gb>

Author

Smith, Luke Alexander

Publication Date

2015

License

<https://creativecommons.org/licenses/by-nc-nd/4.0/> 4.0

Peer reviewed|Thesis/dissertation

UNIVERSITY OF CALIFORNIA,
IRVINE

Refining Multivariate Value Set Bounds

DISSERTATION

submitted in partial satisfaction of the requirements
for the degree of

DOCTOR OF PHILOSOPHY

in Mathematics

by

Luke Alexander Smith

Dissertation Committee:
Professor Daqing Wan, Chair
Professor Alice Silverberg
Professor Karl Rubin

2015

Portions of Introduction and Chapter 1 © 2014 Elsevier Inc.
All other materials © 2015 Luke Alexander Smith

DEDICATION

I would like to dedicate my work to my wife Stacie Ann Smith. I could not imagine my graduate career without her support, sense of pragmatism, organizational ability, and her love and friendship.

I would also like to dedicate my work to my father, Jack Stuart Smith, who always believed in me unconditionally and fostered my education and imagination throughout my earlier years.

TABLE OF CONTENTS

	Page
LIST OF FIGURES	iv
LIST OF TABLES	v
ACKNOWLEDGMENTS	vi
CURRICULUM VITAE	vii
ABSTRACT OF THE DISSERTATION	ix
Introduction	1
0.1 History and Motivation	1
0.2 Recent Multivariate Value Set Theorems	3
0.3 Main Result	5
0.4 The Newton Polytope	6
0.5 The Degree Matrix and Comparison of Constants	8
1 Essential Theorems and Concepts	10
1.1 Single Variable Value Set	10
1.2 From Single Variable to Multivariable	12
1.3 Restatement of First Polytope Bound and Proof	13
2 Refining Cardinality Bounds	17
2.1 A Method to Improve Prior Proofs	17
2.2 Alternate Degree Bound Proof	22
2.3 Improved Newton Polytope Bound	25
2.4 Improved Integral Dilation Bound	27
3 Conclusion	29
3.1 Analysis of Cardinality Bounds	29
3.2 Future Work	30
Bibliography	32

LIST OF FIGURES

		Page
1	The polytopes of $f(x_1, x_2) = x_1 + x_1^3 x_2$ and $h(x_1, x_2) = x_1^4 + x_2^4$, alongside their contractions by $\frac{n}{d} = \frac{2}{4}$. Note that both polynomials are degree 4, $\Delta(f) \cap \mathbb{N}^2 = \{(3, 1)\}$, and $(\frac{2}{4}\Delta(f)) \cap \mathbb{N}^2 = \emptyset$, but $(\frac{2}{4}\Delta(h)) \cap \mathbb{N}^2 = \{(1, 1)\}$. Therefore, $\mu_h = \frac{2}{4} < \mu_f = 1$	9

LIST OF TABLES

	Page
3.1 Examples of polynomials and their rational interpolations over $\mathbb{F}_{2^8}^*$. We denote α to be a multiplicative generator of $\mathbb{F}_{2^8}^*$	31

ACKNOWLEDGMENTS

I would like to express my appreciation and gratitude to the chair of my thesis committee, Professor Daqing Wan, who deepened my appreciation for number theory and provided me the opportunity to explore mathematical questions which sparked my interest. I also owe Daqing Wan for ideas which make up a majority Section 2.1.

I would also like to thank my committee members, Professor Alice Silverberg and Professor Karl Rubin, for their interest in my research and their advice throughout my coursework and my graduate career.

In addition, I thank Elsevier Inc. for permission to include the portions of the Introduction and Chapter 1 which were originally published in their journal *Finite Fields and Their Applications*, Vol. 28.

Personal thanks goes out to Dennis Eichorn, Matt Ketzi, Roger Dellaca and Josh Hill for their guidance throughout the dissertation writing process and for being a part of my brainstorming process.

Financial Support was provided by the University of California, Irvine Mathematics department and Professor Daqing Wan.

Luke Smith

Curriculum Vitae

Education

- June 2015 **Ph.D. Mathematics**, *University of California, Irvine*.
Specialization in Number Theory, advisor Daqing Wan
- March 2012 **M.S. Mathematics**, *University of California, Irvine*.
Specialization in Number Theory
- June 2009 **B.A. Mathematics**, *University of California, San Diego*.
- May 2007 **General Education**, *Mount San Jacinto College, Menifee, CA*.

Teaching Experience

- August 2014 **Instructor**, *University of California, Irvine*.
- Instructor for Integral Calculus during Summer Session II.
- **Responsibilities include:** Preparing and presenting lectures, meeting with teaching assistants to coordinate lecture and discussion sections, setting up gradebook through EEE course management system, managing student homework through WebWork online homework system, composing quizzes, proctoring/grading quizzes and exams, holding weekly office hours for students, corresponding with students through e-mail.
- 2009–March 2015 **Teaching Assistant**, *University of California, Irvine*.
- **Courses taught include:** Differential Calculus (1 quarter), Integral Calculus (3 quarters, 1 summer session), Multivariable Calculus (4 quarters, 1 summer session), Vector Calculus (2 quarters), Linear Algebra and Infinite Series (2 quarters), Linear Algebra (2 quarters, 1 summer session), Differential Equations (3 quarters, 1 summer session), Introduction to Abstract Math (1 summer session), Abstract Algebra (1 quarter), Real Analysis (1 quarter).
- **Responsibilities include:** Preparing and presenting lectures for discussion sections, holding weekly office hours for students, composing quizzes, proctoring/grading quizzes and exams, corresponding with students through e-mail, working in the math department tutoring center.
- 2012–Present **Private Tutoring**.
-Tutored high school students one-on-one in Precalculus, Statistics, Integral and Differential Calculus, and Infinite Series.

Pedagogical Training

- October 2013 **UCI Teaching, Learning & Technology Center Consultation (TLTC)**, *University of California, Irvine*.
- Corresponded with TLTC for a filming of a Linear Algebra discussion to be critiqued.
- Met with a pedagogical consultant who gave constructive feedback after analyzing the discussion.

Research

Publications

- July 2014 **Polytope Bounds on Multivariate Value Sets**, *Finite Fields and their Applications*, 28 (2014), 132-139.
- Sole author of the manuscript.
- ScienceDirect link: <http://dx.doi.org/10.1016/j.ffa.2014.02.001>

Presentations

- January 2015 **Conveying Group Theoretic Concepts to Middle Schoolers**, *Joint Mathematics Meetings at San Antonio*.
- Presented visual representations of groups used to teach group theory to middle school students at the UCI Mathematics Circle. Joint work with Alessandra Pantano and Casey Kelleher.
- December 2014 **Polytope Bounds on Multivariate Value Sets**, *Chapman University*.
- Presented research from my paper **Polytope Bounds on Multivariate Value Sets**, and presented current work and improvements which have been made since publication.
- April 2013 **Mathematics Graduate Student Colloquium**, *University of California, Irvine*.
- Presented a seminar on research which eventually became **Polytope Bounds on Multivariate Value Sets**. The slideshow for the seminar was made in Beamer.
- October 2012 **Polytope Bounds on Multivariate Value Sets, Ph.D Advancement**, *University of California, Irvine*.
- Presented my results from a preprint version of my paper **Polytope Bounds on Multivariate Value Sets** for my Ph.D advancement.

Research Interests

Number Theory.

- Finite Fields, Newton Polytopes, p -adic liftings, Witt Vectors.

Service

- November 2014 - **Activity Facilitator**, *UCI Mathematics Circle*, University of California, Irvine, CA.
Present - Developed mathematics activity to introduce middle school students to group theory concepts.
- Led middle school students in a visual discussion of groups using properties of regular polygons and regular stars.
- Assisted faculty and graduate students in developing and facilitating their own mathematics activities.
- Math Circle webpage: <http://www.math.uci.edu/~mathcircle/>
- August 2014 **Volunteer**, *Math Fair: at the Square Root of Fun! Held by MIND Research Institute*, Bren Events Center, Irvine, CA.
- Demonstrated interesting properties of conic sections in the UCurves exhibit through interactive exploration.
- Demonstrations used 3D-printed manipulatives, lasers, and a parabolic microphone.
- Event webpage: <http://www.mindresearch.net/mathfair/>
- 2013–Present **Referee**, *Journal of Number Theory; Finite Fields and Their Applications*.

Awards and Fellowships

- August 2014 **Associate Mathematician**, *MIND Research Institute*, Irvine, CA.
- Present - Consults with team of mathematicians and content creators to design mathematics games for K-12 students.
- Games are designed to teach without words, both to overcome language barriers and encourage students to experiment and think critically.
- Develops activities for Family Math Nights and facilitates these events by fostering mathematical conversation between children and parents.

ABSTRACT OF THE DISSERTATION

Refining Multivariate Value Set Bounds

By

Luke Alexander Smith

Doctor of Philosophy in Mathematics

University of California, Irvine, 2015

Professor Daqing Wan, Chair

Over finite fields, if the image of a polynomial map is not the entire field, then its cardinality can be bounded above by a significantly smaller value. Earlier results bound the cardinality of the value set using the degree of the polynomial, but more recent results make use of the powers of all monomials.

In this paper, we explore the geometric properties of the Newton polytope and show how they allow for tighter upper bounds on the cardinality of the multivariate value set. We then explore a method which allows for even stronger upper bounds, regardless of whether one uses the multivariate degree or the Newton polytope to bound the value set. Effectively, this provides an alternate proof of Kosters' degree bound, an improved Newton polytope-based bound, and an improvement of a degree matrix-based result given by Zan and Cao.

Introduction

0.1 History and Motivation

For a given polynomial $f(x)$ over a finite field \mathbb{F}_q , let $V_f := \text{Im}(f)$ denote the value set of f . Determining the cardinality and structure of the value set is a problem with a rich history and wide variety of uses in number theory, algebraic geometry, coding theory and cryptography.

Relevant to this paper are theorems which provide upper bounds on the cardinality of our value set when $f(x)$ is not a permutation polynomial.¹ Let $f(x) \in \mathbb{F}_q[x]$ be a single variable polynomial of degree $d > 0$ with $|V_f| < q$. Using the Chebotarev density theorem over rational function fields, S. D. Cohen proved in [6] that there is a finite set of rational numbers $T_d \subset [0, 1]$ (depending on degree d) such that

$$|V_f| = c_f q + O_d(\sqrt{q}) \tag{1}$$

¹Permutation polynomials have also been studied extensively in literature, in view of their application to cryptography and combinatorics. For more information about other ways value sets have been studied historically, please refer to [10].

for some $c_f \in T_d$ depending on $\text{Gal}(f(x) - t)/\mathbb{F}_q(t)$ and $\text{Gal}(f(x) - t)/\overline{\mathbb{F}_q}(t)$. Guralnick and Wan refine this in [9], proving that for $\gcd(d, q) = 1$ and $|V_f| < q$, $|V_f| \leq \frac{47}{63}q + O_d(\sqrt{q})$. In addition, Mullen conjectured the bound

$$|V_f| \leq q - \frac{q-1}{d} \tag{2}$$

for non-permutation polynomials. This was proven by Wan, Shiue and Chen in [19] using p -adic liftings, but Turnwald later averted the use of liftings with a clever proof in [16] using elementary symmetric polynomials. This bound was also proven sharp for any finite field by Cusick and Müller (for $f(x) = (x+1)x^{q-1} \in \mathbb{F}_{q^k}[x]$, $|V_f| = q^k - \frac{q^k-1}{q}$ for all integers k , see [7]). For more sharp examples, see [19].

Despite the interest mathematicians have taken in the value set problem, most of the work in this area has been dedicated towards univariate polynomials. However, In the past 25 or so years, the multivariate value set problem has been addressed in a few different forms. It was first addressed by Serre in 1988 [13] over varieties, in connection with Hilbert's irreducibility theorem and the inverse Galois problem. His theorem, alongside results by Fried [8] and by Guralnick and Wan [9] give us upper bounds on our value set which generalize Cohen's result in (1). Though these results bound $|V_f|$ by some fraction of $|\mathbb{F}_q^n|$, it is important to note that the error terms in both results, though well behaved with respect to q , are exponentially large in terms of the degree d of the map.

0.2 Recent Multivariate Value Set Theorems

A recently published paper by Mullen, Wan, and Wang in 2012 [12] gives another bound on the value set of polynomial maps, one with no error terms:

Theorem 0.1. *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial map over the vector space \mathbb{F}_q^n , and let $\deg f = \max_i \deg f_i$.*

$$\text{If } |V_f| < q^n, \text{ then } |V_f| \leq q^n - \min \left\{ q, \frac{n(q-1)}{\deg f} \right\}.$$

Since the time their paper was published, multiple refinements have been made to this theorem.

One approach towards improving Theorem 0.1 is to replace the term $\frac{n(q-1)}{\deg f}$ by using different properties of the polynomial map f . Note that the degree only takes one monomial of f into account, so it is reasonable to expect tighter bounds on $|V_f|$ if we account for every monomial. In my first paper [15], I improved upon theorem 0.1 by generalizing Mullen, Wan, and Wang's p -adic lifting approach and utilizing the Newton polytope $\Delta(f)$ of the polynomial map f . The Newton polytope is constructed using all monomials of f using discrete geometry, meaning it encodes more information than $\deg f$ and allows for a stronger statement to be made:

Theorem 0.2 (Smith [15], 2014). *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial map over the vector space \mathbb{F}_q^n , let $\Delta(f)$ be the Newton polytope of f , and let μ_f*

be a certain constant (defined explicitly later) dependent on $\Delta(f)$.

$$\text{If } |V_f| < q^n, \text{ then } |V_f| \leq q^n - \min\{q, \mu_f \cdot (q - 1)\},$$

Zan and Cao also refine Theorem 0.1 by using the degree matrix D_f of the polynomial map f in order to account for all of the monomials of f . Their approach generalizes the p -adic lifting technique as well and improves upon my statement in [15]:

Theorem 0.3 (Zan, Cao [20], 2014). *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial map over the vector space \mathbb{F}_q^n and let D_f be the degree matrix of f .*

$$\text{If } |V_f| < q^n, \text{ then } |V_f| \leq q^n - \min\{q, \omega_f\},$$

where the constant ω_f (defined explicitly later) depends on D_f .

Overall, each new refinement gives us stronger bounds, i.e. $\omega_f \geq \mu_f \cdot (q - 1) \geq \frac{n}{\deg f}(q - 1)$ (see [1] and [20]). In addition, in the univariate case, it has been shown that there are instances when ω_f is strictly larger than $\frac{q-1}{\deg f}$ (as opposed to μ_f always being equal to $\frac{1}{\deg f}$ when $n = 1$). However, since each of these bounds are of the form $|V_f| \leq q^n - \min\{C_f, q\}$ with C_f dependent on the theorem, we are limited to removing at most q elements from these cardinality bounds.

Another type of improvement on theorem 0.1 removes this dependence on subtracting the minimum of two constants. Though still dependent on the polynomial map degree, a theorem by Kusters allows for a stronger bound whenever $n > \deg f$:

Theorem 0.4 (Kosters [11], 2014). *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial map over the vector space \mathbb{F}_q^n , and let $\deg f = \max_i \deg f_i$.*

$$\text{If } |V_f| < q^n, \text{ then } |V_f| \leq q^n - \frac{n(q-1)}{\deg f}.$$

In order to achieve this result, Kosters completely averted the use of p -adic liftings, instead using a method more akin to Turnwald's univariate proof in [16].

0.3 Main Result

In this paper, I will refine these multivariate value set bounds even further, removing the minimum condition from theorems 0.2 and 0.3, ultimately proving the following theorem:

Theorem 0.5. *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial map over the vector space \mathbb{F}_q^n and let D_f be the degree matrix of f .*

$$\text{If } |V_f| < q^n, \text{ then } |V_f| \leq q^n - \omega_f,$$

where the constant ω_f depends on D_f .

To properly convey the significance of this bound in relation to prior bounds, we will describe the Newton polytope in Section 0.4 and the degree matrix in Section 0.5. We will also define the constants associated with these objects and connections between the two. This manuscript will also contain portions of my work in [15] which are relevant to the proofs

of theorems in Chapter 2 as well as a proof of the main theorem in [15] to highlight the difference in techniques used.

0.4 The Newton Polytope

Let F be an arbitrary field and let $h \in F[x_1, \dots, x_n]$. If we write h in the form

$$h(x_1, \dots, x_n) = \sum_{j=1}^m a_j X^{D_j}, \quad a_j \in F^* \tag{3}$$

where

$$D_j = (d_{1j}, \dots, d_{nj}) \in \mathbb{Z}_{\geq 0}^n, \quad X^{D_j} = x_1^{d_{1j}} \dots x_n^{d_{nj}}, \tag{4}$$

then we have the following definition:

Definition 0.1 (Newton polytope). *The Newton polytope of polynomial $h \in F[x_1, \dots, x_n]$, $\Delta(h)$, is the convex closure of the set $\{D_1, \dots, D_m\} \cup \{(0, \dots, 0)\}$ in \mathbb{R}^n .*

Geometric properties of the Newton polytope, such as its dilation by $k \in \mathbb{R}$, its volume or its decomposition into other polytopes via Minkowski Sum, are useful tools in discerning properties of their associated polynomials. For more information, see [3], [18], and [17].

The significance of the Newton polytope to the multivariate value set problem comes from the definition of the following quantity:

Definition 0.2 (Minimal dilation factor μ_h). *Let F be a field, let $h \in F[x_1, \dots, x_n]$, and let $\Delta(h)$ be the Newton polytope of h .*

$$\mu_h := \inf\{k \in \mathbb{R}_{>0} \mid k\Delta(h) \cap \mathbb{N}^n \neq \emptyset\}.$$

In other words, μ_h is the infimum of all positive real numbers k such that the dilation of $\Delta(h)$ by k contains a lattice point with strictly positive coordinates, and we define $\mu_h = \infty$ if such a dilation does not exist. For our purposes, since the vertices of our polytopes have integer coordinates, μ_h will always be finite and rational so long as we consider h which is not a polynomial in some proper subset of $\{x_1, \dots, x_n\}$. If h is polynomial in a proper subset of $\{x_1, \dots, x_n\}$, then we may make a linear change of variables $\{z_1, \dots, z_\nu\}$, $\nu < n$, which allows us to consider $\Delta(h(z_1, \dots, z_\nu)) \subset \mathbb{R}^\nu$, where μ_h will be finite.

The quantity μ_f is used by Adolphson and Sperber [1] to put a lower bound on the q -adic valuation ord_q of the number of \mathbb{F}_q -rational points on a variety V , $N(V)$, over \mathbb{F}_q . Namely, let $V = Z(f_1, \dots, f_m)$ be the vanishing set of f_1, \dots, f_m , where $f_i \in \mathbb{F}_q[x_1, \dots, x_n]$. If the collection of polynomials f_1, \dots, f_m is not polynomial in some proper subset of x_1, \dots, x_n , then we have for $f(x_1, \dots, x_n, y_1, \dots, y_m) = f_1(x_1, \dots, x_n)y_1 + \dots + f_m(x_1, \dots, x_n)y_m$,

$$\text{ord}_q(N(V)) \geq \mu_f - m.$$

Note that in the above definitions, the multivariate polynomial h maps the vector space F^n into its base field F . However, for the value set problem, we are interested in studying the polynomial vector $f : \mathbb{F}_q^n \rightarrow \mathbb{F}_q^n$. Fortunately, the definitions we have developed in this

section can be extended to polynomial vectors. If we denote the support of h by $\Gamma(h) := \{D_1, \dots, D_m\}$, then we define $\Delta(f)$ to be the convex closure of $\Gamma(f_1) \cup \dots \cup \Gamma(f_n) \cup \{(0, \dots, 0)\}$ in \mathbb{R}^n .

0.5 The Degree Matrix and Comparison of Constants

For our multivariate polynomial h as in Section 0.4, we define the $n \times m$ degree matrix of h , $D_h := (D_1, \dots, D_m) \in \mathbb{Z}_{\geq 0}^{n \times m}$. The degree matrix has been used by Cao and his collaborators in [2], [4], and [5] in rational point counting and p -adic estimates. In relation to the value set problem, Zan and Cao use the degree matrix in [20] as a succinct way of keeping track of the exponent vectors D_j that does not explicitly rely on a geometry. Using this, they define the following invariant of h .

Definition 0.3 (Integral dilation factor ω_h). *Let F be a field, let $h \in F[x_1, \dots, x_n]$ be as in equation (3).*

$$\omega_h := \min \left\{ \sum_{j=1}^m k_j \mid k_j \in \{0, 1, \dots, q-1\}, \sum_{j=1}^m k_j D_j \in (q-1)\mathbb{N}^n \right\}. \quad (5)$$

This constant can be thought of as the minimal number of exponent vectors (up to $q-1$ duplicates of each) needed to be summed together to reach a lattice point where all coordinates are positive multiples of $q-1$. Again, so long as h is not polynomial in some proper subset of $\{x_1, \dots, x_n\}$, ω_h will always exist.

Though ω_f and μ_f may seem different by their definitions, a lemma in [3] gives us that

$$\mu_f = \min \left\{ \sum_{j=1}^m \alpha_j \mid \alpha_j \in \mathbb{Q}_{\geq 0}, \sum_{j=1}^m \alpha_j D_j \in \mathbb{N}^n \right\}. \quad (6)$$

Intuitively, studying the dilation of $\Delta(f)$ is equivalent to studying linear combinations of the exponent vectors geometrically. Because of similarity, we can use both D_f and $\Delta(f)$ to study μ_f and ω_f .²

In fact, because of this similarity, we have a direct comparison of the two terms proven by [20]. This, alongside a result of Adolphson and Sperber [1], gives us the following inequalities:

Lemma 0.6. $\omega_f \geq \mu_f(q - 1) \geq \frac{n(q-1)}{d}$.

Not only does ω_f provide a better value set bound for nonpermutation polynomials, but [20] gives sharp examples which improve previously known univariate bounds. For an illustration of the proof by Adolphson and Sperber in two dimensions, please refer to Figure 1 given below.




Figure 1: The polytopes of $f(x_1, x_2) = x_1 + x_1^3 x_2$ and $h(x_1, x_2) = x_1^4 + x_2^4$, alongside their contractions by $\frac{n}{d} = \frac{2}{4}$. Note that both polynomials are degree 4, $\Delta(f) \cap \mathbb{N}^2 = \{(3, 1)\}$, and $(\frac{2}{4}\Delta(f)) \cap \mathbb{N}^2 = \emptyset$, but $(\frac{2}{4}\Delta(h)) \cap \mathbb{N}^2 = \{(1, 1)\}$. Therefore, $\mu_h = \frac{2}{4} < \mu_f = 1$.

²Theorems in Chapter 2 which are dependent on these constants both use $\Delta(f)$ in the proofs given.

Chapter 1

Essential Theorems and Concepts

1.1 Single Variable Value Set

To provide insight towards the proof of our main result, we will investigate upper bounds of $|V_f|$ for the case when f is a single variable polynomial. Parts of this proof will generalize to the multivariate case.

Theorem 1.1. *Let $f(x) \in \mathbb{F}_q[x]$ be a single variable polynomial of degree $d > 0$. If $|V_f| < q$, then*

$$|V_f| \leq q - \frac{q-1}{d}.$$

The proof of this theorem relies on the following definition:

Definition 1.1 (The quantity $U(f)$). *Let \mathbb{Z}_q denote the ring of p -adic integers with uniformizer p and residue field \mathbb{F}_q . Also let $\tilde{f}(x) \in \mathbb{Z}_q[x]$ be the lifting of f taking coefficients from the Teichmüller lifting $L_q \subset \mathbb{Z}_q$ of \mathbb{F}_q . Then we define $U(f)$ to be the smallest positive*

integer k such that the sum

$$S_k(f) := \sum_{x \in L_q} \tilde{f}(x)^k \not\equiv 0 \pmod{pk}.$$

By taking into account the following sum,

$$\sum_{x \in L_q} x^k = \begin{cases} 0, & q-1 \nmid k, \\ q-1, & q-1 \mid k, k \neq 0, \\ q, & k = 0, \end{cases} \quad (1.1)$$

and remembering that we are only summing over a finite number of terms, we have that, for f not identically zero, $\frac{q-1}{d} \leq U(f)$. We also have that if f is a permutation polynomial, then $S_k(f) = S_k(x) = \sum_{x \in L_q} x^k$, implying $U(f) = q-1$. The fact that $U(f)$ exists for all nonpermutation polynomials as well is a corollary of lemma 1.2. Overall, the lemma and the above argument give us that

$$\frac{q-1}{d} \leq U(f) \leq q-1.$$

Theorem 1.1 also follows directly from the lemma 1.2:

Lemma 1.2. *If $|V_f| < q$, then*

$$|V_f| \leq q - U(f).$$

The proof of this result is given by Wan, Shiue, and Chen in [19], and their paper also includes more details regarding this lemma. Mullen, Wan, and Wang [12] also describe an alternate proof of this lemma presented to them by Lenstra through private communication.

1.2 From Single Variable to Multivariable

Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial vector, and note $\deg f = \max_i \{\deg f_i\}$. This maps the vector space \mathbb{F}_q^n to itself. Now, take a basis e_1, \dots, e_n of \mathbb{F}_{q^n} over \mathbb{F}_q . Denote $x = x_1e_1 + \dots + x_ne_n$ and define

$$g(x) := f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n)e_n.$$

In this way, we can think of the function g as a non-constant univariate polynomial map from the finite field \mathbb{F}_{q^n} to itself. Even better, we have the equality $|V_f| = |g(\mathbb{F}_{q^n})|$. Therefore, using Lemma 1.2, we know

$$\text{if } |V_f| < q^n, \text{ then } |V_f| \leq q^n - U(g),$$

where g is viewed as a univariate polynomial.

Unfortunately, as a univariate polynomial, we do not have good control of the univariate degree of g in relation to the multivariate degree of f . Even if one were to construct a closed form for $g(x)$ using methods such as Lagrange Interpolation, the degree of g would likely be high enough as to make the resulting upper bound on $|V_f|$ trivial. Because of these issues

with the degree of g , we cannot use the bounds from the previous section directly, and must rely on another method to bound $U(g)$.

Previously, we introduced $g(x)$ as a univariate polynomial. However, using a basis e_1, \dots, e_n of \mathbb{F}_{q^n} over \mathbb{F}_q as before, we can also define a multivariate polynomial

$$g(x_1, \dots, x_n) := f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n)e_n$$

mapping the vector space \mathbb{F}_q^n into the field \mathbb{F}_{q^n} . In this sense, g as a multivariate polynomial shares some important properties with f as a polynomial vector, such as the fact that $\deg(g) = \max_i \{\deg f_i\}$. Whereas the paper by Mullen, Wan, and Wang determine a bound for $U(g)$ relying on the multivariate degree of f , in this paper we will use the Newton polytope of the multivariate polynomial $g(x_1, \dots, x_n)$ to improve upon these bounds. With this in mind, we define $\Delta(f) := \Delta(g(x_1, \dots, x_n))$, $\mu_f := \mu_{g(x_1, \dots, x_n)}$, and prove the original polytope bound in [15].

1.3 Restatement of First Polytope Bound and Proof

Theorem 1.3. *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial vector over the vector space \mathbb{F}_q^n . If $|V_f| < q^n$, then*

$$|V_f| \leq q^n - \min\{q, \mu_f(q - 1)\}.$$

Proof. First, construct g from our polynomial vector f , as we did in Section 1.2. Viewing g as a univariate polynomial $g(x)$, we are allowed to apply Lemma 1.2 to bound $|V_f|$ using $U(g)$. We then consider g as multivariate $g(x_1, \dots, x_n)$, which allows us to define $\Delta(g)$ and μ_g . Noting that $\Delta(f) = \Delta(g)$ and $\mu_f = \mu_g$ by our definition in Section 1.2, it suffices to prove the following lemma on $U(g)$: □

Lemma 1.4. $U(g) \geq \min\{\mu_f(q-1), q\}$.

Proof. Assume the coefficients of $g(x_1, \dots, x_n)$ are lifted to characteristic zero over L_{q^n} , our Teichmüller lifting of \mathbb{F}_{q^n} . Remember that $U(g)$ is defined over univariate polynomials to be the smallest positive integer k such that

$$S_k(g) := \sum_{x \in L_{q^n}} g(x)^k \not\equiv 0 \pmod{pk}.$$

However, using $x = x_1e_1 + \dots + x_ne_n$ as in Section 1.2, we can rewrite $S_k(g)$ in terms of multivariate $g(x_1, \dots, x_n)$. This means $U(g)$ is the smallest positive integer k such that

$$S_k(g) = \sum_{(x_1, \dots, x_n) \in L_q^n} g(x_1, \dots, x_n)^k \not\equiv 0 \pmod{pk}.$$

Let $k \in \mathbb{Z}_{>0}$ be such that $k < \min\{\mu_f(q-1), q\}$. Expand $g(x_1, \dots, x_n)^k = \sum_{j=1}^m a_j X^{V_j}$ as a polynomial in the n variables x_1, \dots, x_n (see (4)). Since $S_k(g)$ is a finite sum, it can be broken up over the monomials of $g(x_1, \dots, x_n)^k$. Therefore, it suffices to prove

$$\sum_{(x_1, \dots, x_n) \in L_q^n} X^{V_j} \equiv 0 \pmod{pk}, \quad 1 \leq j \leq m, \quad m = \# \text{ monomials of } g^k(x_1, \dots, x_n). \quad (1.2)$$

If we denote $\ell_j := \#\{v_{ij}, 1 \leq i \leq n | v_{ij} \neq 0\}$, i.e. ℓ_j denotes the number of nonzero v_{ij} 's with $1 \leq i \leq n$, then we have exactly $n - \ell_j$ zero v_{ij} 's, implying that

$$\sum_{(x_1, \dots, x_n) \in L_q^n} X^{V_j} \equiv 0 \pmod{q^{n-\ell_j}}.$$

Now let v_p denote the p -adic valuation satisfying $v_p(p) = 1$. If the inequality

$$v_p(q)(n - \ell_j) \geq 1 + v_p(k)$$

is satisfied, then (1.2) is true and we are done.

Considering $X^{V_j} = x_1^{v_{1j}} \dots x_n^{v_{nj}}$, the sum on the left side is identically zero if one of the v_{ij} is not divisible by $q - 1$ (see (1.1)). Thus, we shall assume that all v_{ij} 's are divisible by $q - 1$ (Otherwise (1.2) is satisfied and we are done without even using our inequality on k).

Now, the lattice points of g are contained within $\Delta(g)$ by definition, and this implies our lattice points V_j of g^k are contained within $k\Delta(g)$, the dilation of the polytope $\Delta(g)$ by k . But since $(q - 1) | v_{ij}$, we have that $V_j \in (q - 1)\mathbb{Z}_{\geq 0}^n$ as well.

If we further assume that V_j has no zero coordinates, i.e. $\ell_j = n$, this implies

$$\left(\frac{k}{q-1} \Delta(g) \right) \cap \mathbb{Z}_{>0}^n \neq \emptyset.$$

This statement tells us, by the definition of μ_f , that $\frac{k}{q-1} \geq \mu_f$. In other words,

$$k \geq \mu_f(q-1).$$

This contradicts our assumption that $k < \min\{\mu_f(q-1), q\} \leq \mu_f(q-1)$.

Therefore, when $k < \min\{\mu_f(q-1), q\}$, we have that $\ell_j < n$, and $n - \ell_j > 0$. This case, since $k < q$, gives us $q \nmid k$, and

$$1 + v_p(k) \leq v_p(q) \leq v_p(q)(n - \ell_j).$$

This implies that

$$S_k(g) \equiv 0 \pmod{q^{n-\ell_j}} \equiv 0 \pmod{p^{1+v_p(k)}} \equiv 0 \pmod{pk}$$

and we are done. Lemma 1.4 and the main result of [15] are proved. □

Chapter 2

Refining Cardinality Bounds

2.1 A Method to Improve Prior Proofs

One of the major limitations of the use of p -adic liftings in the proof of Lemma 1.4 is that our assumption only allowed us to show $S_k(f) \equiv 0 \pmod{q}$. Indeed, if we immediately split $S_k(f)$ amongst the monomials of the multivariate polynomial $g(x_1, \dots, x_n)^k$, we lose much of the structure and divisibility of each term. Therefore, we will manipulate our summand to leverage a larger p -adic valuation before splitting it into monomials. To do this, we need the following lemma:

Lemma 2.1. *Let x_1, \dots, x_n be in a commutative ring R , and let $e \in \mathbb{N}$. Then*

$$(x_1 + \dots + x_n)^{p^e} = x_1^{p^e} + \dots + x_n^{p^e} + ph_1(x_1^{p^{e-1}}, \dots, x_n^{p^{e-1}}) + p^2h_2(x_1^{p^{e-2}}, \dots, x_n^{p^{e-2}}) \\ + \dots + p^eh_e(x_1, \dots, x_n)$$

where $h_t(x_1^{p^{e-t}}, \dots, x_n^{p^{e-t}}) \in R[x_1, \dots, x_n]$ is such that $\deg h_t(x_1, \dots, x_n) = p^t$.

Proof. We use the multinomial theorem on the left hand side of the above equation.

$$(x_1 + \cdots + x_n)^{p^e} = x_1^{p^e} + \cdots + x_n^{p^e} + \sum_{\substack{a_1 + \cdots + a_n = p^e \\ a_1 \neq p^e, \dots, a_n \neq p^e}} \binom{p^e}{a_1, \dots, a_n} x_1^{a_1} \cdots x_n^{a_n}. \quad (2.1)$$

For simplicity of notation, let

$$A = \binom{p^e}{a_1, \dots, a_n} x_1^{a_1} \cdots x_n^{a_n}.$$

Then the sum in (2.1) can be split as follows:

$$\begin{aligned} \sum_{a_1 + \cdots + a_n = p^e} A &= x_1^{p^e} + \cdots + x_n^{p^e} + \sum_{\substack{a_1 + \cdots + a_n = p^e \\ p^{e-1} \parallel (a_1, \dots, a_n)}} A + \sum_{\substack{a_1 + \cdots + a_n = p^e \\ p^{e-2} \parallel (a_1, \dots, a_n)}} A \\ &\quad + \cdots + \sum_{\substack{a_1 + \cdots + a_n = p^e \\ p \parallel (a_1, \dots, a_n)}} A + \sum_{\substack{a_1 + \cdots + a_n = p^e \\ p \nmid a_\epsilon \text{ for some } \epsilon}} A. \end{aligned}$$

Now, let

$$\sigma_t = \sum_{\substack{a_1 + \cdots + a_n = p^e \\ p^{e-t} \parallel (a_1, \dots, a_n)}} A, \quad 1 \leq t \leq e.$$

If we can show for $1 \leq t \leq e$ that σ_t has the form $p^t h_t(x_1^{p^{e-t}}, \dots, x_n^{p^{e-t}})$ with $\deg h_t(x_1, \dots, x_n) = p^t$, then the proof is done.

Notice that the summand A always has degree $a_1 + \cdots + a_n = p^e$, which means $\deg \sigma_t = p^e$. Since $p^{e-t} \mid a_\epsilon$ for all ϵ between 1 and n , we know that σ_t has the form $\tau_t(x_1^{p^{e-t}}, \dots, x_n^{p^{e-t}}) \in R[x_1, \dots, x_n]$ and $\deg \tau_t(x_1, \dots, x_n) = \frac{p^e}{p^{e-t}} = p^t$.

The fact that $p^t | \binom{p^e}{a_1, \dots, a_n}$ under the conditions that $p^{e-t} | (a_1, \dots, a_n)$ has an elegant proof by Singmaster in [14]. Therefore, we have that $p^t | \tau_t(x_1^{p^{e-t}}, \dots, x_n^{p^{e-t}})$. This tells us σ_t has the form $p^t h_t(x_1^{p^{e-t}}, \dots, x_n^{p^{e-t}})$ with $\deg h_t(x_1, \dots, x_n) = p^t$, and thus the lemma is proved. □

Let f be a polynomial map over \mathbb{F}_q^n , $\text{char } \mathbb{F}_q = p$. Also let e_1, \dots, e_n be a basis of the field \mathbb{F}_{q^n} over \mathbb{F}_q , and let $x = x_1 e_1 + \dots + x_n e_n$ as before in Section 1.2, allowing for the identification of a polynomial map $f(x_1, \dots, x_n) = ((f_1(x_1, \dots, x_n), \dots, (f_n(x_1, \dots, x_n)))$ with the multivariate polynomial $f(x_1, \dots, x_n) = f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n)e_n$ or the univariate polynomial $f(x)$. Also let $S_k(f)$ and $U(f)$ be as in Section 1.1. To improve upon the p -adic lifting method, we will apply Lemma 2.1 to $f(x_1, \dots, x_n)^k$, split $S_k(f)$ amongst these polynomials, and then split the summand polynomials further into monomials.

Write $k = p^e k_1$ with $p \nmid k_1$. For simplicity of notation, assume f has already been lifted with coefficients in L_{q^n} . Then by Lemma 2.1, there exists polynomials $F_0, \dots, F_e \in \mathbb{F}_{q^n}[x_1, \dots, x_n]$ such that

$$\begin{aligned} (f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n)e_n)^{p^e} &= F_0(x_1^{p^e}, \dots, x_n^{p^e}) + pF_1(x_1^{p^{e-1}}, \dots, x_n^{p^{e-1}}) + \dots \\ &\quad + p^e F_e(x_1, \dots, x_n), \end{aligned}$$

where $\deg F_t(x_1, \dots, x_n) \leq dp^t$. This means that

$$\begin{aligned} (f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n))^k &= \left(F_0(x_1^{p^e}, \dots, x_n^{p^e}) + pF_1(x_1^{p^{e-1}}, \dots, x_n^{p^{e-1}}) + \dots \right. \\ &\quad \left. + p^e F_e(x_1, \dots, x_n) \right)^{k_1} \\ &= \sum_{b_0 + \dots + b_e = k_1} \binom{k_1}{b_0, \dots, b_e} p^{b_1 + 2b_2 + \dots + eb_e} F_0(x_1^{p^e}, \dots, x_n^{p^e})^{b_0} \dots F_e(x_1, \dots, x_n)^{b_e}. \end{aligned}$$

Now for fixed b_0, \dots, b_e , let λ be the positive integer such that $b_\lambda \neq 0, b_{\lambda+1} = \dots = b_e = 0$, and let $y_i = x_i^{p^{e-\lambda}}$. This means we can reduce the power and degree of our summand polynomials in the following way:

$$F_0(x_1^{p^e}, \dots, x_n^{p^e})^{b_0} \dots F_\lambda(x_1^{p^{e-\lambda}}, \dots, x_n^{p^{e-\lambda}})^{b_\lambda} = F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \dots F_\lambda(y_1, \dots, y_n)^{b_\lambda}. \quad (2.2)$$

Note that each term may have a different substitution, but we may split $S_k(f)$ amongst each summand to bound the p -divisibility of the entire sum. Using the reduction of $f(x_1, \dots, x_n)^k$ to (2.2), we are given sums of the form

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \dots F_\lambda(y_1, \dots, y_n)^{b_\lambda}. \quad (2.3)$$

Now the fact that $b_\lambda \neq 0$ tells us this sum is divisible by p^λ , i.e. $S_k(f) \equiv 0 \pmod{p^\lambda}$. From here, we must further split this summand product into monomials and determine the p -divisibility of the smaller sums. Let

$$F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \dots F_\lambda(y_1, \dots, y_n)^{b_\lambda} = \sum_{j=1}^m c_j Y^{W_j}, \quad c_j \in \mathbb{F}_{q^n}^*,$$

where

$$W_j = (w_{1j}, \dots, w_{nj}) \in \mathbb{Z}_{\geq 0}^n, \quad Y^{W_j} = y_1^{w_{1j}} \dots y_n^{w_{nj}}. \quad (2.4)$$

This allows the sum in (2.3), and ultimately $S_k(f)$, to be split among the monomials in (2.4) into sums of the form

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} c_j Y^{W_j} = c_j p^{b_1+2b_2+\dots+\lambda b_\lambda} \prod_{i=1}^n \sum_{y_i \in L_q} y_i^{w_{ij}}. \quad (2.5)$$

Let C be an upper bound on k , i.e. $k < C$ (C will depend on which of the theorems in the following sections we are proving). Our goal is to show $C \leq U(f)$ and therefore come up with a nicer bound on $|V_f|$ (thanks to Lemma 1.2). Let v_p be the p -adic valuation with $v_p(p) = 1$, and let ℓ_j be the number of nonzero entries of W_j . We can accomplish our goal by showing the sum in (2.5) is congruent to zero mod $p^\lambda q^{n-\ell_j}$, and that $v_p(p^\lambda q^{n-\ell_j}) \geq v_p(pk)$, i.e.

$$\lambda + (n - \ell_j)v_p(q) \geq e + 1.$$

If this holds true for all monomials, then $S_k(f) \equiv 0 \pmod{pk}$ and $k \leq U(f)$.

Now the sum in (2.5) equals zero if one of the w_{ij} 's is not divisible by $q - 1$, so all that is left is to consider the case when $q - 1 | w_{ij}$ for all i . In this case, since $b_\lambda \neq 0$, and since ℓ_j is the number of nonzero w_{ij} , we have $n - \ell_j$ zero terms, which tells us

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} c_j Y^{W_j} \equiv 0 \pmod{p^\lambda q^{n-\ell_j}}.$$

The above substitution method allows us to refine the recently published results mentioned in Section 0.2, whose proofs simply used the monomials of $f(x_1, \dots, x_n)^k$ directly. These proofs required that $k < q$ to bound the value set, but our proofs do not. The next few sections will show how the added structure our method provides tighter upper bounds on the cardinality of the value set.

2.2 Alternate Degree Bound Proof

Theorem 2.2. *Let f be a polynomial map with $f : \mathbb{F}_q^n \rightarrow \mathbb{F}_q^n$, $\text{char } \mathbb{F}_q = p$,*

$$f(x_1, \dots, x_n) = f_1(x_1, \dots, x_n)e_1 + \dots + f_n(x_1, \dots, x_n),$$

and $d = \max_i \deg f_i$. If $|V_f| < q^n$, then

$$V_f \leq q - \frac{n(q-1)}{d}.$$

Note that this theorem was proven by Kosters in [11], but we provide an alternate proof using the method outlined in Section 2.1.

Proof. If we can show that, for $1 \leq k < \frac{n(q-1)}{d}$ and $k = p^e k_1$ with $p \nmid k_1$,

$$S_k(f) := \sum_{x \in L_{q^n}} \tilde{f}(x)^k = \sum_{x_1, \dots, x_n \in L_q} \left(\tilde{f}_1(x_1, \dots, x_n)\tilde{e}_1 + \dots + \tilde{f}_n(x_1, \dots, x_n)\tilde{e}_n \right)^k \equiv 0 \pmod{pk},$$

then $U(f) \geq \frac{n(q-1)}{d}$ and we are done by Lemma 1.2.

For simplicity of notation, assume f is already lifted to characteristic zero over L_{q^n} . Split $S_k(f)$ into sums of the form (2.3). Notice that, by our substitution and Lemma 2.1, the degree of the summand $F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \dots F_\lambda(y_1, \dots, y_n)^{b_\lambda}$ in (y_1, \dots, y_n) is bounded above by $dp^\lambda a_0 + dp^1 p^{\lambda-1} + \dots + dp^\lambda a_\lambda = dp^\lambda k_1$. When we further split these sums into sums of the form (2.5), we have that

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} c_j Y^{W_j} \equiv 0 \pmod{p^\lambda q^{n-\ell_j}}.$$

Since this sum equals 0 if one of the w_{ij} 's is not divisible by $q - 1$, assume $q - 1 | w_{ij}$ for all i . Using our degree bound we have that

$$(q - 1)\ell_j \leq w_{1j} + \cdots + w_{nj} \leq dp^\lambda k_1,$$

or $\ell_j \leq \left\lfloor \frac{dp^\lambda k_1}{q-1} \right\rfloor$. If we can show that

$$v_p \left(p^\lambda q^{n - \left\lfloor \frac{dk_1 p^\lambda}{q-1} \right\rfloor} \right) \geq v_p(pk),$$

Then we are done. In other words, we must show

$$\lambda + \left(n - \left\lfloor \frac{dk_1 p^\lambda}{q-1} \right\rfloor \right) v_p(q) \geq e + 1.$$

Mullen, Wan, and Wang [12] proved a similar inequality,

$$\left(n - \left\lfloor \frac{dk_1 p^e}{q-1} \right\rfloor \right) v_p(q) \geq e + 1.$$

However, their proof only holds in the cases when:

1. $n \leq d$
2. $n > d$ and $k < q$.

Since their inequality implies ours, we may assume that $n > d$ and $k \geq q$. Let $r = e - \lambda$.

Then it suffices to show that

$$\left(n - \left\lfloor \frac{dk}{p^r(q-1)} \right\rfloor \right) v_p(q) \geq r + 1, \quad 0 \leq r \leq e.$$

Note that $k < \frac{n(q-1)}{d} \leq q^n$ implies that $\frac{dk}{q-1} < n$, which is equivalent to $\frac{dk}{p^r(q-1)} < \frac{n}{p^r}$.

In the case that $n \leq p^r$, we have that $\frac{dk}{p^r(q-1)} < 1$, which implies

$$\left(n - \left\lfloor \frac{dk}{p^r(q-1)} \right\rfloor \right) v_p(q) = nv_p(q) > v_p(k) = e \geq r.$$

In other words,

$$\left(n - \left\lfloor \frac{dk}{p^r(q-1)} \right\rfloor \right) v_p(q) \geq r + 1.$$

Now let us examine the case when $n > p^r$. For $r = 0$,

$$\left(n - \left\lfloor \frac{dk}{(q-1)} \right\rfloor \right) v_p(q) \geq v_p(q) \geq 1.$$

For $r = 1$ and $p = 2$ (implying $n > 2$),

$$\left(n - \left\lfloor \frac{dk}{2(q-1)} \right\rfloor \right) v_p(q) \geq \left(n - \left\lfloor \frac{n}{2} \right\rfloor \right) v_p(q) = \left\lceil \frac{n}{2} \right\rceil v_p(q) \geq 2v_p(q) \geq 2.$$

And finally, when $r \geq 1$,

$$\begin{aligned} \left(n - \left\lfloor \frac{dk}{p^r(q-1)} \right\rfloor \right) v_p(q) &\geq \left(n - \left\lfloor \frac{n}{p^r} \right\rfloor \right) v_p(q) \geq \left(\frac{n(p^r - 1)}{p^r} \right) v_p(q) \\ &\geq \left(\frac{(p^r + 1)(p^r - 1)}{p^r} \right) v_p(q) = \left(p^r - \frac{1}{p^r} \right) v_p(q). \end{aligned}$$

Note that $p^r - \frac{1}{p^r} \geq r + 1$ for all $r \geq 1$ except for when $r = 1$ and $p = 2$ simultaneously.

□

2.3 Improved Newton Polytope Bound

Theorem 2.3. *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial vector over the vector space \mathbb{F}_q^n . Without loss of generality, suppose f is not polynomial in some subset of $\{x_1, \dots, x_n\}$. Let $\Delta(f)$ be the Newton polytope of f , and let μ_f be the minimal dilation factor associated with $\Delta(f)$. If $|V_f| < q^n$, then*

$$|V_f| \leq q^n - \mu_f(q - 1).$$

Proof. If we can show that, for $1 \leq k < \mu_f(q - 1)$ and $k = p^e k_1$ with $p \nmid k_1$,

$$S_k(f) := \sum_{x \in L_{q^n}} \tilde{f}(x)^k = \sum_{x_1, \dots, x_n \in L_q} \left(\tilde{f}_1(x_1, \dots, x_n) \tilde{e}_1 + \dots + \tilde{f}_n(x_1, \dots, x_n) \tilde{e}_n \right)^k \equiv 0 \pmod{pk},$$

then $U(f) \geq \mu_f(q - 1)$ and we are done by Lemma 1.2.

For simplicity of notation, assume f is already lifted to characteristic zero over L_{q^n} . Split $S_k(f)$ into sums of the form (2.3). Notice that, by our substitution and Lemma 2.1, the exponent vectors of the monomials of the product $F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \cdots F_\lambda(y_1, \dots, y_n)^{b_\lambda}$ are contained in $\frac{k}{p^{e-\lambda}} \Delta(f)$. When we further split these sums into sums of the form (2.5), we have that

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} c_j Y^{W_j} \equiv 0 \pmod{p^\lambda q^{n-\ell_j}}.$$

Since this sum equals 0 if one of the w_{ij} 's is not divisible by $q - 1$, assume $q - 1 | w_{ij}$ for all i . By this assumption, we have that

$$W_j \in \frac{k}{p^{e-\lambda}} \Delta(f) \cap (q - 1) \mathbb{Z}_{\geq 0}^n. \tag{2.6}$$

To further develop this proof, we require additional terminology.

Definition 2.1 (The quantity γ).

$$\gamma := \min \left\{ |S| \left| S \subseteq \{W_1, \dots, W_m\}, \sum_{W_j \in S} W_j \in \mathbb{N}^n \right. \right\}.$$

In other words, γ is the size of smallest subset of the exponent vectors, $\{W_1, \dots, W_m\}$, such that the sum of its elements lie in \mathbb{N}^n . Since $f(x_1, \dots, x_n)$ is not polynomial in some proper subset of $\{x_1, \dots, x_n\}$, we have that the polynomials $F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda}), \dots, F_\lambda(y_1, \dots, y_n)$ are not either. This means γ will exist. Also, assume without loss of generality that W_1, \dots, W_γ satisfy the sum property of γ , i.e.

$$W_1 + \dots + W_\gamma \in \mathbb{N}^n.$$

Using this and (2.6), we have that

$$W_1 + \dots + W_\gamma \in \frac{\gamma k}{p^{e-\lambda}} \Delta(f) \cap (q-1)\mathbb{N}^n, \quad (2.7)$$

which means that $\mu_f \leq \frac{\gamma k}{p^{e-\lambda}(q-1)}$. Reorganizing this, and using our assumption on k at the beginning of the proof, we have $\frac{p^{e-\lambda}}{\gamma} \mu_f (q-1) \leq k < \mu_f (q-1)$, or $p^{e-\lambda} < \gamma$. To make use of this inequality, we have the following lemma:

Lemma 2.4. *For all integers $1 \leq j \leq m$, we have*

$$\gamma - 1 \leq n - \ell_j.$$

Proof. Let W_u be such that $\ell_u \geq \ell_j$ for all $1 \leq j \leq m$. If $\ell_u = n$, then $\gamma = 1$ and we are done. If not, W_u has $n - \ell_u$ components which are zero and we can pick elements from $\{W_1, \dots, W_{u-1}, W_{u+1}, \dots, W_m\}$ to add to W_u until the resulting sum is an element of \mathbb{N}^n . This

implies it is possible to pick $n - \ell_u + 1$ vectors from $\{W_1, \dots, W_m\}$ whose sum will lie in \mathbb{N}^n . By the definition of γ , we must have $\gamma \leq n - \ell_u + 1$. But by our assumption on W_u , this means that $\gamma - 1 \leq n - \ell_j$ for all j . \square

With the help of Lemma 2.4 and (2.7), we have that $p^{e-\lambda} \leq \gamma - 1 \leq n - \ell_j$. If we can show that

$$v_p\left(p^\lambda q^{p^{e-\lambda}}\right) \geq v_p(pk),$$

Then $S_k(f) \equiv 0 \pmod{(pk)}$ and we are done. In other words, if $r = e - \lambda$ we must show

$$p^r v_p(q) \geq r + 1. \tag{2.8}$$

Fortunately, this is true for all primes p and all positive integers r .

\square

2.4 Improved Integral Dilation Bound

Theorem 2.5. *Let $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ be a polynomial vector over the vector space \mathbb{F}_q^n . Without loss of generality, suppose f is not polynomial in some subset of $\{x_1, \dots, x_n\}$. Let ω_f be the integral dilation factor associated with $\Delta(f)$. If $|V_f| < q^n$, then*

$$|V_f| \leq q^n - \omega_f.$$

Proof. If we can show that, for $1 \leq k < \omega_f$ and $k = p^e k_1$ with $p \nmid k_1$,

$$S_k(f) := \sum_{x \in L_{q^n}} \tilde{f}(x)^k = \sum_{x_1, \dots, x_n \in L_q} \left(\tilde{f}_1(x_1, \dots, x_n) \tilde{e}_1 + \dots + \tilde{f}_n(x_1, \dots, x_n) \tilde{e}_n \right)^k \equiv 0 \pmod{pk},$$

then $U(f) \geq \omega_f$ and we are done by Lemma 1.2.

For simplicity of notation, assume f is already lifted to characteristic zero over L_{q^n} . Split $S_k(f)$ into sums of the form (2.3). Notice that, by our substitution and Lemma 2.1, the exponent vectors of the monomials of the product $F_0(y_1^{p^\lambda}, \dots, y_n^{p^\lambda})^{b_0} \dots F_\lambda(y_1, \dots, y_n)^{b_\lambda}$ are contained in $\frac{k}{p^{e-\lambda}} \Delta(f)$. When we further split these sums into sums of the form (2.5), we have that

$$p^{b_1+2b_2+\dots+\lambda b_\lambda} \sum_{y_1, \dots, y_n \in L_q} c_j Y^{W_j} \equiv 0 \pmod{p^\lambda q^{n-\ell_j}}.$$

Since this sum equals 0 if one of the w_{ij} 's is not divisible by $q-1$, assume $q-1 \mid w_{ij}$ for all i . By this assumption, we have that

$$W_j \in \frac{k}{p^{e-\lambda}} \Delta(f) \cap (q-1) \mathbb{Z}_{\geq 0}^n.$$

Now let γ be as in Definition 2.1 and, WLOG, let $W_1 + \dots + W_\gamma \in \mathbb{N}^n$. Then by the definition of γ , we have

$$W_1 + \dots + W_\gamma \in \frac{\gamma k}{p^{e-\lambda}} \Delta(f) \cap (q-1) \mathbb{N}^n,$$

and $\omega_f \leq \frac{\gamma k}{p^{e-\lambda}}$. Using this and Lemma 2.4, we have $\frac{p^{e-\lambda}}{\gamma} \omega_f \leq k < \omega_f$, or $p^{e-\lambda} \leq \gamma - 1 \leq n - \ell_j$.

By this inequality and (2.8), we are done. □

Chapter 3

Conclusion

3.1 Analysis of Cardinality Bounds

Each of the bounds given in Chapter 2 are sharp. Let $N(x_1, \dots, x_{n-1})$ be the field norm of $\mathbb{F}_{q^{n-1}}$ over \mathbb{F}_q . Kusters [11] illustrates that Theorem 2.2 is sharp using the map $f(x_1, \dots, x_n) = (x_1, x_2, \dots, N(x_1, \dots, x_{n-1})x_n)$. Based on this example, we give the following sharp example for Theorems 2.3 and 2.5. Let $h(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, N(x_1, \dots, x_{n-1})^a x_n)$ with a in \mathbb{N} . Because $N(x_1, \dots, x_{n-1})$ is a polynomial containing the monomials $x_1^{n-1}, \dots, x_{n-1}^{n-1}$ with nonzero coefficients, we have that $(a, a, \dots, a, 1) \in \Delta(h)$. This explicitly tells us $\Delta(h) \cap \mathbb{N}^n \neq \emptyset$. We also have for all $V = (v_1, \dots, v_n) \in \Delta(h) \cap \mathbb{N}^n$, $v_n = 1$. This implies $\mu_h = 1$ and $\omega_h = q - 1$. In addition, since the preimage $N^{-1}(0) = \{(0, \dots, 0)\}$, we are given $|V_h| = q^n - (q - 1)$. This example highlights the flexibility granted by the use of constants derived from the Newton polytope, since $\deg h = a(n - 1) + 1$ does not allow for a sharp cardinality bound. This flexibility also gives us more freedom to make substitutions when generating more sharp examples. If $z_1(x), \dots, z_{n-1}(x)$ are univariate permutation polynomials in $\mathbb{F}_q[x]$, then the maps $g(x_1, \dots, x_n) = (z_1(x_1), \dots, z_{n-1}(x_{n-1}), N(x_1, \dots, x_{n-1})^a x_n)$

and $h(z_1(x_1), \dots, z_{n-1}(x_{n-1}), x_n)$ will share the same constants and value set cardinality as $h(x_1, \dots, x_n)$.

Using the constant ω_f also has an advantage when determining bounds on univariate value sets. In this case, since $n = 1$, we have that $\mu_f = \frac{1}{\deg f}$ for all $f \in \mathbb{F}_q[x]$, but Zan and Cao [20] give a sharp example which improves upon this for ω_f . If $f(x) = x^7 + ax \in \mathbb{F}_{19}[x]$ with $a \neq 0, 4, 5, 8, 16, 17$, then it is easy to check that $\omega_f = 6$, $|V_f| = 13 = 19 - \omega_f < 19 - \lceil \frac{1}{7}(18) \rceil = 16$.

Note that, in general, it is not immediately clear how large of an improvement the strongest bound in Theorem 2.5 provides over our bounds in Theorems 2.2 and 2.3. I have addressed in [15] that an effective method for calculating μ_f is not directly clear from the definitions given. However, calculation of ω_f should be much more efficient complexity-wise, since only a finite amount of values need to be checked to determine the minimum value. This quantity of values to check by brute force grows with complexity $O(q^n)$ and is therefore polynomial in q (though exponential in n). Therefore, there is much value in the use of ω_f even when it is equal to $\mu_f \cdot (q - 1)$.

3.2 Future Work

It is important to consider whether the results presented in Chapter 2 apply in more general settings. For instance, there are cases when it is more convenient to use rational interpolated form of a map than its polynomial form, especially when the monomials of the rational interpolation have much smaller degree. Even if we strictly considered Laurent polynomials, where we have $f(x) \in \mathbb{F}_q[x, x^{-1}]$ or the Laurent polynomial map $f(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n))$ with $f_i(x_1, \dots, x_n) \in \mathbb{F}_q[x_1, x_2, \dots, x_n, x_1^{-1}, x_2^{-1}, \dots, x_n^{-1}]$, can we

apply the geometry of the Newton polytope to bound their cardinalities? Would such bounds be any stronger than those obtained by using a polynomial-interpolated form of the map?³

Table 3.1: Examples of polynomials and their rational interpolations over $\mathbb{F}_{2^8}^*$. We denote α to be a multiplicative generator of $\mathbb{F}_{2^8}^*$.

Polynomial Interpolation	Rational Interpolation
$x^{18} + 3x^2 + 1$	N/A
$x^{254} + x^{17} + 1$	$(x^{18} + x + 1)/x$
$x^{254} + x^{253} + x^{30}$	$(x^{32} + x + 1)/x^2$
$(\alpha^6 + \alpha^3 + 1)x^{254} + \dots + (\alpha^6 + \alpha^5)$	$x^{32}/(x^2 + \alpha x + \alpha^7)$
$(\alpha^6 + \alpha^5)x^{254} + \dots + (\alpha^7 + \alpha^6 + \alpha^2)$	$(x^{88} + 1)/(x^2 + x + \alpha^5)$

³Thanks to Matt Ketı for compiling the following table.

Bibliography

- [1] A. Adolphson and S. Sperber. p -adic estimates for exponential sums and the theorem of Chevalley-Waring. *Annales scientifiques de l'cole Normale Suprieure*, 20(4):545–556, 1987.
- [2] W. Cao. Smith normal form of augmented degree matrix and its applications. *Linear Algebra and its Applications*, 431(10):1778 – 1784, 2009.
- [3] W. Cao. Dilation of Newton polytope and p -adic estimate. *Discrete and Computational Geometry*, 45(3):522–528, 2011.
- [4] W. Cao and Q. Sun. On a class of equations with special degrees over finite fields. *Acta Arithmetica*, 130:195–202, 2007.
- [5] J. Chen and W. Cao. Degree matrices and divisibility of exponential sums over finite fields. *Archiv der Mathematik*, 94(5):435–441, 2010.
- [6] S. D. Cohen. The distribution of polynomials over finite fields. *Acta Arithmetica*, 17:255–271, 1970.
- [7] T. W. Cusick and P. Müller. Wan’s bound for value sets of polynomials. In *Proceedings of the Third International Conference on Finite Fields and Applications*, FFA ’95, pages 69–72, New York, NY, USA, 1996. Cambridge University Press.
- [8] M. Fried. On Hilbert’s irreducibility theorem. *Journal of Number Theory*, 6(3):211 – 231, 1974.
- [9] R. Guralnick and D. Wan. Bounds for fixed point free elements in a transitive group and applications to curves over finite fields. *Israel Journal of Mathematics*, 101(1):255–287, 1997.
- [10] J. Hill. Weil image sums (and some related problems), <http://untruth.org/s/p9.html>. 2011.
- [11] M. Kusters. Polynomial maps on vector spaces over a finite field. *Finite Fields and Their Applications*, 31(0):1 – 7, 2015.
- [12] G. L. Mullen, D. Wan, and Q. Wang. Value sets of polynomial maps over finite fields. *The Quarterly Journal of Mathematics*, pages 1191 – 1196, 2012.

- [13] J.-P. Serre. *Topics in Galois Theory*. Jones and Bartlett Publishers, 1992.
- [14] D. Singmaster. Divisibility of binomial and multinomial coefficients by primes and prime powers. *A Collection of Manuscripts Related to the Fibonacci Sequence, 18th Anniversary Volume*, pages 98–114, 1980.
- [15] L. Smith. Polytope bounds on multivariate value sets. *Finite Fields and Their Applications*, 28(0):132 – 139, 2014.
- [16] G. Turnwald. A new criterion for permutation polynomials. *Finite Fields and Their Applications*, 1(1):64 – 82, 1995.
- [17] D. Wan. Variation of p -adic Newton polygons for L-functions of exponential sums. *Asian J. Math.*, 8(3):427–472, 09 2004.
- [18] D. Wan. Lectures on zeta functions over finite fields. *Higher-Dimensional Geometry over Finite Fields*, 16:244–268, 2008.
- [19] D. Wan, P. J.-S. Shiue, and C. S. Chen. Value sets of polynomials over finite fields. *Proceedings of the American Mathematical Society*, 119(3):711–717, 1993.
- [20] H. Zan and W. Cao. Powers of polynomials and bounds of value sets. *Journal of Number Theory*, 143:286 – 292, 2014.