

Lawrence Berkeley National Laboratory

Recent Work

Title

BIS (CYCLOPENTADIENYL)DIMETHYLTHORIUM (IV) [BIS (DIMETHYLDIPHOSPHINO) ETHANE] AND BIS (CYCLOPENTADIENYL) DICHLOROTHORIUM(IV) [BIS (DIMETHYLDIPHOSPHINO) ETHANE] (C₅H₅)₂ (CH₃)₂Th. (CH₃)₂PCH₂CH₂P (CH₃)₂' (C₅H₅)₂Th. (CH₃)₂PCH₂CH₂P(CH₃)₂

Permalink

<https://escholarship.org/uc/item/1dc6m1d2>

Authors

Zalkin, A.
Brennan, J.G.
Andersen, R.A.

Publication Date

1986-09-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

Submitted to Acta Crystallographica


BIS(CYCLOPENTADIENYL)DIMETHYLTHORIUM(IV)
[BIS(DIMETHYLDIPHOSPHINO)ETHANE] AND
BIS(CYCLOPENTADIENYL)DICHLOROTHORIUM(IV)
[BIS(DIMETHYLDIPHOSPHINO)ETHANE]
 $(C_5H_5)_2(CH_3)_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2,$
 $(C_5H_5)_2Cl_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$

A. Zalkin, J.G. Brennan, and R.A. Andersen

September 1986

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*




LBL-22313
e.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Bis(cyclopentadienyl)dimethylthorium(IV)[bis(dimethyldiphosphino)ethane]
and Bis(cyclopentadienyl)dichlorothorium(IV)[bis(dimethyldiphosphino)ethane]




By Allan Zalkin, John G. Brennan & Richard A. Andersen

Materials and Molecular Research Division, Lawrence Berkeley Laboratory and
Department of Chemistry, University of California, Berkeley CA 94720

Abstract. $(\text{C}_5\text{H}_5)_2(\text{CH}_3)_2\text{Th}\cdot(\text{CH}_3)_2\text{PCH}_2\text{CH}_2\text{P}(\text{CH}_3)_2$ (1) $M_r = 542.44$, monoclinic, $P2_1/n$, $a = 18.485(4)$, $b = 14.118(4)$, $c = 8.258(2)$ Å, $\beta = 91.05(2)^\circ$, $V = 2154.7$ Å³, $Z = 4$, $D_x = 1.672$ g cm⁻³, $\lambda(\text{MoK}\alpha) = 0.71073$ Å, $\mu = 73.0$ cm⁻¹, $F(000) = 1040$, $T = 296$ K, $R = 0.026$ for 2521 unique reflections with $F^2 > 2\sigma(F^2)$. Distances (Å) are: Th-C(Cp) 2.84 ± 0.03 ; Th-C(methyl) 2.562(8), 2.583(7); Th-P 3.146(2), 3.147(2) Å; Th-Cp 2.59, 2.57 Å.

$(\text{C}_5\text{H}_5)_2\text{Cl}_2\text{Th}\cdot(\text{CH}_3)_2\text{PCH}_2\text{CH}_2\text{P}(\text{CH}_3)_2$ (2) $M_r = 583.28$, monoclinic, $P2_1/n$, $a = 18.268(2)$, $b = 14.237(6)$, $c = 7.996(3)$ Å, $\beta = 92.27(4)^\circ$, $V = 2078.0$ Å³, $Z = 4$, $D_x = 1.864$ g cm⁻³, $\lambda(\text{MoK}\alpha) = 0.71073$ Å, $\mu = 78.3$ cm⁻¹, $F(000) = 1104$, $T = 296$ K, $R = 0.020$ for 2361 unique reflections with $I > \sigma(I)$. Distances (Å) are: Th-C(Cp) 2.80 ± 0.02 ; Th-Cl 2.707(2), 2.708(2); Th-P 3.121(2), 3.122(2) Å; Th-Cp 2.56, 2.54 Å. The structures of (1) and (2) are crystallographically isomorphous and were done to study their six-coordinate stereochemistry. The cyclopentadienyl rings are trans to the bidentate ligand.

Introduction. Compounds of the type $(C_5H_5)_2M(X)_2(L)$ where M is thorium or uranium, X is an anionic ligand such as halide or alkyl, and L is a neutral, bidentate phosphine ligand were prepared in order to examine the details of inter- and intra-molecular ligand exchange in solution. The stereochemistry of these six coordinate compounds (defining the centroid of the cyclopentadienyl ring as occupying a coordination site) in the solid state was essential since two idealized geometries are possible for $MX_2Y_2^-$ (bidentate ligand), that is either the X ligands or the Y ligands are trans to the bidentate ligand. For comparison the X-ray structures of $(C_5H_5)_2Th(X)_2 \cdot [(CH_3)_2PCH_2CH_2P(CH_3)_2]$ where X = Cl, CH₃ (this paper) and where X = CH₂C₆H₅ (Zalkin, Brennan, Andersen, 1986) are described.



Experimental. The dimethyl complex (1) was made by the reaction of (2) with methyl lithium at $-45^\circ C$ (Brennan, 1985). Colorless crystals suitable for X-ray studies were picked from the crystals obtained by crystallization from a toluene:pentane (1:4) solution at $-70^\circ C$. The dichloro complex (2) was synthesized from the reaction of sodium cyclopentadienide with $ThCl_4 \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$ in a tetrahydrofuran solution at $-70^\circ C$ and crystallized from toluene. Crystals suitable for X-ray studies were grown from toluene:diethyl ether (6:4) at $-20^\circ C$. The air sensitive crystals were sealed inside quartz capillaries under argon. Crystal 0.19 x 0.25 x 0.32 mm with 6 faces (1), 0.22 x 0.25 x 0.37 mm with 12 faces (2); modified Picker automatic diffractometer (1), Enraf-Nonius CAD-4 (2); graphite monochromator; cell dimensions from 20 reflections, $20^\circ < 2\theta < 34^\circ$ (1), 24

reflections, $20^\circ < 2\theta < 32^\circ$ (2); analytical absorption correction, range 1.47 to 1.83 (1), 3.26 to 4.66; max. $(\sin\theta)/\lambda = 0.60 \text{ \AA}^{-1}$ (1), 0.54 \AA^{-1} (2); h -22 to 22, k 0 to 16, l -9 to 9 (1); h -19 to 19, k 0 to 15, l 0 to 8 (2); three standard reflections, [0.4%, 0.4%, 0.3% (1)], [1.3%, 1.7%, 1.2% (2)] variation in standards intensities from average, intensities adjusted isotropically; 7611 (1), 2929 [$I > \sigma(I)$] (2) data, 3817 (1), 2361 (2) unique, $R_{\text{int}} = 0.029$ (1), 0.020 ($hk0, \bar{h}k0$) (2); structure solved by Patterson and Fourier methods; refined on F, 190 (1), 188 (2) parameters; 26 hydrogen positions (1) and 30 hydrogen positions (including 4 disordered positions) (2) in calculated locations with fixed isotropic thermal parameters; anisotropic thermal parameters for non-hydrogen atoms except for disordered atoms in (2); $R = 0.060$ for 3817 data (1); $R = 0.026$ for 2521 reflections for which $F^2 > 2\sigma(F^2)$ (1); $R = 0.020$ for 2361 reflections for which $I > \sigma(I)$ (2); $wR = 0.028$ (1), 0.027 (2); $S = 1.1$ (1), 1.6 (2); $w = [\sigma(F)]^{-2}$, $p = 0.035$ (1), 0.020 (2) in calc. of $\sigma(F^2)$; max (shift/ σ) < 0.004 ; empirical extinction correction, $F_{\text{corr}} = (1 + kI)$, $k = 1.7 \times 10^{-9}$ (1), 1.9×10^{-7} (2); max. and min. of ΔF synthesis 0.9 and -0.8 e \AA^{-3} (1), 0.4 and -0.5 e \AA^{-3} (2); atomic f for neutral Th, Cl, P, and C, and spherical bonded H from International Tables (1974); local unpublished programs and ORTEP (Johnson, 1965). Large anisotropic thermal parameters for C(13) and C(14) as well as a shortened C-C distance between these atoms suggests disorder in the backbone of the dimethylphosphinoethane ligand. In the dichloro complex (2) these atoms were treated as half atoms in two distinct positions and refined with restrained distances (Waser, 1963) (P-C $1.86(1)$; C-C $1.54(1) \text{ \AA}$) and with isotropic thermal parameters, whereas in (1) these carbon atoms were refined with anisotropic thermal parameters; either way of treating these atoms resulted in the same agreement factors. The temperature factors

of the phosphorus atoms are a little larger than and about as isotropic as the thorium atoms, which suggests that they are not disordered, however, the terminal phosphorus methyl group carbon atoms have large anisotropic thermal parameters and are likely disordered in a synchronous fashion with C(14) and C(15) through the pivotal phosphorus atom.

Atomic parameters are listed in Table 1, * and distances and angles are listed in Table 2. Figures 1 & 2 show the molecules and numbering schemes.

* Lists of structure factors, anisotropic thermal parameters, calculated hydrogen positions, distances and angles, least-squares planes, and ORTEP drawings have been deposited with the British Library Lending Division as Supplementary Publication No. (39 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Discussion. The geometry of 1 and 2, at first glance, is contra-intuitive since the sterically largest ligands should try to get farthest from each other. The expectation, based upon the concept that the steric size (cone angle) of the $C_5H_5^-$ ligand is larger than that of a chloride or methyl group (Bagnall and Xing-fu, 1982), is that the cyclopentadienyl groups should be trans to each other and the anionic ligands should be trans to the chelating phosphine ligands. This is clearly not the case in 1, 2, and in the benzyl analogue (Zalkin, Brennan & Andersen, 1986). This situation has been analyzed by Keppert (1977). In six coordinate molecules of the type $M(X)_2(Y)_2$ (bidentate ligand) the locations of X relative to Y is determined by the M-X and M-Y distances. Monodentate ligands with the shortest metal-ligand distance will occupy the less sterically crowded sites trans to the

bidentate ligand. In the compounds 2, and the benzyl analogue, the Th-ring centroid distance is shorter than the Th-C(benzyl) or the Th-Cl bond distance. Hence the ligand with the shortest bond distance is located trans to the bidentate ligand in accord with Keppert's prediction. A similar explanation was proposed for the stereochemistry of $UCl_2\{N[Si(CH_3)_3]\}_2 \cdot (CH_3OCH_2CH_2OCH_3)$, (McCullough, Turner, Andersen, Zalkin & Templeton, 1981). A similar geometry is observed for compound 1 even though the Th-ring centroid distance is essentially equal to that of the Th-C(CH₃). This suggests that the application of Keppert's rules to compounds with cyclopentadienyl ligands may not be universally appropriate and the rules should serve only as a guide to stereochemical predictions.

The averaged Th-P distance in 1 and 2 are 3.147 ± 0.001 A and 3.121 ± 0.001 A, respectively. These distances are in the range found in $Th(CH_2C_6H_5)_4 \cdot [(CH_3)_2PCH_2CH_2P(CH_3)_2]$ of 3.17 ± 0.02 A (Edwards, Andersen & Zalkin, 1984).


This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U. S. Department of Energy under Contract No. DE-AC03-76SF00098.

References

- Bagnall, K. W., Xing-fu, L. (1982). J. Chem. Soc. Dalton Trans., 1365-1369.
- Brennan, J. G. (1985). Ph. D. Thesis, University of California at Berkeley.
- Edwards, P. G., Andersen, R. A., & Zalkin, A. (1984) Organometallics 3, 293-298.
- International Tables for X-ray Crystallography (1974). Vol. IV; Table 2.2, pp 71-102. Birmingham: Kynoch Press.
- Johnson, C. K. (1965). ORTEP. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee.
- Keppert, D. L. (1977) Progress in Inorganic Chemistry, 23, 1-65.
- McCullough, L. G., Turner, H. W., Andersen, R. A., Zalkin, A. & Templeton D. H. (1981). Inorg. Chem., 20, 2869-2871.
- Waser, J. (1963). Acta Cryst. 16, 1091-1094.
- Zalkin, A., Brennan, J. G. & Andersen, R. A. (1986) Accompanying paper.

Table 1. Atomic Parameters

$$B_{eq} = \sum_i \sum_j B_{ij} a_i^* a_j^* a_i \cdot a_j / 3, \text{ \AA}^2.$$



Atom	x	y	z	$B_{eq} (\text{\AA}^2)$
Th	0.10278(1)	0.18457(2)	0.17906(3)	3.642(6)
P(1)	0.12141(11)	0.35769(16)	-0.05635(27)	5.37(6)
P(2)	0.25767(10)	0.20883(15)	0.03265(26)	5.08(6)
C(1)	-0.0213(5)	0.3063(7)	0.1659(16)	7.0(3)
C(2)	-0.0446(5)	0.2234(12)	0.1046(13)	8.3(4)
C(3)	-0.0466(5)	0.1579(7)	0.2250(24)	9.0(5)
C(4)	-0.0237(6)	0.2027(10)	0.3626(14)	7.9(4)
C(5)	-0.0073(4)	0.2935(8)	0.3280(15)	6.9(3)
C(6)	0.1985(6)	0.0315(8)	0.2473(17)	7.8(4)
C(7)	0.1891(6)	0.0748(7)	0.3888(16)	7.9(4)
C(8)	0.1183(8)	0.0566(9)	0.4369(12)	8.4(4)
C(9)	0.0871(6)	0.0038(8)	0.3172(20)	8.3(4)
C(10)	0.1362(9)	-0.0112(7)	0.2019(13)	8.5(4)
C(11)	0.0599(6)	0.3730(8)	-0.2315(11)	8.3(4)
C(12)	0.1206(5)	0.4759(6)	0.0293(13)	8.0(3)
C(13)	0.2082(6)	0.3541(8)	-0.1612(14)	9.4(4)
C(14)	0.2681(5)	0.3231(8)	-0.0677(15)	10.3(4)
C(15)	0.2898(5)	0.1255(9)	-0.1190(13)	9.6(4)
C(16)	0.3362(4)	0.2140(7)	0.1685(12)	7.6(3)
C(17)	0.0900(5)	0.1241(6)	-0.1126(10)	6.4(3)
C(18)	0.1712(4)	0.3126(6)	0.3446(10)	6.1(2)

Table 1. Atomic Parameters (continued)

$$(C_5H_5)_2Cl_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$$

Atom	x	y	z	$B_{eq} (A^2)$
Th	0.10512(1)	0.18258(1)	0.16923(2)	3.133(6)
P(1)	0.12195(9)	0.35857(12)	-0.05896(22)	5.09(5)
P(2)	0.25928(8)	0.20910(13)	0.02111(22)	4.99(5)
C(1)	-0.0239(3)	0.2933(5)	0.1423(11)	6.1(2)
C(2)	-0.0466(3)	0.2029(6)	0.0970(11)	6.9(3)
C(3)	-0.0416(4)	0.1506(5)	0.2430(15)	7.3(3)
C(4)	-0.0149(4)	0.2063(8)	0.3699(10)	7.5(3)
C(5)	-0.0059(4)	0.2923(6)	0.3086(11)	6.2(2)
C(6)	0.2047(4)	0.0417(7)	0.2634(15)	8.2(3)
C(7)	0.1782(9)	0.0736(6)	0.4105(16)	10.7(4)
C(8)	0.1072(6)	0.0419(7)	0.4099(11)	8.5(3)
C(9)	0.0928(4)	-0.0040(5)	0.2691(13)	6.9(3)
C(10)	0.1528(6)	-0.0049(5)	0.1802(9)	7.2(3)
C(11)	0.0631(5)	0.3704(6)	-0.2478(9)	8.0(3)
C(12)	0.1122(4)	0.4724(4)	0.0423(11)	6.9(2)
C(13)	0.2044(5)	0.3509(10)	-0.1849(15)	5.17(27)*
C(14)	0.2696(6)	0.3324(7)	-0.0591(15)	4.76(24)*
C(15)	0.2931(4)	0.1194(8)	-0.1165(11)	10.4(4)
C(16)	0.3372(3)	0.2249(6)	0.1667(10)	7.1(2)
Cl(1)	0.08841(9)	0.12385(11)	-0.15195(19)	5.77(5)
Cl(2)	0.17713(9)	0.31184(11)	0.36194(19)	5.19(4)
C(13')	0.2198(6)	0.3772(8)	-0.1076(18)	5.40(28)*
C(14')	0.2560(8)	0.2870(8)	-0.1594(15)	6.3(3)*

* Isotropically refined disordered atoms.

Table 2. Selected distances(A) and angles(°)

Cp ThMe ₂ ·Me ₂ PCH ₂ CH ₂ PMe ₂			Cp ThCl ₂ ·Me ₂ PCH ₂ CH ₂ PMe ₂		
Th	- C(1)	2.866(8)	Th	- C(1)	2.837(6)
Th	- C(2)	2.834(9)	Th	- C(2)	2.824(6)
Th	- C(3)	2.820(9)	Th	- C(3)	2.804(6)
Th	- C(4)	2.821(8)	Th	- C(4)	2.788(6)
Th	- C(5)	2.849(8)	Th	- C(5)	2.824(6)
Th	- C(6)	2.843(9)	Th	- C(6)	2.791(7)
Th	- C(7)	2.802(9)	Th	- C(7)	2.776(7)
Th	- C(8)	2.803(9)	Th	- C(8)	2.777(7)
Th	- C(9)	2.813(10)	Th	- C(9)	2.785(6)
Th	- C(10)	2.838(9)	Th	- C(10)	2.808(6)
Th	- C(18)	2.583(7)	Th	- Cl(1)	2.707(2)
Th	- C(17)	2.562(8)	Th	- Cl(2)	2.708(2)
Th	- CC(1)	2.593	Th	- Cp(1)	2.564
Th	- Cp(1)	2.573	Th	- Cp(2)	2.544
Th	- P(1)	3.146(2)	Th	- P(1)	3.122(2)
Th	- P(2)	3.147(2)	Th	- P(2)	3.121(2)
P(2)	- P(1)	3.352(3)	P(2)	- P(1)	3.332(3)

Cp ₂ ThMe ₂ ·Me ₂ PCH ₂ CH ₂ PMe ₂			Cp ₂ ThCl ₂ ·Cl ₂ PCH ₂ CH ₂ PMe ₂				
P(1)	-Th	-P(2)	64.36(5)	P(1)	-Th	-P(2)	64.52(5)
P(1)	-Th	-C(17)	71.76(20)	P(1)	-Th	-Cl(1)	72.75(5)
P(1)	-Th	-C(18)	74.05(20)	P(1)	-Th	-Cl(2)	74.35(6)
P(2)	-Th	-C(17)	75.18(20)	P(2)	-Th	-Cl(1)	75.33(6)
P(2)	-Th	-C(18)	71.78(18)	P(2)	-Th	-Cl(2)	73.05(5)
C(17)	-Th	-C(18)	140.04(28)	Cl(1)	-Th	-Cl(2)	141.63(5)
Cp(1)	-Th	-Cp(2)	116.1	Cp(1)	-Th	-Cp(2)	113.9
Cp(1)	-Th	-C(18)	98.9	Cp(1)	-Th	-Cl(1)	101.2

Cp(1)-Th	-C(17)	101.4	Cp(1)-Th	-Cl(2)	98.7
Cp(1)-Th	-P(1)	90.4	Cp(1)-Th	-P(1)	91.3
Cp(1)-Th	-P(2)	154.5	Cp(1)-Th	-P(2)	155.6
Cp(2)-Th	-C(17)	99.4	Cp(2)-Th	-Cl(1)	99.6
Cp(2)-Th	-C(18)	102.0	Cp(2)-Th	-Cl(2)	101.8
Cp(2)-Th	-P(1)	153.4	Cp(2)-Th	-P(1)	154.7
Cp(2)-Th	-P(2)	89.3	Cp(2)-Th	-P(2)	90.3
Th	-P(1) -C(11)	120.3(4)	Th	-P(1) -C(11)	119.26(28)
Th	-P(1) -C(12)	118.2(3)	Th	-P(1) -C(12)	116.14(26)
Th	-P(1) -C(13)	112.1(3)	Th	-P(1) -C(13)	111.9(4)
C(11)-P(1) -C(12)		100.9(5)	C(11)-P(1) -C(12)		102.7(4)
C(11)-P(1) -C(13)		99.5(5)	C(11)-P(1) -C(13)		91.2(5)
C(12)-P(1) -C(13)		102.9(5)	C(12)-P(1) -C(13)		113.0(5)
Th	-P(2) -C(14)	112.0(3)	Th	-P(2) -C(14)	110.3(4)
Th	-P(2) -C(15)	120.2(3)	Th	-P(2) -C(15)	118.7(3)
Th	-P(2) -C(16)	119.2(3)	Th	-P(2) -C(16)	117.90(24)
C(14)-P(2) -C(15)		102.7(6)	C(14)-P(2) -C(15)		114.2(5)
C(14)-P(2) -C(16)		98.9(5)	C(14)-P(2) -C(16)		91.0(4)
C(15)-P(2) -C(16)		100.6(5)	C(15)-P(2) -C(16)		101.5(4)
			Th	-P(1) -C(13') ^b	110.7(4)
			C(11)-P(1) -C(13')		110.7(5)
			C(12)-P(1) -C(13')		94.4(4)
			Th	-P(2) -C(14')	111.9(4)
			C(15)-P(2) -C(14')		87.1(5)
			C(16)-P(2) -C(14')		115.6(5)

^a Cp(1) and Cp(2) represent the centroids of cyclopentadienyl atoms C(1)-C(5) and C(6)-C(10) respectively.

^b Primed atoms are the disordered counterparts of C(13) and C(14).

Figure 1. ORTEP (Johnson, 1965) of $(C_5H_5)_2(CH_3)_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$ (1) showing the atomic numbering scheme; 50% probability ellipsoids.





Figure 2. ORTEP (Johnson, 1965) of $(C_5H_5)_2Cl_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$ (2) showing the atomic numbering scheme; 50% probability ellipsoids; disordered atoms C(13') & C(14') not included.



Supplemental Material

for


Bis(cyclopentadienyl)dimethylthorium(IV)bis(dimethyldiphosphino)ethane
and Bis(cyclopentadienyl)dichlorothorium(IV)bis(dimethyldiphosphino)ethane
 $(C_5H_5)_2(CH_3)_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$, $(C_5H_5)_2Cl_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$

By Allan Zalkin, John G. Brennan & Richard A. Andersen


Materials and Molecular Research Division, Lawrence Berkeley Laboratory and
Department of Chemistry, University of California, Berkeley CA 94720

Abstract. $(C_5H_5)_2(CH_3)_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$ (1) $M_r = 542.44$, monoclinic, $P2_1/n$, $a = 18.485(4)$, $b = 14.118(4)$, $c = 8.258(2)$ Å, $\beta = 91.05(2)^\circ$, $V = 2154.7$ Å³, $Z = 4$, $D_x = 1.672$ g cm⁻³, $\lambda(MoK\alpha) = 0.71073$ Å, $\mu = 73.0$ cm⁻¹, $F(000) = 1040$, $T = 296$ K, $R = 0.026$ for 2521 unique reflections with $F^2 > 2\sigma(F^2)$. Distances (Å) are: Th-C(Cp) 2.84 ± 0.03 ; Th-C(methyl) 2.562(8), 2.583(7); Th-P 3.146(2), 3.147(2) Å; Th-Cp 2.59, 2.57 Å.

$(C_5H_5)_2Cl_2Th \cdot (CH_3)_2PCH_2CH_2P(CH_3)_2$ (2) $M_r = 583.28$, monoclinic, $P2_1/n$, $a = 18.268(2)$, $b = 14.237(6)$, $c = 7.996(3)$ Å, $\beta = 92.27(4)^\circ$, $V = 2078.0$ Å³, $Z = 4$, $D_x = 1.864$ g cm⁻³, $\lambda(MoK\alpha) = 0.71073$ Å, $\mu = 78.3$ cm⁻¹, $F(000) = 1104$, $T = 296$ K, $R = 0.020$ for 2361 unique reflections with $I > \sigma(I)$. Distances (Å) are: Th-C(Cp) 2.80 ± 0.02 ; Th-Cl 2.707(2), 2.708(2); Th-P 3.121(2), 3.122(2) Å; Th-Cp 2.56, 2.54 Å. The structures of (1) and (2) are crystallographically isomorphous and were done to study their six-coordinate stereochemistry. The cyclopentadienyl rings are trans to the bidentate ligand.

Supplemental Table 1. Anisotropic Thermal Parameters (\AA^2)^a

Atom	B ₁₁	B ₂₂	B ₃₃	B ₁₂	B ₁₃	B ₂₃
Th	3.101(10)	3.775(11)	4.067(12)	-0.017(12)	0.522(7)	-0.166(13)
P(1)	4.75(10)	5.22(10)	6.14(12)	0.30(8)	0.57(8)	1.12(9)
P(2)	3.73(8)	5.96(12)	5.61(10)	0.46(8)	1.28(7)	0.19(8)
C(1)	4.6(4)	6.6(5)	10.0(7)	1.7(4)	2.5(4)	2.3(6)
C(2)	4.3(5)	12.4(9)	8.1(7)	2.1(5)	-0.7(4)	-2.0(7)
C(3)	4.0(4)	5.5(6)	17.5(12)	-0.7(4)	2.9(6)	-1.7(7)
C(4)	5.9(5)	10.0(8)	7.8(6)	2.8(5)	3.2(5)	2.7(6)
C(5)	4.1(4)	7.4(6)	9.2(7)	1.8(4)	1.5(4)	-2.3(5)
C(6)	6.7(6)	5.7(5)	11.1(8)	1.5(5)	2.5(6)	1.6(5)
C(7)	7.9(6)	5.4(5)	10.3(7)	0.9(5)	-2.9(6)	1.5(5)
C(8)	11.2(8)	8.1(7)	6.0(5)	3.8(6)	3.1(5)	3.3(5)
C(9)	6.6(6)	6.7(6)	11.6(9)	-1.2(5)	-0.3(6)	3.1(6)
C(10)	13.3(9)	4.3(5)	7.9(7)	2.4(6)	-1.5(7)	-0.2(4)
C(11)	10.0(7)	8.8(6)	6.2(5)	1.5(5)	-0.8(5)	1.2(5)
C(12)	7.7(5)	5.5(5)	10.6(7)	-0.2(4)	-2.0(5)	1.6(5)
C(13)	7.6(6)	10.0(7)	10.6(7)	0.5(5)	2.7(5)	4.7(6)
C(14)	4.6(4)	12.6(9)	13.9(9)	0.4(5)	2.9(5)	7.7(7)
C(15)	6.3(5)	12.8(8)	9.7(7)	0.3(5)	2.5(5)	-4.8(6)
C(16)	4.4(4)	9.6(7)	8.8(6)	-0.4(4)	0.3(4)	-0.3(5)
C(17)	7.5(5)	5.9(5)	5.8(4)	-1.0(4)	-0.5(4)	-1.5(4)
C(18)	5.7(4)	6.9(5)	5.6(4)	-1.6(4)	0.4(3)	-2.4(4)

Supplemental Table 2. Anisotropic Thermal Parameters (\AA^2) (continued)

Atom	B_{11}	B_{22}	B_{33}	B_{12}	B_{13}	B_{23}
Th	2.794(11)	3.274(11)	3.371(11)	0.002(6)	0.606(7)	-0.103(7)
Cl(1)	6.62(8)	6.17(9)	4.53(7)	-1.44(7)	0.22(6)	-1.59(7)
Cl(2)	5.32(7)	5.96(8)	4.32(7)	-1.03(6)	0.49(6)	-1.39(6)
P(1)	4.79(8)	4.51(7)	6.02(9)	0.26(6)	0.95(7)	1.20(7)
P(2)	3.56(7)	6.15(8)	5.37(8)	0.61(6)	1.63(6)	0.64(7)
C(1)	3.73(29)	5.9(4)	8.7(5)	1.63(26)	2.0(3)	1.9(4)
C(2)	2.71(26)	9.1(5)	8.8(5)	0.89(29)	-0.14(28)	-2.8(4)
C(3)	3.3(3)	6.0(4)	12.8(7)	-0.07(27)	2.3(4)	2.3(5)
C(4)	4.8(4)	11.3(6)	6.7(4)	2.3(4)	3.3(3)	2.0(5)
C(5)	4.4(3)	7.6(4)	6.7(4)	0.89(29)	1.46(29)	-2.3(4)
C(6)	4.5(3)	6.7(5)	13.2(8)	1.2(3)	0.3(4)	5.1(5)
C(7)	16.5(10)	4.6(4)	10.4(7)	0.3(5)	-9.3(7)	0.8(4)
C(8)	13.1(7)	7.3(5)	5.3(4)	4.2(5)	3.9(5)	3.0(4)
C(9)	5.9(4)	4.4(3)	10.3(6)	-0.17(28)	0.8(4)	2.6(4)
C(10)	9.9(5)	4.6(3)	7.1(4)	2.8(4)	0.9(4)	0.3(3)
C(11)	10.2(5)	7.9(5)	5.9(4)	1.1(4)	-0.2(4)	1.6(4)
C(12)	6.3(4)	4.0(3)	10.2(5)	-0.39(26)	-1.8(3)	0.3(3)
C(15)	6.7(4)	16.1(8)	8.4(5)	1.8(5)	2.1(4)	-5.7(6)
C(16)	4.3(3)	8.3(4)	8.8(5)	-1.1(3)	1.1(3)	-1.5(4)

^a The anisotropic temperature factor has the form:

$$\exp[-0.25(B_{11}h^2a^{*2} + 2B_{12}hka^*b^* + \dots)].$$


Supplemental Table 3. Estimated Hydrogen Positional and Isotropic Thermal

Parameters (Å^2) ^a $(\text{C}_5\text{H}_5)_2(\text{CH}_3)_2\text{Th} \cdot (\text{CH}_3)_2\text{PCH}_2\text{CH}_2\text{P}(\text{CH}_3)_2$

Atom	x	y	z	B
H(1)	-0.0153	0.3643	0.1071	10.0
H(2)	-0.0574	0.2125	-0.0074	10.0
H(3)	-0.0616	0.0928	0.2142	10.0
H(4)	-0.0198	0.1744	0.4682	10.0
H(5)	0.0106	0.3403	0.4030	10.0
H(6)	0.2425	0.0308	0.1875	10.0
H(7)	0.2245	0.1119	0.4468	10.0
H(8)	0.0960	0.0772	0.5350	10.0
H(9)	0.0382	-0.0193	0.3144	10.0
H(10)	0.1282	-0.0463	0.1035	10.0
H(11)	0.0791	0.4209	-0.3018	10.0
H(12)	0.0134	0.3928	-0.1947	10.0
H(13)	0.0556	0.3145	-0.2890	10.0
H(14)	0.0720	0.4919	0.0583	10.0
H(15)	0.1379	0.5200	-0.0487	10.0
H(16)	0.1513	0.4773	0.1243	10.0
H(17)	0.2185	0.4170	-0.1988	10.0
H(18)	0.2028	0.3124	-0.2524	10.0
H(19)	0.3090	0.3184	-0.1376	10.0
H(20)	0.2779	0.3696	0.0146	10.0
H(21)	0.3010	0.0662	-0.0675	10.0
H(22)	0.3323	0.1506	-0.1680	10.0
H(23)	0.2527	0.1161	-0.2000	10.0
H(24)	0.3772	0.2361	0.1102	10.0
H(25)	0.3460	0.1518	0.2107	10.0
H(26)	0.3264	0.2563	0.2562	10.0

^a The isotropic temperature factor has the form $\exp[-B(\sin\theta/\lambda)^2]$.

Supplemental Table 3. (continued)



Atom	x	y	z	B
H(1)	-0.0212	0.3475	0.0675	10.0
H(2)	-0.0630	0.1810	-0.0141	10.0
H(3)	-0.0547	0.0844	0.2539	10.0
H(4)	-0.0042	0.1869	0.4854	10.0
H(5)	0.0112	0.3468	0.3741	10.0
H(6)	0.2544	0.0514	0.2256	10.0
H(7)	0.2043	0.1109	0.4966	10.0
H(8)	0.0729	0.0517	0.4992	10.0
H(9)	0.0458	-0.0325	0.2357	10.0
H(10)	0.1574	-0.0353	0.0716	10.0
H(11)	0.0126	0.3830	-0.2164	10.0
H(12)	0.0805	0.4223	-0.3159	10.0
H(13)	0.0642	0.3119	-0.3126	10.0
H(14)	0.1285	0.5219	-0.0329	10.0
H(15)	0.0607	0.4825	0.0665	10.0
H(16)	0.1421	0.4735	0.1463	10.0
H(17)	0.1992	0.2994	-0.2657	10.0
H(18)	0.2116	0.4101	-0.2445	10.0
H(19)	0.2686	0.3769	0.0335	10.0
H(20)	0.3159	0.3382	-0.1161	10.0
H(21)	0.3368	0.1426	-0.1699	10.0
H(22)	0.3052	0.0630	-0.0513	10.0
H(23)	0.2550	0.1045	-0.2023	10.0
H(24)	0.3482	0.1654	0.2235	10.0
H(25)	0.3797	0.2450	0.1049	10.0
H(26)	0.3255	0.2728	0.2491	10.0
H(17') ^a	0.2225	0.4223	-0.1984	10.0
H(18')	0.2460	0.4012	-0.0071	10.0
H(19')	0.3060	0.2996	-0.1943	10.0
H(20')	0.2278	0.2574	-0.2520	10.0

^a Primed atoms are disordered counterparts to atoms H(17)-H(20).

Supplemental Table 4. Additional Distances(A)

$\text{Cp}_2\text{ThMe}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$			$\text{Cp}_2\text{ThCl}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$		
P(1) - C(11)	1.837(9)		P(1) - C(11)	1.827(8)	
P(1) - C(12)	1.813(9)		P(1) - C(12)	1.824(7)	
P(1) - C(13)	1.837(10)		P(1) - C(13)	1.847(10)	
P(2) - C(14)	1.825(10)		P(2) - C(14)	1.881(9)	
P(2) - C(15)	1.826(10)		P(2) - C(15)	1.810(8)	
P(2) - C(16)	1.820(9)		P(2) - C(16)	1.817(7)	
C(1) - C(2)	1.344(14)		C(1) - C(2)	1.396(9)	
C(1) - C(5)	1.370(14)		C(1) - C(5)	1.357(11)	
C(2) - C(3)	1.358(16)		C(2) - C(3)	1.384(11)	
C(3) - C(4)	1.362(15)		C(3) - C(4)	1.362(12)	
C(4) - C(5)	1.349(13)		C(4) - C(5)	1.332(10)	
C(6) - C(7)	1.333(13)		C(6) - C(7)	1.366(14)	
C(6) - C(10)	1.347(14)		C(6) - C(10)	1.316(11)	
C(7) - C(8)	1.398(14)		C(7) - C(8)	1.374(13)	
C(8) - C(9)	1.358(15)		C(8) - C(9)	1.319(12)	
C(9) - C(10)	1.344(14)		C(9) - C(10)	1.329(11)	
C(13) - C(14)	1.409(14)		C(13) - C(14)	1.551(12)	
			P(1) - C(13') ^a	1.864(10)	
			P(2) - C(14')	1.819(11)	
			C(13') - C(14')	1.511(12)	

Supplemental Table 4. (continued)

$\text{Cp}_2\text{ThMe}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$		$\text{Cp}_2\text{ThCl}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$	
H(1) - C(1)	0.959	H(1) - C(1)	0.979
H(2) - C(2)	0.963	H(2) - C(2)	0.977
H(3) - C(3)	0.963	H(3) - C(3)	0.978
H(4) - C(4)	0.962	H(4) - C(4)	0.977
H(5) - C(5)	0.961	H(5) - C(5)	0.980
H(6) - C(6)	0.960	H(6) - C(6)	0.979
H(7) - C(7)	0.960	H(7) - C(7)	0.978
H(8) - C(8)	0.961	H(8) - C(8)	0.979
H(9) - C(9)	0.961	H(9) - C(9)	0.979
H(10) - C(10)	0.961	H(10) - C(10)	0.977
H(11) - C(11)	0.964	H(11) - C(11)	0.982
H(12) - C(11)	0.958	H(12) - C(11)	0.978
H(13) - C(11)	0.955	H(13) - C(11)	0.981
H(14) - C(12)	0.959	H(14) - C(12)	0.979
H(15) - C(12)	0.957	H(15) - C(12)	0.979
H(16) - C(12)	0.961	H(16) - C(12)	0.978
H(17) - C(13)	0.961	H(17) - C(13)	0.980
H(18) - C(13)	0.960	H(18) - C(13)	0.980
H(19) - C(14)	0.961	H(19) - C(14)	0.975
H(20) - C(14)	0.960	H(20) - C(14)	0.980
H(21) - C(15)	0.960	H(21) - C(15)	0.977
H(22) - C(15)	0.957	H(22) - C(15)	0.978
H(23) - C(15)	0.959	H(23) - C(15)	0.981
H(24) - C(16)	0.958	H(24) - C(16)	0.978
H(25) - C(16)	0.962	H(25) - C(16)	0.980
H(26) - C(16)	0.958	H(26) - C(16)	0.979
		H(17') - C(13') ^a	0.971
		H(18') - C(13')	0.979
		H(19') - C(14')	0.982
		H(20') - C(14')	0.980


^a Primed atoms are disordered counterparts to atoms C(13), C(14) and H(17)-H(20).

Supplemental Table 5. Additional Angles

$\text{Cp}_2\text{ThMe}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$			$\text{Cp}_2\text{ThCl}_2 \cdot \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$		
Atoms		Angle(°)	Atoms		Angle(°)
C(2) - C(1) - C(5)		107.8(10)	C(2) - C(1) - C(5)		107.6(7)
C(1) - C(2) - C(3)		109.3(10)	C(1) - C(2) - C(3)		105.6(7)
C(2) - C(3) - C(4)		106.4(10)	C(2) - C(3) - C(4)		108.9(7)
C(3) - C(4) - C(5)		109.5(10)	C(3) - C(4) - C(5)		107.9(8)
C(1) - C(5) - C(4)		107.1(10)	C(1) - C(5) - C(4)		109.9(7)
C(7) - C(6) - C(10)		109.0(10)	C(7) - C(6) - C(10)		109.3(8)
C(6) - C(7) - C(8)		107.6(10)	C(6) - C(7) - C(8)		104.7(7)
C(7) - C(8) - C(9)		106.3(09)	C(7) - C(8) - C(9)		108.8(8)
C(8) - C(9) - C(10)		108.6(10)	C(8) - C(9) - C(10)		108.7(8)
C(6) - C(10) - C(9)		108.5(10)	C(6) - C(10) - C(9)		108.5(8)
P(1) - C(13) - C(14)		115.8(08)	P(1) - C(13) - C(14)		106.2(8)
P(2) - C(14) - C(13)		115.8(08)	P(2) - C(14) - C(13)		107.1(8)
			P(1) - C(13') - C(14') ^a		111.7(9)
			P(2) - C(14') - C(13')		107.6(9)

^a Primed atoms are counterparts to C(13) & C(14).

Supplemental Table 6. Least-squares Planes



Plane No. 1

Equation of plane: 17.46930 a - 3.71855 b - 1.74192 c = -1.79523

Distance to the plane from atoms


in the plane			not in the plane	
atom	d(A)	$\sigma(d)$	atom	d(A)
C(1)	-0.0042	(169)	Th	2.592(7)
C(2)	0.0039	(218)		
C(3)	0.0018	(238)		
C(4)	-0.0040	(181)		
C(5)	0.0041	(159)		

Plane No. 2

Equation of plane: -5.42084 a 11.74148 b - 3.84877 c = -1.65146

Distance to the plane from atoms

in the plane			not in the plane	
atom	d(A)	$\sigma(d)$	atom	d(A)
C(6)	-0.0068	(177)	Th	2.572(7)
C(7)	0.0075	(173)		
C(8)	-0.0064	(183)		
C(9)	0.0028	(203)		
C(10)	0.0041	(206)		



Plane No. 1

Equation of plane: 17.33518 a - 3.41437 b + -1.93801 c = -1.69297

Distance to the plane from atoms

in the plane			not in the plane	
atom	d(A)	$\sigma(d)$	atom	d(A)
C(6)	0.0131	(138)	Th	2.564(5)
C(7)	-0.0056	(110)		
C(8)	0.0013	(110)		
C(9)	0.0036	(114)		
C(10)	-0.0126	(143)		


Plane No. 2

Equation of plane: - 5.15826 a + 12.07150 b - 3.49565 c = -1.47343


Distance to the plane from atoms

in the plane			not in the plane	
atom	d(A)	$\sigma(d)$	atom	d(A)
C(1)	0.0060	(131)	Th	2.544(6)
C(2)	-0.0070	(145)		
C(3)	0.0077	(189)		
C(4)	0.0003	(156)		
C(5)	-0.0039	(134)		


Supplemental Fig. 1. ORTEP drawing of compound 1 viewed down a line connecting the centroids of the Cp rings.




Supplemental Fig. 2. ORTEP drawing of molecule 2 viewed down a line connecting the centroids of the Cp rings; disordered atoms C(13') & C(14') not included.




Supplemental Fig 3. ORTEP drawing of molecule 1 viewed in a direction edge on to the Cp rings.




Supplemental Fig. 4. ORTEP drawing of molecule 2 viewed in a direction edge on to the Cp rings.



Supplemental Fig. 5. ORTEP drawing of molecule 1 viewed down a pseudo-2-fold axis.



Supplemental Fig 6. ORTEP drawing of molecule 2 viewed down a pseudo-2-fold axis.



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*