UC Irvine

UC Irvine Previously Published Works

Title

Rehabilitation of Cardiovascular Disorders and Sleep Apnea.

Permalink

<https://escholarship.org/uc/item/1fm6d0zh>

Journal

Sleep medicine clinics, 12(2)

ISSN

1556-407X

Author

Jafari, Behrouz

Publication Date

2017-06-01

DOI

10.1016/j.jsmc.2017.01.001

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at

<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Rehabilitation of Cardiovascular Disorders and Sleep Apnea

Behrouz Jafari, MD^{a,b,*}

KEYWORDS

- Obstructive sleep apnea • Sleep disordered breathing • Cardiac rehabilitation
- Stroke rehabilitation • Cardiovascular disease • Stroke

KEY POINTS

- Obstructive sleep apnea (OSA) is a common disorder in patients entering cardiac rehabilitation units.
- Sleep disordered breathing is underdiagnosed in the poststroke period.
- Early screening and treatment of OSA are important in the management of the poststroke period or after cardiac events.
- Failure to treat OSA in cardiac patients and patients with stroke can have negative impacts on outcomes during rehabilitation.

INTRODUCTION

Obstructive sleep apnea (OSA) syndrome is a disorder affecting 2% to 4% of the general population in the United States.^{1–3} It is characterized by recurrent episodes of upper airway collapse during sleep. This condition presents in more than half of patients with acute coronary heart disease who are eligible for enrollment in cardiac rehabilitation programs.⁴ Because of the hemodynamic fluctuations associated with OSA, those patients with concurrent OSA who enter cardiac rehabilitation programs may be placed at greater risk for arrhythmias and exercise-related complications.⁵ This risk can lead to an increase in major adverse cardiac events such as revascularization, heart failure (HF), hospital readmission, functional limitation, and reduced quality of life in cardiac patients.

In contrast, stroke, with an incidence of 2 to 18 per 1000 per year, is the second leading cause

of death worldwide and more than 50% of survivors have mental and physical impairment.^{6,7} Although sleep disordered breathing (SDB) has been recognized in patients with stroke since the early nineteenth century,⁸ over the last 2 decades it has been emerging as an important cardiovascular risk factor.

In the poststroke period, patients with OSA have greater disability and higher mortality than patients without OSA.^{9,10} However, OSA is under-recognized during the poststroke period because of a lack of symptoms or lack of gross obesity, despite evidence that managing this risk factor may benefit those patients.¹¹ There is growing evidence that treatment of coexisting OSA with continuous positive airway pressure (CPAP) or mandibular advancement devices can successfully treat OSA, resulting in improved rehabilitation outcomes and improved health-related quality of life.^{12–15}

Disclosure: The author has nothing to disclose.

^a Section of Pulmonary, Critical Care and Sleep Medicine, School of Medicine, University of California-Irvine, 333 City Boulevard West, Suite 400, Irvine, CA, USA; ^b Sleep Program, VA Long Beach Healthcare System, 5901 East 7th Street, Long Beach, CA 90822, USA

* Sleep Program, VA Long Beach Healthcare System, 5901 East 7th Street, Long Beach, CA.

E-mail address: jafarib@uci.edu

Sleep Med Clin 12 (2017) 193–203

<http://dx.doi.org/10.1016/j.jsmc.2017.01.001>

1556-407X/17/Published by Elsevier Inc.

PREVALENCE OF OBSTRUCTIVE SLEEP APNEA IN PATIENTS ENTERING CARDIAC REHABILITATION PROGRAMS

It is well known that patients with different types of cardiovascular diseases have higher prevalence of SDB.^{16–19} Several studies have shown that OSA is an independent risk factor for development of hypertension and has a dose-response relationship between severity of OSA and incidence of hypertension.^{18,20,21} SDB is also estimated to have a prevalence of 30% to 69% in patients with coronary artery disease (CAD)^{22–24} and 50% to 70% in patients with systolic HF.^{25,26} However, fewer data are available in HF with preserved ejection fraction (EF) than in those with reduced EF. The largest study on this group was done on 244 consecutive patients: 97 patients (39.8%) presented with OSA and 72 patients (29.5%) with central sleep apnea (CSA).²⁷

In the postdischarge period after coronary revascularization procedures, 52% to 64%^{28,29} of patients had moderate to severe OSA. A recent study suggested that, overall, 53% of patients entering cardiac rehabilitation programs are at high risk, or are previously diagnosed with OSA.⁴ However, the experts believe that the prevalence of OSA is substantially higher and is largely under-recognized in patients entering cardiac rehabilitation programs.³⁰

PREVALENCE OF OBSTRUCTIVE SLEEP APNEA IN THE POSTSTROKE PERIOD

In the past decade, longitudinal studies have shown that people with SDB have a greater risk for stroke.^{31–35} The reported prevalence in the literature is very high, ranging from 40% to 90%,^{9,10,34,36–40} likely reflecting different study designs and definitions of SDB.

Yaggi and colleagues⁴¹ followed 1022 patients over 6 years and found that OSA is associated with an increased risk for stroke, transient ischemic attack, and death. This finding was true even after adjusting for age, gender, body mass index (BMI), arterial hypertension, diabetes, atrial fibrillation, hyperlipidemia, and smoking habits. Patients with severe SDB had a higher chance of having stroke (hazard ratio [HR], 3.3; confidence interval [CI], 1.7–6.3). These results were confirmed by other investigators as well.^{35,42–44}

It has been shown that many patients experience a recurrent stroke within 5 years of their first attack⁴⁵ and those with recurrent stroke have a higher chance of having SDB compared with first-time stroke victims.⁴⁶ Note that presence of an apnea-hypopnea index (AHI) greater than 10

per hour is an independent risk factor for stroke recurrence.⁴⁶

The prevalence of sleep apnea is highest early following stroke and may decrease during recovery,^{34,47} reflecting a decrease in sleep apnea in most patients, although it can also be caused by a higher mortality in those with more severe form of disease. Although SDB tends to improve spontaneously several weeks after stroke, approximately 50% of patients still have OSA 3 months after the acute event.^{31,32,34} Szűcs and colleagues⁴⁸ reported that the frequency and severity of sleep apnea were unchanged 3 months later in most patients with ischemic stroke, whereas it was greatly improved in patients with hemorrhagic stroke. Moreover, hemorrhagic strokes lead more often to central apneas.⁴⁹

Both OSA and, less frequently, CSA have been identified as risk factors for stroke.¹⁰ However, only OSA is recognized as a mortality risk for ischemic stroke.^{9,48,49} Periodic breathing (PB) may also develop in patients with stroke in the absence of cardiopulmonary disease, disturbed consciousness, or HF.⁵⁰ During the acute phase of stroke, obstructive apnea is the most predominant type. Central apneas are associated with altered mental status, brain edema, and ischemia of brainstem and their incidence diminishes during recovery after stroke.⁵¹ Rowat and colleagues⁵² investigated the impact of PB in the awake state on mortality in the immediate phase after stroke. They investigated 138 patients with stroke at a median of 4 hours after stroke and found that those with PB were more likely to have severe stroke and more likely to be dead or dependent at 3 months. In contrast, another study showed that CSA was not related to early death among the patients with stroke. Only patients with OSA (AHI >10) had an increased risk of early death.⁹

In a prospective study on 161 patients admitted to a stroke unit, no relationships were found between sleep-related respiratory events and the topographic location of the neurologic lesion or vascular involvement.³⁴

WHY OBSTRUCTIVE SLEEP APNEA IS UNDER-RECOGNIZED IN STROKE AND CARDIAC REHABILITATION UNITS

OSA is rarely considered in rehabilitation units despite evidence that managing OSA, when present, may benefit patients with cardiovascular diseases. It has been shown that, despite higher prevalence of SDB among patients with stroke, none of them had been referred for diagnostic studies because SDB had not been suspected clinically because of less sleepiness and lower BMI.

The late diagnosis of sleep apnea can potentially have a negative impact on the outcomes and rehabilitation processes of these patients.¹¹ Two key clinical symptoms, excessive daytime sleepiness and obesity, do not seem to be prevalent in patients after stroke^{10,13,53,54} or HF.⁵⁵ Bassetti and colleagues⁵³ showed that those with ischemic strokes were neither obese nor sleepy despite having severe OSA. They reported that 26 out of 152 patients with stroke had severe OSA with an AHI greater than or equal to 30, but their mean Epworth Sleepiness Scale (ESS) score was 6 and BMI of 27.9 kg/m² indicated that they were not obese and did not have excessive daytime sleepiness.⁵³ Similarly, Wessendorf and colleagues⁵⁶ showed that, in 105 patients with stroke and moderate to severe OSA (AHI ≥ 15), the mean ESS score was 7 and mean BMI was 28.7 kg/m². Kaneko and colleagues¹⁰ made similar observations as well (mean ESS score of 5 and BMI of 28.8 kg/m²).

The ESS relies on the self-perception of sleepiness. In patients without stroke and cardiovascular disease, the ESS correlates modestly, but significantly, with the AHI and with objective measures of sleepiness such as the Multiple Sleep Latency Test⁵⁷ and the Oxford Sleep Resistance Test.⁵⁵ There are several possibilities that might explain the lack of sleepiness in patients with stroke, including cognitive impairment, low physical activity, or drug-related factors.^{54,58,59} In one study, patients with stroke were only mildly impaired and there was no significant relationship between Mini-Mental State Examination and ESS scores.⁵⁴ This finding indicates that cognitive impairment probably is not the reason for the lack of reported daytime sleepiness.

Another proposed potential explanation for the lower ESS scores in patients with stroke is low level of physical activity, so they may be less likely to feel sleepy.⁵⁴ The investigators suggest that most of their physical needs are met by hospital personnel, and as a result they are less physically active, which might affect subjective sleepiness.⁵⁴ Although it seems a plausible reason, there is no evidence to accept this as the cause of less daytime sleepiness.

Another possible reason is that drugs commonly used in patients with cardiac disease, such as β -receptor blockers, diuretics, statins, thrombocyte aggregation inhibitors, and antiarrhythmics, may cause a reduced perception of daytime sleepiness through a central nervous system-stimulating effect. However, none of them are known to have such effects, and none have been reported to reduce perception of sleepiness.⁶⁰ Therefore, it is unlikely that these drugs could explain lower levels of sleepiness in patients with stroke.

These findings in patients with stroke are similar to those previously described in cardiovascular patients, who also had less daytime sleepiness at any given OSA severity.^{58,59} Arzt and colleagues⁵⁴ reported that patients with HF with any OSA severity had lower mean ESS (7), indicating less sleepiness despite sleeping less. This finding in patients with HF shows that the absence of subjective sleepiness is not a reliable means of ruling out OSA.⁵⁵

Another finding was the lack of an association between BMI and the severity of OSA in patients with stroke, which contrasts with the well-established relationship between increasing BMI and increasing severity of OSA in patients without stroke.^{61,62}

All of the studies discussed earlier showed that lower ESS scores and BMI in patients with stroke or HF and OSA cannot be explained by differences in the severity of OSA or other confounding factors and that those are not sensitive independent predictors of the presence of OSA.⁶³

Therefore, regardless of the underlying mechanisms, lower ESS scores and BMI may make the diagnosis of OSA less likely in the rehabilitation period, and as a result may have a negative impact on clinical course and outcome.

OBSTRUCTIVE SLEEP APNEA AND REHABILITATION OUTCOME

Stroke and cardiac rehabilitation programs combine physical rehabilitation with education, self-management, and psychological support. The goal of postacute rehabilitation is to facilitate return home by assisting patients in achieving independence in personal care activities. In addition, it provides an opportunity to identify any comorbidity such as hypertension, and diabetes that may delay recovery. Therefore, early recognition and treatment of OSA during acute rehabilitation should be important parts of recovery. There is growing evidence that failure to treat OSA in cardiac patients can have a negative effect on postoperative recovery and increase mortality and morbidity, reduce quality of life, and lead to worse outcomes during and after the cardiac rehabilitation.^{64,65} Hargenes and colleagues⁶⁶ studied patients entering an early outpatient cardiac rehabilitation program and found that cardiac patients with OSA had significantly lower cardiac output, stroke volume, and contractility index. They suggested that the decreased cardiac function in the OSA group, likely because of pressure and volume alteration associated with apneic events, may place these patients at a disadvantage in recovering from their cardiac events, and at increased risk for secondary complications.

In a recent study in which 67 consecutive patients referred for coronary artery bypass grafting (CABG) underwent clinical evaluation and were followed for a mean of 4.5 years after CABG, 56% of patients had moderate-severe OSA.⁶⁷ Those with sleep apnea had significantly higher major adverse cardiac or cerebrovascular events (35% vs 16%; $P = .02$) and new revascularization (19% vs 0%; $P = .01$). OSA was an independent factor associated with these events in the multivariate analysis.⁶⁷

In a prospective observational study, patients with acute myocardial infarction (MI) and percutaneous coronary intervention underwent cardiovascular magnetic resonance to define salvaged myocardium and infarct size within 3 to 5 days and at 3 months after acute MI. Of the 56 patients included, 29 (52%) had SDB. Patients with SDB had significantly less salvaged myocardium, smaller reduction in infarct size within 3 months after acute MI, a larger final infarct size, and a lower final left ventricular EF.²⁸ In a multivariate analysis, including established risk factors for large MI, AHI was independently associated with less myocardial salvage and a larger infarct size 3 months after acute MI.²⁸

In contrast, the presence of OSA in the setting of stroke is associated with unfavorable clinical course, including early neurologic worsening, delirium, depressed mood, impaired functional capacity and cognition, longer period of hospitalization and rehabilitation, and increased mortality.^{1,7,9,10,36,37,68–72} In addition, severe OSA not only increases the risk of stroke recurrence but also increases the incidence of fatal and nonfatal cardiovascular events as well as sudden death during sleep.^{44,73–75} The higher blood pressure and its swing during episodes of upper airways obstruction lead to worse outcome following stroke.⁷⁶ Sleep apnea probably promotes this functional impairment through intermittent nocturnal hypoxia, reduced cerebral perfusion, and fragmented sleep.⁷⁷

Sleep quality can influence the rehabilitation process. The development of poor sleep with frequent arousals may be particularly important and persist long after stroke symptoms have resolved. In addition, poor sleep quality might have a negative impact on patients' motivation, level of energy, and participation in rehabilitation. Prior study suggests that SDB and poor sleep during rehabilitation were associated with less functional recovery between admission and discharge.^{37,78} This association can persist up to 3 months after admission⁷⁸ and persisted when analyses were adjusted for other significant independent predictors of functional recovery,

including total hours of therapy received, mental status, acute hospital transfer during the rehabilitation stay, and reason for admission to rehabilitation facility.⁷⁸ This finding is particularly important because, although many other predictors of rehabilitation outcomes are difficult or impossible to change, abnormal sleep pattern can be a modifiable predictor of rehabilitation outcomes. Interventions to improve abnormal sleep/wake patterns and early treatment of SDB during rehabilitation may result in better functional recovery among older people. Studies in other institutional settings (eg, nursing homes, acute care hospitals) suggest that sleep problems are associated with social isolation, poor health, and functional impairment in older people.^{79–83} Those with excessive daytime sleepiness required more assistance with self-care activities, engaged in fewer social interactions,^{82,83} and showed worse health-related quality of life.⁸⁴ In addition, they had slower recovery of functional abilities and longer stay in rehabilitation units.^{10,85} These outcomes can lead to increased caregiving needs at home, nursing home placement, or death,⁸⁶ and may have detrimental effects on rehabilitation^{64,87} and long-term outcome.⁸⁸

Functional independence in personal care activities can be measured using the motor subscale of the Functional Independence Measure (mFIM),⁸⁹ which is widely used in rehabilitation settings to assess functional limitations and changes in functional status with rehabilitation therapy. Prior work suggests that a 1-point improvement in mFIM score is associated with 2.2 fewer minutes of help required from another person each day,⁹⁰ which translates into 15 minutes of caregiving per week for every 1-point change in mFIM. Every 10% reduction in daytime sleepiness is associated with 1-point additional improvement in mFIM during rehabilitation.⁷⁸ This degree of change in daytime sleepiness is achievable by using interventions to improve sleep patterns. It was also associated with increased participation in social and physical activities.⁸² This relationship between more daytime sleepiness and less favorable immediate-term and long-term functional recovery suggests that sleep is an important modifiable predictor of rehabilitation outcomes for this vulnerable patient population.

Good and colleagues³⁷ found that patients with recent stroke and sleep apnea had poorer functional outcomes assessed with the Barthel Activities of Daily Living (ADL) index at discharge from rehabilitation as well as 3 and 12 months after stroke onset. Soon after admission to the stroke rehabilitation unit, these patients are more dependent in their ADLs and have longer latency in

reaction and in response to verbal stimuli, impairing their ability to acquire new skills.¹⁴ In addition, the presence of SDB was associated with a worse Barthel index at 6 months after stroke⁹¹ and predicted a higher mortality during the following 10 years.^{9,53,91} Note that high mortality was found only in patients with OSA but not CSA.⁹

Other symptoms in patients with stroke and cardiac disorders are impaired memory, lack of ability to concentrate, and tiredness, which are common in patients with OSA.^{92,93} These patients are less adherent to rehabilitation programs because of decreased functional capacity⁹⁴ and depression.^{64,87} Poor sleep and fatigue are associated with neuropsychiatric and cognitive disturbances, and result in worse outcomes in rehabilitation and quality of life.⁹⁵ Depression after stroke occurs in 30% to 60% of patients with stroke⁹⁶ and poor sleep strongly correlates with health complaints and depressive symptoms.⁹⁷ Previous studies have shown that depression and cognitive impairment commonly occur in patients with sleep apnea but without stroke, and improve after CPAP treatment.^{98–100} This finding indicates that emotional instability in this population might be sleep related and therefore it is reasonable to consider and investigate the SDB, if other symptoms exist.

Poststroke fatigue is an independent predictor of dependency and death. In a cohort study of 8194 patients with stroke,¹⁰¹ 39% of patients reported fatigue 2 years after their stroke. Patients with fatigue were more likely to be institutionalized and had increased mortality in the subsequent year.¹⁰¹

In some studies, delirium was reported in up to half of patients, especially in the first week after ischemic stroke.^{102,103} This delirium can prolong hospital stay and increases risk of dementia and admission to an institution.^{103,104} Well-known risk factors for delirium are hypoxemia¹⁰⁵ and OSA, which can be reversed by CPAP treatment.^{106–108}

These studies show high prevalence of OSA at different time points from stroke onset or cardiac event and OSA exerts a negative effect on the functional recovery of these patients. Therefore, patients with cardiovascular diseases should be screened for OSA early in the course of their disease.

However, the question remains whether early intervention for treatment of OSA shortly after stroke or cardiac event alters the neurologic recovery and/or outcome.

EARLY TREATMENT OF OBSTRUCTIVE SLEEP APNEA IN STROKE REHABILITATION

The central nervous system is very sensitive to hypoperfusion in the poststroke period. Therefore,

interventions designed to improve poststroke functioning are more effective the earlier they are delivered after symptom onset to reduce mortality and disability. Upper airway obstruction presents a potential therapeutic target to improve outcomes, because control of sleep apnea may prevent hypoxemia, large fluctuations in blood pressure, and cerebral hypoperfusion. To address this, previous studies have examined the potential beneficial effects of OSA treatment on neurologic and cognitive functions of patients during the stable phase of stroke in a rehabilitation setting. Bravata and colleagues⁴⁷ randomized 55 patients to an intervention group (stroke standard of care plus auto-CPAP) and control group (only stroke standard of care). Patients with stroke randomized to the intervention group received 2 nights of auto-CPAP, but only those with evidence of OSA received auto-CPAP for the remainder of the 30-day period. Change in stroke severity was assessed by comparing the National Institutes of Health (NIH) Stroke Scale (NIHSS) score at baseline versus at 30 days. Intervention patients had greater improvements in NIHSS score (−3.0) than control patients (−1.0; $P = .03$). Among patients with OSA, greater improvement was observed with increasing auto-CPAP use.

In another randomized study on 50 patients with stroke,¹⁰⁹ CPAP use within 24 hours of hospitalization resulted in a significant improvement in the NIHSS score. In both trials, greater improvement was noted with increasing CPAP use.

Both sleep apnea and blood pressure variability confer a poor prognosis after stroke and are potentially treatable.⁷⁶ Treatment of coexisting OSA decreases both nocturnal and diurnal blood pressure^{56,110} and as a result reduces risk of stroke recurrence,¹¹¹ reduces mortality,⁷⁴ and improves quality of life.^{56,69} Because SDB improves after the acute phase of stroke, auto-CPAP systems may be preferable.^{47,112}

Among the most prominent potential benefits of CPAP therapy are motor and functional outcomes. However, randomized controlled trials were not strong enough to show this effect. Some studies found that CPAP improved 1 or more neurologic function outcomes, sleepiness, or mood,^{13–15,47} whereas the others found no benefit of CPAP treatment in these areas.^{113–115} However, the negative studies had small sample sizes and patients had low CPAP compliance, which may have affected the results.

One of the obstacles of OSA treatment in the stroke population has been poor compliance. There are several reasons for poor compliance, including spontaneous improvement of SDB, lack of excessive daytime sleepiness in some patients,

and motor (facial and bulbar palsy) and cognitive (confusional states, dementia, aphasia) deficits. Compliance has been reported to be as low as 25% to 50% in the acute^{39,53,113,116,117} or sub-acute^{69,74,114} stroke phase, whereas other groups reported rates as high as 70% in the rehabilitation setting.⁵⁶ This finding indicates the importance of CPAP acclimation by skilled respiratory therapists.

In summary, previous research suggests that treatment of OSA with CPAP improves functional recovery during stroke rehabilitation, but that the effect of CPAP treatment on cognitive function in patients with stroke is still unclear.

EARLY TREATMENT OF OBSTRUCTIVE SLEEP APNEA IN CARDIOVASCULAR REHABILITATION

Before the availability of CPAP, the only known definitive therapy for OSA was tracheostomy. One of the earliest studies that showed the impact of treatment of OSA was a retrospective cohort in the 1970s.¹¹⁸ There were 198 patients followed for 7 years. Seventy-one patients received tracheostomy and 127 received conservative treatment consisting of recommended weight loss. The tracheostomy group (as curative treatment) developed considerably fewer vascular complications.¹¹⁸ Another observational cohort study examined the impact of CPAP on long-term cardiovascular outcomes in patients with sleep apnea and showed that CPAP therapy reduces fatal and nonfatal cardiovascular events and protected against death from cardiovascular disease.^{44,119,120}

Short-term randomized trials have been studying systemic blood pressure as the outcome. Although the results have been different with respect to the magnitude of blood pressure reduction, it seems to be a clinically important blood pressure reduction. Those studies showed that patients who had more severe and symptomatic sleep apnea and hypertension experienced more benefit from the longer use of CPAP.^{110,121,122} Patients less likely to experience benefit were those with asymptomatic mild OSA who had normal blood pressure at baseline.^{123,124} In a double-blind, randomized trial comparing therapeutic airway pressurization with sham CPAP, therapeutic airway pressurization was associated with a reduction of 3 to 7 mm Hg in mean 24-hour systemic arterial blood pressure.¹²² In extrapolating from pharmacologic antihypertensive trials, this blood pressure reduction, in itself, is expected to result in a 20% stroke risk reduction.¹²⁵

To date, there are very few data available about the treatment of OSA in the context of cardiac rehabilitation and, overall, the data supporting

the treatment of OSA in patients with CAD are less robust than for systemic hypertension.

A case-control study of 192 patients with acute MI (63 patients without OSA, 52 untreated patients with OSA, and 71 patients with OSA treated with CPAP), over a 6-year follow-up, showed that the risk of recurrent MI and revascularization procedures was lower in treated than in untreated OSA and was similar to patients without OSA.¹²⁶

In another study, in 390 patients who had undergone percutaneous coronary intervention, over a median follow-up of 4.8 years, untreated OSA was independently associated with a significant increased risk of repeat revascularization that was reduced by CPAP treatment.¹²⁷

SDB, in both forms of OSA and CSA, is common in patients with HF and has been suggested to increase the morbidity and mortality in these patients.

The mainstay of OSA treatment in HF is CPAP. Small studies showed that CPAP in patients with HF and OSA reduced 30-day hospital readmission and emergency visits.¹²⁸ It also improved systolic function, blood pressure, and heart rate.^{129–131} In long-term studies, CPAP decreased mortality and improved hospitalization-free survival in patients with HF and OSA.^{132,133} However, there are no double-blind randomized, controlled trials to evaluate these effects, particularly in the rehabilitation setting.

In contrast, studies in patients with HF and CSA suggested that CPAP also can improve EF^{131,134} but does not show any survival benefit.¹³⁴ A recent randomized controlled trial (Adaptive Servo Ventilation in Patients with Heart Failure [SERVE-HF]) examined the effect of adaptive servoventilation in patients with reduced EF and CSA. This trial showed no significant positive effect, but increased both all-cause (HR, 1.28; 95% CI, 1.06–1.55; $P = .01$) and cardiovascular mortality (HR, 1.34; 95% CI, 1.09–1.65; $P = .006$).^{135,136}

Therefore, until the results from further studies are available, optimization of HF management should be the first-line therapy in patients with concurrent CSA. Although the appropriate therapy for CSA in patients with HF remains controversial, CPAP remains widely accepted for OSA in patients with HF.

All of the studies discussed earlier show the beneficial effect of OSA treatment in the context of cardiac rehabilitation, and those patients should be screened early and considered for CPAP treatment until further robust clinical trials have been done.

SUMMARY

Several cohort studies have shown that OSA significantly increases the risk of cardiovascular

disorders independently of potential confounding risk factors. This finding is particularly important given that sleep apnea is a potentially modifiable risk factor. Randomized controlled trials of CPAP in patients with OSA with follow-up of cerebrovascular and cardiovascular outcomes suggest a clinically significant risk reduction associated with the use of CPAP.

In view of potential benefits of noninvasive ventilation on prognosis and functional outcome with improving the quality of life, cognitive function, and acceleration of recovery in the rehabilitation of patients with cardiovascular diseases, clinicians should have a low threshold to evaluate their patients for SDB.

The reasons discussed earlier support the need for cardiac and stroke rehabilitation units to include sleep apnea screening and treatment as an integral component of their care.

REFERENCES

- Young T, Palta M, Dempsey J, et al. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993;328(17):1230–5.
- Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 2002; 165(9):1217–39.
- Young TB. Epidemiology of daytime sleepiness: definitions, symptomatology, and prevalence. *J Clin Psychiatry* 2004;65(Suppl 16):12–6.
- Sharma S, Parker AT. Prevalence of obstructive sleep apnea in a patient population undergoing cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2011;31(3):188–92.
- Romero-Corral A, Somers VK, Pelikka PA, et al. Decreased right and left ventricular myocardial performance in obstructive sleep apnea. *Chest* 2007;132(6):1863–70.
- Bonita R. Epidemiology of stroke. *Lancet* 1992; 339(8789):342–4.
- Yaggi H, Mohsenin V. Obstructive sleep apnoea and stroke. *Lancet Neurol* 2004;3(6):333–42.
- Cheyne JA. A case of apoplexy in which the fleshy part of the heart was converted into fat. *Dublin Hosp Rep* 1818;2:216–22.
- Sahlin C, Sandberg O, Gustafson Y, et al. Obstructive sleep apnea is a risk factor for death in patients with stroke: a 10-year follow-up. *Arch Intern Med* 2008;168(3):297–301.
- Kaneko Y, Hajek VE, Zivanovic V, et al. Relationship of sleep apnea to functional capacity and length of hospitalization following stroke. *Sleep* 2003;26: 293–7.
- Brooks D, Davis L, Vujovic-Zotovic N, et al. Sleep-disordered breathing in patients enrolled in an inpatient stroke rehabilitation program. *Arch Phys Med Rehabil* 2010;91(4):659–62.
- Bratton DJ, Gaisl T, Wons AM, et al. CPAP vs mandibular advancement devices and blood pressure in patients with obstructive sleep apnea: a systematic review and meta-analysis. *JAMA* 2015; 314:2280–93.
- Ryan CM, Bayley M, Green R, et al. Influence of continuous positive airway pressure on outcomes of rehabilitation in stroke patients with obstructive sleep apnea. *Stroke* 2011;42(4):1062–7.
- Sandberg O, Franklin KA, Bucht G, et al. Sleep apnea, delirium, depressed mood, cognition, and ADL ability after stroke. *J Am Geriatr Soc* 2001; 49(4):391–7.
- Parra O, Sanchez-Armengol A, Bonnin M. Early treatment of obstructive sleep apnoea and stroke outcome: a randomised controlled trial. *Eur Respir J* 2011;011(37):1128–36.
- Javaheri S. Sleep disorders in systolic heart failure: a prospective study of 100 male patients. The final report. *Int J Cardiol* 2006;106(1):21–8.
- Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences. *Lancet* 2009;373(9657):82–93.
- Peppard PE, Young T, Palta M, et al. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000; 342(19):1378–84.
- Porthan KM, Melin JH, Kupila JT, et al. Prevalence of sleep apnea syndrome in lone atrial fibrillation: a case-control study. *Chest* 2004;125(3):879–85.
- Young T, Peppard P, Palta M, et al. Population-based study of sleep-disordered breathing as a risk factor for hypertension. *Arch Intern Med* 1997;157(15):1746–52.
- Nieto F, Young T, Lind B, et al. Association of sleep-disordered breathing, sleep apnea, and hypertension in a large community-based study. *JAMA* 2000;283:1229–36.
- Hung J, Whitford EG, Parsons RW, et al. Association of sleep apnoea with myocardial infarction in men. *Lancet* 1990;336(8710):261–4.
- Schafer H, Koehler U, Ewig S, et al. Obstructive sleep apnea as a risk marker in coronary artery disease. *Cardiology* 1999;92(2):79–84.
- Cepeda-Valery B, Acharjee S, Romero-Corral A, et al. Obstructive sleep apnea and acute coronary syndromes: etiology, risk, and management. *Curr Cardiol Rep* 2014;16(10):535.
- Khayat RN, Jarjoura D, Patt B, et al. In-hospital testing for sleep-disordered breathing in hospitalized patients with decompensated heart failure: report of prevalence and patient characteristics. *J Card Fail* 2009;15(9):739–46.
- Oldenburg O, Lamp B, Faber L, et al. Sleep-disordered breathing in patients with symptomatic heart

- failure: a contemporary study of prevalence in and characteristics of 700 patients. *Eur J Heart Fail* 2007;9(3):251–7.
27. Bitter T, Faber L, Hering D, et al. Sleep-disordered breathing in heart failure with normal left ventricular ejection fraction. *Eur J Heart Fail* 2009;11(6):602–8.
 28. Buchner S, Satzl A, Debl K, et al. Impact of sleep-disordered breathing on myocardial salvage and infarct size in patients with acute myocardial infarction. *Eur Heart J* 2014;35(3):192–9.
 29. Glantz H, Thunstrom E, Herlitz J, et al. Occurrence and predictors of obstructive sleep apnea in a revascularized coronary artery disease cohort. *Ann Am Thorac Soc* 2013;10(4):350–6.
 30. Le Grande MR, Neubeck L, Murphy BM, et al. Screening for obstructive sleep apnoea in cardiac rehabilitation: a position statement from the Australian Centre for Heart Health and the Australian Cardiovascular Health and Rehabilitation Association. *Eur J Prev Cardiol* 2016;23(14):1466–75.
 31. Bassetti C, Aldrich MS. Sleep apnea in acute cerebrovascular diseases: final report on 128 patients. *Sleep* 1999;22(2):217–23.
 32. Bassetti C, Aldrich MS, Chervin RD, et al. Sleep apnea in patients with transient ischemic attack and stroke: a prospective study of 59 patients. *Neurology* 1996;47(5):1167–73.
 33. Bassetti C, Aldrich MS, Quint D. Sleep-disordered breathing in patients with acute supra- and infratentorial strokes. A prospective study of 39 patients. *Stroke* 1997;28(9):1765–72.
 34. Parra O, Arboix A, Bechich S, et al. Time course of sleep-related breathing disorders in first-ever stroke or transient ischemic attack. *Am J Respir Crit Care Med* 2000;161(2 Pt 1):375–80.
 35. Arzt M, Young T, Finn L, et al. Association of sleep-disordered breathing and the occurrence of stroke. *Am J Respir Crit Care Med* 2005;172(11):1447–51.
 36. Dyken ME, Somers VK, Yamada T, et al. Investigating the relationship between stroke and obstructive sleep apnea. *Stroke* 1996;27(3):401–7.
 37. Good DC, Henkle JQ, Gelber D, et al. Sleep-disordered breathing and poor functional outcome after stroke. *Stroke* 1996;27(2):252–9.
 38. Parra O. Sleep-disordered breathing and stroke: is there a rationale for treatment? *Eur Respir J* 2001;18(4):619–22.
 39. Broadley SA, Jorgensen L, Cheek A, et al. Early investigation and treatment of obstructive sleep apnoea after acute stroke. *J Clin Neurosci* 2007;14(4):328–33.
 40. Mohsenin V, Valor R. Sleep apnea in patients with hemispheric stroke. *Arch Phys Med Rehabil* 1995;76(1):71–6.
 41. Yaggi H, Concato J, Kernan W, et al. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med* 2005;353:2034–41.
 42. Munoz R, Duran-Cantolla J, Martinez-Vila E, et al. Severe sleep apnea and risk of ischemic stroke in the elderly. *Stroke* 2006;37(9):2317–21.
 43. Young T, Finn L, Peppard PE, et al. Sleep disordered breathing and mortality: eighteen-year follow-up of the Wisconsin sleep cohort. *Sleep* 2008;31(8):1071–8.
 44. Marin JM, Carrizo SJ, Vicente E, et al. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet* 2005;365(9464):1046–53.
 45. Lai SM, Studenski S, Duncan PW, et al. Persisting consequences of stroke measured by the Stroke Impact Scale. *Stroke* 2002;33(7):1840–4.
 46. Dzierwas R, Humpert M, Hopmann B, et al. Increased prevalence of sleep apnea in patients with recurring ischemic stroke compared with first stroke victims. *J Neurol* 2005;252(11):1394–8.
 47. Bravata DM, Concato J, Fried T, et al. Continuous positive airway pressure: evaluation of a novel therapy for patients with acute ischemic stroke. *Sleep* 2011;34(9):1271–7.
 48. Szűcs A, Vitrai J, Janszky J, et al. Pathological sleep apnoea frequency remains permanent in ischaemic stroke and it is transient in haemorrhagic stroke. *Eur Neurol* 2002;47(1):15–9.
 49. Tosun A, Kokturk O, Karata GK, et al. Obstructive sleep apnea in ischemic stroke patients. *Clinics (Sao Paulo)* 2008;63(5):625–30.
 50. Hermann DM, Siccoli M, Kirov P, et al. Central periodic breathing during sleep in acute ischemic stroke. *Stroke* 2007;38(3):1082–4.
 51. Rola R, Wierzbicka A, Wichniak A, et al. Sleep related breathing disorders in patients with ischemic stroke and transient ischemic attacks: respiratory and clinical correlations. *J Physiol Pharmacol* 2007;58 Suppl 5(Pt 2):575–82.
 52. Rowat AM, Dennis MS, Wardlaw JM. Central periodic breathing observed on hospital admission is associated with an adverse prognosis in conscious acute stroke patients. *Cerebrovasc Dis* 2006;21(5–6):340–7.
 53. Bassetti CL, Milanova M, Gugger M. Sleep-disordered breathing and acute ischemic stroke: diagnosis, risk factors, treatment, evolution, and long-term clinical outcome. *Stroke* 2006;37(4):967–72.
 54. Arzt M, Young T, Peppard PE, et al. Dissociation of obstructive sleep apnea from hypersomnolence and obesity in patients with stroke. *Stroke* 2010;41(3):e129–34.
 55. Bennett LS, Stradling JR, Davies RJ. A behavioural test to assess daytime sleepiness in obstructive sleep apnoea. *J Sleep Res* 1997;6(2):142–5.
 56. Wessendorf TE, Wang YM, Thilmann AF, et al. Treatment of obstructive sleep apnoea with nasal

- continuous positive airway pressure in stroke. *Eur Respir J* 2001;18(4):623–9.
57. Punjabi NM, Bandeen-Roche K, Young T. Predictors of objective sleep tendency in the general population. *Sleep* 2003;26(6):678–83.
 58. Javaheri S, Parker TJ, Liming JD, et al. Sleep apnea in 81 ambulatory male patients with stable heart failure. Types and their prevalences, consequences, and presentations. *Circulation* 1998;97(21):2154–9.
 59. Arzt M, Young T, Finn L, et al. Sleepiness and sleep in patients with both systolic heart failure and obstructive sleep apnea. *Arch Intern Med* 2006;166(16):1716–22.
 60. Kryger M, Roth T, Dement WC, editors. *Principles and practice of sleep medicine*. 4th edition. Philadelphia: Elsevier Saunders; 2005.
 61. Duran J, Esnaola S, Rubio R, et al. Obstructive sleep apnea-hypopnea and related clinical features in a population-based sample of subjects aged 30 to 70 yr. *Am J Respir Crit Care Med* 2001;163(3 Pt 1):685–9.
 62. Young T, Shahar E, Nieto FJ, et al. Predictors of sleep-disordered breathing in community-dwelling adults: the Sleep Heart Health Study. *Arch Intern Med* 2002;162(8):893–900.
 63. Gottlieb DJ, Whitney CW, Bonekat WH, et al. Relation of sleepiness to respiratory disturbance index: the Sleep Heart Health Study. *Am J Respir Crit Care Med* 1999;159(2):502–7.
 64. Banack HR, Holly CD, Lowensteyn I, et al. The association between sleep disturbance, depressive symptoms, and health-related quality of life among cardiac rehabilitation participants. *J cardiopulmonary Rehabil Prev* 2014;34(3):188–94.
 65. Redeker NS, Ruggiero J, Hedges C. Patterns and predictors of sleep pattern disturbance after cardiac surgery. *Res Nurs Health* 2004;27(4):217–24.
 66. Hargens TA, Aron A, Newsome LJ, et al. Effects of obstructive sleep apnea on hemodynamic parameters in patients entering cardiac rehabilitation. *J cardiopulmonary Rehabil Prev* 2015;35(3):181–5.
 67. Uchoa CH, Danzi-Soares Nde J, Nunes FS, et al. Impact of OSA on cardiovascular events after coronary artery bypass surgery. *Chest* 2015;147(5):1352–60.
 68. Decary A, Rouleau I, Montplaisir J. Cognitive deficits associated with sleep apnea syndrome: a proposed neuropsychological test battery. *Sleep* 2000;23(3):369–81.
 69. Sandberg O, Franklin KA, Bucht G, et al. Nasal continuous positive airway pressure in stroke patients with sleep apnoea: a randomized treatment study. *Eur Respir J* 2001;18(4):630–4.
 70. Harbison J, Ford GA, James OF, et al. Sleep-disordered breathing following acute stroke. *QJM* 2002;95(11):741–7.
 71. Iranzo A, Santamaria J, Berenguer J, et al. Prevalence and clinical importance of sleep apnea in the first night after cerebral infarction. *Neurology* 2002;58(6):911–6.
 72. Turkington PM. Effect of upper airway obstruction on blood pressure variability after stroke [comment]. *Clin Sci* 2004;107(1):27–8.
 73. Gami AS, Howard DE, Olson EJ, et al. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med* 2005;352(12):1206–14.
 74. Martinez-Garcia MA, Soler-Cataluna JJ, Ejarque-Martinez L, et al. Continuous positive airway pressure treatment reduces mortality in patients with ischemic stroke and obstructive sleep apnea: a 5-year follow-up study. *Am J Respir Crit Care Med* 2009;180(1):36–41.
 75. Parra O, Arboix A, Montserrat JM, et al. Sleep-related breathing disorders: impact on mortality of cerebrovascular disease. *Eur Respir J* 2004;24(2):267–72.
 76. Turkington PM, Bamford J, Wanklyn P, et al. Effect of upper airway obstruction on blood pressure variability after stroke. *Clin Sci (Lond)* 2004;107(1):75–9.
 77. Balfors EM, Franklin KA. Impairment of cerebral perfusion during obstructive sleep apneas. *Am J Respir Crit Care Med* 1994;150(6 Pt 1):1587–91.
 78. Alessi CA, Martin JL, Webber AP, et al. More daytime sleeping predicts less functional recovery among older people undergoing inpatient post-acute rehabilitation. *Sleep* 2008;31(9):1291–300.
 79. Barbar SI, Enright PL, Boyle P, et al. Sleep disturbances and their correlates in elderly Japanese American men residing in Hawaii. *J Gerontol A Biol Sci Med Sci* 2000;55(7):M406–11.
 80. Kutner NG, Schechtman KB, Ory MG, et al. Older adults' perceptions of their health and functioning in relation to sleep disturbance, falling, and urinary incontinence. FICSIT Group. *J Am Geriatr Soc* 1994;42(7):757–62.
 81. Schmitt FA, Phillips BA, Cook YR, et al. Self report on sleep symptoms in older adults: correlates of daytime sleepiness and health. *Sleep* 1996;19(1):59–64.
 82. Alessi CA, Martin JL, Webber AP, et al. Randomized, controlled trial of a nonpharmacological intervention to improve abnormal sleep/wake patterns in nursing home residents. *J Am Geriatr Soc* 2005;53(5):803–10.
 83. Martin JL, Webber AP, Alam T, et al. Daytime sleeping, sleep disturbance, and circadian rhythms in the nursing home. *Am J Geriatr Psychiatry* 2006;14(2):121–9.
 84. Ohayon MM, Vecchierini MF. Normative sleep data, cognitive function and daily living activities in older adults in the community. *Sleep* 2005;28(8):981–9.

85. Wessendorf TE, Teschler H, Wang YM, et al. Sleep-disordered breathing among patients with first-ever stroke. *J Neurol* 2000;247(1):41–7.
86. Hoogerduijn JG, Schuurmans MJ, Duijnste MS, et al. A systematic review of predictors and screening instruments to identify older hospitalized patients at risk for functional decline. *J Clin Nurs* 2007;16(1):46–57.
87. Hayano J, Carney RM, Watanabe E, et al. Interactive associations of depression and sleep apnea with adverse clinical outcomes after acute myocardial infarction. *Psychosom Med* 2012;74(8):832–9.
88. Le Grande MR, Jackson AC, Murphy BM, et al. Relationship between sleep disturbance, depression and anxiety in the 12 months following a cardiac event. *Psychol Health Med* 2016;21(1):52–9.
89. Hamilton BB, Granger CV, Sherwin FS, et al. A uniform national data system for medical rehabilitation. In: Fuhrer M, editor. *Rehabilitation outcomes: analysis and measurement*. Baltimore (MD): Brookes; 1987. p. 137–47.
90. Granger CV, Cotter AC, Hamilton BB, et al. Functional assessment scales: a study of persons after stroke. *Arch Phys Med Rehabil* 1993;74(2):133–8.
91. Turkington PM, Allgar V, Bamford J, et al. Effect of upper airway obstruction in acute stroke on functional outcome at 6 months. *Thorax* 2004;59(5):367–71.
92. Kales A, Caldwell AB, Cadieux RJ, et al. Severe obstructive sleep apnea-II: associated psychopathology and psychosocial consequences. *J Chronic Dis* 1985;38(5):427–34.
93. Leegaard OF. Diffuse cerebral symptoms in convalescents from cerebral infarction and myocardial infarction. *Acta Neurol Scand* 1983;67(6):348–55.
94. Siddiqui F, Macrea M, Horowitz M, et al. Relationship between excessive daytime sleepiness and 6 minute walk test in patients with coronary artery disease and obstructive sleep apnea. *Chest* 2015;148:1064A.
95. Leppavuori A, Pohjasvaara T, Vataja R, et al. Insomnia in ischemic stroke patients. *Cerebrovasc Dis* 2002;14(2):90–7.
96. Gustafson Y, Nilsson I, Mattsson M, et al. Epidemiology and treatment of post-stroke depression. *Drugs Aging* 1995;7(4):298–309.
97. Foley DJ, Monjan AA, Brown SL, et al. Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 1995;18(6):425–32.
98. Findley LJ, Barth JT, Powers DC, et al. Cognitive impairment in patients with obstructive sleep apnea and associated hypoxemia. *Chest* 1986;90(5):686–90.
99. Millman RP, Fogel BS, McNamara ME, et al. Depression as a manifestation of obstructive sleep apnea: reversal with nasal continuous positive airway pressure. *J Clin Psychiatry* 1989;50(9):348–51.
100. Montplaisir J, Bedard MA, Richer F, et al. Neurobehavioral manifestations in obstructive sleep apnea syndrome before and after treatment with continuous positive airway pressure. *Sleep* 1992;15(6 Suppl):S17–9.
101. Vgontzas AN, Liao D, Bixler EO, et al. Insomnia with objective short sleep duration is associated with a high risk for hypertension. *Sleep* 2009;32(4):491–7.
102. Langhorne P, Stott DJ, Robertson L, et al. Medical complications after stroke: a multicenter study. *Stroke* 2000;31(6):1223–9.
103. McManus J, Pathansali R, Hassan H, et al. The course of delirium in acute stroke. *Age Ageing* 2009;38(4):385–9.
104. McManus J, Pathansali R, Stewart R, et al. Delirium post-stroke. *Age Ageing* 2007;36(6):613–8.
105. Gustafson Y, Olsson T, Eriksson S, et al. Acute confusional states in stroke patients. *Cerebrovasc Dis* 1991;1:257–64.
106. Whitney JF, Gannon DE. Obstructive sleep apnea presenting as acute delirium. *Am J Emerg Med* 1996;14(3):270–1.
107. Lee JW. Recurrent delirium associated with obstructive sleep apnea. *Gen Hosp Psychiatry* 1998;20(2):120–2.
108. Munoz X, Marti S, Sumalla J, et al. Acute delirium as a manifestation of obstructive sleep apnea syndrome. *Am J Respir Crit Care Med* 1998;158(4):1306–7.
109. Minnerup J, Ritter MA, Wersching H, et al. Continuous positive airway pressure ventilation for acute ischemic stroke. A randomized feasibility study. *Stroke* 2012;43:1137–9.
110. Becker H, Jerrentrup A, Ploch T, et al. Effect of nasal continuous positive airway pressure treatment on blood pressure in patients with obstructive sleep apnea. *Circulation* 2003;107:68–73.
111. Martinez-Garcia MA, Galiano-Blancart R, Roman-Sanchez P, et al. Continuous positive airway pressure treatment in sleep apnea prevents new vascular events after ischemic stroke. *Chest* 2005;128(4):2123–9.
112. Bravata DM, Concato J, Fried T, et al. Autotitrating continuous positive airway pressure for patients with acute transient ischemic attack: a randomized feasibility trial. *Stroke* 2010;41(7):1464–70.
113. Hui DS, Choy DK, Wong LK, et al. Prevalence of sleep-disordered breathing and continuous positive airway pressure compliance: results in Chinese patients with first-ever ischemic stroke. *Chest* 2002;122(3):852–60.
114. Hsu CY, Vennelle M, Li HY, et al. Sleep-disordered breathing after stroke: a randomised controlled trial

- of continuous positive airway pressure. *J Neurol Neurosurg Psychiatry* 2006;77(10):1143–9.
115. Brown DL, Chervin RD, Kalbfleisch JD, et al. Sleep Apnea Treatment After Stroke (SATS) trial: is it feasible? *J Stroke Cerebrovasc Dis* 2013;22(8):1216–24.
 116. Palombini L, Guilleminault C. Stroke and treatment with nasal CPAP. *Eur J Neurol* 2006;13(2):198–200.
 117. Scala R, Turkington PM, Wanklyn P, et al. Acceptance, effectiveness and safety of continuous positive airway pressure in acute stroke: a pilot study. *Respir Med* 2009;103(1):59–66.
 118. Partinen M, Guilleminault C. Daytime sleepiness and vascular morbidity at seven-year follow-up in obstructive sleep apnea patients. *Chest* 1990;97(1):27–32.
 119. Marti S, Sampol G, Munoz X, et al. Mortality in severe sleep apnoea/hypopnoea syndrome patients: impact of treatment. *Eur Respir J* 2002;20(6):1511–8.
 120. Doherty L, Kiely J, Swan V, et al. Long-term effects of nasal continuous positive airways pressure therapy on cardiovascular outcomes in sleep apnea syndrome. *Chest* 2005;127(6):2076–84.
 121. Faccenda J, Mackay T, Boon N, et al. Randomized placebo-controlled trial of continuous positive airway pressure on blood pressure in the sleep apnea-hypopnea syndrome. *Am J Respir Crit Care Med* 2001;163:344–8.
 122. Pepperell JC, Ramdassingh-Dow S, Crosthwaite N, et al. Ambulatory blood pressure after therapeutic and subtherapeutic nasal continuous positive airway pressure for obstructive sleep apnoea: a randomised parallel trial. *Lancet* 2002;359(9302):204–10.
 123. Barbe F, Mayoralas LR, Duran J, et al. Treatment with continuous positive airway pressure is not effective in patients with sleep apnea but no daytime sleepiness. a randomized, controlled trial. *Ann Intern Med* 2001;134(11):1015–23.
 124. Monasterio C, Vidal S, Duran J, et al. Effectiveness of continuous positive airway pressure in mild sleep apnea-hypopnea syndrome. *Am J Respir Crit Care Med* 2001;164(6):939–43.
 125. Collins R, Peto R, MacMahon S, et al. Blood pressure, stroke, and coronary heart disease. Part 2, short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335(8693):827–38.
 126. Garcia-Rio F, Alonso-Fernandez A, Armada E, et al. CPAP effect on recurrent episodes in patients with sleep apnea and myocardial infarction. *Int J Cardiol* 2013;168(2):1328–35.
 127. Wu X, Lv S, Yu X, et al. Treatment of OSA reduces the risk of repeat revascularization after percutaneous coronary intervention. *Chest* 2015;147(3):708–18.
 128. Kauta SR, Keenan BT, Goldberg L, et al. Diagnosis and treatment of sleep disordered breathing in hospitalized cardiac patients: a reduction in 30-day hospital readmission rates. *J Clin Sleep Med* 2014;10(10):1051–9.
 129. Kaneko Y, Floras JS, Usui K, et al. Cardiovascular effects of continuous positive airway pressure in patients with heart failure and obstructive sleep apnea. *N Engl J Med* 2003;348(13):1233–41.
 130. Mansfield DR, Gollogly NC, Kaye DM, et al. Controlled trial of continuous positive airway pressure in obstructive sleep apnea and heart failure. *Am J Respir Crit Care Med* 2004;169(3):361–6.
 131. Sun H, Shi J, Li M, et al. Impact of continuous positive airway pressure treatment on left ventricular ejection fraction in patients with obstructive sleep apnea: a meta-analysis of randomized controlled trials. *PLoS One* 2013;8(5):e62298.
 132. Wang H, Parker JD, Newton GE, et al. Influence of obstructive sleep apnea on mortality in patients with heart failure. *J Am Coll Cardiol* 2007;49(15):1625–31.
 133. Kasai T, Narui K, Dohi T, et al. Prognosis of patients with heart failure and obstructive sleep apnea treated with continuous positive airway pressure. *Chest* 2008;133(3):690–6.
 134. Bradley TD, Logan AG, Kimoff RJ, et al. Continuous positive airway pressure for central sleep apnea and heart failure. *N Engl J Med* 2005;353(19):2025–33.
 135. Cowie MR, Wegscheider K, Teschler H. Adaptive servo-ventilation for central sleep apnea in heart failure. *N Engl J Med* 2016;374(7):690–1.
 136. Cowie MR, Woehrle H, Wegscheider K, et al. Adaptive servo-ventilation for central sleep apnea in systolic heart failure. *N Engl J Med* 2015;373(12):1095–105.