

UC Berkeley

Working Papers

Title

ON THE INEVITABILITY OF THE CONSISTENCY OPERATOR

Permalink

<https://escholarship.org/uc/item/1fm9h9bq>

Journal

The Journal of Symbolic Logic, 84(1)

ISSN

0022-4812 1943-5886

Authors

Montalbán, Antonio

Walsh, James

Publication Date

2019-03-14

DOI

10.1017/jsl.2018.65

Peer reviewed

ON THE INEVITABILITY OF THE CONSISTENCY OPERATOR

JAMES WALSH AND ANTONIO MONTALBÁN

ABSTRACT. We examine recursive monotonic functions on the Lindenbaum algebra of EA. We prove that no such function sends every consistent φ to a sentence with deductive strength strictly between φ and $(\varphi \wedge \text{Con}(\varphi))$. We generalize this result to iterates of consistency into the effective transfinite. We then prove that for any recursive monotonic function f , if there is an iterate of Con that bounds f everywhere, then f must be somewhere equal to an iterate of Con.

1. INTRODUCTION

It is a well-known empirical phenomenon that natural axiomatic theories are well-ordered by their consistency strength. However, without a precise mathematical definition of “natural,” it is difficult to explain this observation in a strictly mathematical way. One expression of this phenomenon comes from *ordinal analysis*, a research program whereby recursive ordinals are assigned to theories as a measurement of their consistency strength. One method for calculating the proof-theoretic ordinal of a theory T involves demonstrating that T can be approximated over a weak base theory by a class of formulas that are well understood. In particular, the Π_1^0 fragments of natural theories are often proof-theoretically equivalent to iterated consistency statements over a weak base theory, making these theories amenable to ordinal analysis. For discussion, see, e.g., Beklemishev [4, 5] and Joosten [10].

Why are the Π_1^0 fragments of natural theories proof-theoretically equivalent to iterated consistency statements? Our approach to this question is inspired by Martin’s approach to another famous question from mathematical logic: why are natural Turing degrees well-ordered by Turing reducibility? Martin conjectured that (i) the non-constant degree invariant functions meeting a certain simplicity condition ($f \in L(\mathbb{R})$)¹ are pre-well-ordered by the relation “ $f(a) \leq_T g(a)$ on a cone in the Turing degrees” and (ii) the successor for this well-ordering is induced by the Turing jump. Martin’s conjecture is meant to capture the idea that the Turing jump and its iterates into the transfinite are the only natural non-trivial degree invariant functions.

In this paper we investigate analogous hypotheses concerning jumps on consistent axiomatic theories, namely, consistency statements. We fix elementary arithmetic EA as our base theory. EA is a subsystem of PA that is often used as a base theory in ordinal analysis and in which standard approaches to arithmetization of syntax can be carried out without substantial changes; see [6] for details. We write $[\varphi]$ to denote the equivalence class of φ modulo EA-provable equivalence. We write $\varphi \vdash \psi$

2010 *Mathematics Subject Classification*. Primary 03F40.

¹Martin’s Conjecture is stated under the hypothesis ZF + AD + DC, which is satisfied by $L(\mathbb{R})$ assuming that there are ω many Woodin cardinals with a measurable above them all.

if $\text{EA} \vdash \varphi \rightarrow \psi$ and say that φ *implies* ψ . If $\varphi \vdash \psi$ but $\psi \not\vdash \varphi$ we say that φ *strictly implies* ψ . The *Lindenbaum algebra* of EA is the set of equivalence classes of sentences ordered by \vdash . We focus on recursive functions f that are *monotonic*, i.e.,

$$\text{if } \varphi \vdash \psi, \text{ then } f(\varphi) \vdash f(\psi).$$

We note that (i) a function f is monotonic just in case f preserves implication over EA and (ii) all monotonic functions induce functions on the Lindenbaum algebra of EA. We adopt the convention that all functions named “ f ” in this paper are recursive.

Our goal is to demonstrate that $\varphi \mapsto (\varphi \wedge \text{Con}(\varphi))$ and its iterates into the transfinite are canonical among monotonic functions. Our first theorem to this end is the following.

Theorem 1.1. *Let f be monotonic. Suppose that for all consistent φ ,*
(i) $\varphi \wedge \text{Con}(\varphi)$ implies $f(\varphi)$ and
(ii) $f(\varphi)$ strictly implies φ .
Then for every true φ , there is a true ψ such that $\psi \vdash \varphi$ and $[f(\psi)] = [\psi \wedge \text{Con}(\psi)]$.

Corollary 1.2. *There is no monotonic function f such that for all consistent φ ,*
(i) $\varphi \wedge \text{Con}(\varphi)$ strictly implies $f(\varphi)$ and
(ii) $f(\varphi)$ strictly implies φ .

We note that this result depends essentially on the condition of monotonicity. Shavrukov and Visser [13] studied recursive functions f that are *extensional* over the Lindenbaum algebra of PA, i.e.,

$$\text{if } \text{PA} \vdash (\varphi \leftrightarrow \psi), \text{ then } \text{PA} \vdash (f(\varphi) \leftrightarrow f(\psi)),$$

and proved the following theorem.

Theorem 1.3. *(Shavrukov–Visser) There is a recursive extensional function f such that for all consistent φ ,*
(i) $\varphi \wedge \text{Con}(\varphi)$ strictly implies $f(\varphi)$ and
(ii) $f(\varphi)$ strictly implies φ .

In particular, Shavrukov and Visser proved that for any consistent φ , the sentence

$$\varphi^* := \varphi \wedge \forall x (\text{Con}(I\Sigma_x + \varphi) \rightarrow \text{Con}(I\Sigma_x + \varphi + \text{Con}(I\Sigma_x + \varphi)))$$

has deductive strength strictly between φ and $\varphi \wedge \text{Con}(\varphi)$, and that the map $\varphi \mapsto \varphi^*$ is extensional. By a theorem of Kripke and Pour-El [11], the Lindenbaum algebras of PA and EA are effectively isomorphic, whence Theorem 1.3 also applies to EA. Thus, Corollary 1.2 cannot be strengthened by weakening the hypothesis of monotonicity to the hypothesis of extensionality.

We also note that Friedman, Rathjen, and Weiermann [8] introduced a notion of *slow consistency* with which they produced a Π_1^0 sentence $\text{SlowCon}(\text{PA})$ with deductive strength strictly between PA and $\text{PA} + \text{Con}(\text{PA})$. In general, the statement $\text{SlowCon}(\varphi)$ has the form

$$\forall x (F_{e_0}(x) \downarrow \rightarrow \text{Con}(I\Sigma_x + \varphi))$$

where F_{e_0} is a standard representation of a recursive function that is not provably total in PA. This is not in conflict with Corollary 1.2, however, since $\varphi \wedge \text{Con}(\varphi)$ and $\varphi \wedge \text{SlowCon}(\varphi)$ are provably equivalent for all φ such that $\varphi \vdash \forall x F_{e_0}(x) \downarrow$.

On the other hand, changing the definition of the $\text{SlowCon}(\varphi)$ so that the function in the antecedent varies with the input φ results in a map that is not monotonic.

Theorem 1.1 generalizes to the iterates of Con into the effective transfinite. For an elementary presentation α of a recursive well-ordering (see Definition 3.1) and a sentence φ , we define sentences $\text{Con}^\beta(\varphi)$ for every $\beta < \alpha$.

$$\begin{aligned}\text{Con}^0(\varphi) &:= \top \\ \text{Con}^{\beta+1}(\varphi) &:= \text{Con}(\varphi \wedge \text{Con}^\beta(\varphi)) \\ \text{Con}^\lambda(\varphi) &:= \forall \beta < \lambda (\text{Con}^\beta(\varphi))\end{aligned}$$

For a precise definition using Gödel's fixed point lemma, see Definition 3.2. Note that for every φ , $[\text{Con}^1(\varphi)] = [\text{Con}(\varphi)]$.

Remark 1.4. We warn the reader that there is some discrepancy between our notation and the notation used by other authors. Our iteration scheme $\text{Con}^{\alpha+1}(\varphi) \equiv \text{Con}(\varphi \wedge \text{Con}^\alpha(\varphi))$ is sometimes denoted $\text{Con}((\text{EA} + \varphi)_\alpha)$, e.g., [3]. Moreover, the notation $\text{Con}^{\alpha+1}(\varphi)$ is sometimes used to denote $\text{Con}(\text{Con}^\alpha(\varphi))$, e.g., [2].

With each predicate Con^α we associate a function

$$\varphi \mapsto (\varphi \wedge \text{Con}^\alpha(\varphi)).$$

Theorem 1.1 then generalizes into the effective transfinite as follows.

Theorem 1.5. *Let f be monotonic. Suppose that for all φ ,*

(i) $\varphi \wedge \text{Con}^\alpha(\varphi)$ *implies* $f(\varphi)$,

(ii) *if* $[f(\varphi)] \neq [\perp]$, *then* $f(\varphi)$ *strictly implies* $\varphi \wedge \text{Con}^\beta(\varphi)$ *for all* $\beta < \alpha$.

Then for every true φ , there is a true ψ such that $\psi \vdash \varphi$ and $[f(\psi)] = [\psi \wedge \text{Con}^\alpha(\psi)]$.

Corollary 1.6. *There is no monotonic f such that for all φ , if $[\varphi \wedge \text{Con}^\alpha(\varphi)] \neq [\perp]$, then both*

(i) $\varphi \wedge \text{Con}^\alpha(\varphi)$ *strictly implies* $f(\varphi)$ *and*

(ii) $f(\varphi)$ *strictly implies* $\varphi \wedge \text{Con}^\beta(\varphi)$ *for all* $\beta < \alpha$.

Thus, if the range of a monotonic function f is sufficiently constrained, then for some φ and some α ,

$$[f(\varphi)] = [\varphi \wedge \text{Con}^\alpha(\varphi)] \neq [\perp].$$

This property still holds even when these constraints on the range of f are relaxed considerably. More precisely, if a monotonic function is everywhere bounded by a finite iterate of Con , then it must be somewhere equivalent to an iterate of Con .

Theorem 1.7. *Let $n \in \mathbb{N}$. Let f be a monotonic function such that for every φ ,*

(i) $\varphi \wedge \text{Con}^n(\varphi)$ *implies* $f(\varphi)$ *and*

(ii) $f(\varphi)$ *implies* φ .

Then for some φ and some $k \leq n$, $[f(\varphi)] = [\varphi \wedge \text{Con}^k(\varphi)] \neq [\perp]$.

To generalize this result into the effective transfinite, we focus on a particular class of monotonic functions that we call Π_1^0 .

Definition 1.8. A function f is Π_1^0 if $f(\varphi) \in \Pi_1^0$ for all φ .

Our main theorem is the following: if a monotonic function is everywhere bounded by a transfinite iterate of Con , then it must be somewhere equivalent to an iterate of Con . This is to say that the iterates of the consistency operator are *inevitable*; no

monotonic function that is everywhere bounded by some iterate of Con can avoid all of the iterates of Con .

Theorem 1.9. *Let $\varphi \mapsto f(\varphi)$ be a monotonic Π_1^0 function. Then either*

- (i) *for some $\beta \leq \alpha$ and some φ , $[\varphi \wedge f(\varphi)] = [\varphi \wedge \text{Con}^\beta(\varphi)] \neq [\perp]$ or*
- (ii) *for some φ , $(\varphi \wedge \text{Con}^\alpha(\varphi)) \not\vdash f(\varphi)$.*

The main theorem bears a striking similarity to the following theorem of Slaman and Steel [14].

Theorem 1.10. *(Slaman–Steel) Suppose $f : 2^\omega \rightarrow 2^\omega$ is Borel, order-preserving with respect to \leq_T , and increasing on a cone. Then for any $\alpha < \omega_1$ either*

- (i) *for some $\beta \leq \alpha$, $f(x) \equiv_T x^{(\beta)}$ cofinally or*
- (ii) *$(x^{(\alpha)} <_T f(x))$ cofinally.*

There are two notable disanalogies between Theorem 1.9 and Theorem 1.10. First, Theorem 1.9 guarantees only that sufficiently constrained functions are *some-where* equivalent to an iterate of Con , whereas Theorem 1.10 guarantees *cofinal* equivalence with an iterate of the Turing jump. Second, by assuming AD, Slaman and Steel inferred that this behavior happens not only cofinally but also *on a cone* in the Turing degrees. There is no obvious analogue of AD from which one can infer that if cofinally many Lindenbaum degrees have a property then every element in some non-trivial ideal of Lindenbaum degrees has that property.

We then turn our attention to a generalization of consistency, namely, 1-consistency. Recall that a theory T is *1-consistent* if T is consistent with the true Π_1^0 theory of arithmetic. Just as the Π_1^0 fragments of natural theories are often proof-theoretically equivalent to iterated consistency statements over a weak base theory, the Π_2^0 fragments of natural theories are often proof-theoretically equivalent to iterated 1-consistency statements over a weak base theory.

Conservativity theorems relating 1-consistency and iterated consistency play an important role in the proof-theoretic analysis of arithmetic theories. For instance, it is a consequence of Beklemishev’s *reduction principle* [6] that for any Π_1^0 φ ,

$$\text{EA} + 1\text{Con}(\text{EA}) \vdash \varphi \text{ if and only if } \text{EA} + \{\text{Con}^k(\text{EA}) : k < \omega\} \vdash \varphi.$$

This fact plays an integral role in Beklemishev’s [5] consistency proof of PA. We show that this conservativity result is drastically violated *in the limit*. For functions f and g , we say that f *majorizes* g if there is a consistent φ such that for all ψ , if $\psi \vdash \varphi$ then $f(\psi) \vdash g(\psi)$; if in addition φ is true then we say that f *majorizes* g *on a true ideal*.

Proposition 1.11. *For any elementary presentation α of a recursive well-ordering, 1Con majorizes Con^α on a true ideal.*

It is tempting to conjecture on the basis of this result that 1Con is the weakest monotonic function majorizing each Con^α for α a recursive well-ordering. We prove that this is not the case.

Theorem 1.12. *There are infinitely many monotonic functions f such that for every recursive ordinal α , there is an elementary presentation a of α such that f majorizes Con^a on a true ideal but also 1Con majorizes f on a true ideal.*

Theorem 1.1 demonstrates that for any monotonic f with a sufficiently constrained range, f must agree cofinally with Con . We would like to strengthen

cofinally to on a true ideal. One strategy for establishing this claim would be to show that every set that is closed under EA provable equivalence and that contains cofinally many true sentences also contains every sentence in some true ideal. We show that this strategy fails.

Proposition 1.13. *There is a recursively enumerable set \mathcal{A} that contains arbitrarily strong true sentences and that is closed under EA provable equivalence but does not contain any true ideals.*

It is not clear whether Theorem 1.1 can be strengthened in the desired manner.

2. NO MONOTONIC FUNCTION IS STRICTLY BETWEEN THE IDENTITY AND Con

In this section we prove that no monotonic function sends every consistent φ to a sentence with deductive strength strictly between φ and $(\varphi \wedge \text{Con}(\varphi))$. Most of the work is contained in the proof of the following lemma.

Lemma 2.1. *Let f be a monotonic function such that for all consistent φ , $f(\varphi)$ strictly implies φ . Then for every true sentence φ there is a true sentence θ such that $\theta \vdash \varphi$ and $f(\theta) \vdash (\theta \wedge \text{Con}(\theta))$.*

Proof. Let f be as in the statement of the theorem. By assumption the following statement is true.

$$\chi := \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta)))$$

Let φ be a true sentence. Then the sentence $\psi := \varphi \wedge \chi$ is true. Let

$$\theta := (\psi \wedge (f(\psi) \rightarrow \text{Con}(\psi))).$$

Note that $\theta \vdash \varphi$.

Claim 2.2. $f(\theta) \vdash (\theta \wedge f(\psi))$.

Clearly $\theta \vdash \psi$. So $f(\theta) \vdash f(\psi)$ since f is monotonic. Also $f(\theta) \vdash \theta$ by assumption.

Claim 2.3. $(\theta \wedge f(\psi)) \vdash (\psi \wedge \text{Con}(\psi))$.

Immediate from the definition of θ .

Claim 2.4. $(\psi \wedge \text{Con}(\psi)) \vdash (\theta \wedge \text{Con}(\theta))$.

Clearly $(\psi \wedge \text{Con}(\psi)) \vdash \theta$. It suffices to show that

$$(\psi \wedge \text{Con}(\psi)) \vdash \text{Con}(\theta).$$

We reason as follows.

$$\begin{aligned} (\psi \wedge \text{Con}(\psi)) &\vdash \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))) \text{ by choice of } \psi. \\ &\vdash \text{Con}(\psi) \rightarrow \text{Con}(\psi \wedge \neg f(\psi)) \text{ by instantiation.} \\ &\vdash \text{Con}(\psi \wedge \neg f(\psi)) \text{ by logic.} \\ &\vdash \text{Con}(\theta) \text{ by the definition of } \theta. \end{aligned}$$

It is immediate from the preceding claims that $f(\theta) \vdash (\theta \wedge \text{Con}(\theta))$. \square

A number of results follow immediately from the lemma.

Theorem 2.5 (Restatement of Theorem 1.1). *Let f be monotonic. Suppose that for all consistent φ ,*

- (i) $\varphi \wedge \text{Con}(\varphi)$ implies $f(\varphi)$ and
- (ii) $f(\varphi)$ strictly implies φ .

Then for every true φ , there is a true ψ such that $\psi \vdash \varphi$ and $[f(\psi)] = [\psi \wedge \text{Con}(\psi)]$.

Proof. By the lemma, for every true φ there is a true ψ such that $\psi \vdash \varphi$ and $f(\psi) \vdash (\psi \wedge \text{Con}(\psi))$. Since we are assuming that $(\psi \wedge \text{Con}(\psi)) \vdash f(\psi)$, it follows that $[f(\psi)] = [\psi \wedge \text{Con}(\psi)]$. \square

We note that this theorem applies to a number of previously studied operators. For instance, the theorem applies to the notion of *cut-free consistency*, i.e., consistency with respect to cut-free proofs. EA does not prove the cut-elimination theorem, which is equivalent to the totality of super-exponentiation (over EA), and does not prove the equivalence of cut-free consistency and consistency. Another such operator is the Friedman-Rathjen-Weiermann *slow consistency operator* discussed in §1. Theorem 2.5 implies that these operators exhibit the same behavior as the consistency operator “in the limit.” Indeed, for any φ such that φ proves the cut-elimination theorem, $\varphi \wedge \text{Con}(\varphi)$ and $\varphi \wedge \text{Con}_{\text{CF}}(\varphi)$ are EA-provably equivalent. Likewise, for any φ that proves the totality of F_{e_0} , $\varphi \wedge \text{Con}(\varphi)$ and $\varphi \wedge \text{SlowCon}(\varphi)$ are EA-provably equivalent.

As a corollary of Theorem 2.5 we note that no monotonic function reliably produces sentences strictly between those produced by the identity and by Con.

Corollary 2.6 (Restatement of Corollary 1.2). *There is no monotonic function f such that for all consistent φ ,*

- (i) $\varphi \wedge \text{Con}(\varphi)$ strictly implies $f(\varphi)$ and
- (ii) $f(\varphi)$ strictly implies φ .

Shavrukov and Visser [13] studied functions over Lindenbaum algebras and discovered a recursive *extensional uniform density function* g for the Lindenbaum algebra of EA, i.e., (i) for any φ and ψ such that ψ strictly implies φ , $g(\langle \varphi, \psi \rangle)$ is a sentence with deductive strength strictly between φ and ψ and (ii) if $\text{EA} \vdash (\varphi \leftrightarrow \psi)$ then, for any θ , $[g(\langle \varphi, \theta \rangle)] = [g(\langle \psi, \theta \rangle)]$ and $[g(\langle \theta, \varphi \rangle)] = [g(\langle \theta, \psi \rangle)]$. They asked whether this result could be strengthened by exhibiting a recursive uniform density function that is monotonic in both its coordinates. As a corollary of our theorem we answer their question negatively.

Corollary 2.7. *There is no monotonic uniform density function for the Lindenbaum algebra of EA.*

Proof. Suppose there were such a function g over the Lindenbaum algebra of EA. Then given any input of the form $\langle \varphi, (\varphi \wedge \text{Con}(\varphi)) \rangle$, g would produce a sentence with deductive strength strictly between φ and $(\varphi \wedge \text{Con}(\varphi))$. We then note that $f : \varphi \mapsto g(\langle \varphi, (\varphi \wedge \text{Con}(\varphi)) \rangle)$ is monotonic, but that for every consistent φ , $\varphi \wedge \text{Con}(\varphi)$ strictly implies $f(\varphi)$ and $f(\varphi)$ strictly implies φ , contradicting the previous theorem. \square

Our negative answer to the question raised by Shavrukov and Visser makes use of a Π_2^0 sentence $\forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta)))$. Shavrukov and Visser raised the following question in private communication.

Question 2.8. *Is there a recursive uniform density function for the lattice of Π_1^0 sentences over EA that is monotonic in both its coordinates?*

Remark 2.9. It is clear from the proof of the lemma that any monotonic f meeting the hypotheses of Theorem 2.5 is not only cofinally equivalent to Con ; for every true ψ that implies

$$\chi := \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))),$$

there is a true θ such that $\theta \vdash \psi$ and $[\psi \wedge \text{Con}(\psi)] = [\theta \wedge \text{Con}(\theta)] = [f(\theta)]$.

This observation points the way toward a corollary of our theorem; namely that any monotonic function strictly meeting the hypotheses of the theorem must have the same range as $\varphi \mapsto (\varphi \wedge \text{Con}(\varphi))$ in the limit. To prove this, we first prove a version of jump inversion— $\varphi \mapsto (\varphi \wedge \text{Con}(\varphi))$ inversion—for Lindenbaum algebras. This is to say that the range of Con contains a true ideal in the Lindenbaum algebra. A similar result is established for true Π_2^0 sentences in [1].

Proposition 2.10. *Suppose $\varphi \vdash \text{Con}(\top)$. Then for some ψ , $[\varphi] = [(\psi \wedge \text{Con}(\psi))]$.*

Proof. Let $\psi := \text{Con}(\top) \rightarrow \varphi$.

Claim 2.11. $\varphi \vdash (\psi \wedge \text{Con}(\psi))$.

Trivially, $\varphi \vdash \psi$. Since $\varphi \vdash \text{Con}(\top)$, it follows that from the formalized second incompleteness theorem, i.e., $\text{Con}(\top) \vdash \text{Con}(\neg \text{Con}(\top))$, that $\varphi \vdash \text{Con}(\neg \text{Con}(\top))$. But $\neg \text{Con}(\top)$ is the first disjunct of ψ , so $\varphi \vdash \text{Con}(\psi)$.

Claim 2.12. $(\psi \wedge \text{Con}(\psi)) \vdash \varphi$.

Note that $\text{Con}(\psi) \vdash \text{Con}(\top)$. The claim then follows since clearly $(\psi \wedge \text{Con}(\top)) \vdash \varphi$. \square

Corollary 2.13. *Let f be monotonic. Suppose that for all consistent φ ,*

(i) $\varphi \wedge \text{Con}(\varphi)$ implies $f(\varphi)$ and

(ii) $f(\varphi)$ strictly implies φ .

Then the intersection of the ranges of f and Con in the Lindenbaum algebra contains a true ideal.

Proof. Let φ be a sentence such that $\varphi \vdash \text{Con}(\top)$ and

$$\varphi \vdash \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))).$$

Note that both of these sentences are true, and hence φ is in an element of a true ideal. By the previous proposition, there is a ψ such that $[\psi \wedge \text{Con}(\psi)] = [\varphi]$. By Remark 2.9 there is a θ such that $[f(\theta)] = [\psi \wedge \text{Con}(\psi)]$, that is, φ is in the range of f . \square

3. ITERATING Con INTO THE TRANSFINITE

By analogy with Martin's Conjecture, we would like to show that there is a natural well-ordered hierarchy of monotonic functions and that the successor for this well-ordering is induced by Con . Thus, we define the iterates of Con along elementary presentations of well-orderings.

Definition 3.1. By an *elementary presentation* of a recursive well-ordering we mean a pair $(\mathcal{D}, <)$ of elementary formulas, such that (i) the relation $<$ well-orders \mathcal{D} in the standard model of arithmetic and (ii) EA proves that $<$ linearly orders the elements satisfying \mathcal{D} , (iii) it is elementarily calculable whether an element represents zero or a successor or a limit and (iv) the elementary formulas defining

the set of limit ordinals and the successor relation provably in EA satisfy their corresponding first order definitions in terms of $<$.

Definition 3.2. Given an elementary presentation $\langle \alpha, < \rangle$ of a recursive well-ordering and a sentence φ , we use Gödel's fixed point lemma to define sentences $\mathbf{Con}^*(\varphi, \beta)$ for $\beta < \alpha$ as follows.

$$\mathbf{EA} \vdash \mathbf{Con}^*(\varphi, \beta) \leftrightarrow \forall \gamma < \beta, \mathbf{Con}(\varphi \wedge \mathbf{Con}^*(\varphi, \gamma)).$$

We use the notation $\mathbf{Con}^\beta(\varphi)$ for $\mathbf{Con}^*(\varphi, \beta)$.

Remark 3.3. Note that, since the following clauses are provable in EA.

- $\mathbf{Con}^0(\varphi) \leftrightarrow \top$
- $\mathbf{Con}^{\gamma+1}(\varphi) \leftrightarrow \mathbf{Con}(\varphi \wedge \mathbf{Con}^\gamma(\varphi))$
- $\mathbf{Con}^\lambda(\varphi) \leftrightarrow \forall \gamma < \lambda, \mathbf{Con}^\gamma(\varphi)$ for λ a limit.

Note that this hierarchy is proper for true φ by Gödel's second incompleteness theorem. We need to prove that for transfinite α , \mathbf{Con}^α is monotonic over the Lindenbaum algebra of EA. Before proving this claim we recall Schmerl's [12] technique of *reflexive transfinite induction*. Note that " $\text{Pr}(\varphi)$ " means that φ is provable in EA.

Proposition 3.4. (*Schmerl*) *Suppose that $<$ is an elementary linear order and that $\mathbf{EA} \vdash \forall \alpha (\text{Pr}(\forall \beta < \alpha, A(\beta)) \rightarrow A(\alpha))$. Then $\mathbf{EA} \vdash \forall \alpha A(\alpha)$.*

Proof. From $\mathbf{EA} \vdash \forall \alpha (\text{Pr}(\forall \beta < \alpha, A(\beta)) \rightarrow A(\alpha))$ we infer

$$\begin{aligned} \mathbf{EA} \vdash \text{Pr}(\forall \alpha A(\alpha)) &\rightarrow \forall \alpha \text{Pr}(\forall \beta < \alpha, A(\beta)) \\ &\rightarrow \forall \alpha A(\alpha). \end{aligned}$$

Löb's theorem, i.e.,

$$\text{if } \mathbf{EA} \vdash \text{Pr}(\zeta) \rightarrow \zeta, \text{ then } \mathbf{EA} \vdash \zeta,$$

then yields $\mathbf{EA} \vdash \forall \alpha A(\alpha)$. □

Proposition 3.5. *If $\varphi \vdash \psi$, then $\mathbf{Con}^\alpha(\varphi) \vdash \mathbf{Con}^\alpha(\psi)$.*

Proof. Let $\mathcal{A}(\beta)$ denote the claim that $\mathbf{Con}^\beta(\varphi) \vdash \mathbf{Con}^\beta(\psi)$.

We want to prove that $\mathcal{A}(\alpha)$, without placing any restrictions on α . We prove the equivalent claim that $\mathbf{EA} \vdash \mathcal{A}(\alpha)$. By Proposition 3.4, it suffices to show that

$$\mathbf{EA} \vdash \forall \alpha (\text{Pr}(\forall \beta < \alpha, \mathcal{A}(\beta)) \rightarrow \mathcal{A}(\alpha)).$$

Reason within EA. Suppose that $\text{Pr}(\forall \beta < \alpha, \mathcal{A}(\beta))$, which is to say that

$$\text{Pr}(\forall \beta < \alpha, \text{Pr}(\mathbf{Con}^\beta(\varphi) \rightarrow \mathbf{Con}^\beta(\psi))).$$

Since $\mathbf{Con}^\alpha(\varphi)$ contains EA, we infer that

$$\mathbf{Con}^\alpha(\varphi) \vdash \forall \beta < \alpha \text{Pr}(\mathbf{Con}^\beta(\varphi) \rightarrow \mathbf{Con}^\beta(\psi)).$$

Since $\mathbf{Con}^\alpha(\varphi)$ proves that for all $\beta < \alpha$, $\mathbf{EA} \not\vdash \neg \mathbf{Con}^\beta(\varphi)$ we infer that

$$\mathbf{Con}^\alpha(\varphi) \vdash \forall \beta < \alpha \mathbf{Con}(\mathbf{Con}^\beta(\psi)).$$

Thus,

$$\mathbf{Con}^\alpha(\varphi) \vdash \forall \beta < \alpha (\mathbf{Con}^\beta(\psi)).$$

This concludes the proof of the proposition. □

Thus, for each predicate Con^α the function

$$\varphi \mapsto (\varphi \wedge \text{Con}^\alpha(\varphi))$$

is monotonic over the Lindenbaum algebra of EA.

In this section we show that the functions given by iterated consistency are minimal with respect to each other. We fix an elementary presentation α of a recursive well-ordering. We assume that f is a monotonic function such that for every consistent φ , $f(\varphi)$ strictly implies $\varphi \wedge \text{Con}^\beta(\varphi)$ for all $\beta < \alpha$. We would like to relativize the proof of Lemma 2.1 to Con^β . However, the proof of Lemma 2.1 relied on the truth of the principle

$$\forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))).$$

It is not in general clear that $\text{Con}^\alpha(\varphi)$ implies $\text{Con}^\alpha(\varphi \wedge \neg f(\varphi))$. To solve this problem, we define a sequence of true sentences $(\theta_\beta)_{\beta \leq \alpha}$ such that for every sentence φ , if $\varphi \vdash \theta_\beta$ then $\text{Con}^\beta(\varphi)$ implies $\text{Con}^\beta(\varphi \wedge \neg f(\varphi))$. Thus, we are able to relativize the proof of Lemma 2.1 for Con^β to sentences that imply θ_β .

Definition 3.6. Given an elementary presentation α of a recursive well-ordering, we use Gödel's fixed point lemma to define sentences $\theta^*(\beta)$ for $\beta < \alpha$ as follows.

EA $\vdash \theta^*(\beta) \leftrightarrow$

$$\forall \gamma < \beta (\text{True}_{\Pi_3}(\theta^*(\gamma))) \wedge \forall \zeta \left((\forall \gamma < \beta \text{Pr}(\zeta \rightarrow \theta^*(\gamma))) \rightarrow (\text{Con}^\beta(\zeta) \rightarrow \text{Con}^\beta(\zeta \wedge \neg f(\zeta))) \right).$$

We use the notation θ_β for $\theta^*(\beta)$.

Remark 3.7. Note that every sentence in the sequence $(\theta_\beta)_{\beta \leq \alpha}$ has complexity Π_3^0 . Note moreover that for a successor $\beta + 1$, $\theta_{\beta+1}$ is equivalent to

$$\theta_\beta \wedge \forall \zeta (\text{Pr}(\zeta \rightarrow \theta_\beta) \rightarrow (\text{Con}^{\beta+1}(\zeta) \rightarrow \text{Con}^{\beta+1}(\zeta \wedge \neg f(\zeta)))).$$

Lemma 3.8. Let f be monotonic such that, for all φ ,

(i) $\varphi \wedge \text{Con}^\alpha(\varphi)$ implies $f(\varphi)$,

(ii) if $[f(\varphi)] \neq [\perp]$, then $f(\varphi)$ strictly implies $\varphi \wedge \text{Con}^\beta(\varphi)$ for all $\beta < \alpha$.

Then for each $\beta \leq \alpha$, the sentence θ_β is true.

Proof. Let f be as in the statement of the lemma. We prove the claim by induction on $\beta \leq \alpha$. The **base case** $\beta = 0$ is trivial.

For the **successor case** we assume that $\beta < \alpha$ and that θ_β is true; we want to show that $\theta_{\beta+1}$ is true. So let ζ be a sentence such that $\zeta \vdash \theta_\beta$. We want to show that $\text{Con}^{\beta+1}(\zeta)$ implies $\text{Con}^{\beta+1}(\zeta \wedge \neg f(\zeta))$. We prove the contrapositive, that $\neg \text{Con}^{\beta+1}(\zeta \wedge \neg f(\zeta))$ implies $\neg \text{Con}^{\beta+1}(\zeta)$. So suppose $\neg \text{Con}^{\beta+1}(\zeta \wedge \neg f(\zeta))$, i.e.,

$$(\dagger) \quad \zeta \wedge \neg f(\zeta) \vdash \neg \text{Con}^\beta(\zeta \wedge \neg f(\zeta)).$$

We reason as follows.

Since $\zeta \vdash \theta_\beta$, $\zeta \vdash \forall \gamma < \beta, \text{True}_{\Pi_3}(\theta_\gamma)$. From this we infer

$$(\star) \quad \zeta \vdash \text{Pr}(\zeta \rightarrow \forall \gamma < \beta, \text{True}_{\Pi_3}(\theta_\gamma))$$

by Σ_1^0 completeness. Moreover, since $\zeta \vdash \theta_\beta$,

$$\begin{aligned} & \zeta \vdash \forall \varphi ((\forall \gamma < \beta \text{Pr}(\varphi \rightarrow \theta_\gamma)) \rightarrow (\text{Con}^\beta(\varphi) \rightarrow \text{Con}^\beta(\varphi \wedge \neg f(\varphi)))) \text{ by the definition of } \theta_\beta. \\ & \vdash \forall \gamma < \beta \text{Pr}(\zeta \rightarrow \theta_\gamma) \rightarrow (\text{Con}^\beta(\zeta) \rightarrow \text{Con}^\beta(\zeta \wedge \neg f(\zeta))) \text{ by instantiation.} \\ & \vdash \text{Con}^\beta(\zeta) \rightarrow \text{Con}^\beta(\zeta \wedge \neg f(\zeta)) \text{ by } (\star). \end{aligned}$$

$\zeta \wedge \neg f(\zeta) \vdash \neg \text{Con}^\beta(\zeta \wedge \neg f(\zeta))$ by (\dagger) .

$$\vdash \neg \text{Con}^\beta(\zeta) \text{ by logic.}$$

$$\zeta \vdash \text{Con}^\beta(\zeta) \rightarrow f(\zeta) \text{ by logic.}$$

Thus, $(\zeta \wedge \text{Con}^\beta(\zeta)) \vdash f(\zeta)$. Since $f(\varphi)$ always strictly implies $\varphi \wedge \text{Con}^\beta(\varphi)$, we infer that

$$[\zeta \wedge \text{Con}^\beta(\zeta)] = [\perp].$$

This is to say that $\neg \text{Con}^{\beta+1}(\zeta)$.

For the **limit case** we let β be a limit ordinal and assume that for every $\gamma < \beta$, θ_γ is true. We want to show that θ_β is true. Let ζ be a sentence such that for every $\gamma < \beta$, $\zeta \vdash \theta_\gamma$. We want to show that $\text{Con}^\beta(\zeta)$ implies $\text{Con}^\beta(\zeta \wedge \neg f(\zeta))$. So assume that $\text{Con}^\beta(\zeta)$, i.e., for every $\gamma < \beta$, $\text{Con}^\gamma(\zeta)$. Let $\gamma < \beta$. Since β is a limit ordinal, $\gamma + 1 < \beta$. So by the inductive hypothesis $\theta_{\gamma+1}$ is true. That is, by the definition of $\theta_{\gamma+1}$,

$$\forall \varphi (\text{Pr}(\varphi \rightarrow \theta_\gamma) \rightarrow (\text{Con}^\gamma(\varphi) \rightarrow \text{Con}^\gamma(\varphi \wedge \neg f(\varphi)))).$$

By instantiation, we infer that

$$\text{Pr}(\zeta \rightarrow \theta_\gamma) \rightarrow (\text{Con}^\gamma(\zeta) \rightarrow \text{Con}^\gamma(\zeta \wedge \neg f(\zeta))).$$

Since $\zeta \vdash \theta_\gamma$ and $\text{Con}^\gamma(\zeta)$, this means that $\text{Con}^\gamma(\zeta \wedge \neg f(\zeta))$. Since γ was a generic ordinal less than β , we get that

$$\forall \gamma < \beta, \text{Con}^\gamma(\zeta \wedge \neg f(\zeta)),$$

i.e., $\text{Con}^\beta(\zeta)$. This completes the proof of the lemma. \square

Theorem 3.9 (Restatement of Theorem 1.5). *Let f be monotonic. Suppose that for all φ ,*

(i) $\varphi \wedge \text{Con}^\alpha(\varphi)$ implies $f(\varphi)$,

(ii) if $[f(\varphi)] \neq [\perp]$, then $f(\varphi)$ strictly implies $\varphi \wedge \text{Con}^\beta(\varphi)$ for all $\beta < \alpha$.

Then for every true χ , there is a true ψ such that $\psi \vdash \chi$ and $[f(\psi)] = [\psi \wedge \text{Con}^\alpha(\psi)]$.

Proof. Let χ be a true sentence. By the lemma, θ_α is true. So

$$\varphi := \chi \wedge \theta_\alpha$$

is true. We let

$$\psi := \varphi \wedge (f(\varphi) \rightarrow \text{Con}^\alpha(\varphi)).$$

Note that $\psi \vdash \chi$. We now show that $[\psi \wedge \text{Con}^\alpha(\psi)] = [f(\psi)]$.

Claim 3.10. $f(\psi) \vdash (\psi \wedge f(\varphi))$.

Since f is monotonic.

Claim 3.11. $(\psi \wedge f(\varphi)) \vdash (\varphi \wedge \text{Con}^\alpha(\varphi))$.

By the definition of ψ .

Claim 3.12. $(\varphi \wedge \text{Con}^\alpha(\varphi)) \vdash (\psi \wedge \text{Con}^\alpha(\psi))$.

It is clear from the definition of ψ that $(\varphi \wedge \text{Con}^\alpha(\varphi)) \vdash \psi$. So it suffices to show that $(\varphi \wedge \text{Con}^\alpha(\varphi)) \vdash \text{Con}^\alpha(\psi)$.

$$\begin{aligned} \varphi \wedge \text{Con}^\alpha(\varphi) &\vdash \forall \zeta ((\forall \beta < \alpha \text{Pr}(\zeta \rightarrow \theta_\beta)) \rightarrow (\text{Con}^\alpha(\zeta) \rightarrow \text{Con}^\alpha(\zeta \wedge \neg f(\zeta)))) \text{ by choice of } \varphi. \\ &\vdash \forall \beta < \alpha \text{Pr}(\varphi \rightarrow \theta_\beta) \rightarrow (\text{Con}^\alpha(\varphi) \rightarrow \text{Con}^\alpha(\varphi \wedge \neg f(\varphi))) \text{ by instantiation.} \\ &\vdash \forall \beta < \alpha \text{Pr}(\varphi \rightarrow \theta_\beta) \rightarrow \text{Con}^\alpha(\varphi \wedge \neg f(\varphi)) \text{ by logic.} \end{aligned}$$

Since $\text{Con}^\alpha(\varphi \wedge \neg f(\varphi)) \vdash \text{Con}^\alpha(\psi)$, to prove the desired claim it suffices to show that

$$\varphi \wedge \text{Con}^\alpha(\varphi) \vdash \forall \beta < \alpha \text{Pr}(\varphi \rightarrow \theta_\beta).$$

We reason as follows.

$$\begin{aligned} \varphi &\vdash \theta_\alpha \text{ by choice of } \varphi. \\ &\vdash \forall \beta < \alpha (\text{True}_{\Pi_3} \theta_\beta) \text{ by definition of } \theta_\alpha. \\ &\vdash \text{Pr}(\varphi \rightarrow \forall \beta < \alpha (\text{True}_{\Pi_3} \theta_\beta)) \text{ by } \Sigma_1^0 \text{ completeness.} \\ &\vdash \forall \beta < \alpha \text{Pr}(\varphi \rightarrow \text{True}_{\Pi_3} \theta_\beta) \\ &\vdash \forall \beta < \alpha \text{Pr}(\varphi \rightarrow \theta_\beta) \end{aligned}$$

It is immediate from the preceding claims that $f(\psi) \vdash \psi \wedge \text{Con}^\alpha(\psi)$. By assumption, $\psi + \text{Con}^\alpha(\psi) \vdash f(\psi)$, so it follows that $[f(\psi)] = [\psi \wedge \text{Con}^\alpha(\psi)]$. \square

Corollary 3.13 (Restatement of Corollary 1.6). *There is no monotonic f such that for all φ , if $[\varphi \wedge \text{Con}^\alpha(\varphi)] \neq [\perp]$, then both*

- (i) $\varphi \wedge \text{Con}^\alpha(\varphi)$ strictly implies $f(\varphi)$ and
- (ii) $f(\varphi)$ strictly implies $\varphi \wedge \text{Con}^\beta(\varphi)$ for all $\beta < \alpha$.

4. FINITE ITERATES OF Con ARE INEVITABLE

In this section and the next section we prove that the iterates of Con are, in a sense, inevitable. First we show that, for every natural number n , if a monotonic function f is always bounded by Con^n , then it is somewhere equivalent to Con^k for some $k \leq n$. In §5, we turn to generalizations of this result into the effective transfinite.

Theorem 4.1 (Restatement of Theorem 1.7). *Let $n \in \mathbb{N}$. Let f be a monotonic function such that for every φ ,*

- (i) $\varphi \wedge \text{Con}^n(\varphi)$ implies $f(\varphi)$ and
- (ii) $f(\varphi)$ implies φ .

Then for some φ and some $k \leq n$, $[f(\varphi)] = [\varphi \wedge \text{Con}^k(\varphi)] \neq [\perp]$.

Proof. We suppose, towards a contradiction, that there is no ψ and no $k \leq n$ such that $[f(\psi)] = [\psi \wedge \text{Con}^k(\psi)] \neq [\perp]$. We then let φ_1 be a true statement such that

$$\begin{aligned} \varphi_1 &\vdash \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))) \\ \varphi_1 &\vdash \forall k \forall \zeta (\text{Con}^{k+1}(\zeta) \rightarrow \neg \text{Pr}((\zeta \wedge \text{Con}^k(\zeta)) \leftrightarrow f(\zeta))). \end{aligned}$$

The first condition is that φ_1 proves that for every consistent φ , $f(\varphi)$ strictly implies φ . The second condition is that φ_1 proves that $f(\zeta)$ never coincides with $\zeta \wedge \text{Con}^k(\zeta)$, unless $[\zeta \wedge \text{Con}^k(\zeta)] = [\perp]$.

We define a sequence of statements, starting with φ_1 , as follows:

$$\varphi_{k+1} := \varphi_k \wedge (f(\varphi_k) \rightarrow \text{Con}^k(\varphi_k)).$$

We will use our assumption to show that, for all k , $\varphi_k \wedge \text{Con}^k(\varphi_k) \vdash \text{Con}^k(\varphi_{k+1})$. From this we will deduce that $[f(\varphi_{n+1})] = [\varphi_{n+1} \wedge \text{Con}^n(\varphi_{n+1})] \neq [\perp]$, contradicting the assumption that f and Con^n never coincide. Most of the work is contained in the proof of the following lemma.

Lemma 4.2. *For all k , for all $j \geq k$, $(\varphi_k \wedge \text{Con}^k(\varphi_k)) \vdash \text{Con}^k(\varphi_j)$.*

Proof. We prove the claim by a double induction. The primary induction is on k . For the **base case** $k = 1$, we prove the claim by induction on j . The *base case* $j = 1$ follows trivially. For the *inductive step* we assume that $(\varphi_1 \wedge \text{Con}(\varphi_1)) \vdash \text{Con}(\varphi_j)$ and show that $(\varphi_1 \wedge \text{Con}(\varphi_1)) \vdash \text{Con}(\varphi_{j+1})$.

$$\begin{aligned} \varphi_1 \wedge \text{Con}(\varphi_1) &\vdash \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))) \text{ by choice of } \varphi_1. \\ &\vdash \text{Con}(\varphi_j) \rightarrow \text{Con}(\varphi_j \wedge \neg f(\varphi_j)) \text{ by instantiation.} \\ \varphi_1 \wedge \text{Con}(\varphi_1) &\vdash \text{Con}(\varphi_j) \text{ by the inductive hypothesis.} \\ &\vdash \text{Con}(\varphi_j \wedge \neg f(\varphi_j)) \text{ by logic.} \\ &\vdash \text{Con}(\varphi_{j+1}) \text{ by definition of } \varphi_{j+1}. \end{aligned}$$

For the **inductive step** we assume that the claim is true of $k - 1$, i.e.,

$$\forall j \geq k - 1 ((\varphi_{k-1} \wedge \text{Con}^{k-1}(\varphi_{k-1})) \vdash (\text{Con}^{k-1}(\varphi_j))).$$

We prove the claim for k . Once again, we prove the claim by induction on j . The *base case* $j = k$ follows trivially. For the *inductive step* we assume that $\varphi_k \wedge \text{Con}^k(\varphi_k) \vdash \text{Con}^k(\varphi_j)$. We want to prove that $\varphi_k \wedge \text{Con}^k(\varphi_k) \vdash \text{Con}^k(\varphi_{j+1})$.

$$\begin{aligned} \varphi_k \wedge \text{Con}^k(\varphi_k) &\vdash \forall x \forall \zeta (\text{Con}^{x+1}(\zeta) \rightarrow \neg \text{Pr}((\zeta \wedge \text{Con}^x(\zeta)) \leftrightarrow f(\zeta))) \text{ by choice of } \varphi_1. \\ &\vdash \text{Con}^k(\varphi_j) \rightarrow \neg \text{Pr}((\varphi_j \wedge \text{Con}^{k-1}(\varphi_j)) \leftrightarrow f(\varphi_j)) \text{ by instantiation.} \\ \varphi_k \wedge \text{Con}^k(\varphi_k) &\vdash \text{Con}^k(\varphi_j) \text{ by the inner inductive hypothesis.} \\ &\vdash \neg \text{Pr}((\varphi_j \wedge \text{Con}^{k-1}(\varphi_j)) \leftrightarrow f(\varphi_j)) \text{ by logic.} \end{aligned}$$

Thus, $\varphi_k \wedge \text{Con}^k(\varphi_k)$ proves that one of the following cases holds.

$$(\varphi_j \wedge \text{Con}^{k-1}(\varphi_j)) \not\vdash f(\varphi_j)$$

$$f(\varphi_j) \not\vdash (\varphi_j \wedge \text{Con}^{k-1}(\varphi_j))$$

We now show that $\varphi_k \wedge \text{Con}^k(\varphi_k)$ refutes the second option.

Claim 4.3. $\varphi_k \wedge \text{Con}^k(\varphi_k) \vdash \text{Pr}(f(\varphi_j) \rightarrow (\varphi_j \wedge \text{Con}^{k-1}(\varphi_j)))$.

By the outer inductive hypothesis, EA proves the following conditional:

$$\theta := ((\varphi_{j-1} \wedge \text{Con}^{k-1}(\varphi_{j-1})) \rightarrow (\text{Con}^{k-1}(\varphi_j))).$$

Thus, $f(\varphi_j)$ (which contains EA) also proves θ . We now show that $f(\varphi_j) \vdash \text{Con}^{k-1}(\varphi_j)$.

$$\begin{aligned} f(\varphi_j) &\vdash \varphi_j \wedge f(\varphi_{j-1}) \text{ since } f \text{ is monotonic.} \\ &\vdash (\varphi_{j-1} \wedge (f(\varphi_{j-1}) \rightarrow \text{Con}^{j-1}(\varphi_{j-1}))) \wedge f(\varphi_{j-1}) \text{ by the definition of } \varphi_j. \\ &\vdash \varphi_{j-1} \wedge \text{Con}^{j-1}(\varphi_{j-1}) \text{ by logic.} \\ &\vdash \varphi_{j-1} \wedge \text{Con}^{k-1}(\varphi_{j-1}) \text{ since } j \geq k. \\ &\vdash \text{Con}^{k-1}(\varphi_j) \text{ since } f(\varphi_j) \text{ proves } \theta. \end{aligned}$$

By Σ_1^0 completeness, $(\varphi_k \wedge \text{Con}^k(\varphi_k)) \vdash \text{Pr}(f(\varphi_j) \rightarrow \text{Con}^{k-1}(\varphi_j))$.

Claim 4.4. $(\varphi_k \wedge \text{Con}^k(\varphi_k)) \vdash \text{Con}^k(\varphi_{j+1})$.

We reason as follows.

$$\begin{aligned} (\varphi_k \wedge \text{Con}^k(\varphi_k)) &\vdash \neg \text{Pr}((\varphi_j \wedge \text{Con}^{k-1}(\varphi_j)) \rightarrow f(\varphi_j)) \text{ by the previous claim.} \\ &\vdash \text{Con}(\varphi_j \wedge \neg f(\varphi_j) \wedge \text{Con}^{k-1}(\varphi_j)). \\ &\vdash \text{Con}(\varphi_{j+1} \wedge \text{Con}^{k-1}(\varphi_j)) \text{ by the definition of } \varphi_{j+1}. \\ &\vdash \text{Con}(\varphi_{j+1} \wedge \text{Con}^{k-1}(\varphi_{j+1})) \text{ by the outer inductive hypothesis.} \\ &\vdash \text{Con}^k(\varphi_{j+1}) \text{ by definition of } \text{Con}^k. \end{aligned}$$

This concludes the proof of the lemma. \square

As an instance of the lemma, we get that $(\varphi_n \wedge \text{Con}^n(\varphi_n)) \vdash \text{Con}^n(\varphi_{n+1})$. We reason as follows.

$$\begin{aligned} f(\varphi_{n+1}) &\vdash \varphi_n \wedge (f(\varphi_n) \rightarrow \text{Con}^n(\varphi_n)) \text{ by the definition of } \varphi_{n+1}. \\ f(\varphi_{n+1}) &\vdash f(\varphi_n) \text{ since } f \text{ is monotonic.} \\ &\vdash \text{Con}^n(\varphi_n) \text{ by logic.} \\ &\vdash \text{Con}^n(\varphi_{n+1}) \text{ by the lemma.} \end{aligned}$$

On the other hand, $\varphi_{n+1} \wedge \text{Con}^n(\varphi_{n+1}) \vdash f(\varphi_{n+1})$ since f is everywhere bounded by Con^n . Thus, $[f(\varphi_{n+1})] = [\varphi_{n+1} \wedge \text{Con}^n(\varphi_{n+1})]$, contradicting the assumption that there is no ψ and no $k \leq n$ such that $[f(\psi)] = [\psi \wedge \text{Con}^k(\psi)] \neq [\perp]$. \square

5. TRANSFINITE ITERATES OF CON ARE INEVITABLE.

Generalizing the proof of Theorem 4.1 into the transfinite poses the following difficulty. Recall that the proof of Theorem 4.1 makes use of a sequence of sentences starting with $\varphi_0 := \top$ where

$$\varphi_{k+1} := \varphi_k \wedge (f(\varphi_k) \rightarrow \text{Con}^k(\varphi_k)).$$

It is not clear what the ω th sentence in the sequence should be. A natural idea is that for a limit ordinal λ the corresponding ‘‘limit sentence’’ should quantify over the sentences in the sequence beneath it and express, roughly,

$$\forall \gamma < \lambda (\text{True}(\varphi_\gamma) \wedge (\text{True}(f(\varphi_\gamma)) \rightarrow \text{Con}^\gamma(\varphi_\gamma))).$$

However, if the sentences in the sequence $(\varphi_\gamma)_{\gamma < \lambda}$ have unbounded syntactic complexity, then we are not guaranteed to have a truth-predicate with which we can quantify over them.

Nevertheless, we show that Theorem 4.1 generalizes into the transfinite given an additional assumption on complexity. Note that $\varphi \mapsto (\varphi \wedge \text{Con}(\varphi))$ can be factored into two functions—the identity and $\varphi \mapsto \text{Con}(\varphi)$ —the latter of which always produces a Π_1^0 sentence. For the rest of this section, we will focus on monotonic functions $\varphi \mapsto \varphi \wedge f(\varphi)$ where f is monotonic and also $f(\varphi) \in \Pi_1^0$ for all φ .

Definition 5.1. A function f is Π_1^0 if $f(\varphi) \in \Pi_1^0$ for all φ .

For the next theorem we fix an elementary presentation Γ of a recursive well-ordering. In the statement of the theorem and throughout the proof $\alpha, \beta, \gamma, \delta$, etc. are names of ordinals from the notation system Γ .

Theorem 5.2 (Restatement of Theorem 1.9). *Let f be a monotonic Π_1^0 function. Then either*

- (i) *for some $\beta \leq \alpha$ and some φ , $[\varphi \wedge f(\varphi)] = [\varphi \wedge \text{Con}^\beta(\varphi)] \neq [\perp]$ or*
- (ii) *for some φ , $(\varphi \wedge \text{Con}^\alpha(\varphi)) \not\vdash f(\varphi)$.*

Proof. Let f be a monotonic Π_1^0 function such that for every φ ,

$$(\varphi \wedge \text{Con}^\alpha(\varphi)) \vdash (\varphi \wedge f(\varphi)).$$

We assume, for the sake of contradiction, that there is no sentence ζ and no $\beta \leq \alpha$ such that $[\zeta \wedge \text{Con}^\beta(\zeta)] = [\zeta \wedge f(\zeta)] \neq [\perp]$. We then let φ be the conjunction of the following four sentences.

$$\begin{aligned} & \forall \zeta (\text{Con}(\zeta) \rightarrow \text{Con}(\zeta \wedge \neg f(\zeta))) \\ & \forall \beta \leq \alpha \forall \zeta (\text{Con}^\beta(\zeta) \rightarrow \forall \delta < \beta, \neg \text{Pr}((\zeta \wedge \text{Con}^\delta(\zeta)) \leftrightarrow (\zeta \wedge f(\zeta)))) \\ & \forall \zeta \forall \eta (\text{Pr}(\zeta \rightarrow \eta) \rightarrow \text{Pr}(f(\zeta) \rightarrow f(\eta))) \\ & \forall x (\text{Pr}(\text{True}_{\Pi_2^0}(x)) \rightarrow \text{True}_{\Pi_2^0}(x)) \end{aligned}$$

The first expresses that for every consistent φ , $f(\varphi)$ strictly implies φ . The second sentence expresses that if $\beta < \alpha$, then $f(\zeta)$ and $\zeta \wedge \text{Con}^\beta(\zeta)$ never coincide, unless $[\zeta \wedge \text{Con}^\beta(\zeta)] = [\perp]$. The third sentence expresses the monotonicity of f . The fourth sentence expresses the Π_2^0 soundness of EA. Note that each of these sentences is true, so their conjunction φ is also true. Each of the four sentences is Π_2^0 , whence so is φ .

We are interested in the following sequence $(\varphi_\beta)_{\beta \leq \Gamma}$. Note that the sentences in the sequence $(\varphi_\beta)_{\beta \leq \Gamma}$ all have complexity Π_2^0 . Note moreover that since φ_1 is true, so is φ_β for every β .

$$\varphi_1 := \varphi.$$

$$\varphi_\gamma := \varphi_1 \wedge \forall \delta < \gamma (\text{True}_{\Pi_1}(f(\varphi_\delta)) \rightarrow \text{Con}^\delta(\varphi_\delta)) \text{ for } \gamma > 1.$$

Formally, we define the sequence $(\varphi_\beta)_{\beta \leq \Gamma}$ by Gödel's fixed point lemma as in Definition 3.6.

Remark 5.3. We may assume that the ordinal notation system Γ is provably linear in EA. Thus, $\text{EA} \vdash \forall \beta \leq \alpha, \forall \gamma < \beta (\text{True}_{\Pi_2}(\varphi_\beta) \rightarrow \text{True}_{\Pi_2}(\varphi_\gamma))$.

Our goal is to show that

$$[\varphi_{\alpha+1} \wedge \text{Con}^\alpha(\varphi_{\alpha+1})] = [\varphi_{\alpha+1} \wedge f(\varphi_{\alpha+1})]$$

contradicting the assumption that f and Con^α never coincide. The main lemmas needed to prove this result are the following.

Lemma 5.4. $\text{EA} \vdash \forall \gamma \leq \alpha \text{Pr}((\varphi_\gamma \wedge \neg f(\varphi_\gamma)) \rightarrow \varphi_\alpha)$.

Lemma 5.5. $\text{EA} \vdash \forall \beta \leq \alpha \forall \gamma \leq \beta \text{Pr}((\varphi_\beta \wedge \text{Con}^\gamma(\varphi_\beta)) \rightarrow \text{Con}^\gamma(\varphi_\beta \wedge \neg f(\varphi_\beta)))$.

Lemma 5.4 is needed to derive Lemma 5.5. We now show how we use Lemma 5.5 to derive Theorem 5.2. As an instance of Lemma 5.5, letting $\alpha = \beta = \gamma$, we infer that

$$\text{EA} \vdash \text{Pr}((\varphi_\alpha \wedge \text{Con}^\alpha(\varphi_\alpha)) \rightarrow \text{Con}^\alpha(\varphi_\alpha \wedge \neg f(\varphi_\alpha))).$$

From the soundness of EA, we infer that

$$(\mp) \quad \varphi_\alpha + \text{Con}^\alpha(\varphi_\alpha) \vdash \text{Con}^\alpha(\varphi_\alpha \wedge \neg f(\varphi_\alpha)).$$

We then reason as follows.

$$\begin{aligned} & \varphi_{\alpha+1} \vdash \varphi_\alpha \wedge (f(\varphi_\alpha) \rightarrow \text{Con}^\alpha(\varphi_\alpha)) \text{ by the definition of } \varphi_{\alpha+1}. \\ & f(\varphi_{\alpha+1}) \vdash f(\varphi_\alpha) \text{ since } f \text{ is monotonic.} \\ & \varphi_{\alpha+1} + f(\varphi_{\alpha+1}) \vdash \varphi_\alpha \wedge \text{Con}^\alpha(\varphi_\alpha) \text{ by logic.} \\ & \vdash \text{Con}^\alpha(\varphi_{\alpha+1}) \text{ by } \mp. \end{aligned}$$

On the other hand, $\varphi_{\alpha+1} + \text{Con}^\alpha(\varphi_{\alpha+1}) \vdash f(\varphi_{\alpha+1})$ since f is everywhere bounded by Con^α . Since φ_1 is true, so too is $\varphi_{\alpha+1}$, whence we infer that

$$[\varphi_{\alpha+1} \wedge \text{Con}^\alpha(\varphi_{\alpha+1})] = [\varphi_{\alpha+1} \wedge f(\varphi_{\alpha+1})] \neq [\perp],$$

contradicting the claim that there is no sentence ζ and no $\beta \leq \alpha$ such that $[\zeta \wedge \text{Con}^\beta(\zeta)] = [\zeta \wedge f(\zeta)] \neq [\perp]$. \square

It remains to prove Lemma 5.4 and Lemma 5.5. We devote one subsection to each.

5.1. Proof of Lemma 5.4. In this subsection we prove Lemma 5.4. First we recall the statement of the lemma.

Lemma 5.6 (Restatement of Lemma 5.4).

$$\text{EA} \vdash \forall \gamma \leq \alpha (\text{Pr}(\varphi_\gamma \wedge \neg f(\varphi_\gamma)) \rightarrow \varphi_\alpha).$$

Proof. We reason in EA. Let $\gamma \leq \alpha$. We assume that

$$(\eta) \quad \text{True}_{\Pi_2}(\varphi_\gamma) \wedge \neg \text{True}_{\Pi_1}(f(\varphi_\gamma)).$$

We want to derive φ_α , i.e.

$$\varphi_1 \wedge \forall \sigma < \alpha (\text{True}_{\Pi_1}(f(\varphi_\sigma)) \rightarrow \text{Con}^\sigma(\varphi_\sigma)).$$

The first conjunct follows trivially from the assumption that $\text{True}_{\Pi_2}(\varphi_\gamma)$. We now prove the second conjunct of φ_α in two parts, first for all σ such that $\alpha > \sigma \geq \gamma$ and then for all $\sigma < \gamma$.

$\alpha > \sigma \geq \gamma$: From the assumption that $\text{True}_{\Pi_2}(\varphi_\gamma)$ we infer that φ_1 , whence we infer that f is monotonic. Thus, for all $\delta \geq \gamma$, $f(\varphi_\delta) \vdash f(\varphi_\gamma)$, i.e., $\text{EA} \vdash (f(\varphi_\delta) \rightarrow f(\varphi_\gamma))$. From φ_1 we also infer that EA is Π_2^0 sound, and so we infer that for all $\delta \geq \gamma$, $\text{True}_{\Pi_1}(f(\varphi_\delta)) \rightarrow \text{True}_{\Pi_1}(f(\varphi_\gamma))$. From the assumption that $\neg \text{True}_{\Pi_1}(f(\varphi_\gamma))$ we then infer that for all $\delta \geq \gamma$, $\neg \text{True}_{\Pi_1}(f(\varphi_\delta))$, whence for all $\delta \geq \gamma$, $\text{True}_{\Pi_1}(f(\varphi_\delta)) \rightarrow \text{Con}^\delta(\varphi_\delta)$.

$\sigma < \gamma$: By Remark 5.3, η implies that

$$\forall \sigma < \gamma (\text{True}_{\Pi_1}(f(\varphi_\sigma)) \rightarrow \text{Con}^\sigma(\varphi_\sigma)).$$

This completes the proof of Lemma 5.4. \square

5.2. Proof of Lemma 5.5. In this subsection we prove Lemma 5.5. We recall the statement of Lemma 5.5.

Lemma 5.7 (Restatement of Lemma 5.5).

$$\text{EA} \vdash \forall \beta \leq \alpha \forall \gamma \leq \beta \text{Pr}(\varphi_\beta + \text{Con}^\gamma(\varphi_\beta) \rightarrow \text{Con}^\gamma(\varphi_\beta \wedge \neg f(\varphi_\beta))).$$

The proof of this lemma is importantly different from the proof of Lemma 4.2. In particular, to push the induction through limit stages we need to know not only that the inductive hypothesis is true but also that it is provable in EA. We resolve this issue by using Schmerl’s technique of *reflexive transfinite induction* (see Proposition 3.4).

In the proof of the lemma, we let $\mathcal{C}(\gamma, \delta)$ abbreviate the claim that

$$\varphi_\delta + \text{Con}^\gamma(\varphi_\delta) \vdash \text{Con}^\gamma(\varphi_\delta \wedge \neg f(\varphi_\delta)).$$

Proof. We want to show that

$$\text{EA} \vdash \forall \beta \leq \alpha (\forall \gamma \leq \beta (\mathcal{C}(\gamma, \beta))).$$

By Proposition 3.4 it suffices to show that

$$\text{EA} \vdash \forall \alpha (\text{Pr}(\forall \beta \leq \alpha \forall \gamma \leq \beta \mathcal{C}(\gamma, \beta)) \rightarrow \forall \gamma \leq \alpha \mathcal{C}(\gamma, \alpha)).^2$$

Thus, we **reason in EA** and fix α . We assume that

$$(\Delta) \quad \text{Pr}(\forall \beta \leq \alpha, \forall \gamma \leq \beta, \mathcal{C}(\gamma, \beta)).$$

We let $\gamma \leq \alpha$ and we want to show that $\mathcal{C}(\gamma, \alpha)$.

Since $\varphi_\alpha \vdash \varphi$ we infer that

$$(\sharp) \quad \varphi_\alpha + \text{Con}^\gamma(\varphi_\alpha) \vdash \forall \delta < \gamma, \neg \text{Pr}((\varphi_\alpha \wedge \text{Con}^\delta(\varphi_\alpha)) \leftrightarrow (\varphi_\alpha \wedge f(\varphi_\alpha))).$$

We first note that both

$$\begin{aligned} & \varphi_\alpha \vdash \forall \delta < \gamma (\text{True}_{\Pi_1}(f(\varphi_\delta)) \rightarrow \text{Con}^\delta(\varphi_\delta)) \text{ by the definition of } \varphi_\alpha \text{ and also} \\ & \varphi_\alpha + f(\varphi_\alpha) \vdash \forall \delta < \gamma \text{Pr}(f(\varphi_\alpha) \rightarrow f(\varphi_\delta)) \text{ since } \varphi_1 \text{ proves the monotonicity of } f. \\ & \vdash \forall \delta < \gamma (f(\varphi_\alpha) \rightarrow \text{True}_{\Pi_1}(f(\varphi_\delta))) \text{ since } \varphi_1 \text{ proves the } \Pi_2^0 \text{ soundness of EA.} \\ & \vdash \forall \delta < \gamma, \text{True}_{\Pi_1}(f(\varphi_\delta)) \text{ by logic.} \end{aligned}$$

Thus, we may reason as follows.

$$\begin{aligned} & \varphi_\alpha + f(\varphi_\alpha) \vdash \forall \delta < \gamma, \text{Con}^\delta(\varphi_\delta) \\ & \vdash \forall \delta < \gamma, \text{Con}^\delta(\varphi_\delta \wedge \neg f(\varphi_\delta)) \text{ since } (\Delta) \text{ delivers } \mathcal{C}(\delta, \delta). \\ & \vdash \forall \delta < \gamma, \text{Con}^\delta(\varphi_\alpha) \text{ by Lemma 5.4.} \end{aligned}$$

Thus, by Σ_1^0 completeness,

$$\text{EA} \vdash \forall \delta < \gamma \text{Pr}((\varphi_\alpha \wedge f(\varphi_\alpha)) \rightarrow \text{Con}^\delta(\varphi_\alpha)).$$

²The reader might expect that we need to write “ $\beta < \alpha$ ” instead of “ $\beta \leq \alpha$ ” in the antecedent for this to match the statement of Proposition 3.4. However, it is clear from the proof of Proposition 3.4 that this suffices.

Combined with (\sharp), this delivers

$$\begin{aligned} \varphi_\alpha + \text{Con}^\gamma(\varphi_\alpha) &\vdash \forall \delta < \gamma \neg \text{Pr}((\varphi_\alpha \wedge \text{Con}^\delta(\varphi_\alpha)) \rightarrow f(\varphi_\alpha)). \\ &\vdash \forall \delta < \gamma, \text{Con}(\varphi_\alpha \wedge \neg f(\varphi_\alpha) \wedge \text{Con}^\delta(\varphi_\alpha)). \\ &\vdash \forall \delta < \gamma, \text{Con}(\varphi_\alpha \wedge \neg f(\varphi_\alpha) \wedge \text{Con}^\delta(\varphi_\alpha \wedge \neg f(\varphi_\alpha))) \text{ since } (\Delta) \text{ delivers } \mathcal{C}(\delta, \alpha). \\ &\vdash \text{Con}^\gamma(\varphi_\alpha \wedge \neg f(\varphi_\alpha)). \end{aligned}$$

This completes the proof of Lemma 5.5. \square

Theorem 5.2 shows the inevitability of the consistency operator. For a sufficiently constrained monotonic function f , f must coincide with an iterate of Con on some non-trivial sentence. However, it is not clear from the proofs of Theorem 4.1 or Theorem 5.2 that f must coincide with Con on a *true* sentence.

Question 5.8. *Let f be a monotonic Π_1^0 function. Suppose that for every φ ,*

$$(\varphi \wedge \text{Con}^\alpha(\varphi)) \vdash f(\varphi).$$

*Must there be some $\beta \leq \alpha$ and some **true** φ such that*

$$[\varphi \wedge f(\varphi)] = [\varphi \wedge \text{Con}^\beta(\varphi)]?$$

6. 1-CONSISTENCY AND ITERATED CONSISTENCY

Just as the Π_1^0 fragments of natural theories can often be approximated by iterated consistency statements, the Π_2^0 fragments of natural theories can often be approximated by iterated 1-consistency statements. A theory T is *1-consistent* if $T + \text{Th}_{\Pi_1^0}(\mathbb{N})$ is consistent. The 1-consistency of $\text{EA} + \varphi$ can be expressed by the following Π_2^0 sentence, $1\text{Con}(\varphi)$:

$$\forall x (\text{True}_{\Pi_1^0}(x) \rightarrow \text{Con}(\varphi \wedge \text{True}_{\Pi_1^0}(x))).$$

In this section, we investigate the relationship between 1-consistency and iterated consistency. First, we show that 1Con majorizes every iterate of Con^α .

Proposition 6.1 (Restatement of Proposition 1.11). *For any elementary presentation α of a recursive well ordering, there is a true sentence φ such that for every ψ , if $\psi \vdash \varphi$, then $(\psi \wedge 1\text{Con}(\psi))$ implies $(\psi \wedge \text{Con}^\alpha(\psi))$. Moreover, if $[\psi \wedge \text{Con}^\alpha(\psi)] \neq [\perp]$ then $(\psi \wedge 1\text{Con}(\psi))$ strictly implies $(\psi \wedge \text{Con}^\alpha(\psi))$.*

Proof. Let α be an elementary presentation of a recursive well-ordering. Let φ be a true sentence such that $\varphi \vdash \text{TI}_{\Pi_1^0}^\alpha$, i.e., φ implies the validity of transfinite induction along α for Π_1^0 predicates. We prove that

$$(\varphi \wedge 1\text{Con}(\varphi)) \vdash \text{Con}^{\alpha+1}(\varphi).$$

Since $\varphi \wedge 1\text{Con}(\varphi) \vdash \text{TI}_{\Pi_1^0}^\alpha$, it suffices to show that:

Base case: $(\varphi \wedge 1\text{Con}(\varphi)) \vdash \text{Con}(\varphi)$

Successor case: $(\varphi \wedge 1\text{Con}(\varphi)) \vdash \forall \beta < \alpha (\text{Con}^\beta(\varphi) \rightarrow \text{Con}^{\beta+1}(\varphi))$

Limit case: $(\varphi \wedge 1\text{Con}(\varphi)) \vdash \forall \lambda \left(\text{lim}(\lambda) \rightarrow ((\forall \beta < \lambda \text{Con}^\beta(\varphi)) \rightarrow \text{Con}^\lambda(\varphi)) \right)$

The **base case** and the **limit case** are both trivial. For the **successor case** we first note that by the definition of $1\text{Con}(\varphi)$,

$$1\text{Con}(\varphi) \vdash \forall x (\text{True}_{\Pi_1^0}(x) \rightarrow \text{Con}(\varphi \wedge \text{True}_{\Pi_1^0}(x))),$$

and so by substituting $\text{Con}^\beta(\varphi)$ in for x ,

$$(\oplus) \quad 1\text{Con}(\varphi) \vdash \text{True}_{\Pi_1^0}(\text{Con}^\beta(\varphi)) \rightarrow \text{Con}(\varphi \wedge \text{True}_{\Pi_1^0}(\text{Con}^\beta(\varphi))).$$

Thus, we reason as follows.

$$\begin{aligned} 1\text{Con}(\varphi) \vdash \text{Con}^\beta(\varphi) &\rightarrow \text{Con}(\varphi \wedge \text{True}_{\Pi_1^0}(\text{Con}^\beta(\varphi))) \text{ by } (\oplus). \\ &\rightarrow \text{Con}(\varphi \wedge \text{Con}^\beta(\varphi)). \\ &\rightarrow \text{Con}^{\beta+1}(\varphi) \text{ by the definition of } \text{Con}^{\beta+1}. \end{aligned}$$

It is clear that the implication $\varphi \wedge 1\text{Con}(\varphi) \vdash \varphi \wedge \text{Con}^\alpha(\varphi)$ is strict as long as $[\varphi \wedge \text{Con}^\alpha(\varphi)] \neq [\perp]$. This completes the proof of the proposition. \square

In light of the previous proposition, one might conjecture that 1Con is the weakest monotonic function majorizing every function of the form Con^α for some recursive well-ordering α on true sentences. However, this is not so. To demonstrate this, we use a recursive linear order that has no hyperarithmetic infinite descending sequences. Harrison [9] introduced such an ordering with order-type $\omega_1^{CK} \times (1 + \mathbb{Q})$; see also Feferman and Spector [7] who consider such orderings in the context of iterated reflection principles. We use a presentation \mathcal{H} of Harrison's ordering such satisfying the conditions explicated in Definition 3.1. We note that since \mathcal{H} has no hyperarithmetic descending sequences, transfinite induction along \mathcal{H} for Π_1^0 properties is valid. Our idea is to produce a function stronger than each Con^α but weaker than 1Con by iterating Con along the Harrison linear order.

Theorem 6.2 (Restatement of Theorem 1.12). *There are infinitely many monotonic functions f such that for every recursive ordinal α , there is an elementary presentation a of α such that f majorizes Con^a on a true ideal but also 1Con majorizes f on a true ideal.*

Proof. In Definition 3.2, we used Gödel's fixed point lemma to produce iterates of Con along an elementary well-ordering. We similarly use Gödel's fixed point lemma to define sentences $\mathbf{Con}^*(\varphi, \beta)$ for $\beta \in \mathcal{H}$ as follows.

$$\text{EA} \vdash \mathbf{Con}^*(\varphi, \beta) \leftrightarrow \forall \gamma <_{\mathcal{H}} \beta, \text{Con}(\varphi \wedge \mathbf{Con}^*(\varphi, \gamma)).$$

We use the notation $\text{Con}^\beta(\varphi)$ for $\mathbf{Con}^*(\varphi, \beta)$. Recall that we are assuming that it is elementarily calculable whether an element of \mathcal{H} is zero or a successor or a limit. Thus, the following clauses are provable in EA.

- $\text{Con}^0(\varphi) \leftrightarrow \top$
- $\text{Con}^{\gamma+1}(\varphi) \leftrightarrow \text{Con}(\varphi \wedge \text{Con}^\gamma(\varphi))$
- $\text{Con}^\lambda(\varphi) \leftrightarrow \forall \gamma <_{\mathcal{H}} \lambda, \text{Con}^\gamma(\varphi)$ for λ a limit.

Claim 6.3. *For $\gamma \in \mathcal{H}$, the function $\varphi \mapsto \text{Con}^\gamma(\varphi)$ is monotonic.*

This follows immediately from Proposition 3.5. Note that in the statement of Lemma 3.4 we assume only that $<$ is an elementary *linear* ordering, not a well-ordering.

Claim 6.4. *There are infinitely many monotonic functions f such that for every recursive well-ordering α , there is an elementary presentation a of α such that f majorizes Con^a on true sentences.*

If $x <_{\mathcal{H}} y$ then $\text{Con}^y(\varphi)$ strictly implies $\text{Con}^x(\varphi)$ for every φ such that $\text{Con}^x(\varphi) \neq [\perp]$. Given the order type of \mathcal{H} , this means that for infinitely many γ , for every recursive well-ordering α , Con^γ majorizes Con^a where a represents α in \mathcal{H} .

Claim 6.5. 1Con majorizes Con^a on true sentences for each $a \in \mathcal{H}$.

Since every Π_1^0 definable subset of ω has an \mathcal{H} -least element, the sentence $\text{TI}_{\Pi_1^0}^{\mathcal{H}}$, which expresses the validity of transfinite induction along \mathcal{H} for Π_1^0 predicates, is true. But then if $\varphi \vdash \text{TI}_{\Pi_1^0}^{\mathcal{H}}$, then for any $\gamma \in \mathcal{H}$, $(\varphi \wedge 1\text{Con}(\varphi))$ strictly implies $(\varphi \wedge \text{Con}^\gamma(\varphi))$ as long as $[(\varphi \wedge \text{Con}^\gamma(\varphi))] \neq [\perp]$, as in Proposition 6.1. \square

7. AN UNBOUNDED RECURSIVELY ENUMERABLE SET THAT CONTAINS NO TRUE IDEALS

In this section we prove a limitative result. Theorem 2.5 demonstrates that if f is monotonic and that for all consistent φ , (i) $\varphi \wedge \text{Con}(\varphi)$ implies $f(\varphi)$ and (ii) $f(\varphi)$ strictly implies φ , then for cofinally many true φ , $[f(\varphi)] = [\varphi \wedge \text{Con}(\varphi)]$. It is natural to conjecture that cofinal equivalence with Con be strengthened to equivalence to Con *in the limit*, i.e., on a true ideal. One strategy to strengthen Theorem 2.5 in this way would be to show that every recursively enumerable set that contains arbitrarily strong true sentences and that is closed under provable equivalence contains a true ideal.

We now show that the aforementioned strategy fails. To this end, we define a recursively enumerable set \mathcal{A} that contains arbitrarily strong true sentences and that is closed under provable equivalence but does not contain any true ideals. We are grateful to Matthew Harrison-Trainer for simplifying the proof of the following proposition.

Proposition 7.1 (Restatement of Proposition 1.13). *There is a recursively enumerable set \mathcal{A} that contains arbitrarily strong true sentences and that is closed under EA provable equivalence but does not contain any true ideals.*

Proof. Let $\{\varphi_0, \varphi_1, \dots\}$ be an effective Gödel numbering of the language of arithmetic. We describe the construction of \mathcal{A} in stages. During a stage n we may *activate* a sentence ψ , in which case we say that ψ is *active* until it is *deactivated* at some later stage $n + k$. After describing the construction of \mathcal{A} we verify that \mathcal{A} has the desired properties.

Stage 0: Numerate φ_0 and $\neg\varphi_0$ into \mathcal{A} . Activate the sentences $(\varphi_0 \wedge \text{Con}(\varphi_0))$ and $(\neg\varphi_0 \wedge \text{Con}(\neg\varphi_0))$.

Stage $n+1$: There are finitely many active sentences. For each such sentence ψ , numerate $\theta_0 := (\psi \wedge \varphi_{n+1})$ and $\theta_1 := (\psi \wedge \neg\varphi_{n+1})$ into \mathcal{A} . Deactivate the sentence ψ and activate the sentences $(\theta_0 \wedge \text{Con}(\theta_0))$ and $(\theta_1 \wedge \text{Con}(\theta_1))$.

We dovetail the construction with a search through EA proofs. If we ever see that $\text{EA} \vdash \varphi \leftrightarrow \psi$ for some φ that we have already numerated into \mathcal{A} , then we numerate ψ into \mathcal{A} .

Now we check that \mathcal{A} has the desired properties. It is clear that \mathcal{A} is recursively enumerable and that \mathcal{A} is closed under EA provable equivalence.

Claim 7.2. \mathcal{A} contains arbitrarily strong true sentences. That is, for each true sentence φ , there is a true sentence ψ such that $\psi \vdash \varphi$ and $\psi \in \mathcal{A}$.

At any stage in the construction of \mathcal{A} , there are finitely many active sentences, ψ_0, \dots, ψ_k . An easy induction shows that exactly one of ψ_0, \dots, ψ_k is true. Indeed, exactly one of φ_0 or $\neg\varphi_0$ is true, and hence so is exactly one of $\varphi_0 \wedge \text{Con}(\varphi_0)$ and $\neg\varphi_0 \wedge \text{Con}(\neg\varphi_0)$. And if θ is true, then so is exactly one of $\zeta_0 := \theta \wedge \varphi_k$ and $\zeta_1 := \theta \wedge \neg\varphi_k$, and hence so too is exactly one of $\zeta_0 \wedge \text{Con}(\zeta_0)$ and $\zeta_1 \wedge \text{Con}(\zeta_1)$.

Let φ_k be a true sentence. At stage k in the construction of \mathcal{A} there are only finitely many active sentences ψ_0, \dots, ψ_n . We have already seen that exactly one of ψ_i is true. But then $\varphi_k \wedge \psi_i$ is true, $(\varphi_k \wedge \psi_i \vdash \varphi_k)$, and $(\varphi_k \wedge \psi_i)$ is numerated into \mathcal{A} .

Claim 7.3. *\mathcal{A} contains no true ideals.*

An easy induction shows that if ψ_0 and ψ_1 are both active at the same stage, then for any θ , if θ implies both ψ_0 and ψ_1 then $\theta \in [\perp]$.

Let φ be a true sentence in \mathcal{A} . By the previous remark, the only sentences in \mathcal{A} that strictly imply φ are (i) EA refutable sentences and (ii) sentences that imply $\varphi \wedge \text{Con}(\varphi)$. Since the Lindenbaum algebra of EA is dense, this means there is some ψ such that $(\varphi \wedge \text{Con}(\varphi))$ strictly implies ψ strictly implies φ but $\psi \notin \mathcal{A}$. \square

The following questions remain.

Question 7.4. *Is the relation of cofinal agreement on true sentences an equivalence relation on recursive monotonic operators?*

Question 7.5. *Let f be recursive and monotonic. Suppose that for all consistent φ ,*

(i) $\varphi \wedge \text{Con}(\varphi)$ implies $f(\varphi)$ and

(ii) $f(\varphi)$ implies φ .

Must f be equivalent to the identity or to Con on a true ideal?

REFERENCES

- [1] Uri Andrews, Mingzhong Cai, David Diamondstone, Steffen Lempp, and Joseph S Miller. On the structure of the degrees of relative provability. *Israel Journal of Mathematics*, 207(1):449–478, 2015.
- [2] Lev D Beklemishev. Provability logics for natural turing progressions of arithmetical theories. *Studia Logica*, 50(1):107–128, 1991.
- [3] Lev D Beklemishev. Iterated local reflection versus iterated consistency. *Annals of Pure and Applied Logic*, 75(1-2):25–48, 1995.
- [4] Lev D Beklemishev. Proof-theoretic analysis by iterated reflection. *Archive for Mathematical Logic*, 42(6):515–552, 2003.
- [5] Lev D Beklemishev. Provability algebras and proof-theoretic ordinals, I. *Annals of Pure and Applied Logic*, 128(1-3):103–123, 2004.
- [6] Lev D Beklemishev. Reflection principles and provability algebras in formal arithmetic. *Russian Mathematical Surveys*, 60(2):197, 2005.
- [7] Solomon Feferman and Clifford Spector. Incompleteness along paths in progressions of theories. *The Journal of Symbolic Logic*, 27(4):383–390, 1962.
- [8] Sy-David Friedman, Michael Rathjen, and Andreas Weiermann. Slow consistency. *Annals of Pure and Applied Logic*, 164(3):382–393, 2013.
- [9] Joseph Harrison. Recursive pseudo-well-orderings. *Transactions of the American Mathematical Society*, 131(2):526–543, 1968.
- [10] Joost J Joosten. Turing–Taylor expansions for arithmetic theories. *Studia Logica*, 104(6):1225–1243, 2016.
- [11] Marian Boykan Pour-El and Saul Kripke. Deduction-preserving “recursive isomorphisms” between theories. *Fundamenta Mathematicae*, 61:141–163, 1967.

- [12] Ulf R. Schmerl. A fine structure generated by reflection formulas over primitive recursive arithmetic. *Studies in Logic and the Foundations of Mathematics*, 97:335–350, 1979.
- [13] V Yu Shavrukov and Albert Visser. Uniform density in Lindenbaum algebras. *Notre Dame Journal of Formal Logic*, 55(4):569–582, 2014.
- [14] Theodore A Slaman and John R Steel. Definable functions on degrees. In *Cabal Seminar 81–85*, pages 37–55. Springer, 1988.

GROUP IN LOGIC AND THE METHODOLOGY OF SCIENCE, UNIVERSITY OF CALIFORNIA, BERKELEY
Email address: `walsh@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY
Email address: `antonio@math.berkeley.edu`