Lawrence Berkeley National Laboratory

Recent Work

Title

STUDY OF THE K+p INTERACTION FROM 0.85 TO 1.5 BeV/c

Permalink

https://escholarship.org/uc/item/1fx015d1

Authors

Brown, John L. Bland, Roger W. Bowler, Michael G. <u>et al.</u>

Publication Date

1964-07-22

University of California Ernest O. Lawrence Radiation Laboratory

UCRL 11446

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

STUDY OF THE $K^{+}p$ interaction from 0.85 to 1.5 BeV/c

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California. International Conference on High Energy Physics

1

UCRL-11446

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory Berkeley, California

Contract No. W-7405-eng-48

STUDY OF THE K^{*}p INTERACTION FROM 0.85 TO 1.5 BEV/C

John L. Brown, Roger W. Bland, Michael G. Bowler, Gerson Goldhaber, Sulamith Goldhaber, Allan A. Hirata, John A. Kadyk, Victor H. Seeger, and George H. Trilling

July 22, 1964

(To be presented at the 1964 International Conference on High Energy Physics, Dubna, U.S.S.R. August 5-15, 1964)

UCRL-11446

STUDY OF THE K⁺_p INTERACTION FROM 0.85 TO 1.5 BEV/C^{*}

John L. Brown, Roger W. Bland, Michael G. Bowler, Gerson Goldhaber, Sulamith Goldhaber, Allan A. Hirata, John A. Kadyk, Victor H. Seeger, and George H. Trilling

(Presented by Sulamith Goldhaber)

Lawrence Radiation Laboratory University of California Berkeley, California

July 22, 1964

In an experiment currently in progress at the Bevatron, $K^{+}p$ and $K^{+}d$ reactions in the momentum range 0.86 - 1.58 BeV/c are being studied in a 25-inch hydrogen/deuterium bubble chamber. Preliminary cross sections for the various $K^{+}p$ reactions have been obtained and are presented below. The most notable feature is the large increase in the inelastic cross section in the region of the thresholds for production of the N^{*}(1238) and K^{*}(880) resonances.

For the purposes of this preliminary investigation, approximately 1500 pictures were double-scanned at each of the following five momenta: 0.86, 0.96, 1.20, 1.36 and 1.58 BeV/c. All events were processed through the PACKAGE space reconstruction and kinematic fitting program. Each event was re-examined at the scan table by a physicist to verify the conclusions of the fitting program, and to resolve identification ambiguities, where possible, on the basis of ionization. Residual ambiguous events comprised less than 3% of the total number, and were assigned in the same proportions as the unambiguous events.

* Work sponsored by the U. S. Atomic Energy Commission.

UCRL-11446

At all but the two highest momenta the pion contamination of the K^{T} beam was negligible. To correct for contamination at 1.36 and 1.58 BeV/c, film was also taken with incident positive pions of these momenta. A comparison of the pion and kaon film gave estimates of 7% and 18% respectively for the contamination at the two above momenta. Appropriate corrections have been applied to individual partial cross sections.

The results for the cross sections of the various channels, corrected for pion contamination, are shown in Table I. For all momenta except 1.58 BeV/c the K⁺ flux was determined from a count of the observed τ decays in flight combined with the τ branching ratio of Roe et al.¹ At the highest momentum the τ statistics were insufficient to make this determination meaningful. We also list the results obtained with the total cross sections normalized to the measurements of Cook.² The results of this experiment together with data from other experiments in the same energy range are shown in Figs. 1 and 2. A preliminary study of the three-particle final states indicates that the initial rise in the inelastic channels is accompanied by substantial formation of the N^{*}(1238) resonance. At momenta of 1.20 BeV/c and higher there is in addition substantial production of the K^{*}(880) resonance. The cross section for double pion production remains less than 1/2 mb up to 1600 MeV/c.

REFERENCES

- Byron P. Roe, Daniel Sinclair, John L. Brown, Donald A. Glaser, John A. Kadyk, and George H. Trilling, Phys. Rev. Lett. 7, 346 (1961).
- V. Cook, D. Keefe, L. T. Kerth, D. C. Murphy, W. A. Wentzel, and T. F. Zipf, Phys. Rev. Lett. <u>7</u>, 182 (1961), see also
 Victor Cook (Ph.D. Thesis) "K⁺ Nucleon Interactions in the Momentum Range of 800-2900 MeV/c". Lawrence Radiation Laboratory Report UCRL-10130 (March 1962).
- 3. Brandt Kehoe, Phys. Rev. Lett. <u>11</u>, 93 (1963).

1.

- 4. E. Boldt, J. Duboc, N. H. Duong, P. Eberhard, R. George, V. P. Henri,
 F. Levy, J. Poyen, M. Pripstein, J. Crussard, and A. Tran, Phys. Rev. <u>133 B</u>,
 220 (1964).
- 5. Theodore F. Stubbs, Hugh Bradner, William Chinowsky, Gerson Goldhaber, Sulamith Goldhaber, William Slater, Donald M. Stork, and Harold K. Tikho, Phys. Rev. Lett. 7, 188 (1961).
- 6. Sulamith Goldhaber: "Meson Exchange in the K⁺p Interaction", <u>Proceedings</u> of the Athens Topical Conference on Newly Discovered Resonant Particles, Ohio University, Athens, Ohio (1963).

FIGURE CAPTIONS

- Fig. 1. K⁺p elastic and total cross sections as a function of momentum. In addition to the results of this experiment, data from references 2, 5, and 6 have been included.
 - Fig. 2. K⁺p inelastic partial cross sections as a function of momentum. Data from reference 3 through 6 have been included.


UCRL-11446

			-		: 	·.		
Momentum: 0.86			0.96		1.20		1.36	1.58
Reaction	(1)	(2)	(1)	(2)	(1)	(2)	(1) (2)	(1) (2)
$K^{\dagger}p \rightarrow K^{\dagger}p$	12.0 ± 1.6	12.6 ± 0.7	12.9 ± 2.3	11.5 ± 0.8	12.7 ± 2.1	10.9 ± 0.7	11.5 ± 2.3 8.7 ± 0.6	9.6 ± 1.0
$K^{\dagger}p \rightarrow K^{\dagger}p\pi^{0}$	0.4 ± 0.1	0.4 ± 0.1	0.9 ± 0.3	0.8 ± 0.2	2.4 ± 0.5	2.1 ± 0.3	2.8 ± 0.7 2.1 ± 0.3	0.9 ± 0.3
$K^{\dagger}p \rightarrow K^{0}p\pi^{\dagger}$	0.9 ± 0.2	0.9 ± 0.2	3.1 ± 0.7	2.8 ± 0.4	5.2 ± 0.9	4.4 ± 0.4	7.7 ± 1.6 5.8 ± 0.5	5.6 ± 0.8 ·
$K^+ p \rightarrow K^+ n \pi^+$	0.1 ± 0.1	0.1 ± 0.1	0.2 ± 0.1	0.2 ± 0.1	0.8 ± 0.2	0.7 ± 0.2	1.2 ± 0.4 0.9 ± 0.2	1.3 ± 0.4
Total cross section	13.4 ± 1.8	14.1	17.2 ± 2.9	15.3	21.0 ± 3.4	18.1	23.3 ± 4.6 17.7	17.8
Number of K interactions in sample	431		324		496		474	206
	·							

Table I. Partial K p cross sections (mb) as a function of momentum


(1) Results determined solely from this experiment

(2) Results normalized to the data of Cook et al. Uncertainties in the total cross sections are not included in the quoted errors.



Figure

MUB-3585



ē. 1

Figure

N

MUB-3586

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor. while the second and a first of the second second 不可行了。你也这些新考虑你的自己的意思,我还能够更没有考虑的问题。我有意识的"感问题",她的_是是这些 and the second ا د ه ش in provide the second sec · 医心管 化乙酸盐 医牙外腔 制度的 网络医皮肤酶 化氨酸盐酸盐 化氟化物 a sector a s $\mathbf{e}_{\mathbf{u}}^{\mathbf{u}} = \mathbf{e}_{\mathbf{u}}^{\mathbf{u}} + \mathbf{e}_{\mathbf$ A second s 化输资 机运行性加强的 机运行度 选验 where I was a second of the second of all a start a start a start a start a start a í · · .8 the stand of the second second ار الما بنا ال a te esta l'esta de la sec . ب همی فر د و در 1.00 1. The set of 医弹性间隙 化同应剂 医马克氏结合 医马克马氏 机合金属 医分离器