UC San Diego

UC San Diego Previously Published Works

Title

Specific pain complaints in Iraq and Afghanistan veterans screening positive for post-traumatic stress disorder.

Permalink

<https://escholarship.org/uc/item/1gs8v3qk>

Journal

Psychosomatics, 55(2)

ISSN

0033-3182

Authors

Moeller-Bertram, Tobias
Afari, Niloofar
Mostoufi, Sheeva
et al.

Publication Date

2014-03-01

DOI

10.1016/j.psych.2013.01.011

Peer reviewed

Original Research Reports

Specific Pain Complaints in Iraq and Afghanistan Veterans Screening Positive for Post-Traumatic Stress Disorder

Tobias Moeller-Bertram, M.D., Ph.D., M.A.S., Niloofar Afari, Ph.D., Sheeva Mostoufi, M.S.,
David S. Fink, M.P.H., Lisa Johnson Wright, Ph.D., Dewleen G. Baker, M.D.

Background: Post-traumatic stress disorder (PTSD) and pain are highly comorbid. **Objective:** The purpose of this study was to examine the association of PTSD with specific pain complaints in veterans of Operations Enduring and Iraqi Freedom (OEF/OIF). **Method:** A total of 381 primarily male (88.5%) veterans with a mean age of 30 years completed a battery of self-report questionnaires. A positive PTSD screen was defined as a score of ≥ 40 on the Davidson Trauma Scale. Logistic regression was used to examine the association of positive PTSD screen with specific pain complaints. **Results:** There were no significant demographic or physical and mental health differences between veterans who

screened positive for PTSD only and those with PTSD and at least one pain complaint, although differences on rates of combat injury and depression approached significance. Veterans who screened positive for PTSD were 2 to 3 times more likely to report abdominal pain, muscle aches or cramps, and joint aches, even after controlling for age, gender, combat injury, and depression. **Conclusions:** Similar to findings in other populations, there is a relationship between PTSD and pain complaints in OEF/OIF veterans. Future research should examine the mechanisms that link PTSD with specific pain complaints, especially abdominal pain.

(Psychosomatics 2013; xx:xxx)

Post-traumatic stress disorder (PTSD) is a debilitating illness characterized by symptoms of re-experiencing, avoidance, emotional numbing, and hyperarousal resulting from an emotionally traumatic event with actual or perceived threat.¹ Persons with military combat exposure are at high risk for developing PTSD.^{2,3} Lifetime rates for PTSD range from 18% to 30% in Vietnam veterans.⁴ In addition, as many as 80% of Vietnam veterans with PTSD have been reported to experience chronic pain.⁵ Other studies in Vietnam veterans and those from other conflicts predating Operations Enduring and Iraqi Freedom (OEF/OIF) found increased rates for pain complaints, including overall bodily pain, migraines and headaches,⁶ chronic low back pain, and general chronic pain secondary to systemic disease.⁷

The military operations in Iraq and Afghanistan have been unlike previous conflicts, especially because of the unpredictable nature of combat and guerrilla warfare, mul-

tiples and long deployments, the chronic threat of improvised explosive devices (IEDs), and advances in battlefield medicine that has led to increased survival from injuries with higher degrees of disability. Therefore, a growing body of research has begun to examine the impact of OEF/OIF deployment on the mental and physical health of military personnel and veterans.^{8–10} Recent reports sug-

Received September 21, 2012; revised January 20, 2013; accepted January 21, 2013. From the VA Center of Excellence for Stress and Mental Health and VA San Diego Healthcare System, San Diego, CA (TM-B, NA, SM, DSF, DGB); Department of Anesthesiology, University of California, San Diego, San Diego, CA (TM-B); Department of Psychiatry, University of California, San Diego, San Diego, CA (NA, DGB); SDSU/UCSD Joint Doctoral Program in Clinical Psychology, San Diego, CA (SM); VA Northern California Healthcare System, Oakland, CA (LJW). Send correspondence and reprint requests to Niloofar Afari, Ph.D., Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093; e-mail: nafari@ucsd.edu

Published by Elsevier Inc. on behalf of The Academy of Psychosomatic Medicine.

AQ: 5 Pain and PTSD in OEF/OIF Veterans

gest high rates of mental health disorders, including PTSD and depression, among active duty military personnel and OEF/OIF veterans.^{8,11,12} Further, pain is among the most prevalent complaints of those returning from OEF/OIF,¹³ with reports of pain in up to 80% of OEF/OIF veterans and one-third reporting moderate to severe pain.¹⁴ Another study found that 57% of studied veterans reported moderate to severe pain.¹⁵ In addition, OEF/OIF veterans report high rates of specific pain complaints, including chronic widespread pain,¹³ back and limb pain,¹⁵ and headaches.¹⁶

Previous studies also have reported a high co-occurrence of general pain and PTSD¹⁷ as well as poorer physical health functioning and PTSD¹⁸ in OEF/OIF veterans. However, the rates of specific pain complaints in OEF/OIF veterans with PTSD or the role other factors, such as depression, may play in this relationship have not been fully explored. Further, no studies have examined demographic, physical and mental health differences between individuals with PTSD and those with both pain complaints and PTSD in OEF/OIF veterans.

The purpose of this study was threefold. First, we examined the demographic, physical health, and mental health differences between OEF/OIF veterans who screened positive for PTSD only and those who were positive for both PTSD and at least one pain complaint. Second, we assessed the association of screening positive for PTSD with several specific pain complaints in OEF/OIF veterans. Third, we explored the role of age, gender, combat injury, and depression on the relationship between PTSD and pain complaints.

METHODS

Sample and Procedures

This study was part of a larger cross-sectional evaluation of OEF/OIF veterans registering for healthcare at the VA San Diego Healthcare System (VASDHS) from March to October, 2006. During this period, 459 veterans and reservists agreed to complete a battery of questionnaires at their initial registration visit with Member Services. Of these, 19 (4%) did not take part in OEF or OIF. Of the remaining 440 OEF/OIF veterans, 381 (87%) had completed data and were included in these analyses. The study was approved by the University of California, San Diego Institutional Review Board and by the VA San Diego Research and Development Committee. All authors had access to the data.

Measures

Veterans completed a packet of standardized questionnaires and rating scales to assess demographic, military, and in-theatre factors, mental health symptoms, and health-related diagnoses and symptoms including pain complaints.

Demographic and Military Variables The demographic and military questions included information on age, gender, race/ethnicity, and branch of service.

In-Theatre Variables Veterans completed the Combat Exposure Scale, a 7-item measure of wartime stressors experienced by combatants.¹⁹ A series of dichotomous questions (i.e., yes or no answer choices) asked about (1) experiencing an emotional trauma, including combat or noncombat experiences, (2) combat-related injury, and (3) a history of loss of consciousness due to head injury.

Mental Health Symptoms A battery of commonly used and well-validated questionnaires assessed for mental health symptoms. The Davidson Trauma Scale (DTS) was used to determine the degree of PTSD symptoms.²⁰ The DTS is an often-used measure of the frequency and severity of PTSD symptoms based on the Diagnostic and Statistical Manual of Mental Disorders, 4th edition, criteria.¹ This 17-item measure has good internal consistency and test-retest reliability.²⁰ Screening positive for PTSD was defined as positive endorsement of trauma exposure and a DTS score of 40 or higher. A two-item depression screener asked about feeling down, depressed or hopeless, and little interest or pleasure in doing things for the last month. This depression screener has been used routinely in other studies published on VA populations as a general screen for depression.²¹ The measure is psychometrically sound, useful for detecting depression in primary care settings, and has similar test characteristics to other case-finding instruments.²² A positive depression screen was defined as endorsement of both depressed mood and anhedonia.

The Alcohol Use Disorders Identification Test (AUDIT)²³ and Drug Abuse Screening Test (DAST)²⁴ were included to screen for drug and alcohol problems. Both the AUDIT and the DAST demonstrate strong sensitivity and specificity in identifying problematic alcohol and drug abuse, respectively.^{24,25} Substance abuse was defined as a score of 5 or more on the AUDIT or a score of 2 or more on the DAST.

Pain Complaints Veterans also completed a 52-item checklist of medical symptoms that included items on

TABLE 1. Characteristics of All Veterans and by Post-traumatic Stress Disorder and Pain Status

Characteristic	Overall <i>n</i> = 381	PTSD Only <i>n</i> = 38	PTSD and Pain ^a <i>n</i> = 122
Male gender, %	88.5	86.8	90.2
Age, mean years (SD)	30.3 (8.2)	28.5 (8.5)	30.0 (7.6)
Race, %			
White	44.6	47.4	45.1
Black	16.1	10.5	18.0
Hispanic	21.8	36.8	20.5
Asian	13.4	5.3	12.3
Other	4.2	0.0	4.1
Service branch, %			
Army	22.3	34.2	26.2
Air Force	1.8	5.3	1.6
Marines	33.9	36.8	37.7
Navy	36.8	15.8	27.9
National Guard	5.3	7.9	6.0
Injured during combat, %	20.7	21.1	36.6
CES, mean score (SD)	13.8 (11.1)	20 (9.0)	20 (10.5)
Positive depression, %	44.6	65.8	77.6
Positive substance abuse, %	34.1	44.7	43.3
Loss of consciousness, %	6.8	5.3	13.4

PTSD = post-traumatic stress disorder; CES = Combat Exposure Scale.

^a PTSD and Pain participants had a positive PTSD screen and at least one pain complaint.

RESULTS

Initial analyses indicated that other than branch of service, the 381 OEF/OIF veterans who completed the assessment packet did not differ significantly from the 59 OEF/OIF veterans who were excluded from the analyses because of missing data. A larger proportion of incomplete packets than complete packets were from the Marines (39% vs. 34%, $P < 0.001$). Nonetheless, veterans from the Navy and Marines constituted the largest proportion for both completers and non-completers.

Of the 381 veterans with complete data, 160 (42%) screened positive for PTSD and 221 (58%) did not screen positive for PTSD. Table 1 presents the characteristics of the overall sample as well as for those who screened positive for PTSD only and those who screened positive for both PTSD with at least one pain complaint. A majority of the sample was male (88.5%) with a mean age of 30 years (SD = 8.2). The sample was ethnically diverse and represented primarily the Navy (36.8%), Marines (33.9%), and Army (22.3%). Nearly 21% of the sample reported injury during combat and respondents had a mean score of 13.8 (SD = 11.1) on the Combat Exposure Scale corresponding with light to moderate combat exposure. About 45% and 34% of the sample screened positive for depression and substance abuse, respectively; 7% reported a loss of consciousness. Compared with veterans who were positive for PTSD only ($n = 38$), those with PTSD and at least one pain complaint ($n = 122$) had higher rates of combat injury ($\chi^2 [1, n = 381] = 3.23, P = 0.07$) and depression ($\chi^2 [1, n = 381] = 2.20, P = 0.14$), but these differences approached significance. There were no other differences between groups.

Table 2 presents the unadjusted and adjusted ORs and 95% CIs for the association between positive screen for PTSD and the specific pain complaints. Veterans who were positive for PTSD were significantly more likely to report abdominal pain (OR = 4.5, 95% CI = 2.3–9.1),

stomach or abdominal pain, muscle aches or cramps, joint pain or aches, and back pain or spasms. Veterans indicated if the symptom was currently a problem. This checklist was similar to those used in previous studies of health symptoms in veterans.²⁶

Statistical Analyses

Descriptive analyses, including t -tests for continuous variables and χ^2 for dichotomous variables, were used to examine differences between veterans with complete data and those who were excluded from the analyses because of missing data, and to characterize the sample and examine the demographic and other differences between veterans who screened positive for PTSD only and those who were positive for both PTSD and at least one pain complaint. Unadjusted logistic regression analyses were conducted to examine the association between PTSD and specific pain complaints and to determine odds ratios (OR) and 95% confidence intervals (CI). These were followed by logistic regression analyses that controlled for age, gender, depression, and combat injury. Significant P value was set at 0.05. All analyses were conducted using SAS v. 9.2 (SAS Institute Inc., Cary, NC).

TABLE 2. Unadjusted and Adjusted OR and 95% CI for the Association between PTSD and Pain Complaints

Pain Complaints	Unadjusted	Adjusted ^a
	OR (95% CI)	OR (95% CI)
Stomach or abdominal pain	4.5 (2.3–9.1)	2.8 (1.3–6.4)
Muscle aches/cramps	3.8 (2.3–6.4)	2.7 (1.5–4.9)
Joint aching or pain	2.3 (1.5–3.5)	2.1
Back pain or spasms	2.1 (1.4–3.2)	1.5 (0.9–2.5)

^a Adjusted for age, gender, combat injury, and depression.

Pain and PTSD in OEF/OIF Veterans

muscle aches or cramps (OR = 3.8, 95% CI = 2.3–6.4), joint aches (OR = 2.3, 95% CI = 1.5–3.5), and back pain or spasms (OR = 2.1, 95% CI = 1.4–3.2) than veterans without PTSD. Adjusting for age, gender, combat injury, and depression reduced these effects, however, the association with abdominal pain (OR = 2.8, 95% CI = 1.3–6.3), muscle aches or cramps (OR = 2.7, 95% CI = 1.5–4.9), and joint aches (OR = 2.1, 95% CI = 1.2–3.4) remained significant; the association with back pain or spasms approached significance (OR = 1.5, 95% CI = 0.9–2.5). These findings suggest that age, gender, combat injury, and depression only partially explain the relationship between a positive PTSD screen and these specific pain complaints.

DISCUSSION

This study focused on the relationship between a positive screen for PTSD and specific pain complaints in a cohort of OEF/OIF veterans. We found substantial rates of positive PTSD, depression, and substance abuse screens in the sample. There were few demographic, physical health, and mental health differences between OEF/OIF veterans who screened positive for PTSD only and those who screened positive for both PTSD and at least one pain complaint; the differences on rates of combat injury and positive depression screens approached significance. We also found that a positive PTSD screen was associated with significantly higher odds of abdominal pain, musculoskeletal pain, joint pain, and back pain. Adjusting for age, gender, combat injury, and depression diminished the strength of these associations but aside from the link with back pain, all associations remained significant.

Previous research has found that individuals with PTSD and chronic pain report higher ratings of psychiatric distress, greater disability, and more intense pain compared with those with PTSD or chronic pain only.^{27–29} These previous findings are in contrast to ours and suggest that the more substantial morbidity associated with comorbid PTSD and pain may happen specifically when pain is long-lasting or chronic. We did find that veterans with PTSD and at least one pain complaint were more likely to have experienced combat and to screen positive for depression, but these findings were not statistically significant, likely due to low power. Future research should explore the role of combat as well as depressive symptoms in the link between PTSD and pain, and distinguish between combat-related trauma and noncombat trauma (i.e., military sexual trauma) in theatre. Additional research can

also address the morbidity associated with PTSD and pain as pain complaints develop over time and become more chronic in OEF/OIF veterans.

A large body of research has documented a variety of pain complaints in veterans with PTSD.^{2,5,7,30} In contrast to previous studies in Vietnam veterans but similar to one study in Gulf War veterans,³⁰ we saw an increased association of PTSD with abdominal pain. One possible explanation is the increased experience with blast exposure in OEF/OIF veterans. Blast exposure related to IEDs is specifically related to body cavity pain, including abdominal pain, as well as PTSD and traumatic brain injury.³¹ Alternately, studies, primarily with women and noncombat trauma, have linked PTSD with irritable bowel syndrome, a condition with abdominal pain as a key feature.^{32–34} The link between PTSD and gastrointestinal conditions such as irritable bowel syndrome should be fully explored in OEF/OIF veterans. In addition, future research can better examine the role of blast exposure in the complex relationship between PTSD, traumatic brain injury, and various pain complaints, as well as a comprehensive and interdisciplinary approach to assessment and treatment of these conditions.

We also found that age, gender, depression, and combat injury did not completely account for the relationship between PTSD and pain complaints other than back pain. This is of particular interest since previous studies have found comorbid depression to be either a mediator or moderator of increased pain in PTSD patients.^{35,36} Others also have found that although depression may attenuate the relationship between PTSD and physical health functioning, PTSD remains a significant factor in poorer physical health functioning and greater bodily pain.³⁷ Together with our findings, these discrepant results suggest that further evaluation of the overlapping and unique roles of PTSD and depression in pain is warranted.

Current models aiming to explain the comorbidity of PTSD and chronic pain also acknowledge the intertwined nature of PTSD and pain. The shared vulnerability model suggested by Asmundson et al.³⁸ predicts that individual factors predispose people to develop both PTSD and chronic pain. The mutual maintenance model for PTSD and pain proposes that mechanisms such as depression, anxiety sensitivity, or pain perception may maintain or exacerbate symptoms of both PTSD and chronic pain.³⁹ It is important that future research examine both vulnerability and maintenance factors that may explain the comorbid relationship between PTSD and pain complaints and con-

ditions. Our findings on the potentially different associations of PTSD with various pain complaints suggest that the underlying mechanisms may be unique to different pain complaints and conditions. Understanding the similarities and differences in the mechanisms that underlie the relationship between PTSD and specific pain complaints and conditions will facilitate more tailored prevention and treatment strategies.

Our findings also support the need for a multidisciplinary approach to the assessment and treatment of both mental and physical health symptoms in OEF/OIF veterans. The comprehensive screening and follow-up assessment of both trauma-related symptoms and pain and other physical symptoms in veterans with and without physical injury is likely to facilitate early detection and timely and broad treatment. Although evidence-based treatments are available for both PTSD and pain conditions, only a handful of treatment protocols have been developed that address PTSD and pain in combination.⁴⁰ Clearly, more research is needed to determine optimal treatment of OEF/OIF veterans with these co-occurring symptoms.

This study has several limitations. First, our findings are based on self-reported data and could have been affected by recall bias. Relatedly, the PTSD, depression, and substance abuse measures were brief screening instruments not designed for diagnosis of these conditions but for quick identification of individuals who would benefit from additional assessment. In addition, the measure of pain complaints focused only on self-reported presence of pain and did not provide any information on severity or impact on function. Thus, findings from this study should be replicated using a comprehensive and clinically-based assessment of PTSD, other mental health symptoms, and pain complaints. Second, we had no information on veterans' history of pain symptoms prior to deployment to OEF/OIF, nor the course of their pain complaints. Therefore, we must reiterate that our findings are based on cross-sectional data and do not imply that PTSD causes pain or that preexisting history of pain symptoms predisposes one to the development of PTSD. Third, the data presented here were collected in 2006 and may not fully represent the characteristics of veterans who have served in OEF/OIF since then. Nonetheless, our findings of the link between PTSD and pain complaints are consistent with previous research and further highlight the need to examine the interplay of PTSD and pain in

OEF/OIF veterans. Finally, the sample of OEF/OIF participants was not randomly selected. This point is underscored by differences in the branch of service reported by veterans who completed the assessment packet and those who did not, a variable that may be confounded by clinically relevant exposures. In addition, we obtained data from veterans who were attending a VA Medical Center to register for healthcare. It is likely that once discharged from the military, those who are experiencing mental health symptoms are more likely to seek treatment and register for VA healthcare. Indeed, the substantially high rates of positive PTSD, depression, and substance abuse screens we observed suggest that newly registering veterans may represent a self-selected treatment-seeking sample. Thus, our findings may only be relevant to veterans seeking care at a VA Medical Center.

In conclusion, we found that OEF/OIF veterans who screened positive for PTSD were 2 to 3 times more likely to report a number of pain complaints than those without PTSD, even after controlling for age, gender, combat injury, and depression. Future research can delineate the role of combat vs. noncombat trauma in theater in relation to pain complaints. In addition, other research can focus on the role of blast exposure and depression as well as further examining vulnerability and maintenance factors associated with comorbid PTSD and specific pain complaints and conditions. Veterans could benefit from an interdisciplinary assessment and treatment approach that addresses both mental and physical health symptoms. This line of investigation will be valuable to healthcare providers caring for OEF/OIF veterans to help improve their care.

Disclosure: The authors disclosed no proprietary or commercial interest in any product mentioned or concept discussed in this article.

This material is based upon work supported by the Department of Veterans Affairs Center of Excellence for Stress and Mental Health. This material also is the result of work supported with resources of the VA San Diego Healthcare System. The views expressed in this paper are those of the authors and do not reflect the official policy or position of the Department of Veteran's Affairs or any of the institutions with which the authors are affiliated. The authors thank Michael Kilmer, M.S.W. and Naju J. Madra, M.A. for their support and diligent work on this project.

Pain and PTSD in OEF/OIF Veterans

References

1. APA: Diagnostic and Statistical Manual of Mental Disorders, 4th ed. Washington DC 1994
2. O'Toole BI, Catts SV: Trauma, PTSD, and physical health: an epidemiological study of Australian Vietnam veterans. *J Psychosom Res* 2008; 64:33–40
3. Reeves RR, Parker JD, Konkle-Parker DJ: War-related mental health problems of today's veterans: new clinical awareness. *J Psychosoc Nurs Ment Health Serv* 2005; 43:18–28
4. Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall R: The psychological risks of Vietnam for US veterans: a revisit with new data and methods. *Science* 2006; 313:979–982
5. Beckham JC, Crawford AL, Feldman ME, Kirby AC, Hertzberg MA, Davidson JRT, et al: Chronic posttraumatic stress disorder and chronic pain in Vietnam combat veterans. *J Psychosom Res* 1997; 43:379–389
6. Asmundson GJG, Wright KD, Stein, MB: Stein, MB. Pain and PTSD symptoms in female veterans. *Eur J Pain* 2004; 8:345–350
7. Shipherd JC, Keyes M, Jovanovic T, Ready DJ, Baltzell D, Worley V, et al: Veterans seeking treatment for posttraumatic stress disorder: what about comorbid chronic pain? *J Rehabil Res Dev* 2007; 44:153–166
8. Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL: Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med* 2004; 351:13–22
9. Reger MA, Gahm GA, Swanson RD, Duma SJ: Association between number of deployments to Iraq and mental health screening outcomes in US Army soldiers. *J Clin Psychiatry* 2009; 70:1266–1272
10. Thomas JL, Wilk JE, Riviere LA, McGurk D, Castro CA, Hoge CW: Prevalence of mental health problems and functional impairment among active component and National guard soldiers 3 and 12 months following combat in Iraq. *Arch Gen Psychiatry* 2010; 67:614–623
11. Seal KH, Bertenthal D, Miner CR, Sen S, Marmar C: Bringing the war back home: mental health disorders among 103,788 US veterans returning from Iraq and Afghanistan seen at department of Veterans Affairs facilities. *Arch Intern Med* 2007; 167:476–482
12. Baker DG, Heppner P, Afari N, Nunnink S, Kilmer M, Simmons A, et al: Trauma exposure, branch of service, and physical injury in relation to mental health among U.S. veterans returning from Iraq and Afghanistan. *Mil Med* 2009; 174:773–778
13. Helmer DA, Chandler HK, Quigley KS, Blatt M, Teichman R, Lange G: Chronic widespread pain, mental health, and physical role function in OEF/OIF veterans. *Pain Med* 2009; 10:1174–1182
14. Girona RJ, Clark ME, Massengale JP, Walker RL: Pain among veterans of operations enduring freedom and Iraqi freedom. *Pain Med* 2006; 7:339–343
15. Haskell SG, Brandt CA, Krebs EE, Skanderson M, Kerns RD, Goulet JL: Pain among veterans of operations enduring freedom and Iraqi freedom: do women and men differ? *Pain Med* 2009; 10:1167–1173
16. Afari N, Harder LH, Madra NJ, Heppner PS, Moeller-Bertram T, King C, et al: PTSD, combat injury, and headache in veterans returning from Iraq/Afghanistan. *Headache J Head Face Pain* 2009; 49:1267–1276
17. Stecker T, Fortney J, Owen R, McGovern MP, Williams S: Co-occurring medical, psychiatric, and alcohol-related disorders among veterans returning from Iraq and Afghanistan. *Psychosomatics* 2010; 51:503–507
18. Jakupcak M, Luterek J, Hunt S, Conybeare D, McFall M: Post-traumatic stress and its relationship to physical health functioning in a sample of Iraq and Afghanistan War veterans seeking post-deployment VA health care. *J Nerv Ment Dis* 2008; 196:425–428
19. Keane TM, Fairbank JA, Caddell JM, Zimering RT, Taylor KL, Mora CA: Clinical evaluation of a measure to assess combat exposure. *Psychol Assess* 1989; 1:53–55
20. Davidson JRT, Book SW, Colket JT, Tupler LA, Roth S, David D, et al: Assessment of a new self-rating scale for post-traumatic stress disorder. *Psychol Med* 1997; 27:153–160
21. Seal KH, Bertenthal D, Maguen S, Gima K, Chu A, Marmar CR: Getting beyond “Don't ask, don't tell”: an evaluation of US Veterans Administration post-deployment mental health screening of veterans returning from Iraq and Afghanistan. *Am J Public Health*. 2008; 98(4):714–720
22. Whooley MA, Avins AL, Miranda J, Browner WS: Case-finding instruments for depression. *J Gen Intern Med* 1997; 12:439–445
23. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M: Development of the alcohol use disorders identification test (AUDIT): WHO Collaborative Project on early detection of persons with harmful alcohol consumption–II. *Addiction* 1993; 88(6):791–804
24. Maisto SA, Carey MP, Carey KB, Gordon CM, Gleason JR: Use of the AUDIT and the DAST-10 to identify alcohol and drug use disorders among adults with a severe and persistent mental illness. *Psychol Assess* 2000; 12:186–192
25. Cassidy CM, Schmitz N, Malla A: Validation of the alcohol use disorders identification test and the drug abuse screening test in first episode psychosis. *Can J Psychiatry* 2008; 53:26–33
26. Wolfe J, Proctor SP, Erickson DJ, Heeren T, Friedman MJ, Huang MT, et al: Relationship of psychiatric status to Gulf War veterans' health problems. *Psychosom Med* 1999; 61: 532–540
27. Geisser ME, Roth RS, Bachman JE, Eckert TA: The relationship between symptoms of post-traumatic stress disorder and pain, affective disturbance and disability among patients with accident and non-accident related pain. *Pain* 1996; 66:207–214
28. Palyo SA, Beck JG: Post-traumatic stress disorder symptoms, pain, and perceived life control: associations with psychosocial and physical functioning. *Pain* 2005; 117:121–127
29. Villano CL, Rosenblum A, Magura S, Fong C, Cleland C, Betzler TF: Prevalence and correlates of posttraumatic stress disorder and chronic severe pain in psychiatric outpatients. *J Rehabil Res Dev* 2007; 44:167–178
30. Baker DG, Mendenhall CL, Simbartl LA, Magan LK, Steinberg JL: Relationship between posttraumatic stress disorder and self-reported physical symptoms in Persian Gulf War veterans. *Arch Intern Med* 1997; 157:2076–2078
31. Vanderploeg RD, Belanger HG, Horner RD, Spehard AM, Powell-Cope G, Luther SL, et al: Health outcomes associated with

- military deployment: mild traumatic brain injury, blast, trauma, and combat associations in the Florida national guard. *Arch Phys Med Rehabil* 2012; 93:1887–95
32. Mayer EA, Craske M, Naliboff BD: Depression, anxiety, and the gastrointestinal system. *J Clin Psychiatry* 2001; 62(Suppl 8):28–36; Discussion 37
 33. Schur EA, Afari N, Furberg H, Olarte M, Goldberg J, Sullivan PF, et al: Feeling bad in more ways than one: comorbidity patterns of medically unexplained and psychiatric conditions. *J Gen Intern Med* 2007; 22:818–821
 34. White DL, Savas LS, Daci K, Elserag R, Graham DP, Fitzgerald SJ, et al: Trauma history and risk of the irritable bowel syndrome in women veterans. *Aliment Pharmacol Ther* 2010; 32:551–561
 35. Poundja J, Fikretoglu D, Brunet A: The co-occurrence of post-traumatic stress disorder symptoms and pain: is depression a mediator? *J Trauma Stress* 2006; 19:747–751
 36. Roy-Byrne P, Smith WR, Goldberg J, Afari N, Buchwald D: Post-traumatic stress disorder among patients with chronic pain and chronic fatigue. *Psychol Med* 2004; 34:363–368
 37. Löwe B, Kroenke K, Spitzer RL, Williams JB, Mussell M, Rose M, et al: Trauma exposure and posttraumatic stress disorder in primary care patients: cross-sectional criterion standard study. *J Clin Psychiatry* 2011; 72:304–312
 38. Asmundson GJG, Coons MJ, Taylor S, Katz J: PTSD and the experience of pain: research and clinical implications of shared vulnerability and mutual maintenance models. *Can J Psychiatry* 2002; 47:930–937
 39. Sharp TJ, Harvey AG: Chronic pain and post-traumatic stress disorder: mutual maintenance? *Clin Psychol Rev* 2001; 21:857–877
 40. McGeary D, Moore M, Vriend CA, Peterson AL, Gatchel RJ: The evaluation and treatment of comorbid pain and PTSD in a military setting: an overview. *J Clin Psychol Med Settings* 2011; 18:155–163