

Lawrence Berkeley National Laboratory

Recent Work

Title

A NEW (CARTESIAN) REACTION PATH MODEL FOR DYNAMICS IN POLYATOMIC SYSTEMS WITH APPLICATION TO H-atom TRANSFER IN MALONALDEHYDE

Permalink

<https://escholarship.org/uc/item/1hx2q9rv>

Authors

Ruf, B.A.

Miller, W.H.

Publication Date

1987-08-01

0.2



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Chemical Sciences Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY

MAR 4 1988

LIBRARY AND
DOCUMENTS SECTION

Presented at the Faraday Discussions,
Reading, United Kingdom,
December 14-18, 1987


A New (Cartesian) Reaction Path Model for Dynamics in Polyatomic Systems with Application to H-atom Transfer in Malonaldehyde

B.A. Ruf and W.H. Miller

August 1987

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-24412
0.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

A New (Cartesian) Reaction Path Model for Dynamics in Polyatomic Systems,
with Application to H-atom Transfer in Malonaldehyde

Beverly A. Ruf and William H. Miller

Department of Chemistry, University of California, Berkeley,
and
Materials and Chemical Sciences Division, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720


ABSTRACT

A new kind of reaction path model for describing reactions in polyatomic molecular systems is presented, one which is based on the cartesian coordinates of the atoms. Not only does this lead to a simpler treatment of the interaction between the "system" (i.e., the reaction coordinate) and the "bath" (the other degrees of freedom) than earlier reaction path models based on the (curvilinear) steepest descent reaction path, but in many cases it also provides a more natural description of the dynamics. The resulting Hamiltonian has the standard form of a cartesian "system" linearly coupled to a harmonic "bath", the dynamics of which is treated in this paper by the basis set method of Makri and Miller [J. Chem. Phys. 86, 1451 (1987)]. Application to a model of H-atom transfer in malonaldehyde shows that the overall approach - both the cartesian reaction path model and the basis set method treatment of the dynamics - is quite successful.

I. INTRODUCTION

Over the past seven years or so our research group has developed (and applied) a theoretical description of dynamics in polyatomic molecular systems that is based on the minimum energy reaction path (i.e., the "intrinsic reaction coordinate").¹⁻³ This is the steepest descent path (in mass-weighted cartesian coordinates) from the transition state on a potential energy surface backwards to reactants and forward to products. The picture of the dynamics in this model is that of one-dimensional motion along the reaction path coupled to harmonic vibration about it (in $3N-7$ directions, where N is the number of atoms). Although this model has been useful for a variety of applications, and we believe will continue to be so, there are some aspects of it that are undesirable. The purpose of this paper is to describe a new class of models for describing dynamics in polyatomic systems that seems more advantageous in some respects than that based on the minimum energy reaction path.

One type of process for which the reaction path model has difficulty is hydrogen atom transfer as, for example, in malonaldehyde,^{1,2}



The reactive process is clearly the motion of essentially only one hydrogen atom, H_1 , with the other degrees of freedom playing a modest role. The problem with the reaction path description here is that the reaction path always arrives at the reactant and product wells along the normal mode of

lowest frequency (of the appropriate symmetry); cf., Appendix I. For the above example (1) the lowest frequency is associated with some floppy skeletal vibrational motion that is quite unrelated to the motion of atom H_1 , while the relevant vibration is the $O-H_1$ stretch, which is the highest frequency of the reactant. This means that in going from the transition state to the reactant and product wells the minimum energy path makes many sharp turns in the $(3N-6)$ dimensional space, so that the reaction is in no sensible way well-described as one dimensional motion along this path. (The curvature coupling elements in the reaction path Hamiltonian^{1a} are correspondingly large and difficult to deal with.)

Another drawback of the reaction path model is that the reaction path is mass-dependent. Thus, if one wishes to treat isotopically related reactions, an entire re-calculation of the minimum energy path (and force constant matrix along it) is required for each new isotopic species. One would clearly prefer a model for which the potential energy surface "input" to the dynamical treatment is independent of the nuclear masses.

The new model that we describe in Section II is from the outset much simpler than our earlier reaction path models. Cartesian coordinates of the N atoms are used for all degrees of freedom, and one relies on "chemical intuition" to select which one (or ones) is treated for arbitrarily large displacements, as opposed to the other coordinates which move only slightly. For reaction (1), for example, one notes that it is only the x coordinate of hydrogen atom H_1 that undergoes large displacement; although all the other atoms (and the y and z coordinate of H_1) move, they do not move very much. Thus, the potential energy surface in all these cartesian coordinates can be adequately approximated by a quadratic expansion about some reference configuration. Conservation of total angular momentum is accounted for

approximately (but adequately for H-atom transfer processes).

The Hamiltonian which characterizes this new approach is totally cartesian in structure and thus much simpler to deal with than the reaction path Hamiltonian. In fact, it has the generic form of a "system" which is linearly coupled to a harmonic "bath", as has been so commonly assumed for many model studies.⁴ It is much easier to describe the coupling between the "system" and "bath" here than it is in the reaction path Hamiltonian where the coupling arises from curvature coupling effects in the kinetic energy.^{1a} Finally, the potential energy surface information required for this cartesian model is simpler to generate than that for the reaction path Hamiltonian, and it is also mass-independent so that different isotopes of the same system do not require new potential energy surface calculations.

The cartesian reaction path model is developed in Section II, and Section III describes the theoretical methods used to treat the dynamics. Application to a model of H-atom transfer in malonaldehyde is presented in Section IV.

II. A CARTESIAN MODEL FOR REACTION DYNAMICS

a. Frozen Bath Version

We start with the Hamiltonian for an N atom polyatomic system expressed in the full set of 3N cartesian coordinates.

$$H = \sum_{i=1}^{3N} \frac{p_i^2}{2m_i} + V(\{x_i\}) . \quad (2)$$

The first step in defining the model is to identify which cartesian coordinate(s) one wishes to describe for arbitrarily large displacements. For reaction (1) one could choose only the x coordinate of hydrogen atom H₁; a more accurate model would be to treat both the x and y coordinates of H₁ for arbitrarily large displacement. Let $\underline{r}$ denote the one or two (or maybe three) cartesian coordinates to be described for arbitrary displacements; $\underline{R}$ denotes the remaining 3N-1 or 3N-2 cartesian coordinates that do not move very much during the reaction of interest. The $\underline{R}$ -dependence of the potential energy $V = V(\underline{r}, \underline{R})$ is then expanded in a Taylor series to second order about a reference geometry $\underline{R}_0$:

$$\begin{aligned} V(\underline{r}, \underline{R}) = & V(\underline{r}, \underline{R}_0) + \frac{\partial V(\underline{r}, \underline{R}_0)}{\partial \underline{R}_0} \cdot (\underline{R} - \underline{R}_0) \\ & + \frac{1}{2} (\underline{R} - \underline{R}_0) \cdot \frac{\partial^2 V(\underline{r}, \underline{R}_0)}{\partial \underline{R}_0 \partial \underline{R}_0} \cdot (\underline{R} - \underline{R}_0) . \end{aligned} \quad (3)$$

Equation (3) is the essence of the model, namely exploiting the fact that most

of the coordinates (i.e., atoms) do not move very much during the reaction.

With Eq. (3) the Hamiltonian takes the form

$$\begin{aligned}
 H(\underline{p}, \underline{r}, \underline{P}, \underline{R}) = & \left[\frac{1}{2} \underline{p} \cdot \underline{m}^{-1} \cdot \underline{p} + V_0(\underline{r}) \right] \\
 & + \left[\frac{1}{2} \underline{P} \cdot \underline{M}^{-1} \cdot \underline{P} + \frac{1}{2} (\underline{R} - \underline{R}_0) \cdot \underline{K}(\underline{r}) \cdot (\underline{R} - \underline{R}_0) \right] \\
 & - \underline{f}(\underline{r}) \cdot (\underline{R} - \underline{R}_0) ,
 \end{aligned} \tag{4}$$

which one recognizes as essentially the generic Hamiltonian for a (low-dimensional) "system" - the $\underline{r}$ degree(s) of freedom - coupled to a "bath" of harmonic oscillators, the $\underline{R}$ degrees of freedom. (The quantities in Eq. (4) are clearly recognizable: $V_0(\underline{r}) = V(\underline{r}, \underline{R}_0)$, $\underline{K}(\underline{r}) = \partial^2 V(\underline{r}, \underline{R}_0) / \partial \underline{R}_0 \partial \underline{R}_0$, $\underline{f}(\underline{r}) = -\partial V(\underline{r}, \underline{R}_0) / \partial \underline{R}_0$, and $\underline{m}$ is the diagonal matrix of atomic masses.) The coupling - the last term in Eq. (4) - is linear in the "bath" coordinates $\underline{R}$. This describes "solvent reorganization" effects, i.e., the change in the "bath" coordinates' instantaneous equilibrium positions as the "system" dynamics takes place. There is also another coupling between the "system" and "bath" because the force constant matrix of the bath is a function of the "system" coordinate $\underline{r}$; i.e., the instantaneous vibrational frequencies and normal mode eigenvectors change with the system dynamics. One familiar with the reaction path formalism will immediately recognize how much more readily one can deal with system-bath coupling with this new Hamiltonian, Eq. (4), than with that in the reaction path Hamiltonian.^{1a}

The reader will also note that the required input from ab initio quantum

chemistry calculations is much easier to generate for the Hamiltonian in Eq. (4) than it is for the reaction path Hamiltonian. What is required for Eq. (4) is the energy, gradient, and force constant matrix of the potential energy surface for a frozen bath ($R=R_0$), as a function of r . If r is only one coordinate, for example, this means that one requires an energy, gradient, and force constant matrix at, say, ten predetermined geometries. Geometry optimization is not required.

b. Conservation of Total Angular (and Linear) Momentum

Our initial inclination was to ignore the fact that this model does not conserve total angular and linear momentum (due to the fact that the approximation to the potential surface, Eq. (3), destroys rotational and translational invariance). Earlier, for example, Jaquet and Miller⁵ had used essentially this model to treat H-atom diffusion on a tungsten surface, and there, of course, one does not need to be concerned with rotation and translation of the (infinite) surface of tungsten atoms. In malonaldehyde, Eq. (1), however, the "substrate" is not infinite. It seemed to us that it was sufficiently more massive than the one H atom that tunnels and that one could thus ignore rotation and translation. Unfortunately, test calculations convinced us that this is not the case, at least for malonaldehyde; the low frequencies of the bath - the ones that should be zero, i.e., pure rotations and translations - mix in an unphysical way with the true low vibrational frequencies of the molecule. It was thus deemed necessary to project out six pure rotational and translational degrees of freedom.

To separate off the rotational motion rigorously requires use of curvilinear coordinates (i.e., Euler angles),⁶ thus destroying the cartesian form of the Hamiltonian, Eq. (4). To avoid this, and also because we are

developing the present model for application to large molecular systems for which rotational motion is not of interest, we have chosen to eliminate rotation approximately by projecting out infinitesimal rotation and translation of the N-1 "substrate" atoms.

To be specific, we refer to Eq. (1) with $\underline{r} = x_{H_1}$ chosen as the (one) "system" coordinate. Indices $i = 1, 2, 3$ refer to the three cartesian coordinates of atom H_1 . The matrix

$$\{U_{i,k}\} \quad \begin{array}{l} i = 4, \dots, 3N \\ k = 4, \dots, 3N-6 \end{array} \quad (5)$$

denotes the matrix of eigenvectors of the projected, mass-weighted force constant matrix of the N-1 atom substrate,

$$(\underline{1}-\underline{P}) \cdot \underline{m}^{-1/2} \cdot \underline{K}_O \cdot \underline{m}^{-1/2} \cdot (\underline{1}-\underline{P}) ; \quad (6)$$

here $\underline{K}_O = \underline{K}(x_1^O)$, $\underline{m}$, and $\underline{P}$ are $(3N-3) \times (3N-3)$ matrices for the (N-1) atom substrate, and x_1^O is some convenient intermediate value of the system coordinate x_1 . $\underline{P}$ is the projector onto the 6 degrees of freedom that are infinitesimal rotations and translations of the (N-1) atom substrate; its explicit form has been given before^{1a} and is a function only of the geometry of the substrate. The factor $(\underline{1}-\underline{P})$ in Eq. (6) insures that the projected, mass-weighted force constant matrix will have six zero eigenvalues, corresponding to infinitesimal rotations and translations, and $3(N-1)-6$ non-zero eigenvalues that describe vibration of the (N-1) substrate atoms about their reference positions.

The final matrix $\underline{U}$ is now augmented by the other cartesian coordinates of

atom H_1 that are not taken as system coordinates; e.g., with $\underline{r}$ chosen to be $x_{H_1} = x_1$, the coordinates $y_{H_1} = x_2$ and $z_{H_1} = x_3$ are part of the bath, and the final $\underline{U}$ matrix is

$$\{U_{1,k}\} = \begin{matrix} & \begin{matrix} 2 & 3 & 4 & \dots & 3N-6 \end{matrix} \\ \begin{matrix} 2 \\ 3 \\ 4 \\ \vdots \\ 3N \end{matrix} & \begin{pmatrix} 1 & 0 & & & 0 \\ 0 & 1 & & & \\ \vdots & & \ddots & & \\ 0 & & & \underline{U} \text{ from Eq. (5)} & \\ \vdots & & & & \ddots \end{pmatrix} \end{matrix} \quad (7)$$

The $(3N-1)$ cartesian bath coordinates $\underline{R}$ are now expressed in terms of the $3N-7$ normal mode bath coordinates $\underline{Q}$ which have the six rotations and translations of the substrate eliminated,

$$\underline{R} - \underline{R}_0 = \underline{m}^{-1/2} \cdot \underline{U} \cdot \underline{Q} , \quad (8)$$

with $\underline{U}$ given by Eq. (7). The Hamiltonian of Eq. (4) thus becomes
(with $\underline{r} = x_1$)

$$H(p_1, x_1, \underline{p}, \underline{Q}) = \left[\frac{p_1^2}{2m_1} + v_{\text{eff}}(x_1) \right] + \left[\sum_{k=2}^{3N-6} \frac{1}{2} p_k^2 + \frac{1}{2} \underline{Q} \cdot \underline{K}_{\text{eff}}(x_1) \cdot \underline{Q} \right] - f_{\text{eff}}(x_1) \cdot \underline{Q} , \quad (9a)$$

where $\underline{K}_{\text{eff}}$ and f_{eff} are the $(3N-7)$ matrix and vector

$$\underline{K}_{\text{eff}}(x_1) = \underline{U}^\dagger \cdot \underline{m}^{-1/2} \cdot \underline{K}(x_1) \cdot \underline{m}^{-1/2} \cdot \underline{U} \quad (9b)$$

$$\underline{f}_{\text{eff}}(x_1) = \underline{f}(x_1) \cdot \underline{m}^{-1/2} \cdot \underline{U} \quad (9c)$$

and

$$V_{\text{eff}}(x_1) = V(x_1, \underline{R}_0) . \quad (9d)$$

(Note that $\underline{K}_{\text{eff}}(x_1)$ is not diagonal because $\underline{U}$ was obtained by diagonalizing $\underline{K}(x_1)$ at the fixed value $x_1 = x_1^0$.) This Hamiltonian, Eq. (9), is one for 3N-6 degrees of freedom. It should be clear how Eqs. (5)-(9) are modified if the system coordinate(s) $\underline{r}$ is chosen to be x_1 and $x_2 = y_{H_1}$, say.

The transformations described by Eqs. (5)-(9) for eliminating overall translation and rotation of the substrate must be re-done for different isotopic species; i.e., the matrix $\underline{K}_{\text{eff}}(x_1)$ and coupling vector $\underline{f}_{\text{eff}}(x_1)$ in Eq. (9) depend on the atomic masses. It should be noted, however, that the mass dependence on the projected (3N-3) x (3N-3) substrate portion of $\underline{K}_{\text{eff}}(x_1)$ corresponds to secondary isotope effects. Primary isotopic substitutions are simply incorporated via the identity portion of the matrix $\underline{U}$. Both of these mass variations are simple, though. The important matter is that the original $\underline{K}$ and $\underline{f}$ of Eq. (4), which are obtained from ab initio quantum chemistry calculations, are mass-independent. Also important for treating the dynamics of the "system-bath" Hamiltonian Eq. (9) is that its cartesian and linear coupling form has been maintained by the (approximate) way that we have eliminated overall translation and rotation.

c. Flexible Bath

In some cases one may wish to allow the reference geometry of the "bath" to vary with the system coordinate $\underline{r}$, e.g., so that the equilibrium geometries

of the reactants and products are accurately reproduced by the model. It is actually possible to generalize the above treatment in a simple way to incorporate this and still maintain the simple form of the resulting Hamiltonian.

Thus, let $\underline{R}_0(\underline{r})$ be the reference geometry of the bath as a function of the system coordinate(s) $\underline{r}$. We envision, for example, that $\underline{R}_0(\underline{r})$ may be chosen simply to interpolate between the reactant and product geometries of the bath variables. A Taylor series expansion of $\underline{R}$ about $\underline{R}_0(\underline{r})$, as in Eq. (3), is still possible.

$$V(\underline{r}, \underline{R}) = V(\underline{r}, \underline{R}_0(\underline{r})) + \left(\frac{\partial V(\underline{r}, \underline{R})}{\partial \underline{R}} \right)_{\underline{R}=\underline{R}_0(\underline{r})} \cdot (\underline{R} - \underline{R}_0(\underline{r})) + \frac{1}{2} (\underline{R} - \underline{R}_0(\underline{r})) \cdot \left(\frac{\partial^2 V(\underline{r}, \underline{R})}{\partial \underline{R} \partial \underline{R}} \right)_{\underline{R}=\underline{R}_0(\underline{r})} \cdot (\underline{R} - \underline{R}_0(\underline{r})) , \quad (10)$$

and this can be combined with the cartesian kinetic energy to form a Hamiltonian like that of Eq. (4).

To maintain the simple form of the resulting Hamiltonian it is still necessary to project out rotations and translations of a frozen substrate. Therefore, the transformation matrix $\underline{U}$ is defined as above, Eq. (5)-(7), but where $\underline{K}_0 = \underline{K}(\underline{x}_1^0, \underline{R}_0(\underline{x}_1^0))$, $\underline{x}_1^0$ being an intermediate value of the system coordinate $\underline{x}_1$. The projector $\underline{P}$ is also defined at the frozen substrate geometry $\underline{R}_0(\underline{x}_1^0)$. The relation between the $(3N-1)$ coordinates $\underline{R}$ (for $\underline{r} = \underline{x}_1$) and the $(3N-7)$ coordinates $\underline{Q}$ is thus

$$\underline{R} - \underline{R}_0(\underline{x}_1^0) = \underline{m}^{-1/2} \cdot \underline{U} \cdot \underline{Q} , \quad (11)$$

so that $\underline{R}-\underline{R}_0(x_1)$ in Eq. (10) (with $\underline{r} = x_1$) is given by

$$\underline{R}-\underline{R}_0(x_1) = \underline{R}_0(x_1^0) - \underline{R}_0(x_1) + \underline{m}^{-1/2} \cdot \underline{U} \cdot \underline{Q} . \quad (12)$$

Using Eq. (12) in Eq. (10) then gives the same form of Hamiltonian as before, Eq. (9a), here with

$$\underline{K}_{\text{eff}}(x_1) = \underline{U}^+ \cdot \underline{m}^{-1/2} \cdot \underline{K}(x_1) \cdot \underline{m}^{-1/2} \cdot \underline{U} \quad (9b')$$

$$\underline{f}_{\text{eff}}(x_1) = [\underline{f}(x_1) + (\underline{R}_0(x_1) - \underline{R}_0(x_1^0)) \cdot \underline{K}(x_1)] \cdot \underline{m}^{-1/2} \cdot \underline{U} \quad (9c')$$

$$\underline{V}_{\text{eff}}(x_1) = \underline{V}(x_1, \underline{R}_0(x_1)) + \underline{f}(x_1) \cdot (\underline{R}_0(x_1) - \underline{R}_0(x_1^0)) \quad (9d')$$

$$+ \frac{1}{2} (\underline{R}_0(x_1) - \underline{R}_0(x_1^0)) \cdot \underline{K}(x_1) \cdot (\underline{R}_0(x_1) - \underline{R}_0(x_1^0))$$

where

$$\underline{K}(x_1) = \left(\frac{\partial^2 \underline{V}(x_1, \underline{R})}{\partial \underline{R} \partial \underline{R}} \right)_{\underline{R} = \underline{R}_0(x_1)} ,$$

$$\underline{f}(x_1) = - \left(\frac{\partial \underline{V}(x_1, \underline{R})}{\partial \underline{R}} \right)_{\underline{R} = \underline{R}_0(x_1)} .$$

It is clear that these expressions revert to Eq. (9) in the rigid bath limit, where $\underline{R}_0(x_1) = \underline{R}_0$.

To conclude this description of the model we summarize the advantages that it has over earlier reaction path models. (1) Most important is that it

contains the physically correct picture of the dynamics, clearly identifying the relevant coordinates of the process. The intrinsic reaction path for these heavy-light-heavy mass combinations leads to unphysical reaction paths.⁷ (2) It is considerably easier to treat the dynamics of the resulting Hamiltonian which has the coupling in the potential energy rather than that for the reaction path Hamiltonian that has the coupling in the kinetic energy. (3) It is relatively simple to do calculations for different isotopes. (4) It is much simpler to generate the ab initio quantum chemistry "input" for the model.

Disadvantages of the model, on the other hand, are that we are able to separate off overall rotational motion only approximately if we wish (as we do) to maintain the simple form of the Hamiltonian; this seems to be a minor error, though, particularly so if the "substrate" is large. Finally, the model requires that one invoke "chemical intuition" to choose the cartesian coordinates that constitute the "system"; one may view this as an advantage or a disadvantage.

III. SYSTEM-BATH DYNAMICS

Having defined the model Hamiltonian in the previous section, one is now ready to treat its dynamics. The most elegant and rigorous way to treat these system-harmonic bath Hamiltonians is via Feynman path integral methods.⁸ This permits one to take into account the effect of the "bath" on the "system" exactly. There is currently a great deal of progress being made in this direction,^{5,9-14} but these approaches are at present not available for practical calculations with real chemical systems. In this paper, therefore, we utilize an approximate treatment due to Makri and Miller⁴ that has been shown to do an excellent job for including the effect of coupling to a bath on the tunneling in a double well system, as is the process in Eq. (1). We first summarize the basic ideas of this approach and then describe some necessary modifications for the present application.

The basis set method of Makri and Miller chooses basis functions for the total system-bath Hamiltonian in the form

$$\psi_{i\bar{n}}(r, \underline{Q}) = x_i(r) \phi_{\bar{n}}^i(\underline{Q}) , \quad (13)$$

where $\{x_i(r)\}$ is a set of localized functions in the "system" coordinate, and $\phi_{\bar{n}}^i(\underline{Q})$ is the eigenfunction of the "bath" Hamiltonian that results when the total Hamiltonian is averaged over basis function $x_i(r)$; $\phi_{\bar{n}}^i(\underline{Q})$ is obtainable analytically because the resulting "bath" Hamiltonian is that of linearly coupled harmonic oscillators. (We have taken the set $\{x_i\}$ as the distributed gaussians of Hamilton and Light,¹⁵ as used by Makri and Miller.) The matrix of the total system-bath Hamiltonian, $H_{i'\bar{n}', i\bar{n}}$, is constructed in this basis, and the zeroth order effective system Hamiltonian of ref. 4 is defined by taking the part of the Hamiltonian matrix that is diagonal in the "bath"

quantum numbers, i.e., by setting $\underline{n}' = \underline{n}$. This effective system Hamiltonian matrix has a dimension only of the number of "system" basis functions $\{\chi_i\}$ and is of the form

$$\begin{aligned} H_{i',i}^{\underline{n}} &= H_{i',\underline{n},i,\underline{n}} \\ &= F_{i',i}^{\underline{n}} \times h_{i',i}^{\underline{n}}, \end{aligned} \quad (14)$$

where $F_{i',i}^{\underline{n}}$ is the Franck-Condon factor between the oscillator functions $\phi_{\underline{n}}^i$ and $\phi_{\underline{n}}^{i'}$,

$$F_{i',i}^{\underline{n}} = \int dQ \phi_{\underline{n}}^{i'}(Q) \phi_{\underline{n}}^i(Q), \quad (15)$$

and $h_{i',i}^{\underline{n}}$ is a one-dimensional-like Hamiltonian matrix. Most of the effect of the "bath" on the "system" dynamics is contained in the Franck-Condon factor; i.e., it describes the "solvent relaxation", or polaronic effects of the "bath" on the "system". Makri and Miller found that this approximation, i.e., taking the bath quantum numbers to be diagonal, worked quite well provided that the basic functions $\{\chi_i\}$ are localized. The reader should see ref. 4 for discussion of the reasons for this as well as further aspects of the approach.

Makri and Miller considered a system-bath Hamiltonian with a constant force constant matrix, $\underline{K}$, so their treatment must be generalized in order to apply it to the present Hamiltonian, Eq. (9a). Specifically, the Franck-Condon factor of Eq. (15) in the present case is given more explicitly by

$$F_{i',i}^{\underline{n}} = \int dQ \prod_k \phi_{n_k}^{i'}(q_k^{i'}) \phi_{n_k}^i(q_k^i), \quad (16)$$

where $\phi_{n_k}(q_k)$ are ordinary one-dimensional harmonic oscillator wavefunctions, and the coordinates g^i (and $g^{i'}$) are the linear combinations of Q that diagonalize the oscillator potentials

$$\frac{1}{2} Q \cdot K_1 \cdot Q - f_1 \cdot Q, \quad (17a)$$

where

$$K_1 = \langle \chi_1 | K_{\text{eff}} | \chi_1 \rangle \quad (17b)$$

$$f_1 = \langle \chi_1 | f_{\text{eff}} | \chi_1 \rangle. \quad (17c)$$

More specifically, if L_1 is the $(3N-7) \times (3N-7)$ matrix of eigenvectors of K_1 , then

$$L_1^+ \cdot K_1 \cdot L_1 = (\omega_1)^2 \text{ (diagonal)} \quad (18)$$

are the diagonal frequencies of the bath modes associated with the gaussian χ_1 localized at r_1 , and g^i is given in terms of Q by

$$g^i = L_1^+ \cdot Q - \omega_1^{-2} \cdot L_1^+ \cdot f_1. \quad (19)$$

Because the matrices L_1 and L_1 , are different - which is due to the fact that the force constant matrices K_1 and K_1 , are different - the integrals in Eq. (16) do not factor into a product of one-dimensional integrals; they have the form of a multidimensional gaussian integral (times powers). There have, however, been a number of papers describing the efficient evaluation of these multidimensional harmonic Franck-Condon factors using generating function, recursion, and iterative methods;¹⁶ we used a method similar to that of ref.

16a.

With this more generalized Franck-Condon factor, the effective system Hamiltonian which results from the system-bath Hamiltonian of Eq. (9a) is given explicitly by

$$\begin{aligned}
 H_{1',1}^{\underline{n}} = & F_{1',1}^{\underline{n}} \langle \chi_1, | \frac{p_1^2}{2m_1} + v_{\text{eff}}(x_1) | \chi_1 \rangle \\
 & + \frac{1}{2} \langle \chi_1, | \chi_1 \rangle \langle \phi_{\underline{n}}^{1'} | p_{\underline{n}}^2 | \phi_{\underline{n}}^1 \rangle \\
 & + \frac{1}{2} \langle \phi_{\underline{n}}^1 | \underline{Q} \cdot \underline{K}_{1',1} \cdot \underline{Q} | \phi_{\underline{n}}^1 \rangle \\
 & - \underline{f}_{1',1} \cdot \langle \phi_{\underline{n}}^{1'} | \underline{Q} | \phi_{\underline{n}}^1 \rangle ,
 \end{aligned} \tag{20a}$$

where

$$\underline{K}_{1',1} = \langle \chi_1, | \underline{K}_{\text{eff}}(x_1) | \chi_1 \rangle \tag{20b}$$


$$\underline{f}_{1',1} = \langle \chi_1, | \underline{f}_{\text{eff}}(x_1) | \chi_1 \rangle . \tag{20c}$$

The last three terms in Eq. (20) are multidimensional gaussians times powers and are evaluated by methods similar to those used for the Franck-Condon factor.

IV. APPLICATION TO 3-ATOM MODEL OF MALONALDEHYDE

Before applying the methodology of Sections II and III to an ab initio treatment of malonaldehyde, Eq.(1), we consider here a simple 3-atom model of this reaction that can be treated exactly (because it is a triatomic system). It is important to test both the cartesian reaction path model described in Section II, and also the dynamical treatment summarized in Section III.

The 3-atom model we consider is the O-H-O part of Eq.(1),



with a potential energy function that has the general form of Janoschek, Weideman, and Zundel's¹⁷ double well potential, generalized to include bending motion:

$$V = a_1 + a_2 Q_a^2 + a_3 Q_a^4 + a_4 Q_s^2 + a_5 Q_a^2 Q_s + a_6 Q_s^2 + \frac{1}{2} K_Y (\gamma - \gamma_0)^2, \quad (22)$$

where γ is the angle between the two OH bonds. The coefficients have been chosen to approximate the energetics and geometry of reaction (1):

$$a_1 = 0.01338 \text{ (a.u.)}$$

$$a_2 = -0.03603$$

$$a_3 = 0.02425$$

$$a_4 = -0.02745$$

$$a_5 = 0.03695$$

$$a_6 = 0.07786$$

$$K_Y = 0.20000$$

$$Q_a = \frac{1}{\sqrt{2}} (r_1 - r_2)$$

$$Q_s = \frac{1}{\sqrt{2}} (r_1 + r_2)$$

$$\gamma_0 = 158^\circ$$

The "bare" barrier height (i.e. with no zero point energy corrections) for this potential is 6.88 kcal/mole, in qualitative agreement with what is thought be the correct value for malonaldehyde.^{18,19} Table I compares the equilibrium and transition state geometries for this potential to the corresponding quantities for malonaldehyde.

The cartesian reaction model of Section II was now applied, with the x-coordinate of the H-atom as the "system" coordinate, i.e., the reaction coordinate. Both the "frozen" reference geometry for the bath (Section IIa) and a flexible reference geometry (Section IIc) were tried to test the sensitivity of the results to how well the bath is modeled.

Table II gives the tunneling splitting for the ground vibrational state of this double well potential, as calculated by the basis set method described in Section III. Results in Table II are given for the principal isotope, and also for the deuterated system, each for three different choices of the reference geometry of the bath: "Transition State" refers to a frozen reference geometry of the bath that is that of the transition state; "Equilibrium" also refers to a frozen reference geometry (but one that is the average of the two equilibrium geometries); and "Flexible" refers to the variable reference geometry (Section IIc) that interpolates between the transition state and equilibrium geometries. The "exact" values given in Table II were calculated by the method of Carter and Handy²⁰, which is readily applied to any triatomic system. Finally, the values given in parenthesis in Table II are the results obtained from the one-dimensional vibrationally

adiabatic approximation and are thus a measure of how much coupling to the bath effects the reaction coordinate.

From the results in Table II one sees that both an equilibrium and a flexible reference geometry for the bath do a better job of describing the dynamics of H-atom transfer (i.e., the tunneling splitting) than does a frozen transition state reference geometry; the former results are in quite good agreement with the exact values for this model system. One also sees from Table II that the bath has a very significant effect on the tunneling dynamics; i.e., the values given by the one dimensional vibrationally adiabatic approximation (those in parenthesis) are in error by 50 to 100%. The error for the equilibrium geometry is the smallest while the transition state is the largest. It thus appears that a quadratic expansion of the potential in bath coordinates about a frozen reference geometry can be sufficient if the geometry of the bath does not relax very much during the reaction. A flexible reference is necessary, however, if there is a lot of relaxation in the bath. Fortunately, the flexible bath model of Section IIc is not any more difficult to apply than is the frozen bath treatment of Section IIa.

V. CONCLUDING REMARKS

The cartesian reaction path model presented in Section II provides a much simpler description of the interaction between the "system" (i.e., the reaction coordinate) and the "bath" (the remaining degrees of freedom) than does a reaction path model based on the (curvilinear) minimum energy (i.e., steepest descent) path. As discussed, too, it is often a more physically correct picture of the dynamics (as, for example, in the case of H-atom transfer, where the minimum energy reaction path is very sharply curved). It is also gratifying that the kind of quantum chemistry calculations that are necessary to apply the model in its ab initio mode are simpler to generate than for the steepest descent path.

The Hamiltonian that results from this cartesian reaction path model has the generic form of a cartesian "system" linearly coupled to a harmonic "bath". The most powerful way for treating the dynamics of such a system is Feynman path integral methodology, but it is encouraging to see that the simpler basis set method summarized in Section III does quite a good job of describing how coupling to the bath affects the H-atom transfer dynamics. We thus believe that the overall approach described herein will be useful for treating a variety of polyatomic reactive processes.

Acknowledgments

This research was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy, under Contract Number DE-AC03-76SF00098. The calculations were performed on the Berkeley Theoretical Chemistry computing facility, supported by the National Science Foundation Grant CHE84-16345.

Appendix A

This is a short insert for showing why the reaction coordinate always approaches a stationary point from the lowest frequency mode.

Consider a potential $V = V_0 + \frac{1}{2} \sum_1 \omega_1^2 x_1^2$. For the path of steepest descent

$$\frac{\partial a}{\partial s} = \frac{-\partial V}{\partial \underline{x}} / \left| \frac{\partial V}{\partial \underline{x}} \right| = \frac{\underline{x}'}{|\underline{x}'|}$$

where

$$\underline{x}' = -\frac{\partial V}{\partial \underline{x}}. \quad (\text{A.1})$$

This implies that:

$$\frac{x_1'}{x_j'} = \frac{\partial V}{\partial x_1} / \frac{\partial V}{\partial x_j} = \frac{\omega_1^2 x_1}{\omega_j^2 x_j},$$

$$\int \frac{x_1'}{x_1} dx = \int \left(\frac{\omega_1}{\omega_j} \right)^2 \frac{x_j'}{x_j} dx,$$

and

$$\ln x_1 = \left(\frac{\omega_1}{\omega_j} \right)^2 \ln x_j.$$

therefore

$$x_1 = x_j \left(\frac{\omega_1}{\omega_j} \right)^2 \quad \rightarrow \quad \text{along the path.} \quad (\text{A.2})$$

Examining two possible cases, (a) $\omega_1 > \omega_j$ and (b) $\omega_1 < \omega_j$, one can see that the steepest path approaches the stationary point from the mode of lowest frequency (a) x_j and (b) x_1 .

References

1. (a) W. H. Miller, N. C. Handy, and J. E. Adams, *J. Chem. Phys.* 72, 99 (1980); (b) W. H. Miller, in Potential Energy Surfaces and Dynamics Calculations, edited by D. G. Truhlar (Plenum, New York, 1981), p. 265; (c) C. J. Cerjan, S.-H. Shi, and W. H. Miller, *J. Phys. Chem.* 86, 2244 (1982); (d) S. K. Gray, W. H. Miller, Y. Yamaguchi, and H. F. Schaefer, *J. Chem. Phys.* 73, 2733 (1980); (e) S. K. Gray, W. H. Miller, Y. Yamaguchi, and H. F. Schaefer, *J. Am. Chem. Soc.* 103, 1900 (1981); (f) Y. Osamura, H. F. Schaefer, S. K. Gray, and W. H. Miller, *ibid.* 103, 1094 (1981); (g) B. A. Waite, S. K. Gray, and W. H. Miller, *J. Chem. Phys.* 78, 259 (1983); (h) W. H. Miller, *J. Phys. Chem.* 87, 3811 (1983); (i) T. Carrington, Jr., L. M. Hubbard, H. F. Schaefer, and W. H. Miller, *J. Chem. Phys.* 80, 4347 (1984); (j) T. Carrington, Jr. and W. H. Miller, *J. Chem. Phys.* 81, 3573 (1984); (k) W. H. Miller, in The Theory of Chemical Reaction Dynamics, edited by D. C. Clary (D. Reidel, Boston, 1986), p. 27; (l) T. Carrington, Jr. and W. H. Miller, *J. Chem. Phys.* 84, 4364 (1986).
2. For early work on reaction paths and reaction coordinates, see (a) S. Glasstone, K. J. Laidler, and H. Eyring, The Theory of Rate Processes, (McGraw-Hill, New York, 1941); (b) R. A. Marcus, *J. Chem. Phys.* 45, 4493, 4500 (1966); 49, 2610 (1968); (c) G. L. Hofacker, *Z. Naturforsch. Teil A.* 18, 607 (1963); (d) S. F. Fischer, G. L. Hofacker, and R. Seiler, *J. Chem. Phys.* 51, 3941 (1969); (e) K. Fukui, *J. Phys. Chem.* 74, 4161 (1970).

3. Some of the recent papers by other workers on reaction path models are
(a) S. F. Fischer and M. A. Ratner, J. Chem. Phys. 57, 2769 (1972); (b) P. Russegger and J. Brickman, ibid. 62, 1086 (1975); 60, 1 (1977); (c) M. V. Basilevskii, Chem. Phys. 24, 81 (1977); 67, 337 (1982); M. V. Basilevskii and A. G. Shamov, ibid. 60, 349 (1981); (d) K. Fukui, S. Kato, and H. Fujimoto, J. Am. Chem. Soc. 97, 1 (1975); K. Yamashita, T. Yamabe, and K. Fukui, Chem. Phys. Lett. 84, 123 (1981); A. K. Fukui, Acc. Chem. Res. 14, 363 (1981); (e) K. Ishida, K. Morokuma, and A. Komornicki, J. Chem. Phys. 66, 2153 (1977); (f) A. Nauts and X. Chapuisat, Chem. Phys. Lett. 85, 212 (1982); X. Chapuisat, A. Nauts, and G. Durrand, Chem. Phys. 56, 91 (1981); (g) J. Pancir, Collect. Czech. Commun. 40, 1112 (1975); 42, 16 (1977); (h) G. A. Natanson, Mol. Phys. 46, 481 (1982).
4. N. Makri and W. H. Miller, J. Chem. Phys. 86, 1451 (1987).
5. R. Jaquet and W. H. Miller, J. Phys. Chem. 89, 2139 (1985).
6. See, for example, E. B. Wilson, Jr., J. C. Decius, and P. C. Cross, Molecular Vibrations, McGraw-Hill, N.Y., 1955, pp. 273-279.
7. See, for example, (a) J. Manz and J. Römelt, Chem. Phys. Lett. 81, 179 (1981); (b) J. A. Kaye and A. Kuppermann, ibid. 77, 573 (1981); 78, 546 (1981); (c) V. K. Babamov and R. A. Marcus, J. Chem. Phys. 74, 1790 (1981); (d) C. Hiller, J. Manz, W. H. Miller, and J. Römelt, ibid. 78, 3850 (1983); (e) B. C. Garrett, D. G. Truhlar, A. F. Wagner, and T. H. Dunning, Jr., ibid. 45, 120 (1973).
8. R. P. Feynman and A. R. Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill, N.Y. (1965).

9. (a) W. H. Miller, S. D. Schwartz, and J. W. Tromp, J. Chem. Phys. 79, 4889 (1983); (b) Y. Yamashita and W. H. Miller, J. Chem. Phys. 82, 5475 (1985).
10. (a) A. O. Caldeira and A. J. Leggett, Ann. Phys. (N.Y.) 149, 374 (1983); (b) S. Chakravarty and A. J. Leggett, Phys. Rev. Lett. 52, 5 (1984); (c) H. Grabert and U. Weiss, Phys. Rev. Lett. 54, 1605 (1985); (d) A. T. Dorsey, M. P. A. Fisher, and M. S. Wartak, Phys. Rev. A 33, 1117 (1986).
11. (a) D. Thirumalai and B. J. Berne, J. Chem. Phys. 79, 5029 (1983); (b) D. Thirumalai, E. J. Bruskin, and B. J. Berne, ibid. 79, 5063 (1983); (c) D. Thirumalai and B. J. Berne, ibid. 81, 2512 (1984).
12. (a) J. D. Doll, J. Chem. Phys. 81, 3536 (1984); (b) D. L. Freeman, R. D. Coalson, and J. D. Doll, J. Stat. Phys. 43, 931 (1986).
13. (a) E. C. Behrmann, G. A. Jongeward, and P. G. Wolynes, J. Chem. Phys. 79, 6277 (1983); (b) R. W. Hall and P. G. Wolynes, J. Stat. Phys. 43, 935 (1986).
14. (a) D. Chandler and P. Wolynes, J. Chem. Phys. 74, 4078 (1981); (b) M. Sprik, M. L. Klein, and D. Chandler, J. Chem. Phys. 83, 3042 (1985); (c) M. Sprik, R. W. Impey, and M. L. Klein, J. Chem. Phys. 83, 5802 (1985); (d) C. D. Jonah, C. Romeo, and A. Rahman, Chem. Phys. Lett. 123, 209 (1986); (e) P. J. Rossky, J. Schnitker, and R. A. Kuharski, J. Stat. Phys. 43, 949 (1986); (f) A. Wallquist, D. Thirumalai, B. J. Berne, J. Chem. Phys. 85, 1583 (1986).
15. I. P. Hamilton and J. C. Light, J. Chem. Phys. 84, 306 (1986).
16. (a) D. Gruner and P. Brumer, Chem. Phys. Lett. 138, 310 (1987); (b) M. Kupka and P. H. Cribb, J. Chem. Phys. 85, 1303 (1986); (c) F. T. Chau, J. Mol. Spec. 103, 66 (1984); and references therein.

17. R. Janoschek, E. G. Weidemann, and G. Zundel, J. Chem. Soc. Faraday Trans. 2, 505 (1973).
18. (a) W. F. Rowe, R. W. Duerst, and E. B. Wilson, J. Am. Chem. Soc. 98, 4021 (1976); (b) S. L. Baughcum, R. W. Duerst, W. F. Rowe, A. Smith, and E. B. Wilson, ibid. 103, 6296 (1981); (c) Z. Smith, E. B. Wilson, and R. W. Duerst, Spectrochim. Acta Part A 39, 1117 (1983); (d) S. L. Baughcum, Z. Smith, E. B. Wilson, and R. W. Duerst, J. Am. Chem. Soc. 106, 2260 (1984); (e) P. Turner, S. L. Baughcum, S. L. Coy, and Z. Smith, ibid. 106, 2265 (1984).
19. J. Bicerano, H. F. Schaefer, and W. H. Miller, J. Am. Chem. Soc. 105, 2550 (1983); M. J. Frisch, A. C. Scheiner, and H. F. Schaefer, J. Chem. Phys. 82, 4194 (1985).
20. S. Carter and N. C. Handy, Mol. Phys. 47, 1445 (1982); Mol. Phys. 57, 175 (1986).

Table I. Comparison of Geometrics of the 3-Atom Model to Malonaldehyde.

<u>Internal Coordinates^a</u>	<u>Equilibrium</u>		<u>Transition State</u>	
	<u>3-Atom Model</u>	<u>Malonaldehyde^b</u>	<u>3-Atom Model</u>	<u>Malonaldehyde^b</u>
r_1	0.82 A	0.99	1.20	1.20
r_2	1.46	1.69	1.20	1.20
r_3	2.23	1.22	2.36	2.36
γ	158°	155°	158°	158°

^a r_1 and r_2 are OH bond lengths; r_3 is the O-O bond length; γ is the angle between the two OH bonds

^bSee refs. 18 and 19.

Table II. Tunneling Splitting (cm^{-1}) for the 3-Atom Model of Malonaldehyde.

<u>Isotope</u>	<u>Reference Geometry of Bath</u>			
	<u>Transition State^a</u>	<u>Equilibrium^a</u>	<u>Flexible^b</u>	<u>Exact^c</u>
OHO	81 (156) ^d	76 (63)	66 (95)	76
ODO	23 (30)	9 (18)	7 (22)	9

^aFrozen reference geometry (Section IIa)

^bFlexible reference geometry (Section IIc)

^cExact values of the tunneling splitting (for this model)

^dValues in parentheses are tunneling splittings given by the one-dimension vibrational adiabatic approximation.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*