

Lawrence Berkeley National Laboratory

LBL Publications

Title

SEARCH FOR CHARGE $(4/3)e$ PARTICLES PRODUCED IN e^+e^- ANNIHILATIONS

Permalink

<https://escholarship.org/uc/item/1j73w2mh>

Author

Aihara, H.

Publication Date

1983-10-01

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

Physics, Computer Science &
Mathematics Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY
FEB 1 1984
LIBRARY AND
DOCUMENTS SECTION


Submitted to Physical Review Letters

SEARCH FOR CHARGE $(4/3)e$ PARTICLES PRODUCED IN
 e^+e^- ANNIHILATIONS

PEP-4 Collaboration

October 1983

For Reference
Not to be taken from this room



LBL-16860
C1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Search for Charge $(4/3)e$ Particles Produced in e^+e^- Annihilations

H. Aihara⁶, M. Alston-Garnjost¹, D.H. Badtke⁴, J.A. Bakken⁴, A. Barbaro-Galtieri¹, A.V. Barnes¹, B.A. Barnett⁴, B.J. Blumenfeld⁴, A. Bross¹, C.D. Buchanan², W.C. Carithers¹, O. Chamberlain¹, C-Q. Chen¹, J. Chiba⁶, C-Y. Chien⁴, A.R. Clark¹, O.I. Dahl¹, C.T. Day¹, P. Delpierre¹, K.A. Derby¹, P.H. Eberhard¹, D.L. Fancher¹, H. Fujii⁶, T. Fujii⁶, B. Gabioud¹, J.W. Gary¹, W. Gorn³, W-X. Gu¹, N.J. Hadley¹, J.M. Hauptman², H.J. Hilke¹, W. Hofmann¹, J.E. Huth¹, J. Huyen⁴, H. Iwasaki⁶, T. Kamae⁶, R.W. Kenney¹, L.T. Kerth¹, R.I. Koda², R.R. Kofler⁵, K.K. Kwong³, J.G. Layter³, C.S. Lindsey³, G. London¹, S.C. Loken¹, X-Q. Lu⁴, G.R. Lynch¹, L. Madansky⁴, R.J. Madaras¹, R. Majka⁷, J. Mallet¹, P.S. Martin¹, K. Maruyama⁶, J.N. Marx¹, J.A.J. Matthews⁴, S.O. Melnikoff³, W. Moses¹, P. Nemethy¹, D.R. Nygren¹, P.J. Oddone¹, D. Park², A. Pevsner⁴, M. Pripstein¹, P.R. Robrish¹, M.T. Ronan¹, R.R. Ross¹, F.R. Rouse¹, R.R. Sauerwein¹, G. Shapiro¹, M.D. Shapiro¹, B.C. Shen³, W.E. Slater², M.L. Stevenson¹, D.H. Stork², H.K. Ticho², N. Toge⁶, M. Urban¹, R.F. van Daalen Wetters², G.J. Van Dalen³, R. van Tyen¹, H. Videau¹, M.R. Wayne², W.A. Wenzel¹, M. Yamauchi⁶, M.E. Zeller⁷, and W-M. Zhang⁴

¹Lawrence Berkeley Laboratory, University of California, Berkeley, CA. 94720

²University of California, Los Angeles, Ca 90024

³University of California, Riverside, Ca 92521

⁴Johns Hopkins University, Baltimore, MD 21218

⁵University of Massachusetts, Amherst, MA 01003

⁶University of Tokyo, Tokyo, JAPAN

⁷Yale University, New Haven, CT 06520

PACS numbers: 13.65.+i, 14.80 Dq

In a search for heavy charge $(4/3)e$ particles produced in e^+e^- collisions at a center of mass energy of 29 GeV, no candidate events were found in 22 pb^{-1} of data collected by the PEP-4 TPC. Upper limits are established on the inclusive cross section for the production of charge $(4/3)e$ particles in the mass range of 1-10 GeV/c^2 .

Fractional charge searches at accelerators have largely concentrated on the production of charge $(2/3)e$ and $(1/3)e$ particles. In the framework of Quantum Chromodynamics, the inability to observe free quarks in these searches is considered strong evidence that color triplets are confined. Recently, Slansky, Goldman and Shaw¹ have proposed a model in which color triplets are confined, yet some states with higher order representations of $SU(3)_{\text{color}}$ are unconfined. It is possible in this model to have a free charge $(4/3)e$ diquark (uu). In this letter we report the results of a search for the inclusive production of free charge $(4/3)e$ particles at PEP.

The Time Projection Chamber (TPC) is the central tracking chamber (Fig. 1) of the PEP-4 experiment. The detector system has been described in detail elsewhere². Track ionization formed in the TPC sensitive volume drifts in parallel electric and magnetic fields to arrays of proportional wires. The track position in z (parallel to the beam axis) is found from the arrival time of the ionization (drift velocity ≈ 5 cm/ μ sec) at the proportional wires. Up to 183 samples of ionization per track are collected on the wires. Track coordinates in the bending plane (x,y) are found from signals induced on 15 rows of segmented cathode pads beneath the wires.

The data presented in this letter were collected with the TPC operating at a gas gain of $\sim 10^3$, in an 8.5 atm mixture of 80% Ar and 20% CH_4 . The drift field was 75 kV/m; the magnetic field was 4 kG. The spatial resolution for tracks was 190 μ m in the bending plane (x,y) and 340 μ m in the drift direction (z). The chamber had a momentum resolution, dp/p , of $0.035p$ (p in GeV/c) for particles with momenta greater than 1 GeV/c.

The TPC identifies particle species by a simultaneous measurement of ionization and momentum. To avoid fluctuations in the measurement of track

ionization resulting from close collisions with atomic electrons, the measured value of the energy loss due to ionization, $\langle dE/dx \rangle$, was defined to be the mean of the lowest 65% of the wire pulse heights. Each wire was calibrated with an Fe^{55} source. The $\langle dE/dx \rangle$ resolution, measured with minimum ionizing pions ($\langle dE/dx \rangle = 12$ keV/cm), was 3.9% for tracks with at least 80 samples of ionization.

Figure 2 shows the curves of expected $\langle dE/dx \rangle$ as a function of momentum for pions, muons, protons, kaons, and electrons along with a subset of the data collected. Since the energy loss of a particle scales as its charge squared, one would observe relativistic charge $(4/3)e$ particles as a population in the upper right region of figure 2 at $(16/9)$ times the ionization of a relativistic unit charged particle.

In order to achieve good sensitivity, the search was performed in regions of $\langle dE/dx \rangle$ and momentum (shaded region in figure 2) which were not populated by known stable charge $1e$ particles. The low momentum boundary of the search region was set at the curve of expected $\langle dE/dx \rangle$ for a charge $1e$ particle with a mass of 1.8 GeV/c². The low dE/dx boundary was set at $1.2 \times \langle dE/dx \rangle$ for electrons in the same momentum range. The level at which electronic saturation begins to affect the $\langle dE/dx \rangle$ measurement is at $\langle dE/dx \rangle = 42$ keV/cm. This defines the high $\langle dE/dx \rangle$ boundary of the search region.

The data sample used in the search consisted of 22 pb⁻¹ of integrated luminosity at a center of mass energy of 29 GeV collected during the 1982-83 running cycle of PEP. The events were selected to be one photon annihilations of e^+e^- into hadrons, satisfying the following criteria. The number of charged tracks extrapolated to within 5 cm. in (x,y) and 10 cm

in z of the beam crossing had to be greater than or equal to five. The scalar sum of the visible charged momenta had to be at least 7.25 GeV/c. In each event the momentum imbalance between the forward and backward hemispheres was required to be less than 40% of the visible energy. We estimate the contamination of background events to be 8%, coming from cosmic rays, beam gas, $\tau\bar{\tau}$, and two photon events. In all, 7137 events satisfied the above criteria.

In order to retain good $\langle dE/dx \rangle$ resolution, candidates in the search region were selected only if they had at least 80 samples of ionization with no other ionization detected within 3 cm in z . We also required that candidates have momentum assignment errors, dp/p , estimated from the residuals to the fit, less than or equal to $0.1p$ (p in GeV/c).

After applying these cuts 50 candidate tracks remained. There were two major sources of backgrounds in this sample.

Deuterons and tritons have ionization curves lying partly in the search region. Some of these particles, produced from the interactions of primary tracks with the detector material, will pass the above cuts. Due to finite production angles, however, most of the track fits for these particles will not extrapolate through the beam crossing point. We reduced this background by requiring candidates to have an extrapolated distance of closest approach to the beam crossing point no greater than 5 cm in the bending plane (x,y) and no greater than 10 cm in the drift direction (z). Nine candidate tracks were rejected by imposing this requirement. Eight of these had values of $\langle dE/dx \rangle$ and momentum consistent with deuterons.

The second source of background came from unresolved pairs of nearby tracks. When two high momentum tracks are emitted close together (typically

with an angular separation of 25 mr or less) often only one is reconstructed. The ionization from these tracks adds and the reconstructed track appears to have a large value of $\langle dE/dx \rangle$ and a high momentum.

The fine spatial segmentation of the TPC allowed us to develop a series of cuts to reduce the second background. The cuts were tested on heavily ionizing tracks in hadronic events both to ensure that the requirements were not overly restrictive and to measure the inefficiency created by imposing them. We based these requirements on the expectation that overlapping tracks would produce excessively wide cathode pad and wire ionization clusters and/or nearby clusters not associated with the reconstructed track.

A wire hit is imaged on two to three cathode pads. In the three pad case (2-10 per track) we measure the width, σ , of the ionization cluster ($\langle \sigma \rangle = 3.9$ mm). From $P(\sigma)$, the probability for a cluster on an isolated track to have a width σ , we formed the quantity

$$L \equiv \frac{1}{n} \sum_{i=1}^n \ln (P(\sigma))$$

L is the log of the likelihood for a track to have n clusters with widths coming from the distribution $P(\sigma)$. We required candidates to have values of L consistent with a single track. L has no appreciable $\langle dE/dx \rangle$ dependence.

When only one of a pair of nearby tracks is reconstructed, the second track will often appear as additional ionization clusters near the first. Tracks were eliminated as candidates if they had an unusually large number of nearby pad or wire hits not associated with reconstructed tracks.

In some cases a second track or energetic delta ray may appear to merge with a track for part of its length and give approximately double

normal ionization only for the merged regions. The high density of wire information (up to 183 clusters per track) allowed us to reject these. We required candidates to have statistically uniform amounts of ionization along the track length. Partially merged tracks showed, on many contiguous wire hits, abnormally heavy ionization often correlated with large cluster widths. These were rejected.

These requirements rejected the remaining 41 candidate tracks.

The detection efficiency of the apparatus for charge $(4/3)e$ particles was determined in two steps. First the efficiency due to event topology, momentum distribution, fiducial volume etc. was found using a Monte Carlo calculation. The event generator was a modified LUND³ generator in which a pair of oppositely charged $(4/3)e$ particles were introduced into multihadron events. These events provided input to a detector simulation. The generated $(4/3)e$ particles were given momenta chosen from one of three distributions, $dN/dp \propto p^2/E$, $dN/dp = \text{constant}$, and $dN/dp \propto p^2/E * e^{-3.5E}$. Estimates of the inclusive momentum spectra for heavy fractionally charged particles and heavy hadrons produced in e^+e^- collisions tend to favor the first and second distributions above ^{4,5}, the third is presented as an extreme alternative.

In the detector simulation we assumed that charge $(4/3)e$ particles do not have larger than normal nuclear cross sections. We note that some estimates of nuclear cross sections for free fractionally charged particles⁴ give values 2-3 times that for protons. The amount of material between the beam crossing and the TPC active volume corresponds to 6% of a nuclear interaction length.

We found the efficiency of the cuts used to reject overlapping tracks by applying the same requirements to heavily ionizing charge 1e tracks. This efficiency was 72% with an estimated systematic uncertainty of 10%.

Our limits on $R_Q \equiv \sigma(e^+e^- \rightarrow QQX) / \sigma(e^+e^- \rightarrow \mu^+ \mu^-)$ are plotted in figure 3 along with the results of a similar search reported by JADE⁶.

To conclude, we have seen no evidence for the inclusive production of charge $(4/3)e$ particles in e^+e^- collisions at a 29 GeV center of mass energy. For momentum distributions similar to those for heavy hadrons^{4,5} we set an upper limit on the ratio $R_Q = \sigma(e^+e^- \rightarrow QQX) / \sigma(e^+e^- \rightarrow \mu^+ \mu^-)$ at less than .005 in the mass range 1-8 GeV/c².

We would like to thank the PEP staff for their outstanding work. This work was supported by the Department of Energy under contracts #DE-AC03-76SF00098, and #DE-AM03-76SF00034, the National Science Foundation, and the Joint Japan - U.S. collaboration in High Energy Physics.

REFERENCES

1. R. Slansky, T. Goldman, and G. Shaw, Phys. Rev. Lett. 47, 887 (1981)
See also ref. 4.
2. N. Hadley, Thesis, LBL-16116 (unpublished)
R. Fuzesy, N. Hadley, P. Robrish, LBL-16808 (to be published)
The PEP-4 collaboration, IEEE Trans. Nucl. Sci., NS-30, 63, 76, 162
(1983)
3. B. Andersson, and G. Gustafson, Z. Phys., C3, 223 (1980)
T. Sjöstrand, Comp. Phys. Comm. 27, 243 (1982)
4. A. De Rújula, R. Giles, and R. Jaffe, Phys. Rev. D 17, 285 (1978)
5. J. Bjorken, Phys. Rev. D 17, 171 (1978)
6. W. Bartel et al. (JADE collaboration), Z. Phys., C6, 295 (1980)

FIGURE CAPTIONS

Figure 1

Schematic view of the TPC. Track ionization formed in the sensitive volume drifts to arrays of proportional wires (only one set of wires shown). Position in the bending plane (x,y) is found from signals induced on cathode pads beneath the wires.


Figure 2

Curves of expected $\langle dE/dx \rangle$ vs. momentum for stable charge 1e particles. The points are from a subset of the data collected. The charge

(4/3)e search was performed using tracks found in the cross hatched region.


Figure 3

Limits on the inclusive production cross section for charge (4/3)e particles $R_0 \equiv \sigma(e^+e^- \rightarrow Q\bar{Q}X) / \sigma(e^+e^- \rightarrow \mu^+\mu^-)$ for the momentum distributions: A) $dN/dp \propto P^2/E$, B) $dN/dp = \text{constant}$, and C) $dN/dp \propto P^2/E e^{-3.5E}$. The solid curves are the limits from this search, the dashed curves are the limits from a search performed by the JADE collaboration (see ref. 6).




XBL 8310-754A

FIGURE 1



XBL 8310-753

FIGURE 2



XBL 8310-752

FIGURE 3

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720