Lawrence Berkeley National Laboratory

Recent Work

Title

AN EFFECTIVE-RANGE FORMULA FOR PHOTOPION PRODUCTION FROM PIONS

Permalink

<https://escholarship.org/uc/item/1kg8m0v1>

Author

Wong, How-Sen.

Publication Date

1960-06-14

UNIVERSITY OF
CALIFORNIA

Ernest O. Lawrence

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

For Pub. Phys. Rev. Ltrs

UCRL-9264
Limited Distribution

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

AN EFFECTIVE-RANGE FORMULA
FOR PHOTOPION PRODUCTION FROM PIONS

How-Sen Wong

June 14, 1960

AN EFFECTIVE-RANGE FORMULA FOR PHOTOPION PRODUCTION FROM PIONS*

How-Sen Wong

Lawrence Radiation Laboratory
University of California
Berkeley, California

June 14, 1960

In the Chew-Mandelstam¹ approach to the strong-interaction problem at low energies, the amplitude for $\gamma + \pi \rightarrow 2\pi$ plays a central role in any phenomena involving photons. Recently, Ball² has found that this process produces an additive correction to the formulas of Chew et al.³ for photopion production from nucleons. Compton scattering by pions and by nucleons also involves the $\gamma + \pi \rightarrow 2\pi$ reaction. A simple and reasonably accurate formula for the $\gamma + \pi \rightarrow 2\pi$ amplitude which can be used in any of these applications is given here.

We assume that the simple invariant transition amplitude, $M(s, t, u)$ has the Mandelstam representation⁴

$$M(s, t, u) = \frac{1}{\pi^2} \int_4^\infty \int_4^\infty \rho(x, y) \left[\frac{1}{(x-s)(y-t)} + \frac{1}{(x+t)(y-u)} + \frac{1}{(x-u)(y-s)} \right] dx dy. \quad (1)$$

It is defined through the relation,

$$T_{fi} = \frac{i(2\pi)^4 \delta^4(K + p_1 - p_2 - p_3) \sum \left[\left(-\frac{i}{\sqrt{2}} \right) \epsilon_{\alpha\beta\gamma} \epsilon_{\lambda\sigma\mu\nu} (p_1)_\lambda (p_2)_\sigma K_\mu \epsilon_\nu \right] M(s, t, u)}{(16 p_{10} p_{20} p_{30} K)^{1/2}} \quad (2)$$

*A preliminary account of this work was given at the 1959 Thanksgiving meeting of the American Physical Society, November 27-28, 1959. This work was performed under the auspices of the U.S. Atomic Energy Commission.

where α , β , and γ are the isotopic indices of the pions, and $\epsilon_{\alpha\beta\gamma}$ and $\epsilon_{\lambda\sigma\mu\nu}$ are the conventional antisymmetric tensors of third and fourth rank, respectively. Here s, t , and u are defined⁵ as (see Fig. 1)

$$\begin{aligned} s &= -(K + p_1)^2, \\ t &= -(K - p_2)^2, \end{aligned} \quad (3)$$

and

$$u = -(K - p_3)^2,$$

and are related by the condition

$$s + t + u = 3.$$

When the photon K and meson p_1 are the incoming particles, the variables s, t , and u are related to barycentric system variables

$$\begin{aligned} s &= 4(1 + p^2), \\ t &= 1 - 2kE + 2kp \cos \theta, \end{aligned}$$

and

$$u = 1 - 2kE - 2kp \cos \theta,$$

where p and E are the magnitudes of the outgoing pion momenta and energy, respectively, k is the energy of the photon, and we have $\cos \theta = (\vec{p} \cdot \vec{k})/pk$. The formula for the differential cross section in the barycentric system is

$$\frac{d\sigma}{d\Omega} = \sum \frac{p}{k} \left| \frac{1}{8E} \frac{T_{fi}}{2\pi} \right|^2.$$

MU-20481

Fig. 1. Diagram of $\gamma + \pi \rightarrow 2\pi$.

From the Mandelstam representation, we can locate the singularities of the partial-wave amplitudes, defined through the relation

$$M(s, z) = \sum_{\text{odd } \ell} M_{\ell}(s) P'_{\ell}(z),$$

where $z = \cos \theta$. The M_{ℓ}' s are analytic in the whole complex s plane except for cuts on the real axis. The right-hand cut runs from 4, the physical threshold for two pions, to ∞ ; the left cut runs from 0 to $-\infty$. Using these analytic properties, the unitary condition, and the Omnès⁶-Frazer-Fulco⁷ method, we find

$$M_{\ell}(s) = \frac{1}{\pi D_{\ell}(s)} \int_{-\infty}^0 \frac{D_{\ell}(s') \text{Im } M_{\ell}(s')}{s' - s} ds', \quad (4)$$

where $D_{\ell}(s)$ is the π - π denominator function introduced by Chew and Mandelstam.¹

If we assume that for $s \geq 4$, we have $\text{Im } M \approx \text{Im } M_1$, we can show that

$$M(s, t, u) \approx \frac{1}{\pi} \int_4^{\infty} \text{Im } M_1(x) \left[\frac{1}{x-s} + \frac{1}{x-t} + \frac{1}{x-u} \right] dx. \quad (5)$$

We see that all partial waves for $\ell > 1$ can be obtained once M_1 is known. Formula (5) is the basis of an independent calculation that has been carried out by Gourdin and Martin.⁸

Using crossing symmetry, we can obtain a formula for $\text{Im } M_1(s)$ for $s < 0$ in terms of $\text{Im } M_1(s)$ for $s > 4$. We do not give this formula here because the effect-range approach does not require such detailed

information. In any event, it is clear that for $\ell = 1$, the integral of Eq. (4) is homogeneous, so that its solution is not unique, at least with respect to a multiplicative factor. Therefore an arbitrary parameter must be introduced. In the calculation below, we shall fix this parameter Λ as the value of $M_1(s)$ at $s = 1$. At present, we do not know how to relate Λ to fundamental constants, and for the time being it must be determined from experiment.

To proceed further, we need the denominator function, D_1 , for p-wave pion-pion scattering. The pion-pion calculations of Chew and Mandelstam have not yet reached a conclusive stage but these authors have given an approximate form for D_1 which corresponds to the replacement of the unphysical branch cut in the pion-pion amplitude by a finite number of poles.⁹ Further, they have shown that two poles lead to an accurate approximation in the physical region up to $s \sim 40$. Once the parameters of the two-pole formula have been determined, it will be a straightforward problem to incorporate the information into the amplitude $\gamma + \pi \rightarrow 2\pi$. The determination of the pion-pion parameters is still in progress,¹⁰ but we outline here, for future use, the form of solution of our problem that corresponds to the pole approximation of Chew and Mandelstam.⁹ We shall illustrate the method with the one-pole pion-pion amplitude, for which parameters have been given by Frazer and Fulco.⁷

We propose to replace the left-hand cut of the amplitude M_1 by poles with appropriate positions and residues. The "n-pole formula" thus obtained contains $2n$ parameters. One of these may be identified with Λ ; the rest will be determined from crossing relations.

By representing $\text{Im } M_1$ on the left-hand cut by one or two delta functions, we obtain from Eq. (4) the one-pole formula

$$M_1(s) = \frac{\Lambda(1+a)D_1(1)}{(s+a)D_1(s)} \quad (6)$$

or the two-pole formula

$$M_1(s) = \frac{\Lambda D_1(1)}{D_1(s)} \left[\frac{D_1(-b)}{s+b} + \frac{\Lambda_1 D_1(-c)}{s+c} \right] \left[\frac{D_1(-b)}{1+b} + \frac{\Lambda_1 D_1(-c)}{1+c} \right]^{-1} \quad (7)$$

where Λ and Λ_1 are real, and a , b , and c are real and positive. The parameters Λ_1 , a , b , and c may be determined from the following crossing relations which are derived from Eq. (5):

$$M_1(1) - \frac{3}{2} M_1^L(1) = 0 \quad (8a)$$

$$\frac{\partial}{\partial s} M_1(1) = 0 \quad (8b)$$

$$\frac{\partial^2}{\partial s^2} M_1(1) - 6 \frac{\partial^2}{\partial s^2} M_1^L(1) = 0, \quad (8c)$$

where

$$M_1^L(s) = \frac{1}{\pi} \int_{-\infty}^0 \frac{\text{Im } M_1(s') ds'}{s - s'}$$

$$= \frac{\Lambda (1+a) D_1(1)}{(s+a) D_1(-a)}$$

for the one-pole formula, and

$$= \Lambda D_1(1) \left[\frac{1}{s+b} + \frac{\Lambda_1}{s+c} \right] \left[\frac{D_1(-b)}{1+b} + \frac{\Lambda_1 D_1(-c)}{1+c} \right]^{-1}$$

for the two-pole formula.

If the Frazer-Fulco one-pole $D_1(s)$ function is used for a resonance at $s = 10$, we find $a \approx 5.7$ from the crossing condition (8a). The parameters Λ_1 , b , and c in the two-pole formula may be determined from (8a) to (8c). It turns out that no choice of Λ_1 , b , and c can satisfy all three conditions (8a) to (8c) if the Frazer-Fulco one-pole formula is used. However, the first two conditions of Eqs. (9) can be satisfied with the pair of poles in the range between 0 and -4.93. The p-wave amplitude in the physical region not only is insensitive to the positions of the two poles so long as (8a) and (8b) are satisfied, but also is not much different from the one-pole solutions, which satisfies (8a) alone. Thus we may be confident of the accuracy of the two-pole results once the parameters of the two-pole π - π scattering formula are known. The one-pole formula already obtained is probably adequate for most purposes.

We can estimate the order of magnitude of Λ by considering the decay of neutral pions, assuming that the $\gamma + \pi \rightarrow 2\pi$ reaction plays a prominent role in the intermediate state. We find $|\Lambda| \approx 1.3e$ ($e^2 = 1/137$) if the π^0 lifetime is 4×10^{-16} sec.¹¹ Our calculation is based on the assumption that only the least massive (two-pion) intermediate state contributes to the dispersion integral, but there is no good reason to

neglect the 3π contribution, especially if there is a resonance or a bound state at roughly the same energy as the two-pion resonance.¹² A better estimate of Λ may come from photopion production on nucleons, where Ball has shown that $\gamma + \pi \rightarrow 2\pi$ makes an important and characteristic contribution.²

The author wishes to thank Professor G. F. Chew for suggesting this investigation and for his guidance. He also wishes to thank Dr. D. Y. Wong and Dr. A. Martin for helpful discussions.

REFERENCES

1. G.F. Chew and S. Mandelstam, Theory of the Low-Energy Pion-Pion Interaction, UCRL-8728, April 15, 1959 and Phys. Rev. (to be published).
2. J.S. Ball, The Application of the Mandelstam Representation to Photoproduction of Pions from Nucleons, UCRL-9172, April 11, 1960, and J. S. Ball, Phys. Rev. Letters (following letter).
3. G.F. Chew, M. L. Goldberger, F.E. Low, and Y. Nambu, Phys. Rev. 106, 1337 (1957).
4. S. Mandelstam, Phys. Rev. 112, 1344 (1958); *ibid.* 115, 1741 and 1752 (1959).
5. We use the fundamental metric tensor such that $p_1 \cdot p_2 = \vec{p}_1 \cdot \vec{p}_2 - p_{10}p_{20}$. Units are used in which $\hbar = c = 1$ and $\mu = 1$, where μ is the mass of the pion.
6. R. Omnès, Nuovo cimento 8, 316 (1958).
7. W.R. Frazer and J.R. Fulco, Phys. Rev. Letters 2, 365 (1959); *ibid.*, Phys. Rev. 117, 1909 (1960).
8. M. Gourdin and A. Martin, Photoproduction of Pions on Pions, Preprint CERN 7687 TH/67.
9. G.F. Chew and S. Mandelstam, Theory of the Low-Energy Pion-Pion Interaction, Part II, UCRL-9126, March 24, 1960.
10. J.S. Ball and D.Y. Wong, Lawrence Radiation Laboratory, private communication.
11. G. Harris, J. Orear, and S. Taylor, Phys. Rev. 106, 327 (1957).
12. G.F. Chew, Phys. Rev. Letter 4, 142 (1960). The author wishes to thank Professor R. J. Eden of Cambridge University for indicating this point to him.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.