

# UCSF

## UC San Francisco Previously Published Works

### Title

Moderate-intensity physical activity ameliorates the breast cancer risk in diabetic women.

### Permalink

<https://escholarship.org/uc/item/1ks7j14t>

### Journal

Diabetes reviews (Alexandria, Va.), 35(12)

### Authors

Torres-Mejía, Gabriela  
Angeles-Llerenas, Angélica  
Ortega-Olvera, Carolina  
et al.

### Publication Date

2012-12-01

### DOI

10.2337/dc11-2362

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

# Moderate-Intensity Physical Activity Ameliorates the Breast Cancer Risk in Diabetic Women

GABRIELA TORRES-MEJÍA, PHD<sup>1</sup>  
 ANGÉLICA ANGELES-LLERENAS, MSC<sup>1</sup>  
 CAROLINA ORTEGA-OLVERA, MSC<sup>1</sup>  
 EDUARDO LAZCANO-PONCE, PHD<sup>1</sup>  
 ELAD ZIV, PHD<sup>2,3</sup>

JOSÉ PULIDO-RODRÍGUEZ, MD<sup>4</sup>  
 MANUEL DE JESÚS GARCÍA-SOLÍS, MD<sup>4</sup>  
 EFRÉN MURILLO-ZAMORA, MSC<sup>1,5</sup>  
 JULIA VÁZQUEZ-LARA, MD<sup>6</sup>  
 ISABELLE ROMIEU, PHD<sup>7</sup>

**OBJECTIVE**—To evaluate the association between self-reported diabetes and the risk of breast cancer (BC) and its interaction with moderate-intensity physical activity in pre- and postmenopausal Mexican women.

**RESEARCH DESIGN AND METHODS**—A population-based case-control study was conducted using 1,000 incident case subjects and 1,074 control subjects. Blood samples and information on health, diet, physical activity, and anthropometric measurements were obtained.

**RESULTS**—The association between diabetes and BC risk decreased with increasing tertiles of moderate-intensity physical activity (odds ratio [OR] = 4.9 [95% CI 2.3–10.8]; 3.0 [1.3–6.9]; and 1.0 [0.1–9.2], respectively, for each tertile) (test for interaction = 0.04). Compared with the women in the lowest tertiles, increased risk was observed in those premenopausal women with the highest serum C-peptide, IGF-1, and IGF-1 binding protein 3 levels.

**CONCLUSIONS**—Moderate-intensity physical activity can substantially ameliorate the increased BC risk in diabetic women.

*Diabetes Care* 35:2500–2502, 2012

In Mexico, the prevalence of type 2 diabetes has shown a gradual rise in recent decades (from 6.7% of adults  $\geq 20$  years of age in 1993 to 14.4% in 2006). According to the 2006 National Health and Nutrition Survey (Encuesta Nacional de Salud y Nutrición 2006), its prevalence in women increases with age: 14.2% in women aged 50–59 years and 21.3% in those aged 60–69 years (1).

It has been suggested that diabetes increases the risk of breast cancer (BC) (2); however, physical activity appears to reduce the risk of BC (3) by increasing insulin sensitivity (4). Therefore, the aim of this study was to evaluate the effect of diabetes on the risk of BC in pre- and

postmenopausal Mexican women, the interaction with moderate-intensity physical activity, and the association of several biomarkers, including IGF-1, its binding protein 3 (IGFBP3), and C-peptide with the risk of BC in a subsample of premenopausal women.

## RESEARCH DESIGN AND METHODS

A multicenter, population-based, case-control study was conducted by the National Institute of Public Health in Mexico City, Monterrey, and Veracruz (including their metropolitan areas) from January 2004 through December 2007. Detailed methods and inclusion and exclusion criteria have been described

elsewhere (3). In brief, 1,000 incident BC case subjects and 1,074 control subjects aged 35–69 years were selected. The case subjects were enrolled in the study soon after being informed about of their diagnoses (median = 3 days), were histologically confirmed, and had not been previously treated with chemotherapy or antihormonal therapy.

The control subjects were frequency matched to the case subjects by 5-year age categories, membership in a participating health care institution, and place of residence. They were enrolled at approximately the same time as the BC diagnoses. The control group was selected using a multistage probabilistic design.

After informed consent was reviewed and signed by all of the participants, information on health status, reproductive history, dietary intake, and physical activity was obtained through a questionnaire administered by trained nurses; fasting blood samples were obtained from all of the case and control subjects at recruitment. Information on diet and physical activity was obtained during a typical week prior to the first BC symptoms in order to reduce the possibility of reverse causation bias. Physical activity was estimated using a 7-day recall questionnaire that estimated an individual's time spent performing physical activities (sleep and light-, moderate-, and vigorous-intensity physical activity) (5).

A history of diabetes was self-reported using the following question: Have you ever been diagnosed by a qualified physician as having diabetes (yes/no)? Self-reported diabetes has been used as an accurate disease measure (6,7).

Diet during the year prior to the onset of symptoms was assessed using a 104-item semiquantitative food frequency questionnaire by Willet (8) that had been adapted to the Mexican population and validated in Mexico City (9). Further details have been provided in a previous publication (3).

A random subsample of premenopausal women (244 case subjects and 320 control subjects) was selected for measuring: C-peptide, IGF-1, and IGFBP3.

From the <sup>1</sup>National Institute of Public Health, Center for Population Health Research, Cuernavaca, Morelos, Mexico; the <sup>2</sup>Department of Medicine, Division of General Internal Medicine, Institute for Human Genetics, San Francisco, California; the <sup>3</sup>Department of Epidemiology and Biostatistics, Helen Diller Family Comprehensive Cancer Center, San Francisco, California; the <sup>4</sup>Metropolitan Hospital "Dr. Bernardo Sepúlveda," Nuevo León, Mexico; <sup>5</sup>Hospital No. 4, IMSS, Tecomán, Colima, Mexico; <sup>6</sup>Hospital No. 23 of Gynecology, IMSS, Monterrey, Nuevo León, Mexico; and <sup>7</sup>Nutrition and Metabolism, International Agency for Research on Cancer, Lyon, France.

Corresponding author: Angélica Angeles-Llerenas, aangelica@insp.mx.

Received 5 December 2011 and accepted 25 May 2012.

DOI: 10.2337/dc11-2362

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The blood samples were frozen at temperatures between  $-20$  and  $-70^{\circ}\text{C}$  and processed no later than 4 weeks after the blood was collected. The C-peptide, IGF-1, and IGFBP3 levels were determined using radioimmunoanalysis at Laboratorios Clínicos de Puebla, México (ISO 9001:2008).

This study was approved by the Ethics Committee of the National Institute of Public Health and the local committees from participating hospitals.

### Statistical analyses

To evaluate the effect of moderate-intensity physical activity on BC risk, conditional logistic multiple-regression models were used to estimate odds ratios (ORs) and 95% CI (10). An interaction term

(diabetes and moderate-intensity physical activity) was added to the model in postmenopausal women, and the effect of biomarkers on BC risk was assessed in premenopausal women. We exclude anyone diagnosed before 25 years of age to reduce the probability of including type 1 diabetes women (one pre- and four postmenopausal women). The models were a priori adjusted for 5-year age categories, health institution, site, well-known potential confounders, and European ancestry. Statistical analysis was performed using STATA version 10 (Stata Corp.).

**RESULTS**—Characteristics of study population have been described elsewhere (3). Overall, the women who reported a diabetes diagnosis had an increased risk of BC

(OR = 2.1 [95% CI 1.5–2.8]). This association was statistically significant for the postmenopausal women (2.7 [1.7–4.1]) but not for the premenopausal women (1.4 [0.7–2.6]) (Table 1). Additionally, the effect of diabetes on BC risk in the postmenopausal women decreased with increasing tertiles of moderate-intensity physical activity: 4.9 (2.3–10.8) for the lowest tertile ( $\leq 5$  h/week); 3.0 (1.3–6.9) for the second tertile (5.1 to  $<19$  h/week); and 1.0 (0.1–9.2) for the highest tertile ( $>19$  h/week). The *P* value for the interaction term comparing the lowest two tertiles combined and the highest ( $\leq 19$  vs.  $>19$  h/week) was 0.04.

In the premenopausal subsample, a statistically significant association with BC risk was observed when comparing the highest C-peptide tertile versus the

**Table 1—Diabetes and BC risk in Mexican women residing in Mexico City, Monterrey, and Veracruz. Risk Factors for Breast Cancer in Mexico: Mammographic Patterns, Peptide C, and Growth Factors, a Multi-Center Study, 2004–2007**

	Premenopausal*				Postmenopausal†			
	Case (n)	Control (n)	OR	95% CI	Case (n)	Control (n)	OR	95% CI
<b>Diabetes</b>								
No	264	346	1.0		273	361	1.0	
Yes	26	33	1.4	0.7–2.6	101	85	2.7	1.7–4.1
<b>Moderate-intensity physical activity (hours/week)</b>								
$\leq 5$	122	136	1.0		197	160	1.0	
5.1–19	110	110	1.2	0.8–1.8	142	162	0.7	0.5–1.1
$>19$	58	133	0.5	0.3–0.8	35	124	0.3	0.2–0.4
<i>P</i> for trend			<i>P</i> = 0.002				<i>P</i> < 0.001	
<b>Effect on diabetes of moderate-intensity physical activity (hours/week)‡</b>								
$\leq 5$					197	160	4.9	2.3–10.8
5.1–19					142	162	3.0	1.3–6.9
$>19$					35	124	1.0	0.1–9.2
<b>Biomarkers§</b>								
<b>C-peptide (ng/mL)</b>								
$\leq 1.2$	42	87	1.0					
1.3–2	55	94	2.1	1.1–3.8				
2.1–7.2	91	70	6.5	3.3–12.5				
<b>IGF-1 (ng/mL)</b>								
$\leq 123.8$	31	76	1.0					
123.9–206.5	55	97	1.5	0.8–2.8				
$\geq 206.5$	102	78	2.7	1.5–5.0				
<b>IGFBP3 (ng/mL)</b>								
$\leq 2676$	17	86	1.0					
2677–3451	37	84	2.0	1.0–4.1				
$\geq 3452$	134	81	9.6	4.7–19.6				

\*In a multiple model a priori adjusted for 5-year age categories, health care institution, and site, the model was also adjusted for: European ancestry, socioeconomic status (low, medium, and high), BMI (body weight in kilograms/height in centimeters squared), waist/hip ratio, height in centimeters, parity (number of children born alive), history of BC in first-degree relatives (grandmother, mother, or sisters; yes/no), personal history of benign breast disease (yes/no), smoking at least 100 cigarettes (yes/no), occasional intake of  $>1$  alcohol drinks per month during  $\geq 1$  year (yes/no), ever using oral contraceptives (ever/never), age at menarche (years), moderate-intensity physical activity (hours/week, in tertiles), daily energy intake (kcal) using the residuals method, and daily carbohydrate intake, lipid intake, and protein intake in grams. †In a multiple model adjusted for the same variables used for the premenopausal women, ever using hormones during menopause (yes/no) and age at menopause (years). ‡In the postmenopausal women, the *P* value for interaction was 0.044 when comparing the two lowest tertiles combined to the highest tertile; the *P* value for the interaction was 0.07 when comparing all of the tertiles individually. §In a multiple model, with each biomarker adjusted as in the general premenopausal model.

lowest (OR = 6.5 [95% CI 3.3–12.5]) and the highest IGF-1 and IGFBP3 tertiles versus the lowest tertiles (Table 1).

**CONCLUSIONS**—Our results suggest that moderate-intensity physical activity can substantially ameliorate the increased BC risk observed in diabetic women. These results are consistent with the results of other studies involving Hispanic women residing in the U.S. These studies have observed a protective effect against BC in diabetic women who exercise regularly, possibly through a reduced insulin resistance (11). Data from Mexico, Chile, Peru, Argentina, Brazil, and the multicountry Pan American Health Organization study showed a prevalence of adult physical inactivity from 50–91% (12–14). Given that higher serum levels of C-peptide, IGF-1, and IGFBP3 have been found to increase BC risk, further studies are required to address this issue. Our results suggest that IGF-1 and C-peptide offer a potential biological mechanism that explains the increased BC risk associated with diabetes. There is an urgent need to detect and treat diabetes early and promote physical activity to control weight and improve women's health by improving public policy. Research on chronic disease prevention is a top research priority.

**Acknowledgments**—This study was supported by Consejo Nacional de Ciencia y

Tecnología, the Mexican Research Program (Grant Salud 2002-CO1-7462 to G.T.-M.), and the National Institutes of Health (Grant R01-CA-120120 to E.Z.).

No potential conflicts of interest relevant to this article were reported.

G.T.-M., A.A.-L., C.O.-O., E.L.-P., E.Z., J.P.-R., M.d.J.G.-S., E.M.-Z., J.V.-L., and I.R. read and approved the final version of the manuscript. A.A.-L., C.O.-O., and E.M.-Z. analyzed the data and wrote the draft of the manuscript. E.L.-P., J.P.-R., and M.d.J.G.-S. performed critical review of the manuscript. E.Z., J.V.-L., and I.R. contributed to the discussion of the data and performed critical review of the manuscript. G.T.-M. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

#### References

1. Villalpando S, de la Cruz V, Rojas R, et al. Prevalence and distribution of type 2 diabetes mellitus in Mexican adult population: a probabilistic survey. *Salud Publica Mex* 2010;52(Suppl. 1):S19–S26
2. Larsson SC, Mantzoros CS, Wolk A. Diabetes mellitus and risk of breast cancer: a meta-analysis. *Int J Cancer* 2007;121:856–862
3. Angeles-Llerenas A, Ortega-Olvera C, Pérez-Rodríguez E, et al. Moderate physical activity and breast cancer risk: the effect of menopausal status. *Cancer Causes Control* 2010;21:577–586
4. Colberg SR, Grieco CR. Exercise in the treatment and prevention of diabetes. *Curr Sports Med Rep* 2009;8:169–175
5. Sallis JF, Haskell WL, Wood PD, et al. Physical activity assessment methodology in the Five-City Project. *Am J Epidemiol* 1985;121:91–106
6. Oksanen T, Kivimäki M, Pentti J, Virtanen M, Klaukka T, Vahtera J. Self-report as an indicator of incident disease. *Ann Epidemiol* 2010;20:547–554
7. El Fakiri F, Bruijnzeels MA, Hoes AW. No evidence for marked ethnic differences in accuracy of self-reported diabetes, hypertension, and hypercholesterolemia. *J Clin Epidemiol* 2007;60:1271–1279
8. Willet W. *Nutritional Epidemiology*. 2nd ed. New York, Oxford University Press, 1998
9. Hernández-Avila M, Romieu I, Parra S, Hernández-Avila J, Madrigal H, Willett W. Validity and reproducibility of a food frequency questionnaire to assess dietary intake of women living in Mexico City. *Salud Publica Mex* 1998;40:133–140
10. Hosmer DW, Lemeshow S. Model-building strategies and methods for logistic regression. In *Applied Logistic Regression*. 2nd ed. Canada, John Wiley & Sons, 2000
11. Sanderson M, Peltz G, Perez A, et al. Diabetes, physical activity and breast cancer among Hispanic women. *Cancer Epidemiol* 2010;34:556–561
12. Hernández B, de Haene J, Barquera S, et al. [Factors associated with physical activity among Mexican women of childbearing age]. *Rev Panam Salud Publica* 2003;14:235–245 [in Spanish]
13. Pratt M, Jacoby ER, Neiman A. Promoting physical activity in the Americas. *Food Nutr Bull* 2004;25:183–193
14. Jacoby ER. PAHO regional consultation of the Americas on diet, physical activity and health. *Food Nutr Bull* 2004;25:172–174