Lawrence Berkeley National Laboratory

Recent Work

Title

THE DISPROPORTIONATION OF DIGERMANE IN LIQUID AMMONIA

Permalink

<https://escholarship.org/uc/item/1n08c540>

Authors

Dreyfuss, Robert M.

Jolly, William L.

Publication Date

1968-05-01

cy. 2

RECEIVED
LAWRENCE
RADIATION LABORATORY

OCT 31 1968

LIBRARY AND
DOCUMENTS SECTION

University of California

Ernest O. Lawrence
Radiation Laboratory

THE DISPROPORTIONATION OF DIGERMANE IN LIQUID AMMONIA

Robert M. Dreyfuss and William L. Jolly

May, 1968

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

UCRL-18518
cy 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Inorganic Chemistry
as a Note

UCRL-18518
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

THE DISPROPORTIONATION OF DIGERMANE IN LIQUID AMMONIA

Robert M. Dreyfuss and William L. Jolly

May, 1968

Contribution from the Department of Chemistry of the University
of California and the Inorganic Materials Research
Division of the Lawrence Radiation Laboratory, Berkeley, California 94720

The Disproportionation of Digermene in Liquid Ammonia

By Robert M. Dreyfuss and William L. Jolly

While attempting to measure the acid dissociation constant of digermene in liquid ammonia, we have observed that digermene decomposes in liquid ammonia solution to give germane and a solid germanium hydride of variable composition. Similar base-catalyzed decompositions of digermene have been observed by Bornhorst and Ring,¹ but these workers did not investigate the nature of the solid hydride formed. In view of Glarum and Kraus's earlier report² of the formation of GeH_2 by the reaction of bromobenzene with sodium germyl in ammonia, we decided to determine if the same substance can be prepared by the ammonia-catalyzed disproportionation of digermene.

Experimental Section

Standard greaseless vacuum line techniques were employed. Digermene was prepared according to the method of Jolly and Drake.³ The identity

-2-

and purity of germane and digermane were determined by infrared spectrometry^{4,5} with a Perkin Elmer Model 137 Infracord spectrometer, by mass spectrometry with a Consolidated Engineering Corp. mass spectrometer, Model 21-620, and by vapor pressure measurements.³ Ammonia was dried over sodium before its use.

Reactions were carried out in a vessel equipped with two break-off seals. A measured amount of digermane and approximately 0.5 ml of liquid ammonia were distilled into the vessel at liquid nitrogen temperature. The vessel was then sealed and placed in a bath of appropriate temperature. When a low-temperature bath was used, care was taken to keep the entire vessel below the level of the bath.

After reaction had proceeded, the vessel was attached to the vacuum line, and one of the seals was opened, keeping the vessel in the low temperature bath. The volatile contents were then Toepler-pumped through a liquid nitrogen trap until all volatiles had distilled over. A trace of hydrogen could usually be measured in the buret of the Toepler pump. The vessel was then sealed off again and temporarily set aside. The volatiles were then separated. In runs in which all the digermane had decomposed, it was possible to separate germane and ammonia by means of a -160° trap. However, when digermane was present the ammonia could not be separated by trap-to-trap distillation; it was removed by distilling the ternary mixture very slowly through a trap of anhydrous magnesium perchlorate at 0° . The germane and digermane were then separated by fractional condensation in traps at -112° and -196° ; the amount of each was determined by pressure-volume techniques.

-3-

The residue in the reaction vessel was then pyrolyzed at 400° for two hours. The hydrogen (identified by mass spectrometry) was then Toepler-pumped through the second break-off seal and a -196° trap to remove any ammonia that might have adhered to the solid. The hydrogen was measured and the germanium mirror was either discarded or dissolved in basic hydrogen peroxide and titrated with a pH-meter according to the method of Tchakirian.⁶

Results

The results are presented in Table I. The adequacy of the quantitative measurements is attested by the fact that the hydrogen in the germane produced and that formed in the pyrolysis of the GeH_x accounts for the hydrogen in the consumed digermane with an average discrepancy of $\pm 1.2\%$. And in the four runs where a germanium balance could be calculated, the average discrepancy was $\pm 3.5\%$.

In the -63° runs, the solution was initially colorless, but on standing the solution turned yellow and a yellow precipitate formed. In the -45° run, a yellow precipitate formed quickly. In the room-temperature runs, a blood red precipitate formed immediately.

Discussion

The results indicate that, in liquid ammonia solution, digermane disproportionates as follows:

-4-

Table I.
Quantitative Data for Ge_2H_6 Disproportionations
(All quantities in mmoles)

Temp. °C	Time hr.	$\tilde{A}$	$\tilde{B}$	$\tilde{C}$	$\tilde{D}$	$\tilde{E}$	Calcd. values of x	
		Ge_2H_6 in	Ge_2H_6 out	GeH_4 formed	H_2 from GeH_x pyrolysis	Ge in GeH_x	$2\tilde{D}/\tilde{E}$	$\frac{2\tilde{D}}{2(\tilde{A}-\tilde{B})-\tilde{C}}$
-63	96	0.668	0.273	0.481	0.253	0.264	1.92	1.64 ^a
-63	96	.553	.323	.260	.149	.180	1.66	1.49 ^a
-63	25	.383	.308	.064	.077	.077	2.00	1.79 ^a
-63	25	.536	.479	.0696	.0695	.069	2.01	3.16 ^a
-45	37	.200	0	.220	.169	-	-	1.88
23 ^b	24	.188	0	.244	.0746	-	-	1.13
23	20	.244	0	.334	.0676	-	-	0.88
23 ^c	14	.218	0	.291	.0789	-	-	1.09
23	13	.247	0	.321	.0800	-	-	0.92
23	0.25	.152	.114	.0417	.0316	-	-	1.86 ^a

^a These values are relatively inaccurate because they require the difficult determination of the unreacted Ge_2H_6 ($\tilde{B}$).

^b Ammonia soln. 1 M in NH_4NO_3

^c $\text{NH}_3:\text{Ge}_2\text{H}_6 = 1:1$

-5-

At temperatures above -63° (and even at that temperature on standing), the GeH_2 decomposes to germane and a yellow or red precipitate of GeH_x ($x < 2$). The higher the temperature, the more rapidly the GeH_2 decomposes, the lower the value of x and the darker the color of the precipitated GeH_x . Although Glarum and Kraus² reported similar properties for GeH_2 prepared from bromobenzene and sodium germyl, their substance was more stable inasmuch as its solution in ammonia did not become colored on standing for 24 hours, and it could be isolated for a short time as a white solid at -33° .

In the hope that the solution at -63° contained a soluble GeH_2 species, the proton magnetic resonance spectrum of a 0.5 M Ge_2H_6 solution was obtained. The spectrum showed peaks corresponding to germane, digermane and a series of three smaller peaks from 8 to 25 cycles upfield from germane, perhaps corresponding to a lower hydride. As the solution decomposed, the digermane peak decreased in amplitude, the germane peak increased, and the little peaks began to merge and broaden. By the time the digermane peak had disappeared, the little peaks were gone. Attempts to get better resolution by increasing the digermane concentration were unsuccessful, probably because of limited solubility of the lower hydride.

Efforts to effect reaction between GeH_2 and acetylene or diphenylacetylene at -63° , in analogy with the high-temperature reactions of GeI_2 ,⁷ were unsuccessful.

We have observed that no reaction occurs when both the digermane and the ammonia are completely in the gas phase; however, reaction proceeds in the presence of even traces of liquid ammonia. The basic catalyst

-6-

is ammonia itself and not the amide ion because the reaction goes without noticeable change in 1 M ammonium nitrate liquid ammonia solution. Apparently, relatively weak bases cause the disproportionation of digermane. In addition to ammonia and those bases shown to be effective by Bornhorst and Ring,¹ aqueous sodium hydroxide causes the disproportionation.⁸

Acknowledgment. - This work was supported by the United States Atomic Energy Commission.

References

- (1) W. R. Bornhorst and M. A. Ring, Inorg. Chem., 7, 1009 (1968).
- (2) S. N. Glarum, and C. A. Kraus, J. Am. Chem. Soc., 72, 5398 (1950).
- (3) W. L. Jolly and J. E. Drake, Inorg. Syntheses, 7, 34 (1963).
- (4) For GeH_4 : J. W. Straley, C. H. Tindal, and H. H. Nielsen, Phys. Rev., 62, 161 (1942).
- (5) For Ge_2H_6 : D. A. Dows and R. M. Hexter, J. Chem. Phys., 24, 1029 (1956).
- (6) R. Belcher and C. L. Wilson, New Methods in Analytical Chemistry, Chapman and Hall, London 2nd ed., 1956, p. 232.
- (7) M. E. Volpin, et al., Izvest, Akad. Nauk SSSR Ser. Chim., 1963, 2067.
- (8) L. M. Dennis, R. B. Corey and R. W. Moore, J. Am. Chem. Soc., 46, 657 (1924), and unreported observations of R. Dreyfuss.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.