Lawrence Berkeley National Laboratory

Recent Work

Title

MASS TRANSFER TO THE REAR OF A SPHERE IN STOKES FLOW

Permalink

<https://escholarship.org/uc/item/1n16c049>

Authors

Sin, Ping Huei
Newman, John.

Publication Date

1967-02-01

UCRL-17375

University of California
Ernest O. Lawrence
Radiation Laboratory

MASS TRANSFER TO THE REAR OF A SPHERE IN STOKES FLOW

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to International Journal of
Heat and Mass Transfer

UCRL-17375
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

MASS TRANSFER TO THE REAR OF A SPHERE IN STOKES FLOW

Ping Huei Sih and John Newman

February 1967

Mass Transfer to the Rear of a Sphere in Stokes Flow

Ping Huei Sih and John Newman
Inorganic Materials Research Division,
Lawrence Radiation Laboratory, and
Department of Chemical Engineering,
University of California, Berkeley

February, 1967

Abstract

Mass transfer to a sphere, including the rear region, in Stokes flow and at large Péclet numbers, is investigated. By the singular-perturbation technique, six distinct regions of different mass-transfer mechanisms are found. One of these regions, the diffusion layer, has already been solved by the boundary-layer method. Another area, the region at the rear of the sphere, is solved here. The local Nusselt number at the rear stagnation point is found to be 1.107. To predict the mass transfer rate everywhere on the sphere, a composite solution can be formed from the boundary-layer solution and the rear-region solution.

In heat and mass transfer problems, the method used here complements the boundary-layer methods in predicting the heat or mass transfer rate at the rear of an axisymmetric object.

Introduction

The problem of mass (or heat) transfer to a sphere in Stokes flow has been studied by many people; for example, Levich,¹ Acrivos and Taylor,² and Acrivos and Goddard.³ However, the boundary-layer type of approach used by these authors inevitably breaks down near the rear of the sphere. Although, as pointed out by Acrivos and Goddard, the rate of mass transfer at the rear contributes little to the total mass-transfer rate, it is still physically important in determining the true nature of the processes involved. The method used here is of dual significance. In one sense, it is important for understanding the mass-transfer process of the entire sphere. In another, more general sense, it may be applied to correct boundary-layer solutions of other axisymmetric bodies.

The particular problem investigated here is shown graphically in Figure 1. We shall assume throughout this paper that the velocity field around the sphere is described by Stokes' formula. Due to the concentration difference between the sphere and the surrounding fluid, mass transfer will take place. The convective diffusion equation in spherical coordinates can be written as

$$\frac{Pe}{2} \left[v_r^* \frac{\partial \Theta}{\partial r^*} + \frac{v_\theta^*}{r^*} \frac{\partial \Theta}{\partial \theta} \right] = \frac{\partial^2 \Theta}{\partial r^{*2}} + \frac{2}{r^*} \frac{\partial \Theta}{\partial r^*} + \frac{1}{r^{*2}} \left(\frac{\partial^2 \Theta}{\partial \theta^2} + \frac{\cos \theta}{\sin \theta} \frac{\partial \Theta}{\partial \theta} \right), \quad (1)$$

where $Pe = 2Rv_\infty/D$, the Péclet number; $r^* = r/R$; $\Theta = (c_i - c_o)/(c_\infty - c_o)$; and

$$v_r^* = \left(1 - \frac{3}{2r^*} + \frac{1}{2r^{*3}} \right) \cos \theta ;$$

XBL671-118A

Figure 1. Regions of different mass-transfer mechanisms for mass transfer to a sphere in Stokes flow at high Péclet numbers.

$$v_{\theta}^* = - \left(1 - \frac{3}{4r^*} - \frac{1}{4r^{*3}} \right) \sin \theta .$$

The boundary conditions for equation (1) are:

1. $\Theta = 0$ at $r^* = 1$.
2. $\Theta = 1$ at $r^* = \infty$.
3. $\partial\Theta/\partial\theta = 0$ at $\theta = \pi$, the front axis.
4. $\partial\Theta/\partial\theta = 0$ at $\theta = 0$, the rear axis.

The last two boundary conditions are necessary due to symmetry.

Regions of different mass-transfer mechanisms

In liquid systems where the diffusion coefficient is small, mass transfer usually takes place near the object. On the basis of this concept the authors mentioned previously derived the diffusion-layer equation. The solution indicates that there are two different regions in the mass-transfer process. One is far away from the sphere where the concentration is uniform. The other is very near the surface, where all the concentration variation takes place.

However, in the present work, a more detailed analysis is made. We find that there are four more regions, and each has a different mass-transfer mechanism. The first three regions which are near the sphere are investigated here, and the treatment of the sixth, the so-called "far-wake" region, is relegated to the appendix. A sketch of regions 1 to 5 is shown in figure 1.

Region 1: Region far from the sphere

In this region, the presence of the object of a different concentration is not felt by the fluid, and its concentration is

uniform and equal to the upstream concentration. Consequently, there is no diffusion. Mass is simply carried along by convection.

Region 2: The diffusion (boundary) layer

This is the region solved by the authors mentioned, and only a brief summary shall be given here. In this region, convection and diffusion normal to the surface are important in mass transfer, while diffusion tangential to the surface is relatively unimportant.

Since diffusion is near the surface, a new coördinate normal to the surface is suggested; that is $Y = (r^* - 1)/\epsilon = y/\epsilon$, where $\epsilon = (Pe/2)^{-\frac{1}{3}}$ and $y = r^* - 1$. When equation (1) is expressed in terms of the new variable, the dominant terms for small ϵ yield:

$$\frac{3}{2} Y^2 \cos \theta \frac{\partial \Theta}{\partial Y} - \frac{3}{2} Y \sin \theta \frac{\partial \Theta}{\partial \theta} = \frac{\partial^2 \Theta}{\partial Y^2} \quad (2)$$

with the boundary conditions:

1. $\Theta = 0$ at $Y = 0$.
2. $\Theta = 1$ at $Y = \infty$.
3. $\partial \Theta / \partial \theta = 0$ at $\theta = \pi$.

Since the diffusion term in the θ -direction, $\partial^2 \Theta / \partial \theta^2$ in equation (1), is dropped, only one boundary condition in this direction can be specified for equation (2), namely condition 3.

Equation (2) can be solved by a similarity transformation. The similarity variable is defined by

$$\eta = Y/g(\theta), \quad (3)$$

where

$$g(\theta) = 3^{\frac{1}{3}} \left(\frac{1}{2} \sin 2\theta - \theta + \pi \right)^{\frac{1}{3}} / \sin \theta. \quad (4)$$

The solution obtained is

$$\varepsilon = \frac{1}{\Gamma(4/3)} \int_0^{\eta} e^{-\eta^3} d\eta . \quad (5)$$

The local Nusselt number Nu is

$$Nu_{b\ell}(\theta) = 2 \left. \frac{\partial \Theta}{\partial r^*} \right|_{r^*=1} = \frac{2}{\epsilon \Gamma(4/3) g(\theta)} . \quad (6)$$

As $\theta \rightarrow \pi$, the limit of $g(\theta)$ is $2^{1/3}$, which predicts a finite mass-transfer rate at the front. However, as $\theta \rightarrow 0$, $g(\theta)$ tends toward infinity. This implies an infinitely thick diffusion layer or no mass transfer at the rear stagnation point and contradicts the assumption that mass transfer occurs close to the surface of the sphere. In addition, if one examines the tangential gradient along the rear axis, one will find that this gradient does not vanish, violating boundary condition 4 of equation (1). Thus, it is the objective of this work to eliminate these discrepancies.

Region 3: Convective region

According to the boundary-layer solution, it is apparent that the diffusion layer expands starting from the front to the back of the sphere. Eventually, this layer will expand sufficiently far into the bulk such that the convective velocity becomes large, and the approximate velocity used in the boundary-layer equation becomes invalid. This can be shown by substituting the boundary-layer solution into equation (1) and examining the behavior of the neglected terms near $\theta = 0$:

$$\begin{aligned}
& \eta^2 \Theta' \left[\frac{3}{2} (g^3 \cos \theta + g^2 g' \sin \theta) - \eta g^3 \left(\frac{7}{2} g \cos \theta + \frac{15}{4} g' \sin \theta \right) \epsilon + \right. \\
& \quad \left. + \dots \right] = \underline{\Theta''} + 2\Theta' g(1 - \eta \epsilon g + (\eta \epsilon g)^2 + \dots) \epsilon + \\
& \quad + \left[\eta^2 g'^2 \Theta'' - \eta \Theta' \left(g g'' - 2g'^2 + \frac{\cos \theta}{\sin \theta} g g' \right) \right] (1 - 2\eta \epsilon g + \dots) \epsilon^2 ;
\end{aligned} \tag{7}$$

where the prime designates the derivative with respect to η or θ .

The underlined terms, the terms of the first order of magnitude as $\epsilon \rightarrow 0$, form the boundary-layer equation. However, as $\theta \rightarrow 0$, $g(\theta) \rightarrow O(1/\theta)$, and equation (7) has the form

$$\left(\frac{1}{\theta^3} - \frac{\epsilon}{\theta^4} + \dots \right) = 1 + \frac{\epsilon}{\theta} + \frac{\epsilon^2}{\theta^4} ,$$

with $\eta = O(1)$. The left side represents the convective terms, and the right side, the diffusion terms. It is easy to see that when $\theta = O(\epsilon)$ all the terms in the velocity expansion are of the same order. In other words, as $\theta \rightarrow O(\epsilon)$, the diffusion-layer approximation breaks down.

At the same time, the above analysis suggests that the appropriate variable for this region can be

$$S = \theta/\epsilon .$$

The results of substituting S into equation (1) are

$$\left(1 - \frac{3}{2r^*} + \frac{1}{2r^{*3}} \right) \frac{\partial \Theta}{\partial r^*} - \left(1 - \frac{3}{4r^*} - \frac{1}{4r^{*3}} \right) \frac{S}{r^*} \frac{\partial \Theta}{\partial S} = \frac{\epsilon}{r^{*2}} \left(\frac{\partial^2 \Theta}{\partial S^2} + \frac{1}{S} \frac{\partial \Theta}{\partial S} \right) . \tag{8}$$

The diffusion terms on the right of equation (8), being an order of magnitude smaller, can readily be neglected, and we obtain

$$\left(1 - \frac{3}{2r^*} + \frac{1}{2r^{*3}}\right) \frac{\partial \Theta}{\partial r^*} - \left(1 - \frac{3}{4r^*} - \frac{1}{4r^{*3}}\right) \frac{S}{r^*} \frac{\partial \Theta}{\partial S} = 0. \quad (9)$$

The solution of equation (9) must match with the asymptotic form of the boundary-layer solution at $\theta = 0$.

Equation (9) shows that in this region convection is the dominant mode of mass transfer, and diffusion is negligible. Consequently, Θ is constant along streamlines, $\Theta = \Theta(\Psi)$, where from Stokes' solution one can define a stream function Ψ appropriate to region 3 as

$$\Psi = \frac{S^2}{3^{5/3} \pi^{2/3}} \left(\frac{1}{r^*} - 3r^* + 2r^{*2} \right). \quad (10)$$

The form of $\Theta(\Psi)$ is determined by matching with the boundary-layer solution. For this purpose, it is only necessary to find the asymptotic boundary-layer solution at $\theta = 0$ in terms of the stream function. As $r^* \rightarrow 1$; that is, as r^* asymptotically approaches the boundary layer,

$$\Psi = \theta^2 y^2 / (3\pi)^{2/3} \epsilon^2. \quad (11)$$

On the other hand, as $\theta \rightarrow 0$, from equations (3) and (4) we obtain

$$\eta = \theta y / (3\pi)^{1/3} \epsilon. \quad (12)$$

By comparing equations (11) and (12), we find that in the region of matching the relationship between the similarity variable η and the stream function Ψ is

$$\eta = \sqrt{\Psi}. \quad (13)$$

Thus, the solution in region 3 is

$$\Theta(\Psi) = \frac{1}{\Gamma(4/3)} \int_0^{\sqrt{\Psi}} e^{-x^3} dx, \quad (14)$$

with Ψ defined by equation (10).

Region 4: The rear-axis region

The solution (14) still does not satisfy the boundary condition $\partial\Theta/\partial\theta = 0$ at $\theta = 0$. This can be traced to the neglect of the tangential diffusion terms in equation (8) and indicates the presence of still another region near the rear axis where convection and diffusion in the θ -direction are important, but diffusion in the r -direction is not. At the same time Θ is small, of order $\epsilon^{1/2}$, in this region.

Examination of the terms in equation (1) or (8) neglected in region 3 indicates that the thickness of region 4 is given by $S = O(\sqrt{\epsilon})$. Hence, a suitable variable for this region is

$$s = S/\sqrt{\epsilon} = \theta/\epsilon^{3/2},$$

and the corresponding equation is

$$\begin{aligned} & \left(1 - \frac{3}{2r^*} + \frac{1}{2r^{*3}}\right) \frac{\partial\Theta}{\partial r^*} - \left(1 - \frac{3}{4r^*} - \frac{1}{4r^{*3}}\right) \frac{s}{r^*} \frac{\partial\Theta}{\partial s} \\ & = \frac{1}{r^{*2}} \left(\frac{\partial^2\Theta}{\partial s^2} + \frac{1}{s} \frac{\partial\Theta}{\partial s} \right). \end{aligned} \quad (15)$$

The boundary conditions for equation (15) are

1. $\partial\Theta/\partial s = 0$ at $s = 0$.
2. $\Theta \rightarrow \sqrt{\epsilon} s (2r^{*2} - 3r^* + 1/r^*)^{1/2} / \Gamma(4/3) 3^{5/6} \pi^{1/3}$ as $s \rightarrow \infty$, in order to match with region 3.
3. As $r^* \rightarrow 1$, Θ must match with the solution in region 5 (see equation (19)).

Note that this region, involving the tangential diffusion terms, grows thicker at greater distances from the sphere. Eventually, it blends into a sixth region, the far-wake region,

where the depletion of concentration decays, first, as $1/r^*$ along the rear axis and, second, it decays exponentially in $r^*\theta^2$ in the direction normal to the axis.

Region 5: Region near the rear of the sphere

This region is close to the surface at the rear of the sphere. Being close to the surface, the convective velocity is small. Thus, it is probable that diffusion is comparable to convection. As already shown in region 4, on approaching the rear axis, the tangential diffusion becomes important. In addition, analogous to the diffusion layer concept, the normal diffusion may also be important as the concentration change takes place in a thin region. Order of magnitude comparisons verify these conclusions.

Examination of the radial diffusion terms neglected in regions 3 and 4 show that these terms become important and the treatment of these regions becomes invalid for $y = O(\epsilon)$. Consequently, the appropriate variables in region 5 are

$$Y = y/\epsilon \quad \text{and} \quad S = \theta/\epsilon,$$

and in this region equation (1) reduces for small ϵ to

$$\frac{3}{2} Y^2 \frac{\partial \Theta}{\partial Y} - \frac{3}{2} Y S \frac{\partial \Theta}{\partial S} = \frac{\partial^2 \Theta}{\partial Y^2} + \frac{\partial^2 \Theta}{\partial S^2} + \frac{1}{S} \frac{\partial \Theta}{\partial S}, \quad (16)$$

and one sees that diffusion and convection in both the r - and θ -directions become important. The boundary conditions for equation (16) are

1. $\Theta = 0$ at $Y = 0$.
2. $\partial \Theta / \partial S = 0$ at $S = 0$.
3. $\Theta \rightarrow \eta \Theta' (\eta \rightarrow 0) = \epsilon Y S / \Gamma(4/3) (3\pi)^{2/3}$ as $S \rightarrow \infty$,

in order to match with the boundary-layer solution in region 2.

4. For $Y \rightarrow \infty$, an asymptotic solution for equation (16) is found and will be discussed in the following section.

We have carried the analysis far enough to show that radial diffusion is negligible in regions 3 and 4 and that therefore there can be no upstream propagation of effects from these regions, even though radial diffusion is not negligible in region 5 itself. Thus, another way of stating condition 4 is that the radial diffusion term $\partial^2 C / \partial Y^2$ must become negligible compared to the other terms in equation (16) as $Y \rightarrow \infty$ in order for the solution to match with regions 3 and 4.

In this section we have proceeded logically from one region to another, examining the dominant modes of mass transfer in each region and the limits of validity of the treatment for each region, either by observing whether the boundary conditions are satisfied or by examining the order of magnitude of terms neglected in the equation of convective diffusion. The fact that a limit of validity is found indicates the existence of an adjoining region in which different mechanisms of mass transfer are important. It is not necessary to continue this analysis into the far wake because, as indicated above, radial diffusion is already negligible in regions 3 and 4 and, consequently, there can be no upstream propagation of effects which could affect the mass-transfer rate on the sphere. However, for those who are interested, the far-wake region is discussed in the appendix.

Since region 5 is in contact with the surface, the normal concentration gradient at the surface will yield the rate of mass transfer in the rear region. Thus, it is essential to obtain the solution of equation (16). Condition 3 on equation (16) indicates that $\Theta = O(\epsilon)$ in region 5. Since $y = O(\epsilon)$, the order of the Nusselt number in this region is $Nu = O(\partial\Theta/\partial y) = O(1)$, in contrast to the front part of the sphere where $Nu = O(1/\epsilon)$, as indicated by equation (6).

Solution for the region at the rear of the sphere

To obtain a solution of equation (16), we might first express the boundary condition 4 of equation (16) more explicitly by obtaining the asymptotic solution of region 5 as $Y \rightarrow \infty$. For $Y \rightarrow \infty$, at the boundary joining region 5 to regions 3 and 4, $\partial^2\Theta/\partial Y^2$ should approach zero in order to be consistent with equations (9) and (15) of regions 3 and 4. Imposing this condition on equation (16), we obtain

$$\frac{3}{2} Y^2 \frac{\partial\Theta}{\partial Y} - \frac{3}{2} YS \frac{\partial\Theta}{\partial S} = \frac{\partial^2\Theta}{\partial S^2} + \frac{1}{S} \frac{\partial\Theta}{\partial S} \quad (17)$$

as $Y \rightarrow \infty$. Introduction of the similarity variables

$$\xi = S \sqrt{Y} \quad \text{and} \quad \Theta = \epsilon Y^{\frac{1}{2}} f(\xi)$$

reduces equation (17) to

$$\xi f'' + (1 + 3\xi^2/4)f' - 3\xi f/4 = 0, \quad (18)$$

with the boundary conditions

1. $f \rightarrow \xi/\Gamma(4/3)(3\pi)^{\frac{1}{3}}$ as $\xi \rightarrow \infty$, to match with the boundary-layer solution.
2. $f' = 0$ at $\xi = 0$.

Equation (18) can readily be solved numerically.

To verify that the solution, $\Theta = \epsilon Y^{\frac{1}{2}} f(\xi)$, agrees with the assumption that $\partial^2 \Theta / \partial Y^2 \rightarrow 0$ as $Y \rightarrow \infty$, we calculate $\partial^2 \Theta / \partial Y^2$ and $\partial^2 \Theta / \partial S^2$ from this solution with the result

$$\partial^2 \Theta / \partial Y^2 = \epsilon(-f + \xi f' + \xi^2 f'') / 4Y^{3/2} \quad \text{and} \quad \partial^2 \Theta / \partial S^2 = \epsilon Y^{3/2} f''.$$

It is obvious that the assumption is not violated. Moreover, the function f provides the boundary condition for region 4 when $r^* \rightarrow 1$:

$$\Theta \rightarrow \epsilon^{\frac{1}{2}} \sqrt{r^* - 1} f(s \sqrt{r^* - 1}) \quad \text{as } r^* \rightarrow 1 \quad \text{for region 4.} \quad (19)$$

Thus, region 4 can be solved in connection with region 3 without knowing the solution for all of region 5.

So far, an analytic solution of equation (16) does not seem possible; therefore, a numerical solution is sought. Due to the presence of the convective terms in equation (16), a simple finite-difference method in the Y, S -coördinates does not appear practical. For greater efficiency, two new variables

$$\psi = YS \quad \text{and} \quad X = (Y^2 - S^2)/2$$

are introduced. Note that ψ is proportional to the square root of the stream function in region 5.

Equation (16) thus becomes

$$\frac{3}{2} Y(S^2 + Y^2) \frac{\partial \Theta}{\partial X} = (S^2 + Y^2) \left(\frac{\partial^2 \Theta}{\partial \psi^2} + \frac{\partial^2 \Theta}{\partial X^2} \right) + \frac{Y}{S} \frac{\partial \Theta}{\partial \psi} - \frac{\partial \Theta}{\partial X}. \quad (16')$$

Figure 2 shows regions 2, 3, 4, and 5 in the new coördinate system.

A source of difficulty arises in the numerical calculations because convection along streamlines becomes much larger than

Figure 2. Mass transfer regions in streamline coordinates.

XBL672-827

diffusion along streamlines as Y or S approaches infinity. It is ironic that these diffusion terms cause difficulty in the numerical method in just those regions where they become negligible. This is, in fact, a major reason why it is difficult to obtain numerical solutions of the original problem with equation (1) when the diffusion coefficient is small (another reason, of course, being that the concentration variation takes place very close to the surface of the sphere). The development of perturbation methods in the first place is partly an effort to overcome such difficulties.

In the present problem, one can avoid these difficulties by neglecting explicitly the term for diffusion along streamlines $\partial^2 \Theta / \partial X^2$ in the numerical calculations. In those regions where it is, in fact, negligible. Thus, from the discussion of regions 3 and 4, it can be shown that $\partial^2 \Theta / \partial X^2 \approx 0$ in these regions. Thus, upon approaching regions 3 and 4, the appropriate asymptotic equation can be

$$\frac{3}{2} Y(S^2 + Y^2) \frac{\partial \Theta}{\partial X} = (S^2 + Y^2) \frac{\partial^2 \Theta}{\partial \psi^2} + \frac{Y}{S} \frac{\partial \Theta}{\partial \psi} - \frac{\partial \Theta}{\partial X} \quad (20)$$

Similarly, equation (20) can be used upon reaching region 2. Although equation (20) is not exactly the same as the equations derived for the regions 2, 3, and 4, it approximates the conditions when convection becomes more predominant than diffusion along streamlines. In addition, equation (2), being parabolic, is influenced only by the upstream conditions and can be solved without iteration. Equation (16'), involving a streamline coördinate, makes it easier to drop the term for diffusion along streamlines than would be the case for equation (16).

Equation (16') can be solved by the successive over-relaxation method, and the sequence of computations is as follows (for this purpose, regions 2, 3 and 4 in figure 2 should be regarded as asymptotic parts of region 5 where diffusion along streamlines is negligible):

1. Region 2 and the adjacent region 5 are computed to match the boundary-layer solution.
2. Regions 3 and 5 are solved simultaneously, and no computation in region 4 is needed.

One may already have noted that the origin ($\psi = X = 0$) is a singular point; consequently, it is difficult to obtain the gradient there accurately. An alternative would be to assume first a solution of the form

$$\Theta = \bar{\Theta} + AY, \quad (21)$$

since at small Y , Θ is linearly proportional to Y . First, one can guess a value of A and then back out the term AY from Θ and solve for $\bar{\Theta}$ using equations (16') and (20). Once $\bar{\Theta}$ is found, a new A can be calculated, and the procedure can be repeated until the values for A converge. By using this method, we are able to calculate accurately the gradient at the origin, i.e., $A = 0.5535 \epsilon$.

Results and Discussion

From the calculated concentration distribution in the rear-stagnation region, one can easily obtain the mass-transfer rate along the surface by calculating the normal gradient at the surface. The results are shown in figure 3, where Nu_r is the local Nusselt number and the subscript r denotes the solution for the rear region. Also shown in figure 3 is the local Nusselt

XBL671-119A

Figure 3. Local Nusselt number in the region near the rear of the sphere.

number Nu_{bl} predicted by the boundary-layer solution. It is clear, as one would expect, that there is a large difference at small angles, and yet the solution for region 5 blends into the boundary-layer solution at larger angles.

At the rear stagnation point, $S = 0$, the local Nusselt number is $Nu_r = 1.107$. This is less than the Nusselt number ($Nu = 2$) for the diffusion from a sphere into a stagnant medium. The reason, obviously, is that the convection of depleted solution into the rear region decreases the concentration gradient at the rear.

In order to obtain the mass-transfer rate for the entire sphere, it is necessary to form a composite solution, with the boundary-layer solution as the outer solution and the solution for the rear region as the inner solution. Mathematically, the composite solution for the local Nusselt number can be written as

$$Nu(\theta) = Nu_{bl}(\theta) + Nu_r(S) - 2\theta/\epsilon\Gamma(4/3)(3\pi)^{\frac{1}{3}},$$

where the subscript bl denotes the boundary-layer solution. The last term is the inner limit of the boundary-layer solution, which is the same as the outer limit of the solution for region 5.

Acrivos and Goddard have obtained the asymptotic average Nusselt number for $Pe \rightarrow \infty$, and $Re \rightarrow 0$, as

$$\overline{Nu} = Pe^{\frac{1}{3}} [0.9914 + 0.922 Pe^{-\frac{1}{3}} + O(Re) + O(Pe^{-\frac{2}{3}})],$$

and they also have suggested that the contribution of the rear stagnation region to $\overline{Nu}$ is $O(Pe^{-\frac{1}{3}})$, although it actually is of the order $Pe^{-\frac{2}{3}}$. Therefore, if one wants to include the contribution to $\overline{Nu}$ of the rear stagnation region, one should also carry

out two more higher order approximations in Acrivos and Goddard's analysis for the diffusion layer.

Even though the contribution of the rear of the sphere to the average Nusselt number is very small, the solution in this region is of interest because it can elucidate one example of the failure of boundary-layer methods at the rear of a bluff object. The solution found here for the rear of a sphere in Stokes flow would also be applicable to other problems at high Péclet numbers, first to the sphere at higher Reynolds numbers where Stokes' velocity profile is not applicable and secondly to other axisymmetric bodies in the absence of eddies behind the bodies, as long as the rear of the body is blunt like the sphere and not pointed. The rear region is very small, and in this region the velocity profile for such an axisymmetric body can be expressed as

$$v_y = b y^2 \quad \text{and} \quad v_r = - b r y ,$$

where b is a constant and r denotes the normal distance from the axis of symmetry. In addition one needs the asymptotic form of the mass-transfer rate at the rear of the object as predicted from the diffusion layer on the forward part of the object. Then, by appropriate stretching of the coördinates and concentration, the problem can be reduced to equation (16) and the following boundary conditions, for which the solution has already been given.

Appendix. The far-wake region.

The asymptotic solution for region 4 as $r^* \rightarrow \infty$ is

$$\Theta \rightarrow \sqrt{\epsilon r^*} f(s\sqrt{2r^*/3}), \quad (22)$$

which, we might note, is very similar to the asymptotic solution in the same region as $r^* \rightarrow 1$ (see equation (19)). The treatment of region 4 ceases to be valid when $\Theta = O(1)$, which occurs when $r^* = O(1/\epsilon)$. This then defines a sixth region, that of the far wake. In this region, the appropriate variables are

$$R = \epsilon r^* \quad \text{and} \quad S = \theta/\epsilon^2,$$

and equation (1) reduces to

$$\frac{\partial \Theta}{\partial R} - \frac{S}{R} \frac{\partial \Theta}{\partial S} = \frac{1}{R^2} \left(\frac{\partial^2 \Theta}{\partial S^2} + \frac{1}{S} \frac{\partial \Theta}{\partial S} \right), \quad (23)$$

with the boundary conditions

1. $\Theta \rightarrow 1$ as $S \rightarrow \infty$.
2. $\partial \Theta / \partial S = 0$ at $S = 0$.

$$3. \quad \Theta \rightarrow \frac{1}{\Gamma(4/3)} \int_0^B e^{-x^3} dx + \sqrt{R} f(S\sqrt{2R/3}) - \frac{SR\sqrt{2/3}}{\Gamma(4/3)(3\pi)^{1/3}}$$

as $R \rightarrow 0$, where $B = SR\sqrt{2/3}^{5/6} \pi^{1/3}$.

This last boundary condition comes from matching with both region 3 and region 4 and represents the composite expansion for these two regions as $r^* \rightarrow \infty$.

The asymptotic solution in region 6 as $R \rightarrow \infty$ is

$$\Theta = 1 - (K/R) \exp(-RS^2/4), \quad (24)$$

where the constant K is related to the total rate of mass transfer to the sphere:

$$K = \overline{Nu} / (4Pe)^{\frac{1}{3}} \approx 0.6245 .$$

Thus, although detailed solutions have not been obtained for regions 4 and 6, one does arrive at a reasonably complete picture of the concentration distribution in the entire flow field to the first approximation as $Pe \rightarrow \infty$.

Acknowledgment

This work was supported by the United States Atomic Energy Commission.

Nomenclature

- c_i - concentration of the diffusing species
- c_o - concentration at the surface
- c_∞ - concentration in the bulk solution
- D - diffusion coefficient
- f - asymptotic solution for $\epsilon\Theta/Y^{\frac{1}{2}}$ as $Y \rightarrow \infty$ in region 5
- g - see equation (4)
- Nu - the local Nusselt number
- $Pe = 2Rv_\infty/D$, the Péclet number
- r - radial coördinate
- r^* - dimensionless radial coördinate
- R - radius of the sphere
- $R = \epsilon r^*$, stretched coördinate for region 6
- $s = S/\sqrt{\epsilon} = \theta/\epsilon^{3/2}$, stretched coördinate for region 4
- $S = \theta/\epsilon$, stretched coördinate for regions 3 and 5
- $S = \theta/\epsilon^2$, stretched coördinate for region 6
- v_r^* - dimensionless radial velocity
- v_θ^* - dimensionless tangential velocity
- v_∞ - velocity far from the sphere

- x - distance measured along the rear axis
 $X = (y^2 - s^2)/2$, variable used in numerical solution for region 5
 $y = r^* - 1$, the normal distance from the sphere
 $Y = y/\epsilon$, a stretched diffusion-layer coördinate
 $\Gamma(4/3) = 0.89298$, the gamma function of $4/3$
 $\epsilon = (Pe/2)^{-1/3}$, perturbation parameter
 $\eta = Y/g(\theta)$, similarity variable for diffusion layer
 θ - angular coördinate, measured from rear axis
 $\Theta = (c_1 - c_0)/(c_\infty - c_0)$, dimensionless concentration
 $\xi = SVY$, similarity variable for asymptotic solution in region 5
 $\psi = YS$, stream function variable appropriate to region 5
 Ψ - stream function variable appropriate to region 3 (see equation (10))

References

1. Veniamin G. Levich. Physicochemical Hydrodynamics. Englewood Cliffs, N.J., Prentice-Hall, Inc., 1962, section 14.
2. Andreas Acrivos and Thomas D. Taylor. "Heat and Mass Transfer from Single Spheres in Stokes Flow." The Physics of Fluids, 5, 387-394 (1962).
3. Andreas Acrivos and J. D. Goddard. "Asymptotic expansions for laminar forced-convection heat and mass transfer. Part 1. Low speed flows." Journal of Fluid Mechanics, 23, 273-291 (1965).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000