

eScholarship

International Journal of Comparative Psychology

Title

Journal Publication Trends Regarding Cetaceans Found in Both Wild and Captive Environments: What do we Study and Where do we Publish?

Permalink

<https://escholarship.org/uc/item/1nh5g1f8>

Journal

International Journal of Comparative Psychology, 23(3)

ISSN

0889-3675

Authors

Hill, Heather
Lackups, Monica

Publication Date

2010

DOI

10.46867/ijcp.2010.23.03.01

Copyright Information

Copyright 2010 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Journal Publication Trends Regarding Cetaceans Found in Both Wild and Captive Environments: What do we Study and Where do we Publish?

Heather Hill and Monica Lackups
St. Mary's University, U.S.A.

Scientists conducting research on cetaceans have a variety of publication outlets. However, a formal assessment of those options has not been conducted. To better understand the trends in publications regarding dolphins and whales, we surveyed peer-reviewed articles from 9 different databases. Our survey produced 1,628 unique articles involving 16 cetaceans found both in the wild and in captivity. Each article was coded a variety of information: habitat, geographic location, genus, topic, research design, and journal type. The analyses indicated that 68% studies were conducted with wild populations and 29% were performed with captive populations. A quarter of the journals publishing research on dolphins or whales published almost 80% of all the articles selected for this study. Studies were conducted across many different geographic locations and topics. Other major findings elucidated relationships between various variables. As expected, specific topics were more likely associated with certain research designs, habitats, and journal types. One of the most important findings of this study is the limited publication of research conducted with captive cetaceans. While it is important to continue to examine animals in their natural environments, there is much to be learned from studies conducted with animals in captivity. As a group, we must become cognizant of the publication trends which currently describe our research progress as we integrate our knowledge from captivity and the wild.

Much of the extant knowledge of marine mammals is summarized in large compilations of encyclopedias or books dedicated to specific cetaceans, aspects of their life history, geographic distributions, anatomy and physiology, sensory systems and capabilities, or behavioral and cognitive abilities (e.g., Herman, 1980; Ridgway & Harrison, 1999; Schusterman, Thomas, & Wood, 1986; Thomas & Kastelein, 1990). These comprehensive summaries are instrumental in the integration of past knowledge and current findings. Many of the topics compiled in these comprehensive books include findings from research conducted in both the wild and captive settings. A simple review of this literature suggests that some research topics are better addressed with wild populations (e.g., life history parameters, distributions) while others are more appropriate for captive animals (e.g., sensory system capabilities, cognitive capacities).

It is critical that research continues to integrate information from both wild and captive populations. Each population provides unique contributions to our understanding of the pressures faced by marine mammals. Ten years ago, Mann and her colleagues compiled and published the most comprehensive review of the literature to date on several species of cetaceans in the book, *Cetacean Societies* (Mann, Connor, Tyack, & Whitehead, 2000). The editors of this book provided a wealth of information that integrated different aspects of the lives of four cetaceans. Interestingly, two of the cetaceans are found in both wild and captive settings (i.e., *Tursiops truncatus*, bottlenose dolphin, and *Orcinus orca*, killer

We would like to extend a special thank to Mr. Alonzo Mendoza and Ms. Samantha Lopez for their help in locating and compiling additional sources for this study. Without their help and Ms. Monica Lackups' dedication, this study would not have been possible. Correspondence concerning this article should be addressed to Heather M. Hill, Department of Psychology, St. Mary's University, One Camino Santa Maria, San Antonio, TX, 78228, U.S.A. (hhill1@stmarytx.edu).

whale) and two are found only in the wild (i.e., *Megaptera novaeangliae*, humpback whale, and *Physeter macrocephalus*, sperm whale). Authors of individual chapters compiled information regarding the history of captive research, aspects of life history, behavior, development, relationships, acoustics, and communication abilities of these four cetaceans. One of the particularly exciting components of this edited book was the integration of findings from observations in the wild and observations in captivity. This integration allowed readers to assess the findings of research conducted with wild and captive animals simultaneously. For example, this comprehensive review suggested that wild and captive bottlenose dolphins often exhibit very similar behaviors (e.g., play behaviors, association patterns, affiliative or aggressive interactions) and developmental milestones (e.g., echelon swim versus infant swim patterns, emergence of independence).

With approximately 35 different species of cetaceans in captivity (May, 1998), only a very small number have been studied and compared with their wild counterparts (e.g., *Tursiops*; *Orcinus*). In fact, since the publication of *Cetacean Societies* a decade ago, very few comprehensive reviews regarding the progress made in the knowledge of cetaceans have been published (e.g., Harley, 2008; Shaffer & Costa, 2006). Researchers must comb through and integrate a vast body of published peer-reviewed literature, article by article.

Comprehensive reviews on various topics involving wild and captive cetaceans would allow current researchers opportunities to better understand the areas in which progress has been made as well as identify areas in need of more research. For example, research has progressed in our understanding of the acoustic complexity of a variety of cetaceans, including humpback whales, sperm whales, and killer whales (Deecke, 2006). However, much of this research has been conducted in the wild. In comparison, many improvements have been made in the care of captive animals, resulting in better health and longevity (Wells, 2009). Research regarding the effects of humans on the behavior and physical health of cetaceans has also expanded and incorporated knowledge from research conducted in wild and captive settings (Nowacek, Thorne, Johnston, & Tyack, 2007; Thomas, 2009). Similarly, attempts to examine the cognitive abilities, the importance of play, social interactions, and individual differences of a variety of cetaceans continue to move forward (Harley, 2008; Highfill & Kuczaj, 2007; Kuczaj, Gory, & Xitco, 2009; Paulos, Dudzinski, & Kuczaj, 2008).

A recent report regarding animals in captivity, sponsored by the Humane Society of the United States and the World Society for the Protection of Animals, indicated that research involving captive cetaceans was lacking and had made little progress (Rose, Parsons, & Garinato, 2009). Specifically, the authors indicated that research from United States-based marine mammal facilities was minimal and had produced few significant contributions regarding our understanding of marine mammals. With incredulity, we initiated the current review of the available literature pertaining specifically to cetaceans. Our own experience suggested a different interpretation, and we wished to determine the accuracy of the conclusions drawn by Rose and her colleagues.

A variety of marine mammals have been housed in captive environments for approximately 150 years (Samuels & Tyack, 2000). Given the vast body of published literature on marine mammals from both captive and wild settings, we narrowed the focus of our review to include only journal articles that centered on cetaceans cared for by humans at some point during captivity's documented history. Although this choice biased our understanding of publication trends regarding marine mammals, it allowed for a more comprehensive and interpretable picture of research conducted with one major branch of marine mammals.¹ To better understand the type of research conducted with cetaceans found in both the wild and in captivity and published in journals, we examined the following questions:

1. How are articles on dolphins or whales distributed across a variety of databases?
2. Which journals published research involving dolphins or whales most frequently?
3. What dolphins or whales were represented most frequently?
4. What geographic locations were represented most frequently?
5. From which habitat (wild, captive, or both) were samples derived the most frequently?
6. What topics were represented most frequently?
7. What research designs were used most frequently?
8. Were the articles distributed across the journal types related to the habitat?
9. Were the topics examined related to the habitat from which the samples were derived?
10. Were the habitats related to the type of research design used within the studies?
11. Were the research designs related to the type of journal in which they were published?
12. Were the topics examined related to their research designs?

Method

Sample

We examined a sample of 1,628 unique peer-reviewed journal articles published between 1950 and December 2009. Although there are documented publications dating to 1875, we wanted to focus on articles that were contemporaneous with the current research standards for animal safety and welfare. Our sample was derived from a comprehensive, but not necessarily exhaustive, search of nine databases, using the terms “dolphin” or “whale” as our search terms in *all text available* for the nine search engines: Academic Search Complete, Agricola, Annual Reviews, Biological Abstracts, EbscoHost (E-Journals, TOPICSearch, and PsycArticles), JSTOR, PsychINFO, Psychology & Behavioral Sciences Collection, and Science & Technology Collection. These terms represented the greatest number of response hits as compared to “cetacean” only and even “marine mammal.” We

¹We believe that a similar review of pinnipeds alone would be extremely beneficial for the field. A review of other marine mammals not captured in the current review would also be productive.

recognize that some articles relevant to the current study may not be represented in the current sample.² However, we believe the sample is representative of the research conducted and published.

Procedure

In order to manage the large number of potential sources, we used several exclusionary criteria during the search process. First, we excluded marine mammals that were not cetaceans (e.g., *Ursus maritimus*, polar bears, and *Enhydra lutris kenyoni*, sea otters) and whale or dolphin genera that were found only in the wild (e.g., *Balaenoptera* sp., all baleen whales and sperm whales). Thus, we included cetaceans that had been cared for by humans for some length of time (i.e., beyond animals held temporarily due to strandings or catch-and-release field studies). Second, we excluded all books, chapters, reviews, unpublished dissertations, and popular press references. Only peer-reviewed articles found within the above databases were selected for this study. Third, we excluded articles in which fossils or fossil records comprised the primary sample. Most of the studies excluded using this criterion focused on the paleontology and evolutionary history of cetaceans. While these studies are important, they did not appear to be directly related to the current care and well-being of animals in captivity. Finally, if a genus could not be identified from the title, abstract, or full text, it was coded as “unidentified.”

Using the exclusionary criteria above, the search for articles, from which the current sample was derived, produced over 30,000 hits. Each abstract or article was individually reviewed to determine if it met the inclusion criteria (i.e., the word, dolphin or whale, present somewhere in the abstract or title, the species could be found both in the wild and in captivity, the subjects generally were alive, and the study was a peer-reviewed journal article). Approximately 325 articles were duplicated across different databases. Duplicate articles were excluded from the final sample. Once selected for the sample, each abstract or article was coded for the following information: habitat, geographic location, genus, topic, research design, and journal type. Due to the huge range of geographic location, topics, and journals, we collapsed each category into larger categories to facilitate our statistical analyses. See Table 1 for the final individual designations for each category.

Geographic location was originally designated by United States (U.S.) areas (Northeast, Southwest, etc.), European Union (E.U.), United Kingdom (U.K.), and single countries or locations (e.g., Caribbean, Gulf of Mexico). These areas were then grouped by ocean basin with the exception of the North Atlantic, which was divided into the Canadian North Atlantic, the U.S. North Atlantic and the European North Atlantic. If the specific area could not be determined, then it was coded as a separate category indicating that no location was stated. See Table 1 for a list of collapsed categories of geographic areas.

Each article was also coded for species. Initially, all genus species identifications were made for every species examined in a given study. Species that were not directly identified were placed in an unidentified category. The species categories were then collapsed by genus to facilitate the analyses. Sixteen genera were identified across journal articles (see Table 1).

After identifying over 41 topics across the 1,628 articles, we grouped the topics into 14 broad categories. While a finer level of analysis would be ideal, there were too many categories with less than five cases. This condition violated one of the assumptions for our statistical analyses. Thus, the more specific categories were collapsed and may be found in Table 1. See the appendix for the original categories used to create the broad categories analyzed.

Each study was also coded for the type of research design used to conduct the study. Studies were grouped into one of the following categories: observational, observational-biology, experimental-psychology, experimental-biology, and undetermined or miscellaneous. Observational studies were any studies performed in which no or very little manipulation of the subjects occurred. These studies included population and distribution studies and observational research studies of various behaviors. Observational-biology studies included studies that involved techniques requiring more manipulation or invasion to the animal, such as acquiring biological matter (e.g., lesions,

²Different levels of access are available for various databases. Thus, if the current search was replicated at a different institution or facility with greater levels of access, a different sample would be produced.

stomach contents, genetic material) from stranded animals. Experimental-psychology studies involved any research in which aspects of the environment were manipulated or altered to investigate the effects on cognition or behavior. Experimental-biology studies involved any research in which manipulations occurred and affected the subjects, physiologically. These studies included studies on genetics, physiological studies, sensory system studies, and immunological studies. The categories are also listed in Table 1.

Finally, our initial sample included 290 unique journals. Journals were collapsed into nine categories to better identify any trends in publications. The collapsed categories generally represented journals that shared similar topics or scopes of interest and may be found in Table 1.

Table 1
Categories coded for each article.

Variable	Categories
Habitat	Captivity, Wild, Both
Geographic Location	Arctic, North Atlantic Canada, North Atlantic U.S., North Atlantic Europe, Central Atlantic (Caribbean & Gulf of Mexico), South Atlantic (Antarctic waters), North Pacific, South Pacific, Indian, Mediterranean
Genera	<i>Cephalorhynchus, Delphinapterus, Delphinus, Feresa, Globicephalus, Grampus, Lagenodelphis, Lagenorhynchus, Neophocaena, Orcinus, Phocoena, Pseudorca, Sousa, Stenella, Steno, Tursiops</i> , Unidentified
Topic	Behavior, Biology, Cognition, Conservation, Culture, Development, Echolocation & Sound Detection, Human-Dolphin Interaction-Ecotourism, Human-Dolphin Interaction-Therapy, Maternity/Calf Mortality, Methodology, Noise & Distribution, Social Structure, Sound Production
Journal Type	Animal/Ecology/Zoology, Biology/Chemistry/Sciences, Conservation/Environment, Human Interaction- Tourism & Fisheries, Marine Sciences, Psychology, Sound/Acoustics, Therapy, Other (Technical/Foreign Journals)
Research Type	Observational, Observational-Biology, Experimental-Psychology, Experimental-Biology, Other-Methodology & Technology

Results

Article distribution across databases

The results of a chi-square-goodness-of-fit test indicated that articles on cetaceans were distributed unequally across databases, $\chi^2(8, N = 1628) = 1,242, p < 0.001$. Articles on cetaceans were found most frequently in the EbscoHost, Biological Abstracts, and PsycINFO databases (see Fig. 1).




Figure 1. Distribution of articles per database.

Representation of journals

Approximately a quarter (24.5%) of the journals identified in the current study published 78.5% of the selected journal articles. As seen in Table 2, the greatest number of publications on cetaceans was produced by *Aquatic Mammals* and *Marine Mammal Science*, which specialize in all aspects of marine mammals, and the *Journal of the Acoustical Society of America*, which specializes in the study of sound production and reception. The journals with the next most frequent publications regarding dolphins or whales dropped dramatically in their publication frequency as compared to these three prolific journals. Each of the remaining journals published only 2% or less of the identified articles.

Representation of genera

The results of a chi-square-goodness-of-fit test indicated that the cetaceans were not distributed equally, $\chi^2(16, N = 1865) = 4,555.80, p < 0.001$. The most frequently researched animals were the genus, *Tursiops*, and the genus, *Orcinus*, along with a significant number of articles that did not have an identifiable species. See Table 3 for the distribution of genera.

Representation of geographic location

A chi-square-goodness-of-fit test indicated that the geographic locations were not equally distributed, $\chi^2(11, N = 1677) = 797.21, p < 0.001$. The most frequent location was the North Atlantic Ocean around Canada, followed by the North Pacific Ocean, the North Atlantic Ocean around the United States, and the Indian Ocean. The Arctic Ocean had the fewest studies with less than 2% of the articles. See Table 4 for the distribution of geographic locations.

Table 2
Distribution of articles across journals publishing five or more articles.

Journal	f	%
<i>Aquatic Mammals</i>	250	15.36
<i>Marine Mammal Science</i>	148	9.09
<i>Journal of the Acoustical Society of America</i>	119	7.31
<i>Science</i>	31	1.90
<i>Journal of Mammalogy</i>	31	1.90
<i>Animal Behaviour</i>	30	1.84
<i>Journal of Experimental Biology</i>	27	1.66
<i>Nature</i>	25	1.54
<i>Journal of Comparative Psychology</i>	24	1.47
<i>Canadian Journal of Zoology</i>	24	1.47
<i>Environmental Science & Technology</i>	22	1.35
<i>Molecular Ecology</i>	20	1.23
<i>Marine Biology</i>	19	1.17
<i>Biological Conservation</i>	18	1.11
<i>Annual Reviews</i>	17	1.04
<i>Science of the Total Environment</i>	16	0.98
<i>International Journal of Comparative Psychology</i>	16	0.98
<i>Behavioral Ecology and Sociobiology</i>	16	0.98
<i>Anthrozoos</i>	16	0.98
<i>Veterinary Record</i>	15	0.92
<i>Arctic</i>	15	0.92
<i>Marine Fisheries Review</i>	14	0.86
<i>Chemosphere</i>	14	0.86
<i>Acoustical Physics</i>	14	0.86
<i>Zoo Biology</i>	13	0.80
<i>Wildlife Conservation</i>	12	0.74
<i>Natural History</i>	12	0.74
<i>Bioacoustics</i>	12	0.74
<i>Behavioural Processes</i>	12	0.74
<i>Marine Environmental Research</i>	11	0.68
<i>Proceedings: Biological Sciences</i>	10	0.61
<i>Ethology</i>	10	0.61
<i>Sensory Systems</i>	9	0.55
<i>Fishery Bulletin</i>	9	0.55

Table 2 (cont.)

Journal	f	%
<i>Environmental Toxicology & Chemistry</i>	9	0.55
Ecology	9	0.55
Behaviour	9	0.55
Animal Learning & Behavior	9	0.55
Journal of Animal Ecology	8	0.49
Behavioral & Brain Sciences	8	0.49
Veterinary Immunology & Immunopathology	7	0.43
Journal of Experimental Psychology	7	0.43
Fisheries Science	7	0.43
Pacific Science	6	0.37
New Zealand Journal of Marine & Freshwater Research	6	0.37
Mammal Review	6	0.37
Journal of Wildlife Diseases	6	0.37
Journal of Parasitology	6	0.37
Jn of Comp Physiology B: Biochem, Systemic, & Environ Physiology	6	0.37
ICES Journal of Marine Science	6	0.37
Emerging Infectious Diseases	6	0.37
Ecological Applications	6	0.37
Comparative Biochemistry & Physiology Part B	6	0.37
Brain, Behavior & Evolution	6	0.37
Applied Animal Behaviour Science	6	0.37
Animal Cognition	6	0.37
Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C	6	0.37
Veterinary Microbiology	5	0.31
Polar Biology	5	0.31
Molecular Ecology Notes	5	0.31
Journal of Heredity	5	0.31
Journal of Experimental Marine Biology & Ecology	5	0.31
Journal of Applied Ecology	5	0.31
Environment	5	0.31
Deep-Sea Research Part II, Topical Studies in Oceanography	5	0.31
Conservation Biology	5	0.31
Bioscience	5	0.31
Behavioural Brain Research	5	0.31

Table 2 (cont.)

Journal	f	%
Aquatic Toxicology	5	0.31
American Journal of Veterinary Research	5	0.31
A.N.A.E. Approche Neuropsychologique des Apprentissages chez l'Enfant.	5	0.31

Note: Journals publishing 5 or more cetacean articles are listed in this table. The top 10 journals publishing the most articles are **bolded** and presented in descending order. Journals with 10 or more cetacean articles are *italicized* and presented in descending order. Finally, journals are alphabetized within each number category.

Table 3
The representation of genus categories across articles.

Genus	f	%
<i>Cephalorhynchus</i>	24	1.3
<i>Delphinapterus</i>	104	5.6
<i>Delphinus</i>	86	4.6
<i>Feresa</i>	6	0.3
<i>Globicephala</i>	16	0.9
<i>Grampus</i>	33	1.8
<i>Lagenodelphis</i>	8	0.4
<i>Lagenorhynchus</i>	77	4.1
<i>Neophocaena</i>	32	1.7
<i>Orcinus</i>	170	9.1
<i>Phocoena</i>	65	3.5
<i>Pseudorca</i>	28	1.5
<i>Sousa</i>	49	2.6
<i>Stenella</i>	157	8.4
<i>Steno</i>	14	0.8
<i>Tursiops</i>	743	39.8
Unidentified	253	13.6

Table 4
Distribution of geographic location across articles.

Area	f	%
Arctic	31	1.8
North Atlantic- Canada	318	19.0
North Atlantic- U.S.	235	14.0
North Atlantic- Europe	199	11.9
Central Atlantic	55	5.6
South Atlantic	83	4.9
North Pacific	265	15.8
South Pacific	107	6.4
Indian	212	12.6
Mediterranean	94	5.6
General	54	3.2
None stated	24	1.4

Representation of habitat

A chi-square-goodness-of-fit test indicated that the habitats (wild, captive, or both) were not equally represented across the sample, $\chi^2(2, N = 1628) = 1,062.42, p < 0.001$. Wild habitats were most frequently studied as opposed to captive habitats. A very small number of the articles included samples from both habitats. See Figure 2 for the percentage breakdown of each habitat.




Figure 2. Distribution of articles per habitat.

Representation of topics

A chi-square-goodness-of-fit test indicated that the topics were not equally represented, $\chi^2(13, N = 1631) = 2,034.00, p < 0.001$. As Table 5 demonstrates, biology comprised the most frequently published topic, followed by behavior, and the effects of anthropogenic noise on cetaceans and cetacean distribution. The topics with the fewest articles included culture, human-dolphin interactions involving therapy, and development. See Table 5 for the results regarding the 14 broad topics.

Table 5
Distribution of articles across broad topics.

Topic	f	%
Anthropogenic Noise & Distribution	144	8.8
Behavior	243	14.9
Biology	533	32.7
Calf Mortality & Maternity	33	2.0
Cognition	82	5.0
Conservation, Ecology, & Environment	52	3.2
Culture	11	0.7
Echolocation, Sound Detection	113	6.9
Development	26	1.6
Human-dolphin: Interaction	96	5.9
Human-dolphin: Therapy	24	1.5
Methodology	88	5.4
Sound Production	139	8.5
Social Structure	47	2.9

Representation of research design

A chi-square-goodness-of-fit test indicated that the research designs were not equally utilized, $\chi^2(4, N = 1627) = 830.75, p < 0.001$. The most frequent research design used was observational in nature, followed by studies using an observation-biology based design. See Figure 3 for the percentage breakdown of research designs across articles.

Representation of habitat across journal types

A chi-square-test-of-independence indicated that the journals and habitats were related, $\chi^2(16, N = 1300) = 267.50, p < 0.001$, Cramer's $V = 0.29$. As seen in Table 6, the animal, ecology, and zoology journals published articles investigating wild populations significantly more than captive populations. In contrast, the

marine science journals published articles almost exclusively from wild populations. As might be expected, psychology journals and the sound and echolocation journals published significantly more articles examining data from captive populations. See Table 6 for additional information.




Figure 3. Distribution of articles per research design.

Table 6

Distribution of articles across journal and habitat.

Journal Group	Habitat					
	Wild		Captive		Both	
	f	%	f	%	f	%
Biology/Sciences	313	28.2	<i>133</i>	28.4	11	23.4
Psychology	75	6.8	116	24.8	10	21.3
Animals, Ecology, Zoology	155	14.0	38	8.1	2	4.3
Acoustics	61	5.5	80	17.1	6	12.8
Marine Science	398	35.9	67	14.3	14	29.8
Conservation	74	6.7	3	0.6	0	0
Therapy	1	0.1	8	1.7	0	0
Human-Dolphin Interactions	19	1.7	3	0.6	1	2.1
Other	13	1.2	20	4.3	3	6.4
Total	1,109	100	468	100	47	100

Note: Percentages above 10% and their frequencies are **bolded** to help identify trends. The highest percentage is also *italicized*.

Representation of article topic across habitat

A chi-square-test-of-independence indicated that the topics were related to the habitats from which the samples were taken, $\chi^2 (26, N = 1,624) = 456.10, p < 0.001$, *Cramer's V* = 0.38. As might be expected, research on the effects of anthropogenic noise on cetaceans and cetacean distributions was performed more often in the wild while research on cognition and echolocation or sound detection was conducted most often in captivity. See Table 7 for specific information.

Table 7
Distribution of articles across topic and habitat.

Topic	Habitat					
	Wild		Captive		Both	
	f	%	f	%	f	%
Anthropogenic Noise & Distribution	140	12.6	1	0.2	1	2.1
Behavior	199	17.9	42	9.0	2	4.3
Biology	363	32.7	149	31.8	20	42.6
Calf Mortality & Maternity	12	1.1	20	4.3	0	0
Cognition	8	0.7	73	15.6	1	4.3
Conservation, Ecology, & Environment	46	4.1	4	0.9	2	0
Culture	10	0.9	1	0.2	0	0
Echolocation, Sound Detection	29	2.6	78	16.7	6	12.8
Development	13	1.2	12	2.6	0	0
Human-dolphin Interaction	82	7.4	11	2.4	2	4.3
Human-dolphin: Therapy	1	0.1	23	4.9	0	0
Methodology	70	6.3	13	2.8	5	10.6
Sound Production	92	8.3	38	8.1	8	17.0
Social Structure	44	4.0	3	0.6	0	0
Total	1,109	100	468	100	47	100

Note: Percentages above 10% and their frequencies are **bolded** to help identify trends. The highest percentage is also *italicized*.

Representation of research design across habitats

As indicated by a chi-square-test-of-independence, the two variables were related, $\chi^2 (8, N = 1626) = 366.40, p < 0.001$, *Cramer's V* = 0.34. As expected and seen in Table 8, wild research was more often observational whereas captive research was more experimental in nature.

Table 8
Distribution of articles across research type and habitat.

Research Type	Habitat					
	Wild		Captive		Both	
	f	%	f	%	f	%
Observational	579	52.1	100	21.4	18	38.3
Observational-Biology	359	32.3	117	25.0	12	25.5
Experimental-Psychology	27	2.4	120	25.6	6	12.8
Experimental-Biology	82	7.4	121	25.9	10	21.3
Other	64	5.8	10	2.1	1	2.1
Total	1,111	100	468	100	47	100

Note: Percentages above 10% and their frequencies are **bolded** to help identify trends. The highest percentage is also *italicized*. Research type was equally divided across the major categories so the highest percentage was not notated.

Representation of research design across journal type

A chi-square-test-of-independence indicated that research design and journal type were related, $\chi^2 (32, N = 1624) = 334.20, p < 0.001$, *Cramer's V* = 0.23. As summarized in Table 9, sound and acoustic journals were more likely to publish articles using experimental designs involving clear manipulations while psychology journals were more likely to publish articles using experimental research designs with some manipulation. Biology and science journals were more likely to publish studies that used observational designs involving some invasive techniques as compared to marine science journals, which published more observational research.

Representation of article topic across research designs

A chi-square-test-of-independence suggested that the topics were not equally distributed across the designs, $\chi^2 (52, N = 1624) = 2,233.00, p < 0.001$, *Cramer's V* = 0.59. As expected, research on behavior, the effect of anthropogenic noise on cetaceans, sound production, and social structures incorporated observational research designs more often than expected by chance alone. In contrast, research on biology incorporated more invasive observational and experimental designs. Research on echolocation and sound detection also tended to use experimental designs involving more manipulation. Finally, research on cognition and human-dolphin interactions involving therapy incorporated more experimental-psychology based designs than expected by chance alone. See Table 10 for these trends and others.

Table 9
Distribution of articles across journal and research type.

Journal	Research Type									
	Observational		Observational- Biology		Experimental- Psychology		Experimental- Biology		Other	
	f	%	f	%	f	%	f	%	f	%
Biology/ Sciences	133	29.1	209	45.7	28	6.1	71	15.5	16	3.5
Psychology	101	50.0	20	9.9	49	24.3	27	13.4	5	2.5
Animals, Ecology, Zoology	112	57.4	43	22.1	19	9.7	9	4.6	12	6.2
Acoustics	38	25.9	37	25.2	15	10.2	52	35.4	5	3.4
Marine Science	253	52.9	134	28.0	22	4.6	44	9.2	25	5.2
Conservation	30	39.0	32	41.6	3	3.9	3	3.9	9	11.7
Therapy	5	55.6	0	0	4	44.4	0	0	0	0
Human-Dolphin	17	73.9	3	13.0	2	8.7	1	4.3	0	0
Other	9	25.0	8	22.2	11	30.6	6	16.7	2	5.6

Note: Row percentages add to 100%. Percentages above 10% and their frequencies are **bolded** to help identify trends. The highest percentage is also *italicized*. Note that observational research is generally performed and published most frequently across the different journal categories.

Table 10
Distribution of articles across topic and research type.

Topic	Research Type									
	Observational		Observational Biology		Experimental Psychology		Experimental Biology		Other	
	f	%	f	%	f	%	f	%	f	%
Anthropogenic Noise & Distribution	131	92.3	6	4.2	2	1.4	1	0.7	2	1.4
Behavior	198	81.8	22	9.1	19	7.9	3	1.2	0	0
Biology	35	6.6	361	67.9	6	1.1	124	23.3	6	1.1
Calf mortality & Maternity	16	50.0	6	18.8	6	18.8	4	12.5	0	0
Cognition	15	18.3	5	6.1	57	69.5	5	6.1	0	0
Conservation, Ecology, & Environment	34	65.4	10	19.2	3	5.8	2	3.8	3	5.8
Culture	11	100	0	0	0	0	0	0	0	0
Echolocation, Sound Detection	22	19.5	21	18.6	10	8.8	59	52.2	1	0.9

Table 10 (cont.)

Development	16	61.5	5	19.2	2	7.7	2	7.7	1	3.8
Human-dolphin Interaction	54	56.8	18	18.9	18	18.9	0	0	5	5.3
Human-dolphin: Therapy	6	25.0	1	4.2	17	70.8	0	0	0	0
Methodology	19	21.6	5	5.7	4	4.5	6	6.8	54	61.4
Sound Production	100	72.5	24	17.4	6	4.3	6	4.3	2	1.4
Social Structure	41	87.2	2	4.3	3	6.4	1	2.1	0	0

Note: Row percentages add to 100%. Percentages above 10% and their frequencies are **bolded** to help identify trends. The highest percentage is also *italicized*. Note that observational research is generally performed for most research topics. Some notable exceptions are research in cognition, echolocation, and human-dolphin therapy programs, which tended to incorporate experimental paradigms.

Discussion

The purpose of the current study was to explore the publication trends for research involving dolphin and whale species that may be found in captive and wild conditions. An extensive, although not exhaustive, search for relevant literature was conducted using nine research databases. Despite limitations of specific databases regarding their level of access to published articles, the search produced a representative sample of 1,628 unique articles from 290 journals. Three journals in particular dominated the publication record: *Aquatic Mammals* (18.5% of identified articles), *Marine Mammal Science* (*MMS*, 9% of identified articles), and *Journal of the Acoustical Society of America* (*JASA*, about 7% of identified articles). *Aquatic Mammals*, as the oldest publication dedicated to marine mammals, has published almost twice as many articles as *MMS* or *JASA* and eight times as many articles as *Science*, *Journal of Mammalogy*, *Animal Behavior*, *Journal of Experimental Biology*, *Nature*, and *Journal of Comparative Psychology*, the next most frequent journals.

Roughly 20% of all whale and dolphin species were represented in the sample collected with 16 different genera identified (Leatherwood, Randall, & Foster, 1983). The most frequently cited genus was *Tursiops* (bottlenose dolphins), which accounted for 42.9% of all the articles identified. Considering *Tursiops* has a world-wide distribution with abundant populations and is one of the most frequently occurring genera in captivity, this finding is not surprising (Leatherwood et al., 1983; Mann et al., 2000; May, 1998). Again, with a broad geographic range and widespread presence in captivity, the next most frequent genus was *Orcinus* (killer whales, 9.9%). Interestingly, despite the common presence of these two genera in captivity, most of the published literature included data collected from wild populations (*Tursiops* – 23.5% and *Orcinus* – 8.5%). Thus, captive research with *Tursiops* represented 18.1% of all articles and captive research with *Orcinus*, only 1.2% of all articles. In contrast, the two least represented genera, *Feresa* (pygmy killer whale, 0.3%) and *Lagenodelphis* (Fraser's dolphin, 0.4%), are cetaceans which have relatively limited geographic ranges, numbers, captive presence, and thus available information. Clearly, there is a need to increase our knowledge of animals that are not easily observable in the wild. However, there is also a large need to conduct research with captive populations, such as killer whales, so that we may better understand their cognitive and social needs when in the care of humans.

The sample of articles collected indicated that research from around the world is somewhat equally represented. The North Atlantic-Canada (19.0%), North Pacific (15.9%), North Atlantic-United States (14%), Indian (12.6%), and North Atlantic-Europe (11.9%) oceans were the most frequently represented geographic locations across the articles. The Arctic Ocean was the least represented geographic location, accounting for only 1.8% of the articles examined.

As expected, the majority of the published research was conducted with wild populations (68.4%). A mere 3% of the identified articles examined both wild and captive populations. Although research with captive populations is not

published, or perhaps not conducted, as frequently as research with wild populations, it is nowhere as sparse as suggested by Rose and her colleagues (2009), who concluded that research with captive populations was extremely rare (about 5%). This estimate was derived by Rose and her colleagues by examining the abstracts of research presented at the 16th Biennial Conference for the Biology of Marine Mammals in South Africa during 2007, one of the major international conferences for marine mammals. In direct contrast, our findings indicated that 28.6% of our sample incorporated captive populations. Clearly, additional research involving captive populations is needed, but it is not as meager as implicated by the most current report on the case against captivity.³

The 1,628 articles represented over 40 different topics. As summarized in the appendix, these individual topics were grouped into 14 broad topics. Biology, representing many different aspects of physiology and genetics, comprised the most abundant topic, accounting for a third of the sample. Behavior followed with 15% of the sample. Sound production, particularly sounds used for communication purposes, and the production and reception of echolocation were the next two most frequent categories, accounting for about 15% of the sample when combined. The least represented topic in our sample involved studies examining various aspects of culture across various cetaceans, with less than 1% of the articles dedicated to this topic. This result is intriguing as it is one of the few topics that represents a single category of research. Perhaps, this result should be viewed optimistically as it may indicate a more liberal perspective on the complex behaviors and traditions shared by some non-human species (see Rendell & Whitehead, 2001 for a relatively recent review of this topic).

Surprisingly, less than 4% of the articles investigated topics involving mother-calf interactions and behaviors and only 5% of the articles involved studies pursuing cognitive research. Considering we specifically selected genera that could be found in both wild and captive settings, it is interesting that few studies have examined some of the most important components of an animal's life – development and offspring care and cognitive abilities. The paucity of available publications may be due to a lack in the research itself or in an inability to publish such studies in the peer-reviewed arena. Many of these studies involve relatively small sample sizes examined over long periods of time, and traditionally studies involving small sample sizes can be difficult to publish. Additional research, such as the sample size found across publications, is needed to better understand the reasons behind these less researched areas.

As indicated earlier, the results demonstrated that observational studies were the most abundant type of designs published in our sample, accounting for 43% of the articles. Observational studies involving slightly more invasive or hands-on techniques comprised 30% of the sample. In contrast, studies employing more experimental manipulations or research designs accounted for approximately

³A brief review of the abstracts from the 17th Biennial Conference for the Biology of Marine Mammals in Canada also indicated that 5% of the presented research was conducted solely in captivity.

22.5% of the articles. This trend is not surprising given the prevalence of the topics described above and the difficulties of conducting experimental studies.

Several clear relationships emerged suggesting that the type of research conducted depended on the habitat and that certain categories of journals published specific types of research with regard to habitat and research design. First, most observational studies were conducted in the wild with almost twice as many studies incorporating purely observational techniques (83%) or observational techniques with some invasion or manipulation (74%) as compared to experimental paradigms. In contrast, more experimental designs were conducted in captivity (78.4% - psychology and 56.8% - biological). Interestingly, an equal number of articles existed across the four major research designs for research conducted in captivity whereas the articles involving wild populations clearly emphasized observational forms of research with a limited number of experimental forms of research. This trend is not surprising, given the difficulties of conducting well-constructed experimental studies in the field. However, experimental studies are possible in the field and should be attempted more often (e.g., playback studies).

A second trend indicated that some categories of journals were more likely to publish research conducted in specific habitats. Specifically, journals publishing research on the marine sciences, conservation, and animals, their ecology, and zoology were more likely to publish research conducted in the wild than in captivity. In fact, these three categories of journals published only 108 articles (about 7%) in which the samples were drawn from captive populations only. In comparison, the journals emphasizing the marine sciences, conservation, and animals, their ecology and zoology published five times as many articles (38.5%) involving wild populations. If researchers are interested in publishing research conducted in captivity, then the best opportunities for publication can be found in psychology, acoustic, and miscellaneous journals. In fact, 7% of all the articles involving captive populations were published by psychology journals. Journals publishing acoustic research also published research performed in captivity more often than other journals (i.e., about 5% of all the articles). These trends are not unexpected given the research scopes of each of the journals included in these categories and fit well with the relationship between these journal categories and type of research design used.

As described earlier, observational designs were the most frequent type of design observed across articles. However, the journal categories differed in their representation of the two types of observational research. That is, the topics of marine sciences (15.6% of the articles), animals, ecology, and zoology (about 7% of the articles), psychology (about 6% of the articles), and human-dolphin interactions (about 1% of the articles) most often involved the use of strict observational techniques. The biology/sciences (about 13% of the articles) and conservation (about 2% of the articles) categories of journals most often published observational studies in which more manipulation or invasion occurred. Similarly, experimental studies were most often published by psychology-based (about 3% of

the articles) and acoustic-based (about 3% of the articles) journals, as would be expected.

When the findings associated with journal type and habitat and the findings associated with journal type and research design are examined together, a dilemma for researchers conducting studies with captive populations emerges. First, studies conducted with captive populations are published or conducted half as often as studies conducted with their wild counterparts. Second, journals are more likely to publish experimental research than observational research from captive populations, presumably due to the greater confidence in the research findings under controlled contexts. These two findings may be best explained by the limited opportunities researchers face when conducting research in a captive setting. Research in captivity involves overcoming many competing demands (e.g., availability of animals, training time, and monetary support) and working within the goals of the facility (e.g., education, animal interaction, and entertainment). These limitations pose major obstacles for researchers interested in captive populations and make experimental paradigms very challenging. Given the difficulty of completing an experimental study, an observational research approach seems more plausible and efficient. Unfortunately, the ecological validity of a single study conducted with a population of animals at a particular facility is often questioned.

While many facilities have attempted to provide opportunities to conduct research, only a handful of researchers have successfully created research programs centered on a captive population. What characteristics make these research programs successful? Anecdotal evidence suggests that these captive populations had some animals that were dedicated for research purposes, but a formal assessment is necessary.

As one would expect, some topics were pursued more often with wild populations while other topics were pursued more often with captive populations. For example, the effects of anthropogenic noise on dolphins and whales, the behavior of dolphins or whales, interactions between humans and dolphins, social structure, methodology, and conservation were researched primarily in the wild and represented more than a third of the sample. For research conducted in captivity, the most frequently researched topics included cognition, echolocation production and reception, the use of dolphins as a form of therapy for humans, calf survival and maternal care, and development. These topics accounted for only an eighth of the sample. Perhaps the most interesting finding of this analysis was that research on the production of sounds used for communication was equally represented across the wild and captive habitats and, in fact, had significantly more studies conducted in both habitats within the same study than any other topic. As this type of research is often attempting to determine the function of various sounds within specific contexts, conducting research in both field and controlled settings is extremely important in cross-validating conclusions (Harley, 2008; Mann et al., 2000).

Finally, the analyses corroborated the expectation that certain topics are examined more often with specific research designs. The topics with the greatest

representation of observational techniques were behavior (12% of the articles), the effect of anthropogenic noise on cetaceans and cetacean distributions (8% of the articles), sound production (6% of the articles), and social structure (2.5% of the articles). Cetacean biology research most frequently involved observational techniques requiring more manipulation and invasion, which constituted almost a quarter of the entire sample. Experimental studies using more psychology-based research designs most often included research on cognition (3.5% of the articles) and the use of dolphins as therapy for humans (1% of the articles). Experimental studies from a biological perspective included echolocation (3.6% of the articles) and biological research (7.6% of the articles). While these findings are not surprising, researchers should be aware of the different types of research designs utilized for various topics. Certainly, some topics are better suited for observational research and some topics require more invasive manipulations to fully understand them. However, more progress may be possible if a mixed methods approach was utilized more often. To be successful, cooperation between researchers, facilities, and the federal government is a necessity.

In summary, we explored the trends in the publication of marine mammal research, with an emphasis on dolphin and whale species that can be found in both wild and captive settings. The overall findings suggested that a large number of peer-reviewed articles have been published on a variety of topics in a number of different journals. Marine mammal researchers have a variety of journals to which to submit their research. However, many of these journals publish research conducted in the wild more often than research conducted in captivity. Whether this trend is due to a journal preference for more ecologically valid research, the lack of opportunity to conduct research at captive facilities, the availability of financial resources for different research questions, or simply the preference of researchers themselves, additional research is necessary. It is important to continue to examine animals in their natural environments. However, there is much to be learned from studies conducted with animals in captivity and more efforts should be made by researchers, facilities, and funding agencies alike.

Despite the apparent interest in cetaceans, there are clearly preferences in the pursuit of topics. Unfortunately, some very important topics have been grossly under-represented (e.g., cognition, development, calf survival, social structure, and conservation) while other topics are well-represented (e.g., biological features, behavior, sounds production, echolocation). Again, it is not clear if these trends simply reflect the current interests of scientists or the current sources of funding, but more attention should be focused on the under-represented topics. We experienced the first extinction of an extant cetacean during our lifetime, the baiji, or Yangtze river dolphin (*Lipotes vexillifer*) (see Turvey, 2008 for a review). While this event may not have been preventable, ultimately, perhaps an awareness of the paucity of research in certain arenas will spur individuals within the scientific community to shift their interests (Smith, 2009).

As a group, we must become cognizant of the publication trends which currently describe our research progress so that we may better integrate our knowledge from captivity and the wild. Researchers should consider all

publication sources, such as book chapters, reviews, popular press articles, or unpublished theses and dissertations when developing their studies. We also suggest that those individuals involved in the peer-review process for journals be open-minded to the advantages of both large and small sample studies investigating topics that are less represented. Future studies investigating publication trends should examine the variation in sample sizes and their relation to the journals publishing the studies, the research design used, the population examined, and the topic pursued. While our sample may not contain every study ever published on a dolphin or whale species found in captivity and the wild, we believe that it was representative and provides the field with an accurate description of the current state of peer-reviewed published research. We hope that others may find this review helpful in identifying areas of future research as well as areas that could be improved within the publication process.

References

Note: Bolded references represent citations used to support the premise and findings of the paper. References in regular font include those that were used in the sample from which the statistics were calculated.

- (1987). An idea in space. *Futurist*, 21(1), 32.
- (1988). News and notes. *British Medical Journal (Clinical Research Ed)*, 296, 1742.
- (1992). Dolphin death report raises environmental ire. *Nutrition Week*, 22(10), 2-3.
- (1995). Satellites monitor whale whereabouts. *Science Teacher*, 62(2), 12.
- (1996). Are we being too loud? *Environment*, 38(1), 24.
- (2001). Color blindness in whales. *Science*, 292, 399.
- (2002). Ocean noise affects marine life. *Environment*, 44(5), 4.
- (2003). Flipper or Rambo? *Science Teacher*, 70(5), 12.
- (2004). War's environmental impact. *Alternatives Journal*, 30(4), 26.
- (2005). Dolphin school. *Alternatives Journal*, 31(3), 4.
- (2005). In case you haven't heard.... *Mental Health Weekly*, 15(47), 8.
- (2005). Sleepless in Sea World. *Pediatrics*, 116(3), 648.
- (2006). Beluga whales maintain numbers in Cook Inlet, Alaska. *Oryx*, 40(3), 252.
- (2006). Effects of lactoferrin on immune function in dolphins. *Veterinary Record: Journal of the British Veterinary Association*, 159, 865.
- (2007). Multi-locus phylogeography of the dusky dolphin (*Lagenorhynchus obscurus*): Passive dispersal via the west-wind drift or response to prey species and climate change? *BMC Evolutionary Biology*, 7131-147.
- (2008). Orcas and oil. *Science*, 320, 297.
- (2009). A whale of a fetus. *Science*, 323, 857.
- (2009). Animal acoustics: This whale goes to 11. *Nature*, 457, 515.
doi:10.1038/457515c
- Acevedo-Gutierrez, A. (1999). Aerial behavior is not a social facilitator in bottlenose dolphins hunting in small groups. *Journal of Mammalogy*, 80, 768-776.
- Adams, J., Houde, M., Muir, D., Speakman, T., Bossart, G., & Fair, P. (2008). Land use and the spatial distribution of perfluoroalkyl compounds as measured in the plasma of bottlenose dolphins (*Tursiops truncatus*). *Marine Environmental Research*, 66, 430-437. doi:10.1016/j.marenvres.2008.07.004

- Adams, J. D., Speakman, T., Zolman, E., & Schwacke, L. H. (2006). Automating image matching, cataloging, and analysis for photo-identification research. *Aquatic Mammals*, 32, 374-384.
- Adler, L., & Adler, H. (1990). Matching to sample by Atlantic bottlenose dolphins (*Tursiops truncatus*): Chinese characters and English words. *Annals of the New York Academy of Sciences*, 60, 113-126. doi:10.1111/j.1749-6632.1990.tb22733.x
- Adams, L., & Rosel, P. (2006). Population differentiation of the Atlantic spotted dolphin (*stenella frontalis*) in the western North Atlantic, including the Gulf of Mexico. *Marine Biology*, 148, 671-681. doi:10.1007/s00227-005 0094-2
- Adolphs, R. (2009). The social brain: Neural basis of social knowledge. *Annual Review of Psychology*, 60, 693-716.
- Aguilar, A., & Borrell, A. (2005). DDT and PCB reduction in the western Mediterranean from 1987 to 2002, as shown by levels in striped dolphins (*Stenella coeruleoalba*). *Marine Environmental Research*, 59, 391-404. doi:10.1016/j.marenvres.2004.06.004
- Aguilar Soto, N., Johnson, M., Madsen, P., Díaz, F., Domínguez, I., Brito, A., et al. (2008). Cheetahs of the deep sea: deep foraging sprints in short-finned pilot whales off Tenerife (Canary Islands). *Journal of Animal Ecology*, 77, 936-947. doi:10.1111/j.1365-2656.2008.01393.x
- Ainley, D., Ballard, G., & Olmastroni, S. (2009). An apparent decrease in the prevalence of "Ross Sea Killer Whales" in the Southern Ross Sea. *Aquatic Mammals*, 35, 334-346. doi:10.1578/AM.35.3.2009.334
- Ainley, D. G., Ballard, G., Dugger, K. M. (2006). Competition among penguins and cetaceans reveals trophic cascades in the Western Ross Sea, Antarctica. *Ecology*, 87, 2080-2093.
- Ainley, D. G., Dugger, K. M., Toniolo, V., & Gaffney, I. (2007). Cetacean occurrence patterns in the Amundsen and southern Bellingshausen sea sector, Southern Ocean. *Marine Mammal Science*, 23, 287-305. doi:10.1111/j.1748-7692.2007.00109.x
- Akamatsu, T., Ding, W., & Kexiong, W. (2005). Off-axis sonar beam pattern of free-ranging finless porpoises measured by a stereo pulse event data logger. *Journal of the Acoustical Society of America*, 117, 3325-3330. doi:10.1121/1.1893325
- Akamatsu, T., Nakazawa, I., Tsuchiyama, T., & Kimura, N. (2008). Evidence of nighttime movement of finless porpoises through Kanmon Strait monitored using a stationary acoustic recording device. *Fisheries Science*, 74, 970-975. doi:10.1111/j.1444-2906.2008.01614.x
- Akamatsu, T., Wang, D., Nakamura, K., & Wang, K. (1998). Echolocation range of captive and free-ranging baiji (*Lipotes vexillifer*), finless porpoise (*Neophocaena phocaenoides*), and bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 104, 2511-2516. doi:10.1121/1.423757.
- Akamatsu, T., Wang, D., Wang, K., Li, S., Dong, S., Zhao, X., et al. (2008). Estimation of the detection probability for Yangtze finless porpoises (*Neophocaena phocaenoides asiaeorientalis*) with a passive acoustic method. *Journal of the Acoustical Society of America*, 123, 4403-4411. doi:10.1121/1.2912449
- Allen, K. (1983). Development and application of cetacean population models. *Advance in Applied Biology*, 7, 333-405.
- Allen, M., & Read, A. (2000). Habitat selection of foraging bottlenose dolphins in relation to boat density near Clearwater, Florida. *Marine Mammal Science*, 16, 815-824.

- Altes, R., Dankiewicz, L., Moore, P., & Helweg, D. (2003). Multiecho processing by an echolocating dolphin. *Journal of the Acoustical Society of America*, 114, 1155-1166. doi:10.1121/1.1590969
- Alvarez-Flores, C., & Heide-Jørgensen, M. (2004). A risk assessment of the sustainability of the harvest of beluga (*Delphinapterus leucas* (Pallas 1776)) in West Greenland. *ICES Journal of Marine Science / Journal du Conseil*, 61, 274-286. doi:10.1016/j.icesjms.2003.12.004
- Amaral, A., Sequeira, M., Martínez-Cedeira, J., & Coelho, M. (2007). New insights on population genetic structure of *Delphinus delphis* from the northeast Atlantic and phylogenetic relationships within the genus inferred from two mitochondrial markers. *Marine Biology*, 151, 1967-1976. doi:10.1007/s00227-007-0635-y
- Amato, I. (1993). A sub surveillance network becomes a window on whales. *Science*, 261, 549.
- Ames, A., Van Vleet, E., & Reynolds III, J. (2002). Comparison of lipids in selected tissues of the Florida manatee (Order Sirenia) and bottlenose dolphin (Order Cetacea; Suborder Odontoceti). *Comparative Biochemistry & Physiology Part B*, 132, 625.
- Ames, M. (1991). Saving some cetaceans may require breeding in captivity. *Bioscience*, 41, 746-749.
- Amir, O., Berggren, P., Ndaro, S., & Jiddawi, N. (2005). Feeding ecology of the Indo-Pacific bottlenose dolphin (*Tursiops aduncus*) incidentally caught in the gillnet fisheries off Zanzibar, Tanzania. *Estuarine Coastal & Shelf Science*, 63, 429-437. doi:10.1016/j.ecss.2004.12.006.
- Amundin, M., Starkhammar, J., Evander, M., Almqvist, M., Lindström, K., & Persson, H. (2008). An echolocation visualization and interface system for dolphin research. *Journal of the Acoustical Society of America*, 123, 1188-1194. doi:10.1121/1.2828213
- Andersen, G., Førefid, S., Skaare, J., Jenssen, B., Lydersen, C., & Kovacs, K. (2006). Levels of toxaphene congeners in white whales (*Delphinapterus leucas*) from Svalbard, Norway. *Science of the Total Environment*, 357, 128-137. doi:10.1016/j.scitotenv.2005.05.014
- Anderson, J. (2002). Gone fishing: tool use in animals. *Biologist*, 49, 15.
- Andre, M., Delory, E., Degollada, E., Alonso, J.-M., del Rio, J., van der Schaar, M., et al. (2007). Identifying cetacean hearing impairment at stranding sites. *Aquatic Mammals*, 33, 100-109.
- Andre, M., & Nachtigall, P. E. (2007). Electrophysiological measurements of hearing in marine mammals. *Aquatic Mammals*, 33, 1-5.
- Andrew, J. R., Danielle M. W., Kim W. U., & Dave, S. (2003). Fine-scale behaviour of bottlenose dolphins around gillnets. *Proceedings: Biological Sciences*, 27090.
- Andrews, R., Pitman, R., & Ballance, L. (2008). Satellite tracking reveals distinct movement patterns for Type B and Type C killer whales in the southern Ross Sea, Antarctica. *Polar Biology*, 31, 1461-1468. doi:10.1007/s00300-008-0487-z
- Antonioli, C., & Reveley, M. (2005). Randomised controlled trial of animal facilitated therapy with dolphins in the treatment of depression. *BMJ: British Medical Journal*, 331, 1231-1231. doi:10.1136/bmj.331.7527.1231
- Anwar, S., Jeanneret, C., Parrott, L., & Marceau, D. (2007). Conceptualization and implementation of a multi-agent model to simulate whale-watching tours in the St. Lawrence Estuary in Quebec, Canada. *Environmental Modelling & Software*, 22, 1775-1787. doi:10.1016/j.envsoft.2007.02.007
- Aran Mooney, T., Nachtigall, P., Taylor, K., Rasmussen, M., & Miller, L. (2009). Auditory temporal resolution of a wild white-beaked dolphin (*Lagenorhynchus albirostris*

- albirostris*). *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural & Behavioral Physiology*, 195, 375-384. doi:10.1007/s00359-009 0415-x
- Archer, F., Gerrodette, T., Chivers, S., & Jackson, A. (2004). Annual estimates of the unobserved incidental kill of pantropical spotted dolphin (*Stenella attenuata attenuata*) calves in the tuna purse-seine fishery of the eastern tropical Pacific. *Fishery Bulletin*, 102, 233-244.
- Archer, F., Gerrodette, T., Dizon, A., Abella, K., & Southern, S. (2001). Unobserved kill of nursing dolphin calves in a tuna purse-seine fishery. *Marine Mammal Science*, 17, 540-554.
- Archer, F., & Robertson, K. M. (2004). Age and length at weaning and development of diet of pantropical spotted dolphins, *Stenella attenuata*, from the Eastern tropical Pacific. *Marine Mammal Science*, 20, 232-245.
- Arronte, J. C., Pis-Millan, J. A., & Perez, C. (2005). Injury to an Atlantic white-sided dolphin (*Lagenorhynchus albirostris acutus*) caused by needlefish impalement. *Aquatic Mammals*, 31, 184-186.
- Atem, A., Rasmussen, M., Wahlberg, M., Petersen, H., & Miller, L. (2009). Changes in click source levels with distance to targets: studies of free-ranging white-beaked dolphins *lagenorhynchus albirostris* and captive harbor porpoises *phocoena phocoena*. *Bioacoustics*, 19, 49-65.
- Atkins, S., Pillay, N., & Peddemors, V. M. (2004). Spatial distribution of Indo-Pacific humpback dolphins (*Sousa chinensis*) at Richards Bay, South Africa: Environmental influences and behavioural patterns. *Aquatic Mammals*, 30, 84-93.
- Au, D., & Weihs, D. (1980). At high speeds dolphins save energy by leaping. *Nature*, 284, 548-550. doi:10.1038/284548a0
- Au, W. (1994). Comparison of sonar discrimination: Dolphin and an artificial neural network. *Journal of the Acoustical Society of America*, 95, 2728-2735. doi:10.1121/1.409841
- Au, W. (2004). Echolocation Signals of Wild Dolphins. *Acoustical Physics*, 50, 454-462. doi:10.1134/1.1776224
- Au, W., Andersen, L., Rasmussen, A., & Roitblat, H. (1995). Neural network modeling of a dolphin's sonar discrimination capabilities. *Journal of the Acoustical Society of America*, 98, 43-50. doi:10.1121/1.413700
- Au, W., & Benoit-Bird, K. (2003). Automatic gain control in the echolocation system of dolphins. *Nature*, 423, 861.
- Au, W., & Benoit-Bird, K. (2008). Broadband backscatter from individual Hawaiian mesopelagic boundary community animals with implications for spinner dolphin foraging. *Journal of the Acoustical Society of America*, 123, 2884-2894. doi:10.1121/1.2902187
- Au, W., Branstetter, B., Benoit-Bird, K., & Kastelein, R. (2009). Acoustic basis for fish prey discrimination by echolocating dolphins and porpoises. *Journal of the Acoustical Society of America*, 126, 460-467. doi:10.1121/1.3147497
- Au, W., Ford, J., Horne, J., & Allman, K. (2004). Echolocation signals of free-ranging killer whales (*Orcinus orca*) and modeling of foraging for chinook salmon (*Oncorhynchus tshawytscha*). *Journal of the Acoustical Society of America*, 115, 901-909. doi:10.1121/1.1642628
- Au, W., & Herzing, D. (2003). Echolocation signals of wild Atlantic spotted dolphin (*Stenella frontalis*). *Journal of the Acoustical Society of America*, 113, 598-604. doi:10.1121/1.1518980

- Au, W., Kastelein, R., Benoit-Bird, K., Cranford, T., & McKenna, M. (2006). Acoustic radiation from the head of echolocating harbor porpoises (*Phocoena phocoena*). *Journal of Experimental Biology*, 209, 2726-2733. doi:10.1242/jeb.02306
- Au, W., Lemonds, D., Vlachos, S., Nachtigall, P., & Roitblat, H. (2002). Atlantic bottlenose dolphin (*Tursiops truncatus*) hearing threshold for brief broadband signals. *Journal of Comparative Psychology*, 116, 151-157. doi:10.1037/0735-7036.116.2.151
- Au, W., Moore, P., & Pawloski, D. (1988). Detection of complex echoes in noise by an echolocating dolphin. *Journal of the Acoustical Society of America*, 83, 662-668. doi:10.1121/1.396161
- Au, W., & Pawloski, J. (1989). Detection of noise with rippled spectra by the Atlantic bottlenose dolphin. *Journal of the Acoustical Society of America*, 86, 591-596. doi:10.1121/1.398238
- Au, W., Thomas, J. A., & Ramirez, K. T. (2007). Characteristics of the auditory brainstem evoked potential of a Pacific white-sided dolphin (*Lagenorhynchus albirostris obliquidens*). *Aquatic Mammals*, 33, 76-84.
- Au, W., & Turl, C. (1991). Material composition discrimination of cylinders at different aspect angles by an echolocating dolphin. *Journal of the Acoustical Society of America*, 89, 2448-2451. doi:10.1121/1.400930
- Au, W., Vlachos, S., Lemonds, D., Nachtigall, P., & Roitblat, H. (2002). Atlantic bottlenose dolphin (*Tursiops truncatus*) hearing threshold for brief broadband signals. *Journal of Comparative Psychology*, 116, 151-157.
- Au, W., & Würsig, B. (2004). Echolocation signals of dusky dolphins (*Lagenorhynchus obscurus*) in Kaikoura, New Zealand. *Journal of the Acoustical Society of America*, 115, 2307-2313. doi:10.1121/1.1690082
- Aubauer, R., Au, W., Nachtigall, P., Pawloski, D., & DeLong, C. (2000). Classification of electronically generated phantom targets by an Atlantic bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 107, 2750-2754. doi:10.1121/1.428661
- Azevedo, A., Lailson-Brito, J., Jr., Siciliano, S., Cunha, H. A., & Fragoso, A. B. L. (2003). Colour pattern and external morphology of the Fraser's dolphin (*Lagenodelphis hosei*) in the Southwestern Atlantic. *Aquatic Mammals*, 29, 411-416.
- Azevedo, A., Oliveira, A., Dalla Rosa, L., & Lailson-Brito, J. (2007). Characteristics of whistles from resident bottlenose dolphins (*Tursiops truncatus*) in southern Brazil. *Journal of the Acoustical Society of America*, 121, 2978-2983. doi:10.1121/1.2713726
- Azzellino, A., Gaspari, S., Airoldi, S., & Nani, B. (2008). Habitat use and preferences of cetaceans along the continental slope and the adjacent pelagic waters in the western Ligurian Sea. *Deep-Sea Research Part I, Oceanographic Research Papers*, 55, 296-323. doi:10.1016/j.dsr.2007.11.006
- Bagdonas, A., Bel'kovich, V., & Krushinskaya, N. (1970). Interaction of the analyzers in the white-sided dolphin in discrimination of geometric figures under water. *Zhurnal Vysshhei Nervnoi Deyatel'nosti*, 20, 1070-1075.
- Bailey, H., & Thompson, P. (2006). Quantitative analysis of bottlenose dolphin movement patterns and their relationship with foraging. *Journal of Animal Ecology*, 75, 456-465. doi:10.1111/j.1365-2656.2006.01066.x
- Baird, R., & Gorgone, A. (2005). False killer whale dorsal fin disfigurements as a possible indicator of long-line fishery interactions in Hawaiian Waters. *Pacific Science*, 59, 593-601.

- Baird, R., Gorgone, A. M., McSweeney, D. J., Ligon, A. D., Deakos, M. H., Webster, D. L., et al. (2009). Population structure of island-associated dolphins: Evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian islands. *Marine Mammal Science*, 25, 251-274. doi:10.1111/j.1748-7692.2008.00257.x
- Baird, R., Hanson, M., & Dill, L. (2005). Factors influencing the diving behaviour of fish-eating killer whales: sex differences and diel and interannual variation in diving rates. *Canadian Journal of Zoology*, 83, 257-267.
- Baird, R., Ligon, A., Hooker, S., & Gorgone, A. (2001). Subsurface and nighttime behaviour of pantropical spotted dolphins in Hawai'i. *Canadian Journal of Zoology*, 79, 988-987. doi:10.1139/cjz-79-6-988
- Baird, R., McSweeney, D., Bane, C., Barlow, J., Salden, D., Antoine, L., et al. (2006). Killer whales in Hawaiian Waters: Information on population identity and feeding habits. *Pacific Science*, 60, 523-530.
- Baird, R., Schorr, G., Ligon, A., McSweeney, D., Webster, D., & Mahaffy, S. (2008). Site fidelity and association patterns in a deep-water dolphin: Rough-toothed dolphins (*Steno bredanensis*) in the Hawaiian Archipelago. *Marine Mammal Science*, 24, 535-553.
- Baldwin, R. M., Collins, M., Van Waerebeek, K., & Minton, G. (2004). The Indo-Pacific humpback dolphin of the Arabian region: A status review. *Aquatic Mammals*, 30, 111-124.
- Ballance, L. (1992). Habitat use patterns and ranges of the bottlenose dolphin in the Gulf of California, Mexico. *Marine Mammal Science*, 8, 262-274.
- Ballance, L., & Pitman, R. (1998). Cetaceans of the western tropical Indian ocean: Distribution, relative abundance, and comparisons with cetacean communities of two other tropical ecosystems. *Marine Mammal Science*, 14, 429-459.
- Barco, S., Swingle, W., McLellan, W., Harris, R., & Pabst, D. (1999). Local abundance and distribution of bottlenose dolphins (*Tursiops truncatus*) in the nearshore waters of Virginia Beach, Virginia. *Marine Mammal Science*, 15, 394-408.
- Barlow, J., & Forney, K. (2007). Abundance and population density of cetaceans in the California Current ecosystem. *Fishery Bulletin*, 105, 509-526.
- Baron, S., Martinez, A., Garrison, L., & Keith, E. (2008). Differences in acoustic signals from Delphinids in the western North Atlantic and northern Gulf of Mexico. *Marine Mammal Science*, 24, 42-56.
- Barrett-Lennard, L., Ford, J., & Heise, K. (1996). The mixed blessing of echolocation: Differences in sonar use by fish-eating and mammal-eating killer whales. *Animal Behaviour*, 51, 553-565. doi:10.1006/anbe.1996.0059
- Barros, N. B., Jefferson, T. A., & Parsons, E. C. M. (2004). Feeding habits of Indo-Pacific humpback dolphins (*Sousa chinensis*) stranded in Hong Kong. *Aquatic Mammals*, 30, 179-188.
- Barros, N. B., Parsons, E. C. M., & Jefferson, T. A. (2000). Prey of offshore bottlenose dolphins from the South China Sea. *Aquatic Mammals*, 26, 2-6.
- Barros, N. B., & Wells, R. (1998). Prey and feeding patterns of resident bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, *Journal of Mammalogy*, 79, 1045.
- Barton, R. (2006). Animal communication: Do dolphins have names? *Current Biology*, 16, R598-R599. doi:10.1016/j.cub.2006.07.002
- Bateson, M. (2007). Environmental noise and decision making possible implications of increases in anthropogenic noise for information processing in marine mammals. *International Journal of Comparative Psychology*, 20, 169-178.

- Bauer, G., & Johnson, C. (1994). Trained motor imitation by bottlenose dolphins (*Tursiops truncatus*). *Perceptual and Motor Skills*, 79, 1307-1315.
- Bazua-Duran, C. (2005). The acoustic repertoire of bottlenose dolphins (*Tursiops truncatus*) from the southern Gulf of Mexico. *Journal of the Acoustical Society of America*, 117, 2469.
- Bazua-Duran, C., & Au, W. W. L. (2004). Geographic variations in the whistles of spinner dolphins (*Stenella longirostris*) of the Main Hawaiian Islands. *Journal of the Acoustical Society of America*, 116, 3757-3769.
- Beach, F. (1974). Spatial habit reversal in two species of marine mammals. *The Psychological Record*, 24, 385-391.
- Beach, F., & Herman, L. (1972). Preliminary studies of auditory problem solving and intertask transfer by the bottlenose dolphin. *The Psychological Record*, 22, 49-62.
- Beach, F., & Pepper, R. (1972). Operant responding in the bottlenose dolphin (*Tursiops truncatus*). *Journal of the Experimental Analysis of Behavior*, 17, 159-160. doi:10.1901/jeab.1972.17-159
- Bearzi, M. (2005). Dolphin sympatric ecology. *Marine Biology Research*, 1, 165-175. doi:10.1080/17451000510019132
- Bearzi, G., Fortuna, C., & Reeves, R. (2009). Ecology and conservation of common bottlenose dolphins *Tursiops truncatus* in the Mediterranean Sea. *Mammal Review*, 39, 92-123. doi:10.1111/j.1365-2907.2008.00133.x
- Bearzi, G., Politi, E., Agazzi, S., & Azzellino, A. (2006). Prey depletion caused by overfishing and the decline of marine megafauna in eastern Ionian Sea coastal waters (central Mediterranean). *Biological Conservation*, 127, 373-382. doi:10.1016/j.biocon.2005.08.017
- Bearzi, G., Reeves, R., Notarbartolo-Di-Sciara, G., Politi, E., Cañadas, A., Frantzis, A., et al. (2003). Ecology, status and conservation of short-beaked common dolphins *Delphinus delphis* in the Mediterranean Sea. *Mammal Review*, 33, 224. doi:10.1046/j.1365-2907.2003.00032.x
- Bearzi, M., & Stanford, C. (2007). Dolphins and African apes: comparisons of sympatric socio ecology. *Contributions to Zoology*, 76, 235-254.
- Bearzi, M., Rapoport, S., Chau, J., & Saylan, C. (2009). Skin lesions and physical deformities of coastal and offshore common bottlenose dolphins (*Tursiops truncatus*) in Santa Monica Bay and adjacent areas, California. *AMBIO - A Journal of the Human Environment*, 38, 66-71.
- Beasley, I. L., & Davidson, P. J. A. (2007). Conservation status of marine mammals in Cambodian waters, including seven new cetacean records of occurrence. *Aquatic Mammals*, 33, 368-379.
- Beck, B., & Rice, C. (2003). Serum antibody levels against select bacterial pathogens in Atlantic bottlenose dolphins, *Tursiops truncatus*, from Beaufort, NC, USA and Charleston Harbor, Charleston, SC, USA. *Marine Environmental Research*, 55, 161.
- Bechdel, S., Mazzoil, M., Murdoch, M., Howells, E., Reif, J., McCulloch, S., et al. (2009). Prevalence and impacts of motorized vessels on bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida. *Aquatic Mammals*, 35, 367-377. doi:10.1578/AM.35.3.2009.367
- Becker, P., Krahm, M., Mackey, E., Schantz, M., Epstein, M., Porter, B., et al. (2000). Concentrations of polychlorinated biphenyls (PCB's) chlorinated pesticides and heavy metals and other elements in tissues of belugas from Cook Inlet, Alaska. *Marine Fisheries Review*, 62(3), 81-99.

- Bedjer, L., Samuels, A., Whitehead, H., Gales, N., Mann, J., Connor, R., et al. (2006). Decline in relative abundance of bottlenose dolphins exposed to long-term disturbance. *Conservation Biology*, 20, 1791-1798. doi:10.1111/j.1523-1739.2006.00540.x
- Beedholm, K., & Miller, L. A. (2007). Automatic gain control in harbor porpoises (*Phocoena phocoena*)? Central versus peripheral mechanisms. *Aquatic Mammals*, 33, 69-75.
- Behler, D. (2000). Snow Falling on Water. *Wildlife Conservation*, 103(3), 22.
- Beineke, A., Siebert, U., McLaghlan, M., Bruhn, R., Thron, K., Failing, K., et al. (2005). Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbor porpoises (*Phocoena phocoena*). *Environmental Science & Technology*, 39, 3933-3938. doi:10.1021/es048709j
- Beineke, A., Siebert, U., Müller, G., & Baumgärtner, W. (2007). Increased blood interleukin-10 mRNA levels in diseased free-ranging harbor porpoises (*Phocoena phocoena*). *Veterinary Immunology & Immunopathology*, 115, 100-106. doi:10.1016/j.vetimm.2006.09.006
- Beineke, A., Siebert, U., Stott, J., Müller, G., & Baumgärtner, W. (2007). Phenotypical characterization of changes in thymus and spleen associated with lymphoid depletion in free-ranging harbor porpoises (*Phocoena phocoena*). *Veterinary Immunology & Immunopathology*, 117, 254-265. doi:10.1016/j.vetimm.2007.03.009
- Bejder, L., Fletcher, D., & Bräger, S. (1998). A method for testing association patterns of social animals. *Animal Behaviour*, 56, 719-725. doi:10.1006/anbe.1998.0802
- Bejder, L., Samuels, A., Whitehead, H., & Gales, N. (2006). Interpreting short-term behavioural responses to disturbance within a longitudinal perspective. *Animal Behaviour*, 72, 1149-1158. doi:10.1016/j.anbehav.2006.04.003
- Belikov, R., & Bel'kovich, V. (2006). High-pitched tonal signals of beluga whales (*Delphinapterus leucas*) in a summer assemblage off Solovetskii Island in the White Sea. *Acoustical Physics*, 52, 125-131. doi:10.1134/S1063771006020023
- Belikov, R., & Bel'kovich, V. (2007). Whistles of beluga whales in the reproductive gathering off Solovetskii Island in the White Sea. *Acoustical Physics*, 53, 528-534. doi:10.1134/S1063771007040148
- Belikov, R., & Bel'kovich, V. (2008). Communicative pulsed signals of beluga whales in the reproductive gathering off Solovetskii Island in the White Sea. *Acoustical Physics*, 54, 115-123. doi:10.1007/s11441-008-1016-x
- Bel'kovich, V., & Kreichi, S. (2004). Specific features of vowel like signals of white whales. *Acoustical Physics*, 50, 288-294. doi:10.1134/1.1739497
- Bellante, A., Sprovieri, M., Buscaino, G., Manta, D., Buffa, G., Di Stefano, V., et al. (2009). Trace elements and vanadium in tissues and organs of five species of cetaceans from Italian coasts. *Chemistry & Ecology*, 25, 311-323. doi:10.1080/02757540903193155
- Bender, C., Herzing, D., & Bjorklund, D. (2009). Evidence of teaching in Atlantic spotted dolphins (*Stenella frontalis*) by mother dolphins foraging in the presence of their calves. *Animal Cognition*, 12, 43-53. doi:10.1007/s10071-008-0169-9
- Benoit-Bird, K. (2004). Prey caloric value and predator energy needs: foraging predictions for wild spinner dolphins. *Marine Biology*, 145, 435-444. doi:10.1007/s00227-004-1339-1
- Benoit-Bird, K., & Au, W. (2003). Prey dynamics affect foraging by a pelagic predator (*Stenella longirostris*) over a range of spatial and temporal scales. *Behavioral Ecology and Sociobiology*, 53, 364-373.

- Benoit-Bird, K., & Au, W. (2009a). Cooperative prey herding by the pelagic dolphin, *Stenella longirostris*. *Journal of the Acoustical Society of America*, 125, 125-137. doi:10.1121/1.2967480
- Benoit-Bird, K., & Au, W. (2009b). Phonation behavior of cooperatively foraging spinner dolphins. *Journal of the Acoustical Society of America*, 125, 539-546. doi:10.1121/1.2967477
- Benoit-Bird, K., Au, W., & Kastelein, R. (2006). Testing the odontocete acoustic prey debilitation hypothesis: No stunning results. *Journal of the Acoustical Society of America*, 120, 1118-1123. , doi:10.1121/1.2211508
- Berggren, P., Childerhouse, S., Friday, N., Kasuya, T., Kell, L., Kock, K., et al. (2003). Whaling as science. *Bioscience*, 53, 210.
- Berghan, J., Algie, K., Stockin, K., Wiseman, N., Constantine, R., Tezanos-Pinto, G., et al. (2008). A preliminary photo-identification study of bottlenose dolphin (*Tursiops truncatus*) in Hauraki Gulf, New Zealand. *New Zealand Journal of Marine & Freshwater Research*, 42, 465-472.
- Berghan, J., & Visser, I. N. (2000). Vertebral column malformations in New Zealand delphinids with a review of cases world wide. *Aquatic Mammals*, 26, 17-25.
- Bermudez, L., Van Bressem, M., Reyes-Jaimes, O., Sayegh, A., & Paniz-Mondolfi, A. (2009). Lobomycosis in man and lobomycosis-like disease in bottlenose dolphin, Venezuela. *Emerging Infectious Diseases*, 15, 1301-1303. doi:10.3201/eid1508.090347
- Bernal, R., Olavarria, C., & Moraga, R. (2003). Occurrence and long-term residency of two long-beaked common dolphins, *Delphinus capensis* (Gray 1828), in adjacent small bays on the Chilean central coast. *Aquatic Mammals*, 29, 396-399.
- Besio, K., Johnston, L., & Longhurst, R. (2008). Sexy beasts and devoted mums: Narrating nature through dolphin tourism. *Environment & Planning A*, 40, 1219-1234. doi:10.1068/a38424
- Biancani, B., Da Dalt, L., Lacave, G., Romagnoli, S., & Gabai, G. (2009). Measuring fecal progestogens as a tool to monitor reproductive activity in captive female bottlenose dolphins (*Tursiops truncatus*). *Theriogenology*, 72, 1282-1292. doi:10.1016/j.theriogenology.2009.07.025
- Bibikov, N. (2004). What do evoked potentials tell us about the acoustic system of the harbor porpoise? *Acoustical Physics*, 50, 295-304. doi:10.1134/1.1739498
- Bilger, B. (1994). Gut reaction: a remedy for oil spills may live inside the stomach of a whale.. *Sciences*, 34(6), 12.
- Bilgmann, K., Möller, L., Harcourt, R., Gales, R., & Beheregaray, L. (2008). Common dolphins subject to fisheries impacts in Southern Australia are genetically differentiated: implications for conservation. *Animal Conservation*, 11, 518-528. doi:10.1111/j.1469-1795.2008.00213.x
- Bingyao, C., Dongmei, Z., Guang, Y., Xinrong, X., & Kaiya, Z. (2009). Distribution and conservation of the Indo-Pacific humpback dolphin in China. *Integrative Zoology*, 4, 240-247. doi:10.1111/j.1749-4877.2009.00160.x
- Binns, C. (2006). It's a girl, of course. *Wildlife Conservation*, 109(3), 16.
- Bird, K., Würsig, B., & McFadden, C. (2004). Dusky dolphin (*Lagenorhynchus obscurus*) foraging in two different habitats: Active acoustic detection of dolphins and their prey. *Marine Mammal Science*, 20, 215-231.
- Birkun, A., Kuiken, T., Krivokhizhin, S., Haines, D., Osterhaus, A., Bildt, M., et al. (1999). Epizootic of morbilliviral disease in common dolphins (*Delphinus delphis ponticus*) from the Black Sea. *Veterinary Record: Journal of the British Veterinary Association*, 144, 85-92.

- Bisther, A. (2002). Intergroup interactions among killer whales in Norwegian coastal waters; tolerance vs. aggression at feeding grounds. *Aquatic Mammals*, 28, 14-23.
- Blanchet, M.-A., Nance, T., Ast, C., Wahlberg, M., & Acquarone, M. (2008). First case of a monitored pregnancy of a harbour porpoise (*Phocoena phocoena*) under human care. . *Aquatic Mammals*, 34, 9-20.
- Blomqvist, C., Mello, I., & Amundin, M. (2005). An acoustic play-fight signal in bottlenose dolphins (*Tursiops truncatus*) in human care. *Aquatic Mammals*, 31, 187-194.
- Bloodworth, B., & Marshall, C. (2005). Feeding kinematics of Kogia and *Tursiops* (*Odontoceti: Cetacea*): Characterization of suction and ram feeding. *Journal of Experimental Biology*, 208, 3721-3730. , doi:10.1242/jcb.01807
- Bloodworth, B., & Marshall, C. (2007). A functional comparison of the hyolingual complex in pygmy and dwarf sperm whales (*Kogia breviceps* and *K. sima*), and bottlenose dolphins (*Tursiops truncatus*). *Journal of Anatomy*, 211, 78-91. doi:10.1111/j.1469-7580.2007.00755.x
- Bloom, P., Goodson, A. D., Klinowska, M., & Sturtivant, C. R. (1995). The activities of a wild, solitary bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 21, 19-42.
- Bloom, P., & Jager, M. (1994). The injury and subsequent healing of a serious propeller strike to a wild bottlenose dolphin (*Tursiops truncatus*) resident in cold waters off the Northumberland coast of England. *Aquatic Mammals*, 20, 59-64.
- Boersma, P. (1999). Mystery killer. *Wildlife Conservation*, 102(2), 17.
- Boisseau, O. (2005). Quantifying the acoustic repertoire of a population: The vocalizations of free-ranging bottlenose dolphins in Fiordland, New Zealand. *Journal of the Acoustical Society of America*, 117, 2318-2329. doi:10.1121/1.1861692
- Booth, W. (1988). The social lives of dolphins. *Science*, 240, 1273-1274.
- Boreman, S. (1992). Dolphin-safe tuna: What's in a label? The killing of dolphins in the Eastern tropical Pacific and the case for an international legal solution. *Natural Resources Journal*, 32, 425-447.
- Borrell, A., & Aguilar, A. (1990). Loss of organochlorine compounds in the tissues of a decomposing stranded dolphin. *Bulletin of Environmental Contamination and Toxicology*, 45, 46-53.
- Borrell, A., & Aguilar, A. (2005a). Differences in DDT and PCB residues between common and striped dolphins from the southwestern Mediterranean. *Archives of Environmental Contamination and Toxicology*, 48, 501-508.
- Borrell, A., & Aguilar, A. (2005b). Mother-calf transfer of organochlorine compounds in the common dolphin (*Delphinus delphis*). *Bulletin of Environmental Contamination and Toxicology*, 75, 149-156.
- Borrell, A., & Aguilar, A. (2007). Organochlorine concentrations declined during 1987–2002 in western Mediterranean bottlenose dolphins, a coastal top predator. *Chemosphere*, 66, 347-352. doi:10.1016/j.chemosphere.2006.04.074
- Borrell, A., Aguilar, A., Tornero, V., Sequeira, M., Fernandez, G., & Alis, S. (2006). Organochlorine compounds and stable isotopes indicate bottlenose dolphin subpopulation structure around the Iberian Peninsula. *Environment International*, 32, 516-523. doi:10.1016/j.envint.2005.12.001
- Borrell, A., Cantos, G., Pastor, T., & Aguilar, A. (2004). Levels of organochlorine compounds in spotted dolphins from the Coiba archipelago, Panama. *Chemosphere*, 54, 669. doi:10.1016/j.chemosphere.2003.09.011
- Bortolotto, A., Casini, L., & Stanzani, L. A. (1992). Dolphin mortality along the southern Italian Coast (June-September 1991). *Aquatic Mammals*, 18, 56-60.

- Bossart, G., Ewing, R., Herron, A., Cray, C., Mase, B., Decker, S., et al. (1997). Immunoblastic malignant lymphoma in dolphins: histologic, ultrastructural, and immunohistochemical features. *Journal of Veterinary Diagnostic Investigation : Official Publication of the American Association of Veterinary Laboratory Diagnosticicians, Inc*, 9(4), 454-458.
- Bossart, G., Ghim, S.-j., Rehtanz, M., Goldstein, J., Varela, R., Ewing, R. Y., et al. (2005). Orogenital neoplasia in Atlantic bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 31, 473-480.
- Bossart, G., Romano, T. A., Peden-Adams, M. M., Rice, C. D., Fair, P. A., Goldstein, J. D., et al. (2008). Hematological, biochemical, and immunological findings in Atlantic bottlenose dolphins (*Tursiops truncatus*) with orogenital papillomas. *Aquatic Mammals*, 34, 166-177.
- Bracht, A., Brudek, R., Ewing, R., Manire, C., Burek, K., Rosa, C., et al. (2006). Genetic identification of novel poxviruses of cetaceans and pinnipeds. *Archives of Virology*, 151, 423-438. doi:10.1007/s00705-005-0679-6
- Bradshaw, C., Evans, K., & Hindell, M. (2006). Mass cetacean strandings—a plea for empiricism. *Conservation Biology*, 20, 584-586. doi:10.1111/j.1523-1739.2006.00329.x
- Bräger, S. (1993). Diurnal and seasonal behavior patterns of bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science*, 9, 434-438.
- Bräger, S. (1998). Feeding associations between White-fronted Terns and Hector's dolphins in New Zealand. *Condor*, 100, 560.
- Bräger, S., Dawson, S., Slooten, E., Smith, S., Stone, G., & Yoshinaga, A. (2002). Site fidelity and along-shore range in Hector's dolphin, an endangered marine dolphin from New Zealand. *Biological Conservation*, 108, 281-287.
- Bräger, S., Harraway, J., & Manly, B. (2003). Habitat selection in a coastal dolphin species (*Cephalorhynchus hectori*). *Marine Biology*, 143, 233-244. doi:10.1007/s00227-003-1068-x
- Bräger, S., & Würsig, B. (1994). Association patterns of bottlenose dolphins (*Tursiops truncatus*) in Galveston Bay, Texas. *Journal of Mammalogy*, 75, 431.
- Branstetter, B., & Finneran, J. (2008). Comodulation masking release in bottlenose dolphins (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 124, 625-633. doi:10.1121/1.2918545
- Branstetter, B., Finneran, J., & Houser, D. (2008). Frequency and level dependent masking of the multiple auditory steady-state response in the bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 123, 2928-2935. doi:10.1121/1.2896753
- Branstetter, B., & Mercado III, E. (2006). Sound localization by cetaceans. *International Journal of Comparative Psychology*, 19, 26-61.
- Branstetter, B., Mevissen, S., Herman, L., Pack, A., & Roberts, S. (2003). Horizontal angular discrimination by an echolocating bottlenose dolphin *Tursiops truncatus*. *Bioacoustics*, 14, 15-34.
- Branstetter, B., Mevissen, S., Pack, A., Herman, L., Roberts, S., & Carsrud, L. (2007). Dolphin (*Tursiops truncatus*) echoic angular discrimination: Effects of object separation and complexity. *Journal of the Acoustical Society of America*, 121, 626-635. doi:10.1121/1.2400664
- Brault, S., & Caswell, H. (1993). Pod-specific demography of killer whales (*Orcinus orca*). *Ecology*, 74, 1444-1454.
- Breitenbach, E., Stumpft, E., Fersen, L., & Ebert, H. (2009). Dolphin-assisted therapy: Changes in interaction and communication between children with severe

- disabilities and their caregivers. *Anthrozoös*, 22, 277-289. doi:10.2752/175303709X457612
- Brensing, K., & Linke, K. (2003). Behavior of dolphins towards adults and children during swim-with-dolphin programs and towards children with disabilities during therapy sessions. *Anthrozoös*, 16, 315-331. doi:10.2752/089279303786992035
- Brensing, K., Linke, K., Busch, M., Matthes, I., & van der Woude, S. (2005). Impact of different group of swimmers on dolphins in swim-with-the-dolphin programs in two settings. *Anthrozoös*, 18, 409-429. doi:10.2752/089279305785593956
- Brensing, K., Linke, K., & Todt, D. (2003). Can dolphins heal by ultrasound? *Journal of Theoretical Biology*, 225, 99. doi:10.1016/S0022-5193(03)00225-X
- Bressem, M., Waerebeek, K., Jepson, P., Raga, J., Duignan, P., Nielsen, O., et al. (2001). An insight into the epidemiology of dolphin morbillivirus worldwide. *Veterinary Microbiology*, 81, 287-304.
- Brill, R., & Harder, P. (1991). The effects of attenuating returning echolocation signals at the lower jaw of a dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 89, 2851-2857. doi:10.1121/1.400723
- Brill, R., Pawloski, J., Helweg, D., & Au, W. (1992). Target detection, shape discrimination, and signal characteristics of an echolocating false killer whale (*Pseudorca crassidens*). *Journal of the Acoustical Society of America*, 92, 1324-1330. doi:10.1121/1.403926
- Brill, R., Sevenich, M., Sullivan, T., Sustman, J., & Witt, R. (1988). Behavioral evidence for hearing through the lower jaw by an echolocating dolphin (*Tursiops truncatus*). *Marine Mammal Science*, 4, 223-230.
- Bristow, T. (2004). Changes in coastal site usage by bottlenose dolphins (*Tursiops truncatus*) in Cardigan Bay, Wales. *Aquatic Mammals*, 30, 398-404.
- Bristow, T., Glanville, N., & Hopkins, J. (2001). Shore-based monitoring of bottlenose dolphins (*Tursiops truncatus*) by trained volunteers in Cardigan Bay, Wales. *Aquatic Mammals*, 27, 115-120.
- Bristow, T., & Rees, E. I. S. (2001). Site fidelity and behaviour of bottlenose dolphins (*Tursiops truncatus*) in Cardigan Bay, Wales. *Aquatic Mammals*, 27, 1-10.
- Broadhurst, M. (1998). Bottlenose dolphins, *Tursiops truncatus*, removing by-catch from prawn-trawl codends, during fishing in New South Wales, Australia. *Marine Fisheries Review*, 60(3), 9.
- Bröker, K., & Ilangakoon, A. (2008). Occurrence and conservation needs of cetaceans in and around the Bar Reef Marine Sanctuary, Sri Lanka. *Oryx*, 42, 286-291. doi:10.1017/S0030605308006728
- Brook, F. M., & Kinoshita, R. E. (2005). Controlled unassisted breeding of captive indo-pacific bottlenose dolphins, *Tursiops aduncus*, using ultrasonography. *Aquatic Mammals*, 31, 89-95.
- Brook, F., Lim, E. H. T., Chua, F. H. C., & Mackay, B. (2004). Assessment of the reproductive cycle of the Indo-Pacific humpback dolphin, *Sousa chinensis*, using ultrasonography. *Aquatic Mammals*, 30, 137-148.
- Brotons, J., Grau, A., & Rendell, L. (2008). Estimating the impact of interactions between bottlenose dolphins and artisanal fisheries around the Balearic Islands. *Marine Mammal Science*, 24, 112-127.
- Brown, J., Hodgins-Davis, A., & Miller, P. (2006). Classification of vocalizations of killer whales using dynamic time warping. *Journal of the Acoustical Society of America*, 119, EL34-EL40. doi:10.1121/1.2166949

- Brown Gladden, J., Ferguson, M., & Clayton, J. (1997). Matriarchal genetic population structure of North American beluga whales *Delphinapterus leucas* (Cetacea: *Monodontidae*). *Molecular Ecology*, 6, 1033-1046.
- Brown Gladden, J. G., Ferguson, M. M., Friesen, M. K., & Clayton, J. W. (1999). Population structure of North American beluga whales (*Delphinapterus leucas*) based on nuclear DNA microsatellite variation and contrasted with the population structure revealed by mitochondrial DNA variation. *Molecular Ecology*, 8, 347-363. doi:10.1046/j.1365-294X.1999.00559.x.
- Bruemmer, F. (1986). White whales on holiday. *Natural History*, 95(1), 40.
- Bryan, C., Christopher, S., Balmer, B., & Wells, R. (2007). Establishing baseline levels of trace elements in blood and skin of bottlenose dolphins in Sarasota Bay, Florida: Implications for non-invasive monitoring. *Science of the Total Environment*, 388, 325-342.
- Buchholtz, E., & Schurt, S. (2004). Vertebral osteology in Delphinidae (Cetacea). *Zoological Journal of the Linnean Society*, 140, 383-401. doi:10.1111/j.1096-3642.2003.00105.x
- Buchholtz, E., Wolkovich, E. M., & Cleary, R. J. (2005). Vertebral osteology and complexity in *Lagenorhynchus acutus* (Delphinidae) with comparison to other Delphinoid genera. *Marine Mammal Science*, 21, 411-428.
- Buckstaff, K. C. (2004). Effects of watercraft noise on the acoustic behavior of bottlenose dolphins, *Tursiops truncatus*, in Sarasota Bay, Florida. *Marine Mammal Science*, 20, 709-725.
- Buddington, K., Holmes, W., Clemons-Chevis, C., Solangi, M., Vanderpool, D., & Buddington, R. (2006). Oral administration of sucrose solutions and measurement of serum sucrose concentrations to evaluate gastric permeability in adult bottlenose dolphins (*Tursiops truncatus*). *American Journal of Veterinary Research*, 67, 931-935.
- Buscaino, G., Buffa, G., Sarà, G., Bellante, A., Tonello, A., Hardt, F., et al. (2009). Pinger affects fish catch efficiency and damage to bottom gill nets related to bottlenose dolphins. *Fisheries Science*, 75, 537-544. doi:10.1007/s12562-009 0059-3
- Butti, C., Corain, L., Cozzi, B., Podestà, M., Pirone, A., Affronte, M., et al. (2007). Age estimation in the Mediterranean bottlenose dolphin *Tursiops truncatus* (Montagu 1821) by bone density of the thoracic limb. *Journal of Anatomy*, 211, 639-646. doi:10.1111/j.1469-7580.2007.00805.x
- Byrd, B., Hohn, A., Munden, F., Lovewell, G., & Lo Piccolo, R. (2008). Effects of commercial fishing regulations on stranding rates of bottlenose dolphin (*Tursiops truncatus*). *Fishery Bulletin*, 106, 72-81.
- Cabezon, O., Resendes, A., Domingo, M., Raga, J., Agusti, C., Alegre, F., et al. (2004). Seroprevalence of Toxoplasma gondii antibodies in wild dolphins from the Spanish Mediterranean coast. *Journal of Parasitology*, 90, 643-644.
- Caldwell, D. (1956). Intentional removal of a disturbing object by an Atlantic Bottlenosed Dolphin. *Journal of Mammalogy*, 37, 454-455.
- Caldwell, M., & Caldwell, D. (1968). Vocalization of naïve captive dolphins in small groups. *Science*, 159(3819), 1121-1123. doi:10.1126/science.159.3819.1121
- Camargo, F., Rollo Jr., M., Giampaoli, V., & Bellini, C. (2006). Whistle variability in South Atlantic spinner dolphins from the Fernando de Noronha Archipelago off Brazil. *Journal of the Acoustical Society of America*, 120, 4071-4079. doi:10.1121/1.2359704

- Campbell, G. S., Bilgre, B. A., & Defran, R. H. (2002). Bottlenose dolphins (*Tursiops truncatus*) in Turneffe Atoll, Belize: Occurrence, site fidelity, group size, and abundance. *Aquatic Mammals*, 28, 170-180.
- Cañadas, A., Sagarminaga, R., & García-Tiscar, S. (2002). Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. *Deep-Sea Research Part I, Oceanographic Research Papers*, 49, 2053-2063.
- Capaldo, T. (1989). Animal welfare tests the waters of a human-dolphin bond project. *PsyETA – Psychologists for the Ethical Treatment of Animals Bulletin*, 8(2), 7-8.
- Capus, C., Pailhas, Y., Brown, K., Lane, D.M., Moore, P.W., & Houser, D. (2007). Bio-inspired wideband sonar signals based on observations of the bottlenose dolphin (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 121, 594-604.
- Cardelluccio, N., Decataldo, A., Di Leo, A., & Giandomenico, S. (2002). Trace elements in organs and tissues of striped dolphins (*Stenella coeruleoalba*) from the Mediterranean sea (Southern Italy). *Chemosphere*, 49, 85-90.
- Carretta, J., Price, T., Petersen, D., & Read, R. (2004). Estimates of marine mammal, sea turtle, and seabird mortality in the California drift gillnet fishery for swordfish and thresher shark, 1996-2002. *Marine Fisheries Review*, 66(2), 21-30.
- Carroll, G., Suydam, R., Hazard, K., Withrow, D., George, J., & Philo, L. (1994). Frequency of killer whale (*Orcinus orca*) attacks and ship collisions based on scarring on bowhead whales (*Balaena mysticetus*) of the Bering-Chukchi-Beaufort Seas stock. *Arctic*, 47, 247.
- Carter, N. (1982). Effects of psycho-physiological stress on captive dolphins. *International Journal for the Study of Animal Problems*, 3, 193-198.
- Cassens, I., van Waerebeek, K., Best, P., Tzika, A., van Helden, A., Crespo, E., et al. (2005). Evidence for male dispersal along the coasts but no migration in pelagic waters in dusky dolphins (*Lagenorhynchus albirostris obscurus*). *Molecular Ecology*, 14, 107-121. doi:10.1111/j.1365-294X.2004.02407.x
- Caswell, H., Brault, S., Read, A. J., & Smith, T. D. (1998). Harbor porpoise and fisheries: An uncertainty analysis of incidental mortality. *Ecological Applications*, 8, 1226-1238.
- Certain, G., Ridoux, V., Van Canneyt, O., & Bretagnolle, V. (2008). Delphinid spatial distribution and abundance estimates over the shelf of the Bay of Biscay. *ICES Journal of Marine Science / Journal du Conseil*, 65, 656-666. doi:10.1093/icesjms/fsn046
- Charkovsky, I. (1986). 'Water babies' and dolphins. *PSI Research*, 5, 214-222.
- Chechina, O. N. (2007). Dynamics of development of locomotion of the bottlenose dolphin (*Tursiops truncatus ponticus baravash*, 1940) in early ontogenesis. *Journal of Evolutionary Biochemistry & Physiology*, 43, 427-432.
- Chechina, O. (2009). Formation of behavior in bottle-nosed dolphin newborns under conditions of an oceanarium: Heterochrony of involvement of various types of afferentiation. *Biology Bulletin*, 36, 489-492. doi:10.1134/S1062359009050094
- Chen, L., Bruford, M., & Yang, G. (2007). Isolation and characterization of microsatellite loci in the finless porpoise (*Neophocaena phocaenoides*). *Molecular Ecology Notes*, 7, 1129-1131. doi:10.1111/j.1471-8286.2007.01803.x
- Chen, D., Hao, Y., Zhao, Q., & Wang, D. (2006). Reproductive seasonality and maturity of male *Neophocaena phocaenoides asiaeorientalis* in captivity: A case study based on the hormone evidence. *Marine & Freshwater Behaviour & Physiology*, 39, 163-173. doi:10.1080/10236240600563396

- Chen, B., Zheng, D., Zhai, F., Xu, X., Sun, P., Wang, Q., et al. (2008). Abundance, distribution and conservation of Chinese White Dolphins (*Sousa chinensis*) in Xiamen, China. *Mammalian Biology*, 73, 156-164. doi:10.1016/j.mambio.2006.12.002
- Chivers, S., Baird, R., McSweeney, D., Webster, D., Hedrick, N., & Salinas, J. (2007). Genetic variation and evidence for population structure in eastern North Pacific false killer whales (*Pseudorca crassidens*). *Canadian Journal of Zoology*, 85, 783-794. doi:10.1139/Z07-059
- Chilvers, B., & Corkeron, P. (2002). Association patterns of bottlenose dolphins (*Tursiops aduncus*) off Point Lookout, Queensland, Australia. *Canadian Journal of Zoology*, 80, 973-979.
- Chilvers, B., Corkeron, P. J., Blanshard, W. H., Long, T. R., & Martin, A. R. (2001). A new VHF tag and attachment technique for small cetaceans. *Aquatic Mammals*, 27, 11-15.
- Chilvers, B., Corkeron, P., & Puotinen, M. (2003). Influence of trawling on the behaviour and spatial distribution of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) in Moreton Bay, Australia. *Canadian Journal of Zoology*, 81, 1947-1955.
- Chivers, S., & Danil, K. (2007). Growth and reproduction of female short-beaked common dolphins, *Delphinus delphis*, in the eastern tropical Pacific. *Canadian Journal of Zoology*, 85, 108-121.
- Chivers, S., & DeMaster, D. (1994). Evaluation of biological indices for three eastern Pacific dolphin species. *Journal of Wildlife Management*, 58, 470.
- Choi, Y., Kang, M., Sohn, H., & Kim, D. (2003). Disseminated ciliated protozoan infection in a Pacific dolphin (*Tursiops gilli*). *Veterinary Record: Journal of the British Veterinary Association*, 153, 714-715.
- Christensen, A., Rowe, S., & Needham, M. (2007). Value orientations, awareness of consequences, and participation in a whale watching education program in Oregon. *Human Dimensions of Wildlife*, 12, 275-279. doi:10.1080/10871200701442999
- Christensen, V., & Pauly, D. (1998). Changes in models of aquatic ecosystems approaching carrying capacity. *Ecological Applications*, 8, S104-S109.
- Clark, L., Cowan, D., & Pfeiffer, D. (2008). A morphological and histological examination of the Pan-tropical Spotted dolphin (*Stenella attenuata*) and the Spinner dolphin (*Stenella longirostris*) adrenal gland. *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 37, 153-159. doi:10.1111/j.1439-0264.2007.00821.x
- Clark, S., & Odell, D. (1999a). Allometric relationships and sexual dimorphism in captive killer whales (*Orcinus orca*). *Journal of Mammalogy*, 80, 777.
- Clark, S., & Odell, D. (1999b). Nursing parameters in captive killer whales (*Orcinus orca*). *Zoo Biology*, 18, 373-384.
- Clark, S., Odell, D., & Lacinak, C. (2000). Aspects of growth in captive killer whales (*Orcinus orca*). *Marine Mammal Science*, 16, 110-123.
- Clark, S. T., & Odell, D. K. (1999). Nursing behavior in captive false killer whales (*Pseudorca crassidens*). *Aquatic Mammals*, 25, 183-191.
- Clark, L., Pfeiffer, D., & Cowan, D. (2005). Morphology and histology of the Atlantic Bottlenose dolphin (*Tursiops truncatus*) adrenal gland with emphasis on the medulla. *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 34, 132-140. doi:10.1111/j.1439-0264.2004.00600.x

- Cloke, P., & Perkins, H. (2005). Cetacean performance and tourism in Kaikoura, New Zealand. *Environment & Planning D: Society & Space*, 23, 903-924. doi:10.1068/d57j
- Coan Jr., A., & Wallace, K. (1992). Differences in dolphin mortality rates in night and day sets for the U.S. eastern tropical. *Marine Fisheries Review*, 54(2), 7.
- Cole, B. (2000). poetry IN MOTION. *Wildlife Conservation*, 103(6), 52.
- CoŁmara Pellissod, S., SoŁnchez-VizcaUño, J., Carballo, M., & Mułoz, M. (2008). Determination of the immunotoxic potential of heavy metals on the functional activity of bottlenose dolphin leukocytes in vitro. *Veterinary Immunology and Immunopathology*, 121, 189-198.
- Connor, R. (2006). Male coercion of females in bottlenose dolphins. *International Journal of Primatology*, 27, 145-146.
- Connor, R., Heithaus, M., & Barre, L. (1999). Superalliance of bottlenose dolphins. *Nature*, 397, 571-572. doi:10.1038/17501
- Connor, R., Heithaus, M., Berggren, P., & Miksis, J. (2000). "Kerplunking": Surface fluke-splashes during shallow-water bottom foraging by bottlenose dolphins. *Marine Mammal Science*, 16, 646-653.
- Connor, R., Mann, J., & Watson-Capps, J. (2006). A Sex-Specific Affiliative Contact Behavior in Indian Ocean Bottlenose Dolphins, *Tursiops* sp. *Ethology*, 112, 631-638.
- Connor, R., Richards, A., Smolker, R., & Mann, J. (1996). Patterns of female attractiveness in Indian Ocean bottlenose dolphins. *Behaviour*, 133, 37-69. doi:10.1163/156853996X00026
- Connor, R., & Smolker, R. (1995). Seasonal changes in the stability of male-male bonds in Indian Ocean bottlenose dolphins (*Tursiops* sp.). *Aquatic Mammals*, 21, 213-216.
- Connor, R., & Smolker, R. (1996). 'Pop' goes the dolphin: A vocalization male bottlenose dolphins produce during consortships. *Behaviour*, 133, 643-662. doi:10.1163/156853996X00404
- Connor, R., Smolker, R., & Bejder, L. (2006). Synchrony, social behaviour and alliance affiliation in Indian Ocean bottlenose dolphins, *Tursiops aduncus*. *Animal Behaviour*, 72, 1371-1378.
- Constantine, R. (2001). Increased avoidance of swimmers by wild bottlenose dolphins (*Tursiops truncatus*) due to long-term exposure to swim-with-dolphin tourism. *Marine Mammal Science*, 17, 689-702.
- Constantine, R., Brunton, D., & Dennis, T. (2004). Dolphin-watching tour boats change bottlenose dolphin (*Tursiops truncatus*) behaviour. *Biological Conservation*, 117, 299. doi:10.1016/j.biocon.2003.12.009
- Constantine, R., Visser, I., Buurman, D., Buurman, R., & McFadden, B. (1998). Killer whale (*Orcinus orca*) predation on dusky dolphins (*Lagenorhynchus obscurus*) in Kaikoura, New Zealand. *Marine Mammal Science*, 14, 324-330.
- Conti, J., & Frohlich, R. (1984). Gastrointestinal parasitism in pygmy killer whales. *Proceedings - Helminthological Society of Washington*, 51, 364-365.
- Cook, M., Manire, C., & Mann, D. (2005). Auditory evoked potential (AEP) measurements in stranded rough-toothed dolphins (*Steno bredanensis*). *Journal of the Acoustical Society of America*, 117, 2441.
- Cook, M., Sayigh, L. S., Blum, J. E., & Wells, R. S. (2004). Signature-whistle production in undisturbed free-ranging bottlenose dolphins (*Tursiops truncatus*). *Proceedings: Biological Sciences*, 271, 1043-1049.
- Cook, T. D., & Shadish, W. R. (1994). Social experiments: some developments over the past fifteen years. *Annual Review of Psychology*, 45, 545-580.

- Corkeron, P., & Van Parijs, S. (2001). Vocalizations of eastern Australian Risso's dolphins, *Grampus griseus*. *Canadian Journal of Zoology*, 79, 160-165. doi:10.1139/cjz-79-1-160
- Cornaglia, E., Rebora, L., Gili, C., & G. (2000). Histopathological and immunohistochemical studies on cetaceans found stranded on the coast of Italy between 1990 and 1997. *Journal of Veterinary Medicine Series A*, 47, 129-142. doi:10.1046/j.1439-0442.2000.00268.x
- Corpa, J., Peris, B., Ribes, V., Palacio, J., & Liste, F. (2004). Hydrocephalus in a newborn bottlenosed dolphin (*Tursiops truncatus*). *Veterinary Record: Journal of the British Veterinary Association*, 155, 208-210.
- Coscarella, M., & Crespo, E. (2010). Feeding aggregation and aggressive interaction between bottlenose (*Tursiops truncatus*) and Commerson's dolphins (*Cephalorhynchus commersonii*) in Patagonia, Argentina. *Journal of Ethology*, 28, 183-187.
- Cosier, S. (2007). Winter's tail. *Wildlife Conservation*, 110(2), 15.
- Coughlan, J., Mirimin, L., Dillane, E., Rogan, E., & Cross, T. (2006). Isolation and characterization of novel microsatellite loci for the short-beaked common dolphin (*Delphinus delphis*) and cross-amplification in other cetacean species. *Molecular Ecology Notes*, 6, 490-492. doi:10.1111/j.1471-8286.2006.01284.x
- Courbis, S., & Timmel, G. (2009). Effects of vessels and swimmers on behavior of Hawaiian spinner dolphins (*Stenella longirostris*) in Kealake'akua, Honaunau, and Kauhako Bays, Hawai'i. *Marine Mammal Science*, 25, 430-440. doi:10.1111/j.1748-7692.2008.00254.x
- Cousteau, J. (1994). Free willy--free them all. *Animals' Agenda*, 14(1), 20-23.
- Cox, T., Read, A., Swanner, D., Urian, K., & Waples, D. (2004). Behavioral responses of bottlenose dolphins, *Tursiops truncatus*, to gillnets and acoustic alarms. *Biological Conservation*, 115, 203. doi:10.1016/S0006 3207(03)00108-3
- Craik, J., Young, J., & Cheeseman, C. (1998). GLUT-1 mediation of rapid glucose transport in dolphin (*Tursiops truncatus*) red blood cells. *American Journal of Physiology: Regulatory, Integrative & Comparative Physiology*, 275, R112.
- Crissey, S., & Wells, R. (1999). Serum alpha- and gamma-tocopherols, retinol, retinyl palmitate, and carotenoid concentrations in captive and free-ranging bottlenose dolphins (*Tursiops truncatus*). *Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology*, 124B, 391-396.
- Cronin, T., Fasick, J., & Howland, H. (1998). Video photoretinoscopy of the eyes of the small odontocetes *Tursiops truncatus*, *Phocoena phocoena*, and *Kogia breviceps*. *Marine Mammal Science*, 14, 584-590.
- Cubero-Pardo, P. (2007a). Distribución y condiciones ambientales asociadas al comportamiento del delfín bufeo (*Tursiops truncatus*) y el delfín manchado (*Stenella attenuata*) (Cetacea: *Delphinidae*) en el Golfo Dulce, Costa Rica. (Spanish). *Revista de Biología Tropical*, 55, 549-557.
- Cubero-Pardo, P. (2007b). Environmental factors governing the distribution of the bottlenose (*Tursiops truncatus*) and the spotted dolphin (*Stenella attenuata*) in Golfo Dulce, South Pacific, off Costa Rica. *Investigaciones Marinas*, 35, 15-23.
- Cullet, P., & Kameri-Mbote, A. (1996). Dolphin bycatches in tuna fisheries: A smokescreen hiding the real issues? *Ocean Development & International Law*, 27, 333.
- Cunningham-Smith, P., Colbert, D. E., Wells, R. S., & Speakman, T. (2006). Evaluation of human interactions with a provisioned wild bottlenose dolphin (*Tursiops*

- truncatus*) near Sarasota Bay, Florida, and efforts to curtail the interactions. *Aquatic Mammals*, 32, 346-356.
- Curren, K., Bose, N., & Lien, J. (1994). Swimming kinematics of a harbor porpoise (*Phocoena phocoena*) and an Atlantic white-side dolphin (*Lagenorhynchus acutus*). *Marine Mammal Science*, 10, 485-492.
- Currey, R., Slooten, E., & Dawson, S. (2009). An approach for regional threat assessment under IUCN Red List criteria that is robust to uncertainty: The Fiordland bottlenose dolphins are critically endangered. *Biological Conservation*, 142, 1570-1579.
- Curtin, S. (2006). Swimming with dolphins: A phenomenological exploration of tourist recollections. *International Journal of Tourism Research*, 8, 301-315.
- Curtin, S., & Wilkes, K. (2007). Swimming with captive dolphins: Current debates and post-experience dissonance. *International Journal of Tourism Research*, 9, 131-146. doi:10.1002/jtr.599
- Cyranoski, D. (2002). Satellite set to keep track of whales. *Nature*, 415, 354.
- Dabin, W., Cesarini, C., Clemenceau, I., Dhermain, F., Jauniaux, T., Van Canneyt, O., et al. (2004). Double-faced monster in the bottlenosed dolphin (*Tursiops truncatus*) found in the Mediterranean sea. *Veterinary Record: Journal of the British Veterinary Association*, 154, 306-308.
- Dabin, W., Cossais, F., Pierce, G., & Ridoux, V. (2008). Do ovarian scars persist with age in all Cetaceans: new insight from the short-beaked common dolphin (*Delphinus delphis linnaeus*, 1758). *Marine Biology*, 156, 127-139. doi:10.1007/s00227-008-1070-4
- Dagleish, M., Barley, J., Howie, F., Reid, R., Herman, J., & Foster, G. (2007). Isolation of Brucella species from a diseased atlanto-occipital joint of an Atlantic white-sided dolphin (*Lagenorhynchus albirostris acutus*). *Veterinary Record: Journal of the British Veterinary Association*, 160, 876-878.
- Dahlheim, M. E., Schulman-Janiger, A., Black, N., Ternullo, R., Ellifrit, D., & Balcomb III, K. C. (2008). Eastern temperate north pacific offshore killer whales (*Orcinus orca*): Occurrence, movements, and insights into feeding ecology. *Marine Mammal Science*, 24, 719-729. doi:10.1111/j.1748-7692.2008.00206.x
- Danil, K., Maldini, D., & Marten, K. (2005). Patterns of use of Maku'a beach, O'ahu, Hawai'i, by spinner dolphins (*Stenella longirostris*) and potential effects of swimmers on their behavior. *Aquatic Mammals*, 31, 403-412.
- Dankiewicz, L., Helweg, D., Moore, P., & Zafran, J. (2002). Discrimination of amplitude-modulated synthetic echo trains by an echolocating bottlenose dolphin. *Journal of the Acoustical Society of America*, 112, 1702-1708. doi:10.1121/1.1504856
- Dans, S., Crespo, E., Pedraza, S., & Koen Alonso, M. (1997). Notes on the reproductive biology of female dusky dolphins (*Lagenorhynchus obscurus*) off the Patagonian coast. *Marine Mammal Science*, 13, 303-307.
- Dans, S. L., Alonso, M. K., Pedraza, S. N., & Crespo, E. A. (2003). Incidental catch of dolphins in trawling fisheries off Patagonia, Argentina: can populations persist? *Ecological Applications*, 13, 754-762.
- Das, K., Jacob, V., & Bouquegneau, J. (2002). White-sided dolphin metallothioneins: Purification, characterisation and potential role. *Comparative Biochemistry & Physiology Part C: Toxicology & Pharmacology*, 131, 245-251
- Datta, S., & Sturtivant, C. (2002). Dolphin whistle classification for determining group identities. *Signal Processing*, 82, 251.
- Davidson, S. (2002). Do dolphins have names? *Ecos*, 111, 36.

- Davis, L., Gallman, R., & Hutchins, T. (1988). The decline of U.S. whaling: Was the stock of whales running out?. *Business History Review*, 62, 569.
- Davis, S. (2009). Sounding off [marine animal detection]. *Engineering & Technology* (17509637), 4(14), 74-75. doi:10.1049/et.2009.1419
- Davison, N., Cranwell, M., Perrett, L., Dawson, C., Deaville, R., Stubberfield, E., et al. (2009). Meningoencephalitis associated with Brucella species in a live-stranded striped dolphin (*Stenella coeruleoalba*) in south-west England. *Veterinary Record: Journal of the British Veterinary Association*, 165, 87-89.
- Dawson, C., Perrott, L., Young, E., Davison, N., & Monies, R. (2006). Isolation of Brucella species from a bottlenosed dolphin (*Tursiops truncatus*). *Veterinary Record: Journal of the British Veterinary Association*, 158, 831-832.
- Dawson, S. (1991). Clicks and communication: The behavioural and social contexts of Hector's dolphin vocalizations. *Ethology*, 88, 265-276.
- Dawson, S., Slooten, E., DuFresne, S., Wade, P., & Clement, D. (2004). Small-boat surveys for coastal dolphins: line-transect surveys for Hector's dolphins (*Cephalorhynchus hectori*). *Fishery Bulletin*, 102, 441-451.
- Dawson, S., Wade, P., Slooten, E., & Barlow, J. (2008). Design and field methods for sighting surveys of cetaceans in coastal and riverine habitats. *Mammal Review*, 38, 19-49. doi:10.1111/j.1365-2907.2008.00119.x
- de Bruyn, P., Hofmeyr, G., & de Villiers, M. (2006). First record of a vagrant Commerson's dolphin, *Cephalorhynchus commersonii*, at the southern African continental shelf. *African Zoology*, 41, 131-133.
- De Kock, A., & Lord, D. (1989). Predicting the fate and effects of chlorinated hydrocarbons in a coastal marine system--south-east Indian Ocean. *International Journal of Environmental Analytical Chemistry*, 36, 133-138.
- de Oliveira Santos, M. C., Rosso, S., dos Santos, R. A., Lucato, S. H. B., & Bassoi, M. (2002). Insights on small cetacean feeding habits in southeastern Brazil. *Aquatic Mammals*, 28, 38-45.
- De Silva, A., Muir, D., & Mabury, S. (2009). Distribution of perfluorocarboxylate isomers in select samples from the North American environment. *Environmental Toxicology & Chemistry*, 28, 1801-1814.
- Deecke, V. (2006). Studying marine mammal cognition in the wild: A review of four decades of playback experiments. *Aquatic Mammals*, 32, 461-482.
- Deecke, V., Ford, J., & Slater, P. (2005). The vocal behaviour of mammal-eating killer whales: Communicating with costly calls. *Animal Behaviour*, 69, 395-405. doi:10.1016/j.anbehav.2004.04.014
- Deecke, V., Ford, J., & Spong, P. (1999). Quantifying complex patterns of bioacoustic variation: Use of a neural network to compare killer whale (*Orcinus orca*) dialects. *Journal of the Acoustical Society of America*, 105, 2499-2507. doi:10.1121/1.426853
- Deecke, V., Ford, J., & Spong, P. (2000). Dialect change in resident killer whales: Implications for vocal learning and cultural transmission. *Animal Behaviour*, 60, 629-638. doi:10.1006/anbe.2000.1454
- Deecke, V., & Janik, V. (2006). Automated categorization of bioacoustic signals: Avoiding perceptual pitfalls. *Journal of the Acoustical Society of America*, 119, 645-653. doi:10.1121/1.2139067
- Deecke, V., Slater, P., & Ford, J. (2002). Selective habituation shapes acoustic predatory recognition in harbour seals. *Nature*, 420, 171-173. doi:10.1038/nature01030

- Defran, R., & Weller, D. (1999). Occurrence, distribution, site fidelity, and school size of bottlenose dolphins (*Tursiops truncatus*) off San Diego, California. *Marine Mammal Science*, 15, 366-380.
- Defran, R., Weller, D., Kelly, D., & Espinosa, M. (1999). Range characteristics of Pacific coast bottlenose dolphins (*Tursiops truncatus*) in the Southern California Bight. *Marine Mammal Science*, 15, 381-393.
- Degrati, M., Dans, S., Pedraza, S., Crespo, E., & Garaffo, G. (2008). Diurnal behavior of dusky dolphins, *Lagenorhynchus albirostris obscurus*, in Golfo Nuevo, Argentina. *Journal of Mammalogy*, 89, 1241-1247.
- Delfour, F. (1997). Bubbleblow in beluga whales (*Delphinapterus leucas*): A play activity?. *Behavioural Processes*, 40, 183-186. doi:10.1016/S0376-6357(97)00782-1
- Delfour, F. (2006). Marine mammals in front of the mirror-body experiences to self-recognition: A cognitive ethological methodology combined with phenomenological questioning. *Aquatic Mammals*, 32, 517-527.
- Delfour, F. (2008). Ethologie et bien-être des animaux utilisés dans les thérapies assistées par l'animal. *A.N.A.E. Approche Neuropsychologique des Apprentissages chez l'Enfant*, 20, 141-142.
- Delfour, F., & Marten, K. (2001). Mirror image processing in three marine mammal species: Killer whales (*Orcinus orca*), false killer whales (*Pseudorca crassidens*) and California sea lions (*Zalophus californianus*). *Behavioural Processes*, 53, 181-190. doi:10.1016/S0376-6357(01)00134-6
- Delfour, F., & Marten, K. (2005). Inter-modal learning task in bottlenosed dolphins (*Tursiops truncatus*): A preliminary study showed that social factors might influence learning strategies. *Acta Ethologica*, 8, 57-64. doi:10.1007/s10211-005-0110-z
- Delfour, F., & Marten, K. (2006). Lateralized visual behavior in bottlenose dolphins (*Tursiops truncatus*) performing audio-visual tasks: The right visual field advantage. *Behavioural Processes*, 71, 41-50. doi:10.1016/j.beproc.2005.09.005
- DeLong, C., Au, W., Harley, H., Roitblat, H., & Pytka, L. (2007). Human listeners provide insights into echo features used by dolphins (*Tursiops truncatus*) to discriminate among objects. *Journal of Comparative Psychology*, 121, 306-319. doi:10.1037/0735-7036.121.3.306
- DeLong, C., Au, W., Lemonds, D., Harley, H., & Roitblat, H. (2006). Acoustic features of objects matched by an echolocating bottlenose dolphin. *Journal of the Acoustical Society of America*, 119, 1867-1879. doi:10.1121/1.2161434
- DeMaster, D., Moore, S., & Dayton, P. (2000). Cetacean habitat selection in the Alaskan Arctic during summer and autumn. *Arctic*, 53, 432.
- DeMaster, D., Trites, A., Clapham, P., Mizroch, S., Wade, P., Small, R., et al. (2006). The sequential megafaunal collapse hypothesis: Testing with existing data. *Progress in Oceanography*, 68, 329-342. doi:10.1016/j.pocean.2006.02.007
- DeRuiter, S., Bahr, A., Blanchet, M., Hansen, S., Kristensen, J., Madsen, P., et al. (2009). Acoustic behaviour of echolocating porpoises during prey capture. *Journal of Experimental Biology*, 212, 3100-3107. doi:10.1242/jeb.030825
- Denardo, C., Dougherty, M., Hastie, G., Leaper, R., Wilson, B., & Thompson, P. (2001). A new technique to measure spatial relationships within groups of free-ranging coastal cetaceans. *Journal of Applied Ecology*, 38, 888-895. doi:10.1046/j.1365-2664.2001.00639.x
- Desportes, G., Buholzer, L., Anderson-Hansen, K., Blanchet, M.-A., Acquarone, M., Shephard, G., et al. (2007). Decrease stress; train your animals: The effect of

- handling methods on cortisol levels in harbour porpoises (*Phocoena phocoena*) under human care. *Aquatic Mammals*, 33, 286-292.
- Dewsbury, D.A. (1989). Comparative psychology, ethology and animal behavior. *Annual Review of Psychology*, 40, 581-602.
- Di Benedetto, A. P. M., Ramos, R. M. A., Siciliano, S., Bastos, R. A. d. S. G., & Fagundes-Netto, E. (2001). Stomach contents of delphinids from Rio de Janeiro, southeastern Brazil. *Aquatic Mammals*, 27, 24-28.
- Díaz López, B. (2006). Interactions between Mediterranean bottlenose dolphins (*Tursiops truncatus*) and gillnets off Sardinia, Italy. *ICES Journal of Marine Science / Journal du Conseil*, 63, 946-951. doi:10.1016/j.icesjms.2005.06.012
- Díaz López, B., & Bernal Shirai, J. (2008). Marine aquaculture and bottlenose dolphins' (*Tursiops truncatus*) social structure. *Behavioral Ecology and Sociobiology*, 62, 887- 894. doi:10.1007/s00265-007-0512-1
- Dill, L., & Baird, R. (1996). Ecological and social determinants of group size in transient killerwhales. *Behavioral Ecology*, 7, 408.
- Dill, L., Dill, E., & Charles, D. (2003). Feeding preferences of the Monkey Mia dolphins: Results from a simultaneous choice protocol. *Marine Mammal Science*, 19, 650-660.
- Dill, L., Heithaus, M. R., Walters, C. J. (2003). Behaviorally mediated indirect interactions in marine communities and their conservation implications. *Ecology*, 84, 1151-1157.
- Di-Meglio, N., Romero-Alvarez, R., & Collet, A. (1996). Growth comparison in striped dolphins, *Stenella coeruleoalba*, from the Atlantic and Mediterranean coasts of France. *Aquatic Mammals*, 22, 11-19.
- Ding, W., Wuersig, B., & Evans, W. E. (1995). Whistles of bottlenose dolphins: Comparisons among populations. *Aquatic Mammals*, 21, 65-77.
- DiPaola, S., Akai, C., & Kraus, B. (2007). Experiencing belugas: Action selection for an interactive aquarium exhibit. *Adaptive Behavior*, 15, 99-113. doi:10.1177/1059712306076251
- Doksæter, L., Olsen, E., Nøttestad, L., & Fernö, A. (2008). Distribution and feeding ecology of dolphins along the Mid-Atlantic Ridge between Iceland and the Azores. *Deep-Sea Research Part II, Topical Studies in Oceanography*, 55, 243-253. doi:10.1016/j.dsrr.2007.09.009
- Dolar, M., Walker, W., Kooyman, G., & Perrin, W. (2003). Comparative feeding ecology of spinner dolphins (*Stenella longirostris*) and Fraser's dolphins (*Lagenodelphis hosei*) in the Sulu Sea. *Marine Mammal Science*, 19, 1-19.
- Domenici, P., & Batty, R. (2000). Killer whales (*Orcinus orca*) feeding on schooling herring (*Clupea harengus*) using underwater tail-slaps: Kinematic analyses of field observations. *Journal of Experimental Biology*, 203, 283-294.
- Domingo-Roura, X., Shinohara, M., & Takenaka, O. (1997). Microsatellites in the bottlenose dolphin *Tursiops truncatus*. *Molecular Ecology*, 6, 695.
- Domning, D. (1991). A former dolphin-human fishing cooperative in Australia. *Marine Mammal Science*, 7, 94-96.
- Donio, S. (2008). Expérience de soutien à l'aide des dauphins. *A.N.A.E. Approche Neuropsychologique des Apprentissages chez l'Enfant*, 20, 155-156.
- Dorneles, P., Lailson-Brito, J., Santos, R., Costa, P., Malm, O., Azevedo, A., et al. (2007). Cephalopods and cetaceans as indicators of offshore bioavailability of cadmium off Central South Brazil Bight. *Environmental Pollution*, 148, 352-359.

- Dubey, J. P., Mergl, J., Gehring, E., Sundar, N., Velmurugan, G. V., Kwok, O. C. H., et al. (2009). Toxoplasmosis in captive dolphins (*Tursiops truncatus*) and walrus (*Odobenus rosmarus*). *Journal of Parasitology*, 95, 82-85.
- Dubey, J., McFee, W., Majumdar, D., Su, C., Kwok, O., Fair, P., et al. (2008). Isolation of toxoplasma gondii from bottlenose dolphins (*Tursiops truncatus*). *Journal of Parasitology*, 94, 821-823.
- Dubey, J., Morales, J., Sundar, N., Velmurugan, G., González-Barrientos, C., Hernández-Mora, G., et al. (2007). Isolation and genetic characterization of Toxoplasma gondii from striped dolphin (*Stenella coeruleoalba*) from Costa Rica. *Journal of Parasitology*, 93, 710-711.
- Dubey, J. P., Fair, P. A., Bossart, G. D., Hill, D., Fayer, R., Sreekumar, C., et al. (2005). A comparison of several serologic tests to detect antibodies to Toxoplasma gondii in naturally exposed bottlenose dolphins (*Tursiops truncatus*). *Journal of Parasitology*, 91, 1074-1081.
- Dubey, J., Zarnke, R., Thomas, N., Wong, S., Van Bonn, W., Briggs, M., et al. (2003). Toxoplasma gondii, Neospora caninum, Sarcocystis neurona, and Sarcocystis canis-like infections in marine mammals. *Veterinary Parasitology*, 116, 275. doi:10.1016/S0304-4017(03)00263-2
- Dubrovsky, N. (2004). Echolocation system of the bottlenose dolphin. *Acoustical Physics*, 50, 305-317. doi:10.1134/1.1739499
- Dubrovsky, N., & Sukhoruchenko, M. (2002). Audibility thresholds for a pure tone in the presence of a pulse jam in the bottlenose dolphin. *Acoustical Physics*, 48, 441-447.
- Dubrovsky, N., Zorikov, T., Kvizhinadze, O., & Kuratashvili, M. (1992). Feature description of acoustic pulses in their auditory discrimination by *Tursiops truncatus*. *Sensory Systems*, 6, 27-36.
- Dudzinski, K. (1998). Contact behavior and signal exchange in Atlantic spotted dolphins (*Stenella frontalis*). *Aquatic Mammals*, 24, 129-142.
- Dudzinski, K., Clark, C. W., & Würsig, B. (1995). A mobile video/acoustic system for simultaneous underwater recording of dolphin interactions. *Aquatic Mammals*, 21, 187-193.
- Dudzinski, K., Frohoff, T. G., & Crane, N. L. (1995). Behavior of a lone female bottlenose dolphin (*Tursiops truncatus*) with humans off the coast of Belize. *Aquatic Mammals*, 21, 149-153.
- Dudzinski, K., Gregg, J., Ribic, C., & Kuczaj, S. (2009). A comparison of pectoral fin contact between two different wild dolphin populations. *Behavioural Processes*, 80, 182-190. doi:10.1016/j.beproc.2008.11.011
- Duffield, D., Odell, D., McBain, J., & Andrews, B. (1995). Killer whale (*Orcinus orca*) reproduction at Sea World. *Zoo Biology*, 14, 417-430.
- Duggan, Z., Koopman, H., & Budge, S. (2009). Distribution and development of the highly specialized lipids in the sound reception systems of dolphins. *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 179, 783-798. doi:10.1007/s00360-009-0360-6
- Dunbar, M. (1987). Arctic seas that never freeze. *Natural History*, 96(4), 50-53.
- Dunkin, R., Mclellan, W., Blum, J., & Pabst, A. (2005). The ontogenetic changes in the thermal properties of blubber from Atlantic bottlenose dolphin *Tursiops truncatus*. *Journal of Experimental Biology*, 208, 1469-1480. doi:10.1242/jeb.01559
- Durban, J., Parsons, K., Claridge, D., & Balcomb, K. (2000). Quantifying dolphin occupancy patterns. *Marine Mammal Science*, 16, 825-828.

- Durden, W., Stolen, M., Adams, D., & Stolen, E. (2007). Mercury and selenium concentrations in stranded bottlenose dolphins from the Indian River Lagoon system. *Bulletin of Marine Science*, 81, 37-54.
- Durden, W. (2005). The harmful effects of inadvertently conditioning a wild Bottlenose dolphin (*Tursiops truncatus*) to interact with fishing vessels in the Indian River Lagoon, Florida, USA. *Aquatic Mammals*, 31, 413-419.
- Earthwatch, L. L. C. (1993). Ceasing cetacean snares. *Environment*, 35, 21.
- Edwards, E. (2006). Duration of unassisted swimming activity for spotted dolphin (*Stenella attenuata*) calves: implications for mother-calf separation during tuna purse-seine sets. *Fishery Bulletin*, 104, 125-135.
- Edwards, E. (2007). Fishery effects on dolphins targeted by tuna purse-seiners in the Eastern tropical Pacific Ocean. *International Journal of Comparative Psychology*, 20, 217- 227.
- Eisenfeld, M. (2008). Ohaleï Yacoov—Le silence des justes: Une porte sur la vie. *A.N.A.E. Approche Neuropsychologique des Apprentissages chez l'Enfant*, 20, 157-158.
- Elk, C., Bildt, M., Martina, B., Osterhaus, A., & Kuiken, T. (2007). Escherichia coli septicemia associated with lack of maternally acquired immunity in a bottlenose dolphin calf. *Veterinary Pathology*, 44, 88-92.
- Elk, C., Epping, N., & Gans, S. (2001). Pulmonary function measurements in dolphins using capnography. *Veterinary Record: Journal of the British Veterinary Association*, 149, 308-309.
- Ellis, B., Gattoni-Celli, S., Mancia, A., & Kindy, M. (2009). The vitamin D3 transcriptomic response in skin cells derived from the Atlantic bottlenose dolphin. *Developmental & Comparative Immunology*, 33, 901-912. doi:10.1016/j.dci.2009.02.008
- Emery, N. J., & Clayton, N. S. (2008). Comparative social cognition. *Annual Review of Psychology*, 60, 87-113.
- Endo, T., Kimura, O., Hisamichi, Y., Minoshima, Y., & Haraguchi, K. (2007). Age-dependent accumulation of heavy metals in a pod of killer whales (*Orcinus orca*) stranded in the northern area of Japan. *Chemosphere*, 67, 51-59. doi:10.1016/j.chemosphere.2006.09.086
- Erbe, C. (2000). Detection of whale calls in noise: Performance comparison between a beluga whale, human listeners, and a neural network. *Journal of the Acoustical Society of America*, 108, 297-303. doi:10.1121/1.429465
- Erbe, C. (2002). Underwater noise of whale-watching boats and potential effects on killer whales (*Orcinus orca*), based on an acoustic impact model. *Marine Mammal Science*, 18, 394-418.
- Erbe, C. (2008). Critical ratios of beluga whales (*Delphinapterus leucas*) and masked signal duration. *Journal of the Acoustical Society of America*, 124, 2216-2223. doi:10.1121/1.2970094
- Erbe, C., & Farmer, D. (1998). Masked hearing thresholds of a beluga whale (*Delphinapterus leucas*) in icebreaker noise. *Deep-Sea Research Part II, Topical Studies in Oceanography*, 45, 1373-1388.
- Esch, H., Sayigh, L., Blum, J., & Wells, R. (2009). Whistles as potential indicators of stress in bottlenose dolphins (*Tursiops truncatus*). *Journal of Mammalogy*, 90, 638-650.
- Esch, H., Wells, R., & Sayigh, L. (2009). Quantifying parameters of bottlenose dolphin signature whistles. *Marine Mammal Science*, 25, 976-986.
- Escorza-Treviño, S., Archer, F., Rosales, M., Lang, A., & Dizon, A. (2005). Genetic differentiation and intraspecific structure of Eastern Tropical Pacific spotted

- dolphins, *Stenella attenuata*, revealed by DNA analyses. *Conservation Genetics*, 6, 587-600.
- Escorza-Treviño, S., Pastene, L., & Dizon, A. (2004). Molecular analyses of the truei and dalli morphotypes of dalls porpoise (*Phocoenoides dalli*). *Journal of Mammalogy*, 85, 347-355.
- Essington, T. E., Schindler, D. E., Olson, R. J., Kitchell, J. F., Boggs, C., & Hilborn, R. (2002). Alternative fisheries and the predation rate of yellowfin tuna in the Eastern Pacific ocean. *Ecological Applications*, 12, 724-734.
- Essumang, D. (2008). Bioaccumulation of platinum group metals in dolphins, *Stenella* sp., caught off Ghana. *African Journal of Aquatic Science*, 33, 255-259. doi:10.2989/AJAS.2008.33.3.8.620
- Etnier, S., McLeIIan, W., BIum, J., & Pabst, D. (2008). Ontogenetic changes in the structural stiffness of the tailstock of bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Biology*, 211, 3205-3213. , doi:10.1242/jeb.012468
- Evans, D. (1987). Dolphins as beaters for gulls? *Bird Behavior*, 7(1), 47-48.
- Evans, J., Al-Abiani, S., Klesius, P., & Pasnik, D. (2006). First report of *Streptococcus agalactiae* and *Lactococcus garvieae* from a wild bottlenose dolphin (*Tursiops truncatus*). *Journal of Wildlife Diseases*, 42, 561-569.
- Evans, P., & Hammond, P. (2004). Monitoring cetaceans in European waters. *Mammal Review*, 34, 131-156. doi:10.1046/j.0305-1838.2003.00027.x
- Evans, W. (1973). Echolocation by marine delphinids and one species of fresh-water dolphin. *Journal of the Acoustical Society of America*, 54, 191-199. doi:10.1121/1.1913562
- Fahlman, A., Olszowka, A., Bostrom, B., & Jones, D. (2006). Deep diving mammals: Dive behavior and circulatory adjustments contribute to bends avoidance. *Respiratory Physiology & Neurobiology*, 153, 66-77. doi:10.1016/j.resp.2005.09.014
- Fair, P., Bossart, G., Alaee, M., Muir, D., Henry, N., Adams, J., et al. (2009). Occurrence of triclosan in plasma of wild Atlantic bottlenose dolphins (*Tursiops truncatus*) and in their environment [electronic resource]. *Environmental Pollution*, 157, 2248-2254.
- Fair, P., Mitchum, G., Hulsey, T., Adams, J., Zolman, E., McFee, W., et al. (2007). Polybrominated diphenyl ethers (PBDEs) in blubber of free-ranging bottlenose dolphins (*Tursiops truncatus*) from two Southeast Atlantic estuarine areas [electronic resource]. *Archives of Environmental Contamination and Toxicology*, 53, 483-494.
- Fair, P., Hulsey, T. C., Varela, R. A., Goldstein, J. D., Adams, J., Zolman, E. S., et al. (2006). Hematology, serum chemistry, and cytology findings from apparently healthy Atlantic bottlenose dolphins (*Tursiops truncatus*) inhabiting the estuarine waters of Charleston, South Carolina. *Aquatic Mammals*, 32, 182-195.
- Farley, J. & Alkon, D. L. (1985). Cellular mechanisms of learning, memory and information storage. *Annual Review of Psychology*, 36, 419-494.
- Fazioli, K. L., Hofmann, S., & Wells, R. S. (2006). Use of Gulf of Mexico coastal waters by distinct assemblages of bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 32, 212-222.
- Fedorowicz, S. M., Beard, D. A., & Connor, R. C. (2003). Food sharing in wild bottlenose dolphins. *Aquatic Mammals*, 29, 355-359.
- Feinholz, D. M., & Atkinson, S. (2000). Possible aetiologies of yellow coloration in dolphin calves. *Aquatic Mammals*, 26, 191-195.

- Felix, F. (1997). Organization and social structure of the coastal bottlenose dolphin *Tursiops truncatus* in the Gulf de Guayaquil, Ecuador. *Aquatic Mammals*, 23, 1-16.
- Feng, A. S., & Ratnam, R. (2000). Neural basis of hearing in real-world situations. *Annual Review of Psychology*, 51, 699-725.
- Ferber, D. (2005). Whaling endangers more than whales. *Science*, 307(5713), 1190-1191.
- Ferrer-y-Cancho, R., & McCowan, B. (2009). A law of word meaning in dolphin whistle types. *Entropy*, 11, 688-701. doi:10.3390/e11040688
- Ferrero, R., Moore, S., & Hobbs, R. (2000). Development of beluga capture and satellite tagging protocol in Cook Inlet, Alaska. *Marine Fisheries Review*, 62(3), 112-123.
- Fertl, D., Acevedo-Gutierrez, A., & Darby, F. (1996). A report of killer whales (*Orcinus orca*) feeding on a carcharhinid shark in Costa Rica. *Marine Mammal Science*, 12, 606-611.
- Fertl, D., Jefferson, T., Moreno, I., Zerbini, A., & Mullin, K. (2003). Distribution of the Clymene dolphin (*Stenella clymene*). *Mammal Review*, 33, 253. doi:10.1046/j.1365-2907.2003.00033.x
- Fertl, D., Pusser, L., & Long, J. (1999). First record of an albino bottlenose dolphin (*Tursiops truncatus*) in the Gulf of Mexico, with a review of anomalously white cetaceans. *Marine Mammal Science*, 15, 227-234.
- Fertl, D., & Schiro, A. (1994). Carrying of dead calves by free-ranging Texas bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 20, 53-56.
- Fertl, D., Schiro, A. J., & Peake, D. (1997). Coordinated feeding by Clymene dolphins (*Stenella clymene*) in the Gulf of Mexico. *Aquatic Mammals*, 23, 111-112.
- Fertl, D., & Wilson, B. (1997). Bubble use during prey capture by a lone bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 23, 113-114.
- Fertl, D., & Wuersig, B. (1995). Coordinated feeding by Atlantic spotted dolphins (*Stenella frontalis*) in the Gulf of Mexico. *Aquatic Mammals*, 21, 3-5.
- Filatova, O., Fedutin, I., Burdin, A., & Hoyt, E. (2006). Using a mobile hydrophone stereo system for real-time acoustic localization of killer whales (*Orcinus orca*). *Applied Acoustics*, 67, 1243-1248. doi:10.1016/j.apacoust.2006.05.004
- Filatova, O., Fedutin, I., Ivkovich, T., Nagaylik, M., Burdin, A., & Hoyt, E. (2009). The function of multi-pod aggregations of fish-eating killer whales (*Orcinus orca*) in Kamchatka, Far East Russia. *Journal of Ethology*, 27, 333-341. doi:10.1007/s10164-008-0124-x
- Finneran, J. (2003). Whole-lung resonance in a bottlenose dolphin (*Tursiops truncatus*) and white whale (*Delphinapterus leucas*). *Journal of the Acoustical Society of America*, 114, 529-535. doi:10.1121/1.1575747
- Finneran, J. (2008). Modified variance ratio for objective detection of transient evoked potentials in bottlenose dolphins (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 124, 4069-4082. doi:10.1121/1.2996320
- Finneran, J., & Houser, D. (2006). Comparison of in-air evoked potential and underwater behavioral hearing thresholds in four bottlenose dolphins (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 119, 3181-3192. doi:10.1121/1.2180208
- Finneran, J., & Houser, D. (2007). Bottlenose dolphin (*Tursiops truncatus*) steady-state evoked responses to multiple simultaneous sinusoidal amplitude modulated tones. *Journal of the Acoustical Society of America*, 121, 1775-1782. doi:10.1121/1.2431330
- Finneran, J., Carder, D., Dear, R., Belting, T., McBain, J., Dalton, L., et al. (2005). Pure tone audiograms and possible aminoglycoside-induced hearing loss in belugas

- (*Delphinapterus leucas*). *Journal of the Acoustical Society of America*, 117, 3936-3943. doi:10.1121/1.1893354
- Finneran, J., Carder, D., & Ridgway, S. (2002). Low-frequency acoustic pressure, velocity, and intensity thresholds in a bottlenose dolphin (*Tursiops truncatus*) and white whale (*Delphinapterus leucas*). *Journal of the Acoustical Society of America*, 111, 447-456. doi:10.1121/1.1423925
- Finneran, J., Carder, D., Schlundt, C., & Ridgway, S. (2005). Temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. *Journal of the Acoustical Society of America*, 118, 2696-2705. doi:10.1121/1.2032087
- Finneran, J., Houser, D., Blasko, D., Hicks, C., Hudson, J., & Osborn, M. (2008). Estimating bottlenose dolphin (*Tursiops truncatus*) hearing thresholds from single and multiple simultaneous auditory evoked potentials. *Journal of the Acoustical Society of America*, 123, 542-551. doi:10.1121/1.2812595
- Finneran, J., Houser, D. S., & Schlundt, C. E. (2007). Objective detection of bottlenose dolphin (*Tursiops truncatus*) steady-state auditory evoked potentials in response to AM/FM tones. *Aquatic Mammals*, 33, 43-54.
- Finneran, J., London, H., & Houser, D. (2007). Modulation rate transfer functions in bottlenose dolphins (*Tursiops truncatus*) with normal hearing and high-frequency hearing loss. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural & Behavioral Physiology*, 193, 835-843. doi:10.1007/s00359-007-0238-6
- Finneran, J., Oliver, C., Schaefer, K., & Ridgway, S. (2000). Source levels and estimated yellowfin tuna (*Thunnus albacares*) detection ranges for dolphin jaw pops, breaches, and tail slaps. *Journal of the Acoustical Society of America*, 107, 649-656. doi:10.1121/1.428330
- Finneran, J., & Schlundt, C. (2007). Underwater sound pressure variation and bottlenose dolphin (*Tursiops truncatus*) hearing thresholds in a small pool. *Journal of the Acoustical Society of America*, 122, 606-614. doi:10.1121/1.2743158
- Finneran, J., Schlundt, C., Branstetter, B., & Dear, R. (2007). Assessing temporary threshold shift in a bottlenose dolphin (*Tursiops truncatus*) using multiple simultaneous auditory evoked potentials. *Journal of the Acoustical Society of America*, 122, 1249-1264. doi:10.1121/1.2749447
- Finneran, J., Schlundt, C., Carder, D., Clark, J., Young, J., Gaspin, J., et al. (2000). Auditory and behavioral responses of Bottlenose dolphins (*Tursiops truncatus*) and a beluga whale (*Delphinapterus leucas*) to impulsive sounds resembling distant signatures of underwater explosions. *Journal of the Acoustical Society of America*, 108, 417-431. doi:10.1121/1.429475
- Finneran, J., Schlundt, C., Carder, D., & Ridgway, S. (2002). Auditory filter shapes for the bottlenose dolphin (*Tursiops truncatus*) and the white whale (*Delphinapterus leucas*) derived with notched noise. *Journal of the Acoustical Society of America*, 112, 322-328. doi:10.1121/1.1488652
- Fire, S., Fauquier, D., Flewelling, L., Henry, M., Naar, J., Pierce, R., et al. (2007). Brevetoxin exposure in bottlenose dolphins (*Tursiops truncatus*) associated with Karenia brevis blooms in Sarasota Bay, Florida. *Marine Biology*, 152, 827-834. doi:10.1007/s00227007-0733-x
- Fire, S., Flewelling, L. J., Wang, Z., Naar, J., Henry, M. S., Pierce, R. H., et al. (2008). Florida red tide and brevetoxins: Association and exposure in live resident bottlenose dolphins (*Tursiops truncatus*) in the eastern Gulf of Mexico, U.S.A. *Marine Mammal Science*, 24, 831-844.
- Fish, F. (1993). Power output and propulsive efficiency of swimming bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Biology*, 185, 179-193.

- Fish, F. (1998). Comparative kinematics and hydrodynamics of odontocete cetaceans: Morphological and ecological. *Journal of Experimental Biology*, 201, 2867.
- Fish, F., Nicastro, A., & Weihs, D. (2006). Dynamics of the aerial maneuvers of spinner dolphins. *Journal of Experimental Biology*, 209, 590-598. doi:10.1242/jeb.02034
- Fléss, D., Starodubtsev, Y., & Starodubtseva, E. (1990). On the capability of bottle-nosed dolphins to generalize according to a relative sign. *Zhurnal Vysshhei Nervnoi Deyatel'nosti*, 40(1), 44-51.
- Fléss, D., Starodubtsev, Y., & Starodubtseva, E. (1991). Capacity of bottle-nosed dolphins for generalization based on a relative sign. *Neuroscience and Behavioral Physiology*, 21, 166-172. doi:10.1007/BF01182894
- Flores, G. (2005). Flipper fashion. *Natural History*, 114(7), 12.
- Fontaine, M. C., Galan, M., Bouquegneau, J.-M., & Michaux, J. R. (2006). Efficiency of fluorescent multiplex polymerase chain reactions (PCRs) for rapid genotyping of harbour porpoises (*Phocoena phocoena*) with 11 microsatellite loci. *Aquatic Mammals*, 32, 301-304.
- Foote, A., Newton, J., Piertney, S., Willerslev, E., & Gilbert, M. (2009). Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology*, 18, 5207-5217. doi:10.1111/j.1365-294X.2009.04407.x
- Foote, A., & Nystuen, J. (2008). Variation in call pitch among killer whale ecotypes. *Journal of the Acoustical Society of America*, 123, 1747-1752. doi:10.1121/1.2836752
- Foote, A., Osborne, R., & Hoelzel, A. (2004). Whale-call response to masking boat noise. *Nature*, 428, 910-910. doi:10.1038/428910a
- Foote, A., & Hoezel, R. (2004). A little bit louder now. *Environment*, 46(6), 7-8.
- Foote, A., Osborne, R., & Hoelzel, A. (2008). Temporal and contextual patterns of killer whale (*Orcinus orca*) call type production. *Ethology*, 114, 599-606. doi:10.1111/j.1439-0310.2008.01496.x
- Ford, J. (1991). Family fugues. *Natural History*, 100(3), 68.
- Ford, J., & Reeves, R. (2008). Fight or flight: Antipredator strategies of baleen whales. *Mammal Review*, 38, 50-86.
- Ford, J., Ellis, G. M., Matkin, D. R., Balcomb, K. C., Briggs, D., & Morton, A. B. (2005). Killer whale attacks on minke whales: Prey capture and antipredator tactics. *Marine Mammal Science*, 21, 603-618.
- Forestell, P., & Herman, L. (1988). Delayed matching of visual materials by a bottlenosed dolphin aided by auditory symbols. *Animal Learning & Behavior*, 16, 137-146.
- Fossi, M., Casini, S., Bucalossi, D., & Marsili, L. (2008). First detection of CYP1A1 and CYP2B induction in Mediterranean cetacean skin biopsies and cultured fibroblasts by Western blot analysis. *Marine Environmental Research*, 66, 3-6. doi:10.1016/j.marenvres.2008.02.006
- Fossi, M., Marsili, L., Lauriano, G., Fortuna, C., Canese, S., Ancora, S., et al. (2004). Assessment of toxicological status of a SW Mediterranean segment population of striped dolphin (*Stenella coeruleoalba*) using skin biopsy. *Marine Environmental Research*, 58, 269-274. doi:10.1016/j.marenvres.2004.03.070
- Frantzis, A., & Herzing, D. L. (2002). Mixed-species associations of striped dolphins (*Stenella coeruleoalba*), short-beaked common dolphins (*Delphinus delphis*), and Risso's dolphins (*Grampus griseus*) in the Gulf of Corinth (Greece, Mediterranean Sea). *Aquatic Mammals*, 28, 188-197.
- Fripp, D. (2005). Bubblestream whistles are not representative of a bottlenose dolphin's vocal repertoire. *Marine Mammal Science*, 21, 29-44.

- Fripp, D., & Tyack, P. (2008). Postpartum whistle production in bottlenose dolphins. *Marine Mammal Science*, 24, 479-502.
- Fripp, D., Owen, C., Quintana-Rizzo, E., Shapiro, A., Buckstaff, K., Jankowski, K., et al. (2005). Bottlenose dolphin (*Tursiops truncatus*) calves appear to model their signature whistles on the signature whistles of community members. *Animal Cognition*, 8, 17- 26. doi:10.1007/s10071-004-0225-z
- Frodello, J., & Marchand, B. (2001). Cadmium, copper, lead, and zinc in five toothed whale species of the Mediterranean Sea. *International Journal of Toxicology*, 20, 339-343. doi:10.1080/10915810175333613
- Frohoff, T., & Packard, J. (1995). Human interactions with free-ranging and captive bottlenose dolphins. *Anthrozoös*, 8, 44-53. doi:10.2752/089279395787156527
- Frost, K., Russell, R., & Lowry, L. (1992). Killer whales, *Orcinus orca*, in the Southeastern Bering Sea: Recent sightings and predation on other marine mammals. *Marine Mammal Science*, 8, 110-119.
- Funke, C., King, D., McBain, J., Adelung, D., & Stott, J. (2003). Expression and functional characterization of killer whale (*Orcinus orca*) interleukin-6 (IL-6) and development of a competitive immunoassay. *Veterinary Immunology & Immunopathology*, 93, 69. doi:10.1016/S0165-2427(03)00055-2
- Furutani, R. (2008). Laminar and cytoarchitectonic features of the cerebral cortex in the Risso's dolphin (*Grampus griseus*), striped dolphin (*Stenella coeruleoalba*), and bottlenose dolphin (*Tursiops truncatus*). *Journal of Anatomy*, 213, 241-248. doi:10.1111/j.1469-7580.2008.00936.x
- Galatius, A. (2005). Sexually dimorphic proportions of the harbour porpoise (*Phocoena phocoena*) skeleton. *Journal of Anatomy*, 206, 141-154. doi:10.1111/j.1469-7580.2005.00381.x
- Galatius, A. (2006). Bilateral directional asymmetry of the appendicular skeleton of the white-beaked dolphin (*Lagenorhynchus albirostris albirostris*). *Aquatic Mammals*, 32, 232-235.
- Galatius, A., Andersen, M., Haugan, B., Langhoff, H., & Jespersen, Å. (2006). Timing of epiphyseal development in the flipper skeleton of the harbour porpoise (*Phocoena phocoena*) as an indicator of paedomorphosis. *Acta Zoologica*, 87, 77-82. doi:10.1111/j.1463-6395.2006.00223.x
- Galhardo, L., Appleby, M., Waran, N., & Santos, M. (1996). Spontaneous activities of captive performing bottlenose dolphins (*Tursiops truncatus*) *Animal Welfare*, 5, 373-389.
- Galindo, J. A., Serrano, A., Vazquez-Castan, L., Gonzalez-Gandara, C., & Lopez-Ortega, M. (2009). Cetacean diversity, distribution, and abundance in northern Veracruz, Mexico. *Aquatic Mammals*, 35, 12-18.
- Gallup, G. (1995). Mirrors, minds, and cetaceans. *Consciousness and Cognition: An International Journal*, 4, 226-228. doi:10.1006/ccog.1995.1028
- Gannier, A. (2000). Distribution of cetaceans off the Society Islands (French Polynesia) as obtained from dedicated surveys. *Aquatic Mammals*, 26, 111-126.
- Gannier, A. (2002). Cetaceans of the Marquesas Islands (French Polynesia): Distribution and relative abundance as obtained from a small boat dedicated survey. *Aquatic Mammals*, 28, 198-210.
- Gannier, A., & Petiau, E. (2006). Environmental variables affecting the residence of spinner dolphins (*Stenella longirostris*) in a bay of Tahiti (French Polynesia). *Aquatic Mammals*, 32, 202-211.

- Gannier, A., & West, K. (2005). Distribution of the rough-toothed dolphin (*Steno bredanensis*) around the Windward Islands (French Polynesia). *Pacific Science*, 59, 17-24.
- Gannon, D., Barros, N., Nowacek, D., Read, A., Waples, D., & Wells, R. (2005). Prey detection by bottlenose dolphins, *Tursiops truncatus*: An experimental test of the passive listening hypothesis. *Animal Behaviour*, 69, 709-720. doi:10.1016/j.anbehav.2004.06.020
- Gannon, D., & Waples, D. M. (2004). Diets of coastal bottlenose dolphins from the U.S. mid-Atlantic coast differ by habitat. *Marine Mammal Science*, 20, 527-545.
- Gao, A., Zhou, K., & Wang, Y. (1995). Geographical variation in morphology of bottlenose dolphins (*Tursiops* sp.) in Chinese waters. *Aquatic Mammals*, 21, 121-135.
- García-Martínez, J., Moya, A., Raga, J., & Latorre, A. (1999). Genetic differentiation in the striped dolphin *Stenella coeruleoalba* from European waters according to mitochondrial DNA (mtDNA) restriction analysis. *Molecular Ecology*, 8, 1069-1073. doi:10.1046/j.1365-294X.1999.00672.x
- Garrick, R., Woodin, B., Wilson, J., Middlebrooks, B., & Stegeman, J. (2006). Cytochrome P4501A is induced in endothelial cell lines from the kidney and lung of the bottlenose dolphin, *Tursiops truncatus*. *Aquatic Toxicology*, 76, 295-305. doi:10.1016/j.aquatox.2005.10.005
- Garrison, L. (2007). Interactions between marine mammals and pelagic longline fishing gear in the U.S. Atlantic Ocean between 1992 and 2004. *Fishery Bulletin*, 105, 408-417.
- Gaspar, C., Lenzi, R., Reddy, M., & Sweeney, J. (2000). Spontaneous lactation by an adult *Tursiops truncatus* in response to a stranded *Steno bredanensis* calf. *Marine Mammal Science*, 16, 653-658.
- Gaspari, S., Azzellino, A., Airoldi, S., & Hoelzel, A. (2007). Social kin associations and genetic structuring of striped dolphin populations (*Stenella coeruleoalba*) in the Mediterranean Sea. *Molecular Ecology*, 16, 2922-2933. doi:10.1111/j.1365-294X.2007.03295.x
- Gaunaurd, G., Brill, D., Huang, H., Moore, P., & Strifors, H. (1998). Signal processing of the echo signatures returned by submerged shells insonified by dolphin 'clicks': Active classification. *Journal of the Acoustical Society of America*, 103, 1547-1557. doi:10.1121/1.421302
- Gaydos, J., Balcomb III, K., Osborne, R., & Dierauf, L. (2004). Evaluating potential infectious disease threats for southern resident killer whales, *Orcinus orca*: A model for endangered species. *Biological Conservation*, 117, 253. doi:10.1016/j.biocon.2003.07.004
- George, J., & Suydam, R. (1998). Observations of killer whale (*Orcinus orca*) predation in the northeastern Chukchi and western Beaufort seas. *Marine Mammal Science*, 14, 330-332.
- Gerber, L., & DeMaster, D. (2000). Measuring success in conservation. *American Scientist*, 88, 316.
- Gerber, L., Morissette, L., Kaschner, K., & Pauly, D. (2009). Should whales be culled to increase fishery yield? *Science*, 323, 880-881.
- Gero, S., Bejder, L., Whitehead, H., Mann, J., & Connor, R. (2005). Behaviourally specific preferred associations in bottlenose dolphins, *Tursiops* spp. *Canadian Journal of Zoology*, 83, 1566-1573. doi:10.1139/Z05-155

- Gibson, Q., & Mann, J. (2008a). Early social development in wild bottlenose dolphins: Sex differences, individual variation and maternal influence. *Animal Behaviour*, 76, 375-387. doi:10.1016/j.anbehav.2008.01.021
- Gibson, Q., & Mann, J. (2008b). The size, composition and function of wild bottlenose dolphin (*Tursiops* sp.) mother-calf groups in Shark Bay, Australia. *Animal Behaviour*, 76, 389-405. doi:10.1016/j.anbehav.2008.01.022
- Gibson, Q., & Mann, J. (2009). Do sampling method and sample size affect basic measures of dolphin sociality? *Marine Mammal Science*, 25, 187-198.
- Giesy, J., Tanabe, S., Loganathan, B., Kannan, K., Odell, D., Watanabe, M., et al. (2000). Polychlorinated biphenyls, organochlorine pesticides, tris(4-chlorophenyl) methane, and tris(4-chlorophenyl) methanol in livers of small cetaceans stranded along Florida coastal waters, USA. *Environmental Toxicology & Chemistry*, 19, 1566.
- Gladden, J., Brodie, P., & Clayton, J. (1999). Mitochondrial DNA used to identify an extralimital beluga whale (*Delphinapterus leucas*) from Nova Scotia as originating from the St. Lawrence population. *Marine Mammal Science*, 15, 556-558.
- Gnone, G., Benoldi, C., Bonsignori, B., & Fognani, P. (2001a). Observations of rest behaviours in captive bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 27, 29-33.
- Gnone, G., Caresano, F., Cosmai, T., & Gnone, E. (2008). The orientation of cutaneous ridges influences the appearance of longitudinal stripes on the skin surface of bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science*, 24, 711-718. doi:10.1111/j.1748-7692.2008.00204.x
- Gnone, G., Moriconi, T., & Gambini, G. (2006). Sleep behaviour: Sleep in continuously active dolphins; Activity and sleep in dolphins (Reply). *Nature*, 441, E11. doi:10.1038/nature04900
- Goldsmith, M., & Korit, A. (2003). Dolphins on the witness stand? The comparative psychology of strategic memory regulation. *Behavioral & Brain Sciences*, 26, 345-346.
- Goldstein, H. (2005). ROCKY. *Wildlife Conservation*, 108, 29-35.
- Goley, P. (1999). Behavioral aspects of sleep in pacific white-sided dolphins (*Lagenorhynchus obliquidens*, Gill 1865). *Marine Mammal Science*, 15, 1054-1064.
- Gomercic, M. D., Galov, A., Gomercic, T., Skrtic, D., Curkovic, S., Lucic, H., et al. (2009). Bottlenose dolphin (*Tursiops truncatus*) depredation resulting in larynx strangulation with gill-net parts. *Marine Mammal Science*, 25, 392-401. doi:10.1111/j.1748-7692.2008.00259.x
- Gómez de Segura, A., Crespo, E., Pedraza, S., Hammond, P., & Raga, J. (2006). Abundance of small cetaceans in waters of the central Spanish Mediterranean. *Marine Biology*, 150, 149-160. doi:10.1007/s00227-006-0334-0
- Gonzalez, L., Capella, J., & Rosenbaum, H. (1994). Attack of killer whales (*Orcinus orca*) on humpback whales (*Megaptera novaeangliae*) on a South American Pacific breeding ground. *Marine Mammal Science*, 10, 218-222.
- Gonzalvo, J., Valls, M., Cardona, L., & Aguilar, A. (2008). Factors determining the interaction between common bottlenose dolphins and bottom trawlers off the Balearic Archipelago (western Mediterranean Sea). *Journal of Experimental Marine Biology & Ecology*, 367, 47-52. doi:10.1016/j.jembe.2008.08.013
- Goodson, A., & Sturtivant, C. (1996). Sonar characteristics of the harbour porpoise (*Phocoena phocoena*): source levels and spectrum. *ICES Journal of Marine*

- Science / Journal du Conseil*, 53, 465-472 Goodwin, L. (2008). Diurnal and tidal variations in habitat use of the harbour porpoise (*Phocoena phocoena*) in southwest Britain. *Aquatic Mammals*, 34, 44-53.
- Goodwin, L., & Cotton, P. A. (2004). Effects of boat traffic on the behaviour of bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 30, 279-283.
- Goodyear, J. (1993). A sonic/radio tag for monitoring dive depths and underwater movements of whales. *Journal of Wildlife Management*, 57, 503.
- Goold, J. C., & Jefferson, T. A. (2004). A note on clicks recorded from free-ranging Indo-Pacific humpback dolphins, *Sousa chinensis*. *Aquatic Mammals*, 30, 175-178.
- Gorgone, A., Haase, P., Griffith, E., & Hohn, A. (2008). Modeling Response of Target and Nontarget Dolphins to Biopsy Darting. *Journal of Wildlife Management*, 72, 926-932. doi:10.2193/2007-202
- Gorman, J. (1980). Bells toll warning to whales. *Natural History*, 89(6), 6.
- Gouabault, E. (2008). Un couple magique: Le dauphin et l'autiste»: Analyse socio anthropologique d'une figure de l'imaginaire contemporain. *A.N.A.E. Approche Neuropsychologique des Apprentissages chez l'Enfant*, 20, 146-154.
- Gouteux, B., Lebeuf, M., Muir, D., & Gagné, J. (2003). Levels and temporal trends of toxaphene congeners in beluga whales (*Delphinapterus leucas*) from the St. Lawrence Estuary, Canada. *Environmental Science & Technology*, 37, 4603-4609.
- Gowans, S., Würsig, B., & Karczmarski, L. (2007). The social structure and strategies of delphinids: Predictions based on an ecological framework. *Advances in Marine Biology*, 54, 195-294. doi:10.1016/S0065-2881(07)53003-8
- Green, M., Herzing, D., & Baldwin, J. (2007). Noninvasive methodology for the sampling and extraction of DNA from free-ranging Atlantic spotted dolphins (*Stenella frontalis*). *Molecular Ecology Notes*, 7, 1287-1292. doi:10.1111/j.1471-8286.2007.01858.x
- Gregg, J., Dudzinski, K., & Smith, H. (2007). Do dolphins eavesdrop on the echolocation signals of conspecifics?. *International Journal of Comparative Psychology*, 20, 65-88.
- Gregg, J., Dudzinski, K., & Smith, H. (2008). 3D MASC: A method for estimating relative head angle and spatial distance of Dolphins from underwater video footage. *Animal Behaviour*, 75, 1181-1186. doi:10.1016/j.anbehav.2007.09.009
- Gregory, P. R., & Rowden, A. A. (2001). Behaviour patterns of bottlenose dolphins (*Tursiops truncatus*) relative to tidal state, time-of-day, and boat traffic in Cardigan Bay, West Wales. *Aquatic Mammals*, 27, 105-113.
- Greig, T. W., Bemiss, J. A., Lyon, B. R., Bossart, G. D., & Fair, P. A. (2007). Prevalence and diversity of antibiotic resistant *Escherichia coli* in Bottlenose dolphins (*Tursiops truncatus*) from the Indian River Lagoon, Florida, and Charleston Harbor Area, South Carolina. *Aquatic Mammals*, 33, 185-194.
- Grellier, K., Hammond, P., Wilson, B., Sanders-Reed, C., & Thompson, P. (2003). Use of photo identification data to quantify mother-calf association patterns in bottlenose dolphins. *Canadian Journal of Zoology*, 81, 1421-1427.
- Grellier, K., & Wilson, B. (2003). Bottlenose dolphins using the Sound of Barra, Scotland. *Aquatic Mammals*, 29, 378-382.
- Griebel, U., & Schmid, A. (2002). Spectral sensitivity and color vision in the bottlenose dolphin (*Tursiops truncatus*). *Marine and Freshwater Behaviour and Physiology*, 35, 129-137. doi:10.1080/1023624021000014716
- Griffin, R. B., & Griffin, N. J. (2004). Temporal variation in Atlantic spotted dolphin (*Stenella frontalis*) and bottlenose dolphin (*Tursiops truncatus*) densities on the west Florida continental shelf. *Aquatic Mammals*, 30, 380-390.

- Gross, A., Kiszka, J., Van Canneyt, O., Richard, P., & Ridoux, V. (2009). A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, southwest Indian Ocean. *Estuarine Coastal & Shelf Science*, 84, 367-374. doi:10.1016/j.ecss.2009.05.017
- Gubbins, C. (2002a). Association patterns of resident bottlenose dolphins (*Tursiops truncatus*) in a South Carolina estuary. *Aquatic Mammals*, 28, 24-31.
- Gubbins, C. (2002b). Use of home ranges by resident bottlenose dolphins (*Tursiops truncatus*) in a South Carolina estuary. *Journal of Mammalogy*, 83, 178.
- Gubbins, C., McCowan, B., Lynn, S., Hooper, S., & Reiss, D. (1999). Mother-infant spatial relations in captive bottlenose dolphins, *Tursiops truncatus*. *Marine Mammal Science*, 15, 751-765.
- Guerrero-Ruiz, M., Garcia-Godos, I., & Urban R, J. (2005). Photographic match of a killer whale (*Orcinus orca*) between Peruvian and Mexican waters. *Aquatic Mammals*, 31, 438-441.
- Guerrero-Ruiz, M., Perez-Cortes M, H., Salinas Z, M., & Urban R, J. (2006). First mass stranding of killer whales (*Orcinus orca*) in the Gulf of California, Mexico. *Aquatic Mammals*, 32, 265-272.
- Guerrero-Ruiz, M., & Urban R, J. (2000). First report of remoras on two killer whales (*Orcinus orca*) in the Gulf of California, Mexico. *Aquatic Mammals*, 26, 148-150.
- Guinet, C., Barrett-Lennard, L., & Loyer, B. (2000). Co-ordinated attack behavior and prey sharing by killer whales at Crozet archipelago: Strategies for feeding on negatively-buoyant prey. *Marine Mammal Science*, 16, 829-834.
- Guise, S., Martineau, D., Beland, P., & Fournier, M. (1998). Effects of in vitro exposure of Beluga whale leukocytes to selected organochlorines. *Journal of Toxicology & Environmental Health: Part A*, 55, 479-493. doi:10.1080/009841098158287
- Guissamulo, A., & Cockcroft, V. G. (2004). Ecology and population estimates of Indo-Pacific humpback dolphins (*Sousa chinensis*) in Maputo Bay, Mozambique. *Aquatic Mammals*, 30, 94-102.
- Guitart, R., Martinez Silvestre, A., Guerrero, X., & Mateo, R. (1999). Comparative study on the fatty acid composition of two marine vertebrates: Striped dolphins and loggerhead turtles. *Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology*, 124B, 439-443.
- Gutierrez, A., Brennan, B., Rodriguez, P., & Thomas, M. (1997). Resightings and behavior of false killer whales (*Pseudorca crassidens*) in Costa Rica. *Marine Mammal Science*, 13, 307-314.
- Gutierrez, A., & Parker, N. (2000). Surface behavior of bottlenose dolphins is related to spatial arrangement of prey. *Marine Mammal Science*, 16, 287-298.
- Gygax, L. (1993). Spatial movement patterns and behaviour of two captive bottlenose dolphins (*Tursiops truncatus*): Absence of stereotyped behaviour or lack of definition? *Applied Animal Behaviour Science*, 38, 337-344. doi:10.1016/0168-1591(93)90031-J
- Gygax, L. (2002). Evolution of group size in the dolphins and porpoises: Interspecific consistency of intraspecific patterns. *Behavioral Ecology*, 13, 583-590.
- Haarr, M. L., Charlton, L. D., Terhune, J. M., & Trippel, E. A. (2009). Harbour porpoise (*Phocoena phocoena*) presence patterns at an aquaculture cage site in the Bay of Fundy, Canada. *Aquatic Mammals*, 35, 203-211.
- Haider, M., & Lindsley, D. (1964). Microvibrations in man and dolphin. *Science*, 146, 1181-1182. doi:10.1126/science.146.3648.1181

- Hale, P. T., Barreto, A. S., & Ross, G. J. B. (2000). Comparative morphology and distribution of the aduncus and *truncatus* forms of bottlenose dolphin *Tursiops* in the Indian and Western Pacific Oceans. *Aquatic Mammals*, 26, 101-110.
- Hall, A., Hugunin, K., Deaville, R., Law, R., Allchin, C., & Jepson, P. (2006). The risk of infection from polychlorinated biphenyl exposure in the harbor porpoise (*Phocoena phocoena*): A case - control approach. *Environmental Health Perspectives*, 114, 704-711. doi:10.1289/ehp.8222
- Hall, A., McConnell, B., Rowles, T., Aguilar, A., Borrell, A., Schwacke, L., et al. (2006). Individual-based model framework to assess population consequences of polychlorinated biphenyl exposure in bottlenose dolphins. *Environmental Health Perspectives*, 114, 60-64. doi:10.1289/ehp.8053
- Halsey, L. (2002). Ultimate divers. *Biologist*, 49(4), 161.
- Hamer, D., Ward, T., & McGarvey, R. (2008). Measurement, management and mitigation of operational interactions between the South Australian Sardine Fishery and short-beaked common dolphins (*Delphinus delphis*). *Biological Conservation*, 141, 2865-2878. doi:10.1016/j.biocon.2008.08.024
- Hamer, D., Ward, T., & McGarvey, R. (2009). Objective reporting of scientific results is critical for maintaining relationships with industry and achieving conservation outcomes for fisheries. *Animal Conservation*, 12, 287-288. doi:10.1111/j.1469-1795.2009.00271.x
- Hammond, P., Berggren, P., Benke, H., Borchers, D., Collet, A., Heide-Jørgensen, M., et al. (2002). Abundance of harbour porpoise and other cetaceans in the North Sea and adjacent waters. *Journal of Applied Ecology*, 39, 361-376. doi:10.1046/j.1365-2664.2002.00713.x
- Hansen, L., Schwacke, L., Mitchum, G., Hohn, A., Wells, R., Zolman, E., et al. (2004). Geographic variation in polychlorinated biphenyl and organochlorine pesticide concentrations in the blubber of bottlenose dolphins from the US Atlantic coast. *Science of the Total Environment*, 319, 147. doi:10.1016/S0048-9697(03)00371-1
- Hansen, M., Wahlberg, M., & Madsen, P. (2008). Low-frequency components in harbor porpoise (*Phocoena phocoena*) clicks: Communication signal, by-products, or artifacts? *Journal of the Acoustical Society of America*, 124, 4059-4068. doi:10.1121/1.2945154
- Hanson, M. T., & Defran, R. H. (1993). The behaviour and feeding ecology of the Pacific coast bottlenose dolphin, *Tursiops truncatus*. *Aquatic Mammals*, 19, 127-142.
- Haraguchi, K., Hisamichi, Y., & Endo, T. (2009). Accumulation and mother-to-calf transfer of anthropogenic and natural organohalogens in killer whales (*Orcinus orca*) stranded on the Pacific coast of Japan. *Science of the Total Environment*, 407, 2853-2859. doi:10.1016/j.scitotenv.2009.01.003.
- Hare, M., Cipriano, F., & Palumbi, S. (2002). Genetic evidence on the demography of speciation in allopatric dolphin species. *Evolution*, 56, 804-816.
- Harley, H. (2008). Whistle discrimination and categorization by the Atlantic bottlenose dolphin (*Tursiops truncatus*): A review of the signature whistle framework and a perceptual test. *Behavioural Processes*, 77, 243-268. doi:10.1016/j.beproc.2007.11.002**
- Harley, H., & DeLong, C. (2008). Echoic object recognition by the bottlenose dolphin. *Comparative Cognition & Behavior Reviews*, 3, 346-365. doi:10.3819/ccbr.2008.30003
- Harley, H., Putman, E., & Roitblat, H. (2003). Bottlenose dolphins perceive object features through echolocation. *Nature*, 424, 667-669. doi:10.1038/nature01846

- Harley, H., Roitblat, H., & Nachtigall, P. (1996). Object representation in the bottlenose dolphin (*Tursiops truncatus*): Integration of visual and echoic information. *Journal of Experimental Psychology: Animal Behavior Processes*, 22, 164-174. doi:10.1037/0097-7403.22.2.164
- Harlin, A., Markowitz, T., Baker, C., Würsig, B., & Honeycutt, R. (2003). Genetic structure, diversity, and historical demography of New Zealand's dusky dolphins (*Lagenorhynchus albirostris obscurus*). *Journal of Mammalogy*, 84, 702-718.
- Harper, C., & Dangler, C. (2000). Isolation and characterization of a Helicobacter sp. from the gastric mucosa of dolphins. *Applied & Environmental Microbiology*, 66, 4751.
- Hartman, K., Visser, F., & Hendriks, A. (2008). Social structure of Risso's dolphins (*Grampus griseus*) at the Azores: a stratified community based on highly associated social units. *Canadian Journal of Zoology*, 86, 294-306. doi:10.1139/Z07-138
- Hartmann, M. G (2000). The European studbook of bottlenose dolphins (*Tursiops truncatus*): 1998 Survey results. *Aquatic Mammals*, 26, 95-100.
- Harwood, L., & Smith, T. (2002). Whales of the Inuvialuit settlement region in Canada's Western Arctic: An overview and outlook. *Arctic*, 55, 77-93.
- Harzen, S. (1998). Habitat use by the bottlenose dolphin (*Tursiops truncatus*) in the Sado estuary, Portugal. *Aquatic Mammals*, 24, 117-128.
- Harzen, S., & Brunnick, B. J. (1997). Skin disorders in bottlenose dolphins (*Tursiops truncatus*), resident in the Sado estuary, Portugal. *Aquatic Mammals*, 23, 59-68.
- Harzen, S., & Santos, M. E. D. (1992). Three encounters with wild bottlenose dolphins (*Tursiops truncatus*) carrying dead calves. *Aquatic Mammals*, 18, 49-55.
- Harzen, S. E. (2002). Use of an electronic theodolite in the study of movements of the bottlenose dolphin (*Tursiops truncatus*) in the Sado estuary, Portugal. *Aquatic Mammals*, 28, 251-260.
- Hastie, G., Swift, R., Slesser, G., Thompson, P., & Turrell, W. (2005). Environmental models for predicting oceanic dolphin habitat in the Northeast Atlantic. *ICES Journal of Marine Science / Journal du Conseil*, 62, 760-770. doi:10.1016/j.icesjms.2005.02.004
- Hastie, G., Wilson, B., & Thompson, P. (2003). Fine-scale habitat selection by coastal bottlenose dolphins: Application of a new land-based video-montage technique. *Canadian Journal of Zoology*, 81, 469-479.
- Hastie, G., Wilson, B., & Thompson, P. (2006). Diving deep in a foraging hotspot: Acoustic insights into bottlenose dolphin dive depths and feeding behaviour. *Marine Biology*, 148, 1181-1188. doi:10.1007/s00227-005-0143-x
- Hastie, G., Wilson, B., Tufft, L., & Thompson, P. (2003). Bottlenose dolphins increase breathing synchrony in response to boat traffic. *Marine Mammal Science*, 19, 74-84.
- Hastie, G., Wilson, B., Wilson, L., Parsons, K., & Thompson, P. (2004). Functional mechanisms underlying cetacean distribution patterns: hotspots for bottlenose dolphins are linked to foraging. *Marine Biology*, 144, 397-403. doi:10.1007/s00227-003-1195-4
- Hatfield, B., Marks, D., Tinker, M., Nolan, K., & Peirce, J. (1998). Attacks on sea otters by killer whales. *Marine Mammal Science*, 14, 888-894.
- Hawkins, E. R., & Gartside, D. F. (2009a). Interactive behaviours of bottlenose dolphins (*Tursiops aduncus*) during encounters with vessels. *Aquatic Mammals*, 35, 259-268.

- Hawkins, E. R., & Gartside, D. F. (2009b). Patterns of whistles emitted by wild Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) during a provisioning program. *Aquatic Mammals*, 35, 171-186.
- Hayes, W. (1959). Wave-riding dolphins. *Science*, 130, 1657-1658.
- Heath, M., & Ridgway, S. (1999). How dolphins use their blubber to avoid heat stress during encounters with warm water. *American Journal of Physiology*, 276, R1188.
- Heide-Jørgensen, M., Hammeken, N., Dietz, R., Richard, P., & Orr, J. (2001). Surfacing times and dive rates for narwhals (*Monodon monoceros*) and belugas (*Delphinapterus leucas*). *Arctic*, 54, 284.
- Heide-Jørgensen, M., & Reeves, R. (1996). Evidence of a decline in beluga, *Delphinapterus leucas*, abundance off West Greenland. *ICES Journal of Marine Science / Journal du Conseil*, 53, 61-72.
- Heide-Jørgensen, M., Richard, P., Dietz, R., Laidre, K., Orr, J., & Schmidt, H. (2003). An estimate of the fraction of belugas (*Delphinapterus leucas*) in the Canadian high Arctic that winter in West Greenland. *Polar Biology*, 26, 318-326. doi:10.1007/s00300-003-0488-x
- Heimeier, D., Hutt, A., Stone, G., Duignan, P., Baker, C., & Russell, K. (2009). Confirmed expression of MHC class I and class II genes in the New Zealand endemic Hector's dolphin (*Cephalorhynchus hectori*). *Marine Mammal Science*, 25, 68-90.
- Heise, K., Barrett-Lennard, L. G., Saulitis, E., Matkin, C., & Bain, D. (2003). Examining the evidence for killer whale predation on Steller sea lions in British Columbia and Alaska. *Aquatic Mammals*, 29, 325-334.
- Heithaus, M. (2001). Shark attacks on bottlenose dolphins (*Tursiops aduncus*) in Shark Bay, Western Australia: Attack rate, bite scar frequencies, and attack seasonality. *Marine Mammal Science*, 17, 526-539.
- Heithaus, M., & Dill, L. M. (2002). Food availability and tiger shark predation risk influence bottlenose dolphin habitat use. *Ecology*, 83, 480-491.
- Heithaus, M., & Dill, L. M. (2006). Does tiger shark predation risk influence foraging habitat use by bottlenose dolphins at multiple spatial scales? *Oikos*, 114, 257-264. doi:10.1111/j.2006.0030-1299.14443.x
- Helweg, D., Au, W., Roitblat, H., & Nachtigall, P. (1996). Acoustic basis for recognition of aspect-dependent three-dimensional targets by an echolocating bottlenose dolphin. *Journal of the Acoustical Society of America*, 99, 2409-2420. doi:10.1121/1.415429
- Helweg, D., & Hautus, M. (1996). Recognition of aspect-dependent three-dimensional objects by an echolocating Atlantic bottlenose.. *Journal of Experimental Psychology: Animal Behavior Processes*, 22, 19-31.
- Helweg, D., Moore, P., Dankiewicz, L., Zafran, J., & Brill, R. (2003). Discrimination of complex synthetic echoes by an echolocating bottlenose dolphin. *Journal of the Acoustical Society of America*, 113, 1138-1144. doi:10.1121/1.1531175
- Helweg, D., Roitblat, H., Nachtigall, P., & Hautus, M. (1996). Recognition of aspect-dependent three-dimensional objects by an echolocating Atlantic bottlenose dolphin. *Journal of Experimental Psychology: Animal Behavior Processes*, 22, 19-31. doi:10.1037/0097-7403.22.1.19
- Henley, D. (1992). Facilitating artistic expression in captive mammals: Implications for art therapy and art empathicism. *Art Therapy*, 9, 178-192.
- Herman, L. (1975). Interference and auditory short-term memory in the bottlenosed dolphin. *Animal Learning & Behavior*, 3, 43-48.
- Herman, L. (1980). *Cetacean behavior: Mechanisms and functions*. New York: Wiley-Interscience.**

- Herman, L., Abichandani, S., Elhajj, A., Herman, E., Sanchez, J., & Pack, A. (1999). Dolphins (*Tursiops truncatus*) comprehend the referential character of the human pointing gesture. *Journal of Comparative Psychology*, 113, 347-364. doi:10.1037/0735-7036.113.4.347
- Herman, L., & Arbeit, W. (1972). Frequency difference limens in the bottlenose dolphin: 1-70 Kc/s. *Journal of Auditory Research*, 12, 109-120.
- Herman, L., & Arbeit, W. (1973). Stimulus control and auditory discrimination learning sets in the bottlenose dolphin. *Journal of the Experimental Analysis of Behavior*, 19, 379- 394. doi:10.1901/jeab.1973.19-379
- Herman, L., Beach, F., Pepper, R., & Stallings, R. (1969). Learning-set formation in the bottlenose dolphin. *Psychonomic Science*, 14(3), 98-99.
- Herman, L., & Forestell, P. (1985). Reporting presence or absence of named objects by a language-trained dolphin. *Neuroscience & Biobehavioral Reviews*, 9, 667-681. doi:10.1016/0149-7634(85)90013-2
- Herman, L., & Gordon, J. (1974). Auditory delayed matching in the bottlenose dolphin. *Journal of the Experimental Analysis of Behavior*, 21, 19-26. doi:10.1901/jeab.1974.21-19
- Herman, L., Hovancik, J., Gory, J., & Bradshaw, G. (1989). Generalization of visual matching by a bottlenosed dolphin (*Tursiops truncatus*): Evidence for invariance of cognitive performance with visual and auditory materials. *Journal of Experimental Psychology: Animal Behavior Processes*, 15, 124-136. doi:10.1037/0097-7403.15.2.124
- Herman, L., Kuczaj, S., & Holder, M. (1993). Responses to anomalous gestural sequences by a language-trained dolphin: Evidence for processing of semantic relations and syntactic information. *Journal of Experimental Psychology: General*, 122, 184-194. doi:10.1037/0096-3445.122.2.184
- Herman, L., Matus, D., Herman, E., Ivancic, M., & Pack, A. (2001). The bottlenosed dolphin's (*Tursiops truncatus*) understanding of gestures as symbolic representations of its body parts. *Animal Learning & Behavior*, 29, 250-264.
- Herman, L., Morrel-Samuels, P., & Pack, A. (1990). Bottlenosed dolphin and human recognition of veridical and degraded video displays of an artificial gestural language. *Journal of Experimental Psychology: General*, 119, 215-230. doi:10.1037/0096-3445.119.2.215
- Herman, L., Pack, A., & Hoffmann-Kuhnt, M. (1998). Seeing through sound: Dolphins (*Tursiops truncatus*) perceive the spatial structure of objects through echolocation. *Journal of Comparative Psychology*, 112, 292-305. doi:10.1037/0735-7036.112.3.292
- Herman, L., Pack, A., & Wood, A. (1994). Bottlenose dolphins can generalize rules and develop abstract concepts. *Marine Mammal Science*, 10, 70-80.
- Herman, L., Richards, D., & Wolz, J. (1984). Comprehension of sentences by bottlenosed dolphins. *Cognition*, 16, 129-219. doi:10.1016/0010-0277(84)90003-9
- Herman, L., & Thompson, R. (1982). Symbolic, identity, and probe delayed matching of sounds by the bottlenosed dolphin. *Animal Learning & Behavior*, 10, 22-34.
- Herman, L., & Uyeyama, R. (1999). The dolphin's grammatical competency: Comments on Kako (1999). *Animal Learning & Behavior*, 27, 18-23.
- Herman, L., Uyeyama, R., & Pack, A. (2008). Bottlenose dolphins understand relationships between concepts. *Behavioral & Brain Sciences*, 31, 139-140. doi:10.1017/S0140525X08003658

- Hernandez, E. N., Kuczaj, S., Houser, D. S., & Finneran, J. J. (2007). Middle- and long-latency auditory evoked potentials in bottlenose dolphins (*Tursiops truncatus*) resulting from frequent and oddball stimuli. *Aquatic Mammals*, 33, 34-42.
- Hernandez-Milian, G., Goetz, S., Varela-Dopico, C., Rodriguez-Gutierrez, J., Romón-Olea, J., Fuertes-Gamundi, J., et al. (2008). Results of a short study of interactions of cetaceans and longline fisheries in Atlantic waters: Environmental correlates of catches and depredation events. *Hydrobiologia*, 612, 251-268. doi:10.1007/s10750-008-9501-2
- Hernández-Mora, G., González-Barrientos, R., Morales, J., Chaves-Olarte, E., Guzmán-Verri, C., Baquero Calvo, E., et al. (2008). Neurobrucellosis in stranded dolphins, Costa Rica. *Emerging Infectious Diseases*, 14, 1430-1433. doi:10.3201/eid1409.071056
- Herraez, P., Sierra, E., Arbelo, M., Jaber, J., Espinosa de los Monteros, A., & Fernandez, A. (2007). Rhabdomyolysis and myoglobinuric nephrosis (capture myopathy) in a striped dolphin. *Journal of Wildlife Diseases*, 43, 770-774.
- Herring, P. (1998). Dolphins glow with the flow. *Nature*, 393, 731-733.
- Herzing, D. L. (1996). Vocalizations and associated underwater behavior of free-ranging Atlantic spotted dolphins, *Stenella frontalis*, and bottlenose dolphins, *Tursiops truncatus*. *Aquatic Mammals*, 22, 61-79.
- Herzing, D. L. (2006). The currency of cognition: Assessing tools, techniques, and media for complex behavioral analysis. *Aquatic Mammals*, 32, 544-553.
- Herzing, D. L., & Brunnick, B. J. (1997). Coefficients of association of reproductively active female Atlantic spotted dolphins, *Stenella frontalis*. *Aquatic Mammals*, 23, 155-162.
- Herzing, D. L., & Johnson, C. M. (1997). Interspecific interactions between Atlantic spotted dolphins (*Stenella frontalis*) and bottlenose dolphins (*Tursiops truncatus*) in the Bahamas, 1985-1995. *Aquatic Mammals*, 23, 85-99.
- Herzing, D. L., Moewe, K., & Brunnick, B. J. (2003). Interspecies interactions between Atlantic spotted dolphins, *Stenella frontalis*, and bottlenose dolphins, *Tursiops truncatus*, on Great Bahama Bank, Bahamas. *Aquatic Mammals*, 29, 335-341.
- Hess, E. H. (1953). Comparative psychology. *Annual Review of Psychology*, 4, 239-254.
- Heyning, J. (1988). Presence of solid food in a young calf killer whale (*Orcinus orca*). *Marine Mammal Science*, 4, 68-71.
- Hickie, B., Ross, P., MacDonald, R., & Ford, J. K. B. (2007). Killer whales (*Orcinus orca*) face protracted health risks associated with lifetime exposure to PCBs. *Environmental Science & Technology*, 41, 6613-6619. doi:10.1021/es0702519
- Hicks, B., St. Aubin, D., Geraci, J., & Brown, W. (1985). Epidermal growth in the Bottlenose dolphin, *Tursiops truncatus*. *Journal of Investigative Dermatology*, 85, 60-63. doi:10.1111/1523-1747.ep12275348
- Highfill, L. E., & Kuczaj, S. A., II. (2007). Do bottlenose dolphins (*Tursiops truncatus*) have distinct and stable personalities? *Aquatic Mammals*, 33, 380-389.**
- Hill, H. (2009). The behavioral development of two beluga calves during the first year of life. *International Journal of Comparative Psychology*, 22, 234-253.
- Hill, H., Carder, D., & Ridgway, S. (2008). Vigilance in female bottlenose dolphins (*Tursiops* sp.) before and after calving. *International Journal of Comparative Psychology*, 21, 35-57.
- Hill, H., Greer, T., Solangi, M., & Kuczaj II, S. (2007). All mothers are not the same: Maternal styles in bottlenose dolphins (*Tursiops truncatus*). *International Journal of Comparative Psychology*, 20, 35-54.

- Hobbs, R., Laidre, K., Vos, D., Mahoney, B., & Eagleton, M. (2005). Movements and area use of belugas, *Delphinapterus leucas*, in a Subarctic Alaskan Estuary. *Arctic*, 58, 331-340.
- Hobbs, R., Rugh, D., & Demaster, D. (2000). Abundance of belugas, in Cook Inlet, Alaska, 1994-2000. *Marine Fisheries Review*, 62(3), 37-46.
- Hobbs, R., Waite, J., & Rugh, D. (2000). Beluga, group sizes in Cook Inlet, Alaska, based on observer counts and aerial video. *Marine Fisheries Review*, 62(3), 46-59.
- Hoelzel, A. (1991). Killer whale predation on marine mammals at Punta Norte, Argentina: Food sharing, provisioning and foraging strategy. *Behavioral Ecology and Sociobiology*, 29, 197-204. doi:10.1007/BF00166401
- Hoelzel, A. (1993). Foraging behaviour and social group dynamics in Puget Sound killer whales. *Animal Behaviour*, 45, 581-591. doi:10.1006/anbe.1993.1068
- Hoelzel, A. (1994). Genetics and ecology of whales and dolphins. *Annual Review of Ecology and Systematics*, 25, 377-399.
- Hoelzel, A., Dahlheim, M., & Stern, S. (1998). Low genetic variation among killer whales (*Orcinus orca*) in the eastern North Pacific and.. *Journal of Heredity*, 89, 121.
- Hoelzel, A., Ford, J. K. B., & Dover, G. (1991). A paternity test case for the killer whale (*Orcinus orca*) by DNA fingerprinting. *Marine Mammal Science*, 7, 35-43.
- Hof, P., Glezer, L., Nimchinsky, E., & Erwin, J. (2000). Neurochemical and cellular specializations in the mammalian neocortex reflect phylogenetic relationships: Evidence from primates, cetaceans, and artiodactyls. *Brain, Behavior & Evolution*, 55, 300. doi:10.1159/000006665
- Hogg, C., Vickers, E., & Rogers, T. (2005). Determination of testosterone in saliva and blow of bottlenose dolphins (*Tursiops truncatus*) using liquid chromatography–mass spectrometry. *Journal of Chromatography B: Analytical Technologies in the Biomedical & Life Sciences*, 814, 339-346. doi:10.1016/j.jchromb.2004.10.058
- Holden, C. (1993). Whale alarm. *Science*, 260, 896.
- Holden, C. (1998). Tales of springy whales. *Science*, 279, 329.
- Holden, C. (1999). Flipper makes a pass. *Science*, 284, 737.
- Holden, C. (2004). Flipping over a new tail. *Science*, 306, 1678.
- Holden, C. (2005). Cetacean culture? *Science*, 308, 1545.
- Holden, C. (2006). Avoiding whale-ship hits. *Science*, 314, 1663.
- Holt, M., Veirs, V., & Veirs, S. (2008). Noise effects on the call amplitude of Southern resident killer whales (*Orcinus orca*). *Bioacoustics*, 17, 164-166.
- Honma, Y., Ushiki, T., Hashizume, H., Takeda, M., Matsuishi, T., & Honno, Y. (2004). Histological observations on the reproductive organs of harbor porpoises *Phocoena phocoena* incidentally caught in a set net installed off Usujiri, southern Hokkaido. *Fisheries Science*, 70, 94-99. doi:10.1111/j.1444-2906.2003.00776.x
- Hooker, S., & Baird, R. (2001). Diving and ranging behaviour of odontocetes: A methodological review and critique. *Mammal Review*, 31, 81-105. doi:10.1046/j.1365-2907.2001.00080.x
- Hooper, S., Reiss, D., Carter, M., & McCowan, B. (2006). Importance of contextual saliency on vocal imitation by bottlenose dolphins. *International Journal of Comparative Psychology*, 19, 116-128.
- Houde, M., Bujas, T., Small, J., Wells, R., Fair, P., Bossart, G., et al. (2006). Biomagnification of perfluoroalkyl compounds in the bottlenose dolphin (*Tursiops truncatus*) food web. *Environmental Science & Technology*, 40, 4138-4144.

- Houde, M., Measures, L., & Huot, J. (2003). Lungworm (*Pharurus pallasii*: Metastrongyloidea: *Pseudaliidae*) infection in the endangered St. Lawrence beluga whale (*Delphinapterus leucas*). *Canadian Journal of Zoology*, 81, 543.
- Houde, M., Pacepavicius, G., Darling, C., Fair, P., Alaee, M., Bossart, G., et al. (2009). Polybrominated diphenyl ethers and their hydroxylated analogs in plasma of bottlenose dolphins (*Tursiops truncatus*) from the United States East coast. *Environmental Toxicology & Chemistry*, 28, 2061-2068.
- Houde, M., Pacepavicius, G., Wells, R., Fair, P., Letcher, R., Alaee, M., et al. (2006). Polychlorinated biphenyls and hydroxylated polychlorinated biphenyls in plasma of bottlenose dolphins (*Tursiops truncatus*) from the Western Atlantic and the Gulf of Mexico. *Environmental Science & Technology*, 40, 5860-5866. doi:10.1021/es060629n
- Houde, M., Wells, R., Fair, P., Bossart, G., Hohn, A., Rowles, T., et al. (2005). Polyfluoroalkyl compounds in free-ranging bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Mexico and the Atlantic Ocean. *Environmental Science & Technology*, 39, 6591-6598.
- Houser, D., Dankiewicz, L., Stockard, T., & Ponganis, P. (2008). Ultrasound inspection for intravascular bubbles in a repetitively diving dolphin. *Bioacoustics*, 17, 310-312.
- Houser, D., & Finneran, J. (2005). Auditory evoked potentials (AEP) methods for population level assessment of hearing sensitivity in bottlenose dolphins. *Journal of the Acoustical Society of America*, 117, 2408.
- Houser, D., & Finneran, J. (2006). Variation in the hearing sensitivity of a dolphin population determined through the use of evoked potential audiometry. *Journal of the Acoustical Society of America*, 120, 4090-4099. doi:10.1121/1.2357993
- Houser, D., Finneran, J., Carder, D., Van Bonn, W., Smith, C., Hoh, C., et al. (2004). Structural and functional imaging of bottlenose dolphin (*Tursiops truncatus*) cranial anatomy. *Journal of Experimental Biology*, 207, 3657-3665. doi:10.1242/jeb.01207
- Houser, D., Gomez-Rubio, A., & Finneran, J. (2008). Evoked potential audiometry of 13 Pacific bottlenose dolphins (*Tursiops truncatus gilli*). *Marine Mammal Science*, 24, 28-41.
- Houser, D., Martin, S., Bauer, E., Phillips, M., Herrin, T., Cross, M., et al. (2005). Echolocation characteristics of free-swimming bottlenose dolphins during object detection and identification. *Journal of the Acoustical Society of America*, 117, 2308-2317. doi:10.1121/1.1867912
- Houser, D., & Finneran, J. J. (2006). A comparison of underwater hearing sensitivity in bottlenose dolphins (*Tursiops truncatus*) determined by electrophysiological and behavioral methods. *The Journal of the Acoustical Society of America*, 120, 1713-1722.
- Howells, E. M., Reif, J. S., Bechdel, S. E., Murdoch, M. E., Bossart, G. D., McCulloch, S. D., et al. (2009). A novel case of non-offspring adoption in a free-ranging atlantic bottlenose dolphin (*Tursiops truncatus*) inhabiting the Indian River Lagoon, Florida. *Aquatic Mammals*, 35, 43-47.
- Hubard, C. W., Maze-Foley, K., Mullin, K. D., & Schroeder, W. W. (2004). Seasonal abundance and site fidelity of bottlenose dolphins (*Tursiops truncatus*) in Mississippi sound. *Aquatic Mammals*, 30, 299-310.
- Hubbard, J., Hansen, D., & Mahoney, B. (1999). Winter sighting of beluga whales (*Delphinapterus leucas*) in Yakutat-Disenchantment Bay, Alaska. *Arctic*, 52, 411-412.

- Hui, C. (1994). Lack of association between magnetic patterns and the distribution of free ranging dolphins. *Journal of Mammalogy*, 75, 399-405.
- Hung, C., Lau, R., Lam, J., Jefferson, T., Hung, S., Lam, M., et al. (2007). Risk assessment of trace elements in the stomach contents of Indo-Pacific Humpback Dolphins and Finless Porpoises in Hong Kong waters. *Chemosphere*, 66, 1175-1182. doi:10.1016/j.chemosphere.2006.08.005
- Hung, C., So, M., Connell, D., Fung, C., Lam, M., Nicholson, S., et al. (2004). A preliminary risk assessment of trace elements accumulated in fish to the Indo-Pacific Humpback dolphin (*Sousa chinensis*) in the Northwestern waters of Hong Kong. *Chemosphere*, 56, 643-651. doi:10.1016/j.chemosphere.2004.04.016
- Hung, C., Xu, Y., Lam, J., Jefferson, T., Hung, S., Yeung, L., et al. (2006). An assessment of the risks associated with polychlorinated biphenyls found in the stomach contents of stranded Indo-Pacific Humpback Dolphins (*Sousa chinensis*) and Finless Porpoises (*Neophocaena phocaenoides*) from Hong Kong waters. *Chemosphere*, 63, 845-852. doi:10.1016/j.chemosphere.2005.07.059
- Hung, S. K., & Jefferson, T. A. (2004). Ranging patterns of Indo-Pacific humpback dolphins (*Sousa chinensis*) in the Pearl River Estuary, People's Republic of China. *Aquatic Mammals*, 30, 159-174.
- Huntington, H. (2000). Traditional knowledge of the ecology of belugas, in Cook Inlet, Alaska. *Marine Fisheries Review*, 62(3), 134-140.
- Hurtley, S. (2005). Dolphin Culture. *Science*, 309, 222.
- Ibsen, S., Au, W., Nachtigall, P., DeLong, C., & Breese, M. (2007). Changes in signal parameters over time for an echolocating Atlantic bottlenose dolphin performing the same target discrimination task. *Journal of the Acoustical Society of America*, 122, 2446-2450. doi:10.1121/1.2772213
- Iglesias, E. (1982). Falklands war threatens whales. *Environment*, 24(5), 24.
- Ilyukhina, V., Krivoshchapova, M., & Manzhosova, G. (2008). Psychophysiological study of the effects of adaptation to tactile interaction with dolphins in six- to seven-year-old children. *Human Physiology*, 34, 421-430. doi:10.1134/S0362119708040051
- Iniguez, M. A. (2001). Seasonal distribution of killer whales (*Orcinus orca*) in Northern Patagonia, Argentina. *Aquatic Mammals*, 27, 154-161.
- Iniguez, M. A., & De Haro, J. C. (1994). Preliminary report on the feeding habits of the Peale's dolphin (*Lagenorhynchus albirostris australis*) in southern Argentina. *Aquatic Mammals*, 20, 35-37.
- Iniguez, M. A., & Tossenberger, V. P. (2007). Commerson's dolphins (*Cephalorhynchus commersonii*) off Ria Deseado, Patagonia, Argentina. *Aquatic Mammals*, 33, 276-285.
- Innes, S., Lavigne, D.M., Earle, W.M. & Kovacs, K.M. (1987). Feeding rates of seal sand whales. *Journal of Animal Ecology*, 56, 115-130.
- Inoue, Y., Itou, T., Jimbo, T., Sakai, T., Ueda, K., & Imajoh-Ohmi, S. (2001). Molecular cloning and identification of bottle-nosed dolphin p40phox, p47phox and p67phox. *Veterinary Immunology and Immunopathology*, 78, 21-33.
- International Whaling Commission. (1989). Whale numbers down in the Southern Hemisphere. *Bioscience*, 39, 662.
- Irvine, A., Scott, M., Wells, R., & Kaufmann, J. (1981). Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fishery Bulletin United States, National Marine Fisheries Service*, 79, 671-688.
- Irwin, L. J. (2005). Marine toxins: Adverse health effects and biomonitoring with resident coastal dolphins. *Aquatic Mammals*, 31, 195-225.

- Ivanov, M. (2004). Dolphin's echolocation signals in a complicated acoustic environment. *Acoustical Physics*, 50, 469-479. doi:10.1134/1.1776226
- Iwanami, K., Mita, H., Yamamoto, Y., Fujise, Y., Yamada, T., & Suzuki, T. (2006). cDNA-derived amino acid sequences of myoglobins from nine species of whales and dolphins. *Comparative Biochemistry & Physiology Part B*, 145, 249-256. doi:10.1016/j.cbpb.2006.08.002
- Kumagai, J. (2006). How much sound is too much? *IEEE Spectrum*, 43, 60.
- Jaakkola, K., Fellner, W., Erb, L., Rodriguez, M., & Guarino, E. (2005). Understanding of the concept of numerically "less" by bottlenose dolphins (*Tursiops truncatus*). *Journal of Comparative Psychology*, 119, 296-303. doi:10.1037/0735-7036.119.3.296
- Jaber, J., Fernández, A., Herráez, P., Espinosa de los Monteros, A., Ramírez, G., García, P., et al. (2003). Cross-reactivity of human and bovine antibodies in striped dolphin paraffin wax embedded tissues. *Veterinary Immunology & Immunopathology*, 96, 65. doi:10.1016/S0165-2427(03)00158-2
- Jackson, J., Patenaude, N., Carroll, E., & Baker, C. (2008). How few whales were there after whaling? Inference from contemporary mtDNA diversity. *Molecular Ecology*, 17, 236-251. doi:10.1111/j.1365-294X.2007.03497.x
- Jacobsen, T., Mayntz, M., & Amundin, M. (2003). Splitting suckling data of bottlenose dolphin (*Tursiops truncatus*) neonates in human care into suckling bouts. *Zoo Biology*, 22, 477-488. doi:10.1002/zoo.10094
- Janik, V. (1999). Pitfalls in the categorization of behaviour: A comparison of dolphin whistle classification methods. *Animal Behaviour*, 57, 133-143. doi:10.1006/anbe.1998.0923
- Janik, V. (2000). Whistle matching in wild bottlenose dolphins (*Tursiops truncatus*). *Science*, 289, 1355-1357. doi:10.1126/science.289.5483.1355
- Janik, V., Dehnhardt, G., & Todt, D. (1994). Signature whistle variations in a bottlenosed dolphin, *Tursiops truncatus*. *Behavioral Ecology and Sociobiology*, 35, 243-248. doi:10.1007/BF00170704
- Janik, V., Sayigh, L., & Wells, R. (2006). Signature whistle shape conveys identity information to bottlenose dolphins. *PNAS Proceedings of the National Academy of Sciences of the United States of America*, 103, 8293-8297. doi:10.1073/pnas.0509918103
- Janik, V., & Slater, P. (1998). Context-specific use suggests that bottlenose dolphin signature whistles are cohesion calls. *Animal Behaviour*, 56, 829-838. doi:10.1006/anbe.1998.0881
- Janik, V., Van Parijs, S., & Thompson, P. (2000). A two-dimensional acoustic localization system for marine mammals. *Marine Mammal Science*, 16, 437-447.
- Jefferson, T. A. (1996). Morphology of the Clymene dolphin (*Stenella clymene*) in the northern Gulf of Mexico. *Aquatic Mammals*, 22, 35-43.
- Jefferson, T. A. (2000). Population biology of the Indo-Pacific Hump-backed dolphin in Hong Kong. *Wildlife Monographs*, 144, 1-66.
- Jefferson, T. A., Fertl, D., Bolaños-Jiménez, J., & Zerbini, A. (2009). Distribution of common dolphins (*Delphinus spp.*) in the western Atlantic Ocean: A critical re-examination. *Marine Biology*, 156, 1109-1124. doi:10.1007/s00227-009-1152-y
- Jefferson, T. A., & Hung, S. K. (2004). A review of the status of the Indo-Pacific humpback dolphin (*Sousa chinensis*) in Chinese waters. *Aquatic Mammals*, 30, 149-158.

- Jefferson, T. A., & Hung, S. K. (2008). Effects of biopsy sampling on Indo-Pacific humpback dolphins (*Sousa chinensis*) in a polluted coastal environment. *Aquatic Mammals*, 34, 310-316.
- Jefferson, T. A., & Van Waerebeek, K. (2004). Geographic variation in skull morphology of humpback dolphins (*Sousa* spp.). *Aquatic Mammals*, 30, 3-17.
- Jelinski, D., Krueger, C., & Duffus, D. (2002). Geostatistical analyses of interactions between killer whales (*Orcinus orca*) and recreational whale-watching boats. *Applied Geography*, 22, 393-412.
- Jensen, F., Bejder, L., Wahlberg, M., & Madsen, R. (2009). Biosonar adjustments to target range of echolocating bottlenose dolphins (*Tursiops* sp.) in the wild. *Journal of Experimental Biology*, 212, 1078-1086. doi:10.1242/jeb.025619
- Jepson, P., Arbelo, M., Deaville, R., Patterson, I., Castro, P., Baker, J., et al. (2003). Gas-bubble lesions in stranded cetaceans. *Nature*, 425(6958), 575-576. doi:10.1038/425575a
- Jianqiang, X., & Ding, W. (2006). Respiratory pattern of captive Yangtze finless porpoises (*Neophocaena phocaenoides asiaeorientalis*). *Journal of Ethology*, 24, 205-212. doi:10.1007/s10164-005-0181-3
- Joanna Y., W., Randall, W., Alex, A., Asuncion, B., Victoria, T., Peter, R., et al. (2007). Correlates of cytochrome P450 1A1 expression in Bottlenose dolphin (*Tursiops truncatus*) integument biopsies. *Toxicological Sciences*, 97, 111-119.
- Johnson, C., McManus, M., & Skaar, D. (1989). Masked tonal hearing thresholds in the beluga whale. *Journal of the Acoustical Society of America*, 85, 2651-2654. doi:10.1121/1.397759
- Johnson, M. (2005). Playing tag with whales. *Oceanus*, 43(2), 57-58.
- Johnson, S., Venn-Watson, S., Cassle, S., Smith, C., Jensen, E., & Ridgway, S. (2009). Use of phlebotomy treatment in Atlantic bottlenose dolphins with iron overload. *Journal of the American Veterinary Medical Association*, 235, 194-200.
- Johnson-Restrepo, B., Adams, D., & Kannan, K. (2008). Tetrabromobisphenol A (TBBPA) and hexabromocyclododecanes (HBCDs) in tissues of humans, dolphins, and sharks from the United States. *Chemosphere*, 70, 1935-1944. doi:10.1016/j.chemosphere.2007.10.002
- Johnson-Restrepo, B., Kannan, K., Addink, R., & Adams, D. (2005). Polybrominated diphenyl ethers and polychlorinated biphenyls in a marine foodweb of coastal Florida. *Environmental Science & Technology*, 39, 8243-8250. doi:10.1021/es051551y
- Johnston, D. W., & Woodley, T. H. (1998). A survey of acoustic harassment device (AHD) use in the Bay of Fundy, NB, Canada. *Aquatic Mammals*, 24, 51-61.
- Jones, F. M., & Pfeiffer, C. J. (1994). Morphometric comparison of the epidermis in several cetacean species. *Aquatic Mammals*, 20, 29-34.
- Jones, G., & Sayigh, L. (2002). Geographic variation in rates of vocal production of free-ranging bottlenose dolphins. *Marine Mammal Science*, 18, 374-393.
- Kajiwara, N., Kamikawa, S., Ramu, K., Ueno, D., Yamada, T., Subramanian, A., et al. (2006). Geographical distribution of polybrominated diphenyl ethers (PBDEs) and organochlorines in small cetaceans from Asian waters. *Chemosphere*, 64, 287-295. doi:10.1016/j.chemosphere.2005.12.013
- Kakuschke, A., & Prange, A. (2007). The influence of metal pollution on the immune system a potential stressor for marine mammals in the North Sea. *International Journal of Comparative Psychology*, 20, 179-193.

- Kalinichenko, S., & Pushchin, I. (2008). Calcium-binding proteins in the cerebellar cortex of the bottlenose dolphin and harbour porpoise. *Journal of Chemical Neuroanatomy*, 35, 364-370. doi:10.1016/j.jchemneu.2008.03.003
- Kamminga, C., Stuart, A. C., & Silber, G. K. (1996). Investigations on cetacean sonar: XI. Intrinsic comparison of the wave shapes of some members of the Phocoenidae family. *Aquatic Mammals*, 22, 45-55.
- Kannan, K., & Senthilkumar, K. (1997). Elevated accumulation of tributyltin and its breakdown products in bottlenose dolphins (*Tursiops* sp.). *Environmental Science & Technology*, 31, 296.
- Kaplan, J. D., & Connor, R. (2007). A preliminary examination of sex differences in tactile interactions among juvenile Atlantic spotted dolphins (*Stenella frontalis*) *Marine Mammal Science*, 23, 943-953.
- Karandeeva, O. (1969). Features of dolphin behavior in captivity and their training. *Zhurnal Vysshei Nervnoi Deyatel'nosti*, 19(1), 55-58.
- Karczmarski, L. (2000). Conservation and management of humpback dolphins: the South African perspective. *Oryx*, 34, 207-216.
- Karczmarski, L., & Cockcroft, V. G. (1998). Matrix photo-identification technique applied in studies of free-ranging bottlenose and humpback dolphins. *Aquatic Mammals*, 24, 143-147.
- Karczmarski, L., Cockcroft, V., & McLachlan, A. (2000). Habitat use and preferences of Indo-Pacific humpback dolphins *Sousa chinensis* in Algoa Bay, South Africa. *Marine Mammal Science*, 16, 65-79.
- Karczmarski, L., Thornton, M., & Cockcroft, V. G. (1997). Description of selected behaviours of humpback dolphins *Sousa chinensis*. *Aquatic Mammals*, 23, 127-133.
- Karczmarski, L., Thornton, M., & Cockcroft, V. (2000). Daylight occurrence of humpback dolphins *Sousa chinensis* in Algoa Bay, South Africa. *African Journal of Ecology*, 38, 86-90. doi:10.1046/j.1365-2028.2000.00215.x
- Karczmarski, L., Würsig, B., Gailey, G., Larson, K., & Vanderlip, C. (2005). Spinner dolphins in a remote Hawaiian atoll: Social grouping and population structure. *Behavioral Ecology*, 16, 675-685. doi:10.1093/beheco/ari028
- Kasamatsu, M., Kawauchi, R., Tsunokawa, M., Ueda, K., Uchida, E., Oikawa, S., et al. (2009). Comparison of serum lipid compositions, lipid peroxide, α -tocopherol and lipoproteins in captive marine mammals (bottlenose dolphins, spotted seals and West Indian manatees) and terrestrial mammals. *Research in Veterinary Science*, 86, 216-222. doi:10.1016/j.rvsc.2008.07.006
- Kasamatsu, M., Tsunokawa, M., Taki, M., Higuchi, H., & Nagahata, H. (2001). Serum lipid peroxide and alpha-tocopherol concentrations and superoxide dismutase activity in captive bottle-nosed dolphins. *American Journal of Veterinary Research*, 62, 1952-1956.
- Kastelein, R., De Haan, D., & Verboom, W. (2007). The influence of signal parameters on the sound source localization ability of a harbor porpoise (*Phocoena phocoena*). *Journal of the Acoustical Society of America*, 122, 1238-1248. doi:10.1121/1.2747202
- Kastelein, R. A., Dokter, T., & Hilgenkamp, J. (1995). A swimming support for dolphins undergoing veterinary care. *Aquatic Mammals*, 21, 155-159.
- Kastelein, R. A., Hagedoorn, M., Au, W., & de Haan, D. (2003). Audiogram of a striped dolphin (*Stenella coeruleoalba*). *Journal of the Acoustical Society of America*, 113, 1130-1137. doi:10.1121/1.1532310

- Kastelein, R. A., Janssen, M., Verboom, W., & De Haan, D. (2005). Receiving beam patterns in the horizontal plane of a harbor porpoise (*Phocoena phocoena*). *Journal of the Acoustical Society of America*, 118, 1172-1179. doi:10.1121/1.1945565
- Kastelein, R. A., Jennings, N., Verboom, W., de Haan, D., & Schooneman, N. (2006). Differences in the response of a striped dolphin (*Stenella coeruleoalba*) and a harbour porpoise (*Phocoena phocoena*) to an acoustic alarm. *Marine Environmental Research*, 61, 363-378. doi:10.1016/j.marenvres.2005.11.005
- Kastelein, R. A., & Lavaleije, M. S. S. (1992). Foreign bodies in the stomach of a female Harbour porpoise (*Phocoena phocoena*) from the North Sea. *Aquatic Mammals*, 18, 40-46.
- Kastelein, R. A., & Mosterd, J. (1995). Improving parental care of a female bottlenose dolphin (*Tursiops truncatus*) by training. *Aquatic Mammals*, 21, 165-169.
- Kastelein, R. A., Mosterd, J., Schooneman, N. M., & Wiepkema, P. R. (2000). Food consumption, growth, body dimensions, and respiration rates of captive false killer whales (*Pseudorca crassidens*). *Aquatic Mammals*, 26, 33-44.
- Kastelein, R. A., van Dooren, M. F., & Tibboel, D. (2009). A case study of congenital diaphragmatic hernia in a juvenile striped dolphin (*Stenella coeruleoalba*). *Aquatic Mammals*, 35, 32-35.
- Kastelein, R., Vaughan, N., Walton, S., & Wiepkema, P. (2002). Food intake and body measurements of Atlantic bottlenose dolphins (*Tursiops truncatus*) in captivity. *Marine Environmental Research*, 53, 199-218.
- Kastelein, R. A., Verboom, W., Muijsers, M., Jennings, N., & van der Heul, S. (2005). The influence of acoustic emissions for underwater data transmission on the behaviour of harbour porpoises (*Phocoena phocoena*) in a floating pen. *Marine Environmental Research*, 59, 287-307. doi:10.1016/j.marenvres.2004.05.005
- Kastelein, R. A., Verlaan, M., & Jennings, N. (2008). Number and duration of echolocation click trains produced by a harbor porpoise (*Phocoena phocoena*) in relation to target and performance. *Journal of the Acoustical Society of America*, 124, 40-43. doi:10.1121/1.2924132
- Kasuya, T. (2007). Japanese whaling and other cetacean fisheries. *Environmental Science and Pollution Research International*, 14, 39-48.
- Katsufumi, S., Yutaka, W., Akinori, T., Hideji, T., Ryo, K., Paul J., P., et al. (2007). Stroke frequency, but not swimming speed, is related to body size in free-ranging seabirds, pinnipeds and cetaceans. *Proceedings: Biological Sciences*, 274, 471-477.
- Kawano, M., Falandysz, J., & Morita, M. (2008). Instrumental neutron activation analysis of extractable organohalogens in marine mammal, harbour porpoise (*Phocoena phocoena*) and its feed, Atlantic herring (*Clupea harengus*), from the Baltic Sea. *Journal of Radioanalytical & Nuclear Chemistry*, 278, 263-266. doi:10.1007/s10967-008-0401-8
- Kehrig, H., Seixas, T., Palermo, E., Di Beneditto, A., Souza, C., & Malm, O. (2008). Different species of mercury in the livers of tropical dolphins. *Analytical Letters*, 41, 1691-1699. doi:10.1080/00032710802122677
- Keith, M., Bester, M., Bartlett, P., & Baker, D. (2001). Killer whales (*Orcinus orca*) at Marion Island, Southern Ocean. *African Zoology*, 36, 161-173.
- Keith, M., Peddemors, V., Bester, M., & Ferguson, J. (2002). Population characteristics of Indo Pacific humpback dolphins at Richards Bay, South Africa: Implications for incidental capture in shark nets. *South African Journal of Wildlife Research*, 32, 153-162.

- Kellar, N., Danil, K., Archer, F., Chivers, S., Trego, M., & Marks, C. (2009). Blubber testosterone: A potential marker of male reproductive status in short-beaked common dolphins. *Marine Mammal Science*, 25, 507-522.
- Kellogg, W. N. (1958). On the psychological study of small whales. *Journal of Psychology: Interdisciplinary and Applied*, 46(97), 100.
- Kellogg, W., & Rice, C. (1963). Visual discrimination in a bottlenose porpoise. *The Psychological Record*, 13, 483-498.
- Kellogg, W., & Rice, C. (1964). Visual problem solving in a bottlenose dolphin. *Science*, 143, 1053-1055.
- Kern, A., Seidel, K., & Oelschläger, H. (2009). The central vestibular complex in dolphins and humans: Functional implications of Deiters' nucleus. *Brain, Behavior & Evolution*, 73, 102-110. doi:10.1159/000213646
- Ketten, D., & Madin, K. (2004). How to see what whales hear. *Oceanus*, 43(2), 59-62.
- Kilborn, S. (1994). Object carrying in a captive beluga whale (*Delphinapterus leucas*) as possible surrogate behavior. *Marine Mammal Science*, 10, 496-501.
- Kilian, A., von Fersen, L., & Güntürkün, O. (2000). Lateralization of visuospatial processing in the bottlenose dolphin (*Tursiops truncatus*). *Behavioural Brain Research*, 116, 211-215. doi:10.1016/S0166-4328(00)00273-4
- Kilian, A., von Fersen, L., & Güntürkün, O. (2005). Left hemispheric advantage for numerical abilities in the bottlenose dolphin. *Behavioural Processes*, 68, 179-184. doi:10.1016/j.beproc.2004.11.003
- Kilian, A., Yaman, S., von Fersen, L., & Güntürkün, O. (2003). A bottlenose dolphin discriminates visual stimuli differing in numerosity. *Learning & Behavior*, 31, 133-142.
- Killebrew, D. A., Mercado, E., III, Herman, L. M., & Pack, A. A. (2001). Sound production of a neonate bottlenose dolphin. *Aquatic Mammals*, 27, 34-44.
- Kim, G., & Tanabe, S. (1996). Accumulation of butyltin compounds in Risso's dolphin (*Grampus griseus*) from the Pacific Coast of. *Environmental Science & Technology*, 30(8), 2620-2625.
- Kimura, S., Akamatsu, T., Wang, K., Wang, D., Li, S., Dong, S., et al. (2009). Comparison of stationary acoustic monitoring and visual observation of finless porpoises. *Journal of the Acoustical Society of America*, 125, 547-553. doi:10.1121/1.3021302
- Kinghorn, A., Humphries, M., Outridge, P., & Chan, H. (2008). Teeth as biomonitor of selenium concentrations in tissues of beluga whales (*Delphinapterus leucas*). *Science of the Total Environment*, 402, 43-50. doi:10.1016/j.scitotenv.2008.04.031
- Kingsley, M. (1998). Population index estimates for the St. Lawrence belugas, 1973-1995. *Marine Mammal Science*, 14, 508-530.
- Kingston, S., & Rosel, P. (2004). Genetic differentiation among recently diverged delphinid taxa determined using AFLP markers. *Journal of Heredity*, 95, 1-10. doi:10.1093/jhered/esh010
- Klatsky, L., Wells, R., & Sweeney, J. (2007). Offshore bottlenose dolphins (*Tursiops truncatus*): Movement and dive behavior near the Bermuda Pedestal. *Journal of Mammalogy*, 88, 59-66.
- Klissarov, D. V. Z., & Milushev, G. (1992). Successful therapy on upper-jaw fracture in a pregnant bottlenose dolphin (*Tursiops truncatus*) in the Varna Dolphinarium. *Aquatic Mammals*, 18, 47-48.

- Knieriem, A., & Hartmann, M. G. (2001). Comparative histopathology of lungs from by-caught Atlantic white-sided dolphins (*Leucopleurus acutus*). *Aquatic Mammals*, 27, 73-81.
- Knoff, A., Hohn, A., & Macko, S. (2008). Ontogenetic diet changes in bottlenose dolphins (*Tursiops truncatus*) reflected through stable isotopes. *Marine Mammal Science*, 24, 128-137.
- Knowlton, N. (1993). Sibling species in the sea. *Annual Review of Ecology and Systematics*, 24, 189-216.
- Kock, K., Purves, M., & Duhamel, G. (2006). Interactions between Cetacean and Fisheries in the Southern Ocean. *Polar Biology*, 29, 379-388. doi:10.1007/s00300-005-0067-4
- Kogi, K., Hishii, T., Imamura, A., Iwatani, T., & Dudzinski, K. M. (2004). Demographic parameters of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) around Mikura Island, Japan. *Marine Mammal Science*, 20, 510-526.
- Kon-No, A., Ashahina, K., Yumioka, C., Kobayashi, H., Arahata, T., & Murayama, T. (2005). Discrimination of achromatic colors in a bottlenose dolphin. *Japanese Journal of Animal Psychology*, 55, 59-64. doi:10.2502/janip.55.59
- Koopman, H. (2007). Phylogenetic, ecological, and ontogenetic factors influencing the biochemical structure of the blubber of odontocetes. *Marine Biology*, 151, 277-291. doi:10.1007/s00227-006-0489-8
- Koopman, H., Iverson, S., & Read, A. (2003). High concentrations of isovaleric acid in the fats of odontocetes: variation and patterns of accumulation in blubber vs. stability in the melon. *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 173, 247-261. doi:10.1007/s00360-003-0329-9
- Koopman, H., Pabst, D., McLellan, W., Dillaman, R., & Read, A. (2002). Changes in blubber distribution and morphology associated with starvation in the Harbor porpoise (*Phocoena phocoena*): Evidence for regional differences in blubber structure and function. *Physiological & Biochemical Zoology*, 75, 498-512.
- Krahn, M., Herman, D., Matkin, C., Durban, J., Barrett-Lennard, L., Burrows, D., et al. (2007). Use of chemical tracers in assessing the diet and foraging regions of eastern North Pacific killer whales. *Marine Environmental Research*, 63, 91-114. doi:10.1016/j.marenvres.2006.07.002
- Krasnova, V., Bel'kovich, V., & Chernetsky, A. (2006). Mother-infant spatial relations in wild beluga (*Delphinapterus leucas*) during postnatal development under natural conditions. *Biology Bulletin*, 33, 53-58. doi:10.1134/S1062359006010079
- Kroeger, R. H. H., & Kirschfeld, K. (1994). Refractive index in the cornea of a harbor porpoise (*Phocoena phocoena*) measured by two-wavelengths laser-interferometry. *Aquatic Mammals*, 20, 99-107.
- Krützen, M., Barré, L., Connor, R., Mann, J., & Sherwin, W. (2004). ‘O father: Where art thou?’— Paternity assessment in an open fission–fusion society of wild bottlenose dolphins (*Tursiops* sp.) in Shark Bay, Western Australia. *Molecular Ecology*, 13, 1975-1990. doi:10.1111/j.1365-294X.2004.02192.x
- Krützen, M., Barré, L., Moeller, L., Heithaus, M., Simms, C., & Sherwin, W. (2002). A biopsy system for small cetaceans: Darting success and wound healing in *Tursiops* spp. *Marine Mammal Science*, 18, 863-878.
- Krützen, M., Mann, J., Heithaus, M., Connor, R., Bejder, L., & Sherwin, W. (2005). Cultural transmission of tool use in bottlenose dolphins. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 8939-8943. doi:10.1073/pnas.0500232102

- Krützen, M., Sherwin, W., Berggren, P., & Gales, N. (2004). Population structure in an inshore cetacean revealed by microsatellite and mtDNA analysis: Bottlenose dolphins (*Tursiops* sp.) in Shark Bay, Western Australia. *Marine Mammal Science*, 20, 28-47.
- Krushinskii, L., Dashevskii, B., Krushinskaya, N., & Dmitrieva, I. (1972). An investigation of the ability of *Tursiops truncatus* (Montagu) dolphins to utilize empirical figural metrics. *Doklady Akademii Nauk SSSR*, 204, 755-758.
- Kuczaj, S., Gory, J., & Xitco, M. (2009). How intelligent are dolphins? A partial answer based on their ability to plan their behavior when confronted with novel problems. *Japanese Journal of Animal Psychology*, 59, 99-115. doi:10.2502/janip.59.1.9**
- Kuczaj, S., & Highfill, L. (2005). Dolphin play: Evidence for cooperation and culture?. *Behavioral & Brain Sciences*, 28, 705-706.
- Kuczaj, S., Makecha, R., Trone, M., Paulos, R., & Ramos, J. (2006). Role of peers in cultural innovation and cultural transmission: Evidence from the play of dolphin calves. *International Journal of Comparative Psychology*, 19, 223-240.
- Kuczaj, S., Solangi, M., Hoffland, T., & Romagnoli, M. (2008). Recognition and discrimination of human actions across the senses of echolocation and vision in the bottlenose dolphin: Evidence for dolphin cross-modal integration of dynamic information. *International Journal of Comparative Psychology*, 21, 84-95.
- Kuehl, D., Haebler, R., Potter, C., Lahvis, G., Donahue, M., & Regal, R. (2009). PFOS and PFOSA in bottlenose dolphins: An investigation into two unusually high mortality epizootics. *Reproductive Toxicology*, 27, 421. doi:10.1016/j.reprotox.2008.11.070
- Kurihara, N., & Oda, S.-i. (2006). Cranial variation and taxonomic revision of bottlenose dolphins (*Tursiops* spp.) from Japanese waters. *Aquatic Mammals*, 32, 289-300.
- Kuttin, E. S., & Kaller, A. (1996). Cystoisospora delhini n. Sp. causing enteritis in a bottlenosed dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 22, 57-59.
- Kyhn, L., Tougaard, J., Jensen, F., Wahlberg, M., Stone, G., Yoshinaga, A., et al. (2009). Feeding at a high pitch: Source parameters of narrow band, high-frequency clicks from echolocating off-shore hourglass dolphins and coastal Hector's dolphins. *Journal of the Acoustical Society of America*, 125, 1783-1791. doi:10.1121/1.3075600
- Kyngdon, D., Minot, E., & Stafford, K. (2003). Behavioural responses of captive common dolphins *Delphinus delphis* to a 'Swim-with-Dolphin' programme. *Applied Animal Behaviour Science*, 81, 163-171. doi:10.1016/S0168-1591(02)00255-1
- Lacave, G., Eggermont, M., Verslycke, T., Brook, F., Salbany, A., Roque, L., et al. (2004). Prediction from ultrasonographic measurements of the expected delivery date in two species of bottlenosed dolphin (*Tursiops truncatus* and *Tursiops aduncus*). *Veterinary Record: Journal of the British Veterinary Association*, 154, 228-233.
- Lacoboni, M. (2009). Imitation, empathy, and mirror neurons. *Annual Review of Psychology*, 60, 653-670.
- Ladygina, T., & Supin, A. (1970). On the acoustic projection in the cortex of the dolphin brain. *Fiziologicheskii Zhurnal SSSR*, 56, 1554-1560.
- Lahaye, V., Bustamante, P., Dabin, W., Churlaud, C., & Caurant, F. (2007). Trace element levels in foetus-mother pairs of short-beaked common dolphins (*Delphinus delphis*) stranded along the French coasts. *Environment International*, 33, 1021-1028. doi:10.1016/j.envint.2007.05.008
- Laidre, K., Shelden, K., Rugh, D., & Mahoney, B. (2000). Beluga, distribution and survey effort in the Gulf of Alaska. *Marine Fisheries Review*, 62(3), 27-37.

- Lam, J., Lau, R., Murphy, M., & Lam, P. (2009). Temporal trends of hexabromocyclododecanes (hbcds) and polybrominated diphenyl ethers (pbdes) and detection of two novel flame retardants in marine mammals from Hong Kong, South China. *Environmental Science & Technology*, 43, 6944-6949.
- LaMere, S., Leger, J., Schrenzel, M., Anthony, S., Rideout, B., & Salomon, D. (2009). Molecular characterization of a novel gammaretrovirus in killer whales (*Orcinus orca*). *Journal of Virology*, 83, 12956-12967. doi:10.1128/JVI.01354-09
- Lammers, M., & Au, W. (2003). Directionality in the whistles of Hawaiian spinner dolphins (*Stenella longirostris*): A signal feature to cue direction of movement? *Marine Mammal Science*, 19, 249-264.
- Lammers, M., Au, W., & Herzing, D. (2003). The broadband social acoustic signaling behavior of spinner and spotted dolphins. *Journal of the Acoustical Society of America*, 114, 1629-1639. doi:10.1121/1.1596173
- Lammers, M., Schotten, M., & Au, W. (2006). The spatial context of free-ranging Hawaiian spinner dolphins (*Stenella longirostris*) producing acoustic signals. *Journal of the Acoustical Society of America*, 119, 1244-1250. doi:10.1121/1.2151804
- Lammers, M. (2004). Occurrence and behavior of Hawaiian spinner dolphins (*Stenella longirostris*) along Oahu's leeward and south shores. *Aquatic Mammals*, 30, 237-250.
- Lang, T., & Smith, H. (1965). Communications between dolphins in separate tanks by way of an electronic acoustic link. *Science*, 150, 1839-1844. doi:10.1126/science.150.3705.1839
- Larese, J., & Chivers, S. (2009). Growth and reproduction of female eastern and whitebelly spinner dolphins incidentally killed in the eastern tropical Pacific tuna purse-seine fishery. *Canadian Journal of Zoology*, 87, 537-552. doi:10.1139/Z09-038
- Larsen, F., Eigaard, O., & Tougaard, J. (2007). Reduction of harbour porpoise (*Phocoena phocoena*) bycatch by iron-oxide gillnets. *Fisheries Research*, 85, 270-278. doi:10.1016/j.fishres.2007.02.011
- Laura, S., López-Sánchez, J., Aguayo-Lobo, A., & Medrano-González, L. (2004). Morphometry and sexual dimorphism of the coastal spotted dolphin, *Stenella attenuata* graffmani, from Bahía de Banderas, Mexico. *Acta Zoologica*, 85, 223-232. doi:10.1111/j.0001-7272.2004.00175.x
- Lavery, T., Butterfield, N., Kemper, C., Reid, R., & Sanderson, K. (2008). Metals and selenium in the liver and bone of three dolphin species from South Australia, 1988–2004. *Science of the Total Environment*, 390, 77-85. doi:10.1016/j.scitotenv.2007.09.016
- Lavery, T., Kemper, C., Sanderson, K., Schultz, C., Coyle, P., Mitchell, J., et al. (2009). Heavy metal toxicity of kidney and bone tissues in South Australian adult bottlenose dolphins (*Tursiops aduncus*). *Marine Environmental Research*, 67, 1-7. doi:10.1016/j.marenvres.2008.09.005
- Law, R., Allchin, C., & Morris, R. (1995). Uptake of organochlorines (Chlorobiphenyls, Dieldrin; total PCB & DDT) in bottlenose dolphins (*Tursiops truncatus*) from Cardigan Bay, West Wales. *Chemosphere*, 30, 547-560.
- Leatherwood, S., & Reeves, R. (1990). *The bottlenose dolphin*. Chicago: University of Chicago.
- Leatherwood, S., & Reeves, R. R. (1994). River dolphins: A review of activities and plans of the cetacean specialist group. *Aquatic Mammals*, 20, 137-154.
- Leatherwood, S., Reeves, R., & Foster, L. (1983). *Sierra Club handbook of whales and dolphins*.** San Francisco: Sierra Club Books.

- Lebeuf, M., Noël, M., Trottier, S., & Measures, L. (2007). Temporal trends (1987–2002) of persistent, bioaccumulative and toxic (PBT) chemicals in beluga whales (*Delphinapterus leucas*) from the St. Lawrence Estuary, Canada. *Science of the Total Environment*, 383, 216-231. doi:10.1016/j.scitotenv.2007.04.026
- Lefebvre, L., Marino, L., Sol, D., Lemieux-Lefebvre, S., & Arshad, S. (2006). Large brains and lengthened life history periods in Odontocetes. *Brain, Behavior And Evolution*, 68, 218-228.
- Lemon, M., Cato, D., Lynch, T., & Harcourt, R. (2008). Short-term behavioural response of bottlenose dolphins, *Tursiops aduncus*, to recreational powerboats. *Bioacoustics*, 17, 171-173.
- Lemon, M., Lynch, T., Cato, D., & Harcourt, R. (2006). Response of travelling bottlenose dolphins (*Tursiops aduncus*) to experimental approaches by a powerboat in Jervis Bay, New South Wales, Australia. *Biological Conservation*, 127, 363-372. doi:10.1016/j.biocon.2005.08.016
- Lende, R., & Welker, W. (1972). An unusual sensory area in the cerebral neocortex of the bottlenose dolphin (*Tursiops truncatus*). *Brain Research*, 45, 555-560. doi:10.1016/0006-8993(72)90482-9
- Lennert-Cody, C., & Berk, R. (2007). Statistical learning procedures for monitoring regulatory compliance: an application to fisheries data. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 170, 671-689.
- Lerczak, J., Shelden, K., & Hobbs, R. (2000). Application of suction-cup-attached VHF transmitters to the study of beluga surfacing behavior in Cook Inlet, Alaska. *Marine Fisheries Review*, 62(3), 99-111.
- Lesage, V., Barrette, C., Kingsley, M., & Sjare, B. (1999). The effect of vessel noise on the vocal behavior of belugas in the St. Lawrence River Estuary, Canada. *Marine Mammal Science*, 15, 65-84.
- Letcher, R., Norstrom, R. J., Muir, D., Sandau, C., Koczanski, K., Michaud, R., et al. (2000). Methylsulfone polychlorinated biphenyl and 2,2-bis(chlorophenyl)-1,1-dichloroethylene metabolites in beluga whale (*Delphinapterus leucas*) from the St. Lawrence River estuary and western Hudson Bay, Canada. *Environmental Toxicology & Chemistry*, 19, 1378.
- Levin, M., Morsey, B., & De Guise, S. (2007). Non-coplanar PCBs induce calcium mobilization in Bottlenose dolphin and Beluga whale, but not in mouse leukocytes. *Journal of Toxicology & Environmental Health: Part A*, 70, 1220-1231. doi:10.1080/15287390701380898
- Levin, M., Morsey, B., Mori, C., & De Guise, S. (2004). Specific non-coplanar PCB-mediated modulation of bottlenose dolphin and beluga whale phagocytosis upon in vitro exposure. *Journal of Toxicology & Environmental Health: Part A*, 67, 1517-1535.
- Levy, B. (1992). Psychoaesthetics dolphin project. *Art Therapy*, 9(4), 193-197.
- Lewis, A., Hammill, M., Power, M., Dodge, D., & Lesage, V. (2009). Movement and aggregation of Eastern Hudson Bay beluga Whales (*Delphinapterus leucas*): A comparison of patterns found through satellite telemetry and Nunavik traditional ecological knowledge. *Arctic*, 62, 13-24.
- Lilly, J. (1962). Vocal behavior of the bottlenose dolphin. *Proceedings of the American Philosophical Society*, 106, 520-529.
- Lilly, J., & Miller, A. (1962). Operant conditioning of the bottlenose dolphin with electrical stimulation of the brain. *Journal of Comparative and Physiological Psychology*, 55, 73-79. doi:10.1037/h0043540

- Lilly, J., Miller, A., & Truby, H. (1968). Reprogramming of the sonic output of the dolphin: Sonic burst count matching. *Journal of the Acoustical Society of America*, 43, 1412- 1424. doi:10.1121/1.1911001
- Link, W. A., Doherty, P. F. (2002). Scaling in sensitivity analysis. *Ecology*, 83, 3299-3305.
- Litwiler, T., & Cronin, T. (2001). No evidence of accommodation in the eyes of the bottlenose dolphin, *Tursiops truncatus*. *Marine Mammal Science*, 17, 508-525.
- Litz, J., Garrison, L., Fieber, L., Martinez, A., Contillo, J., & Kucklick, J. (2007). Fine-scale spatial variation of persistent organic pollutants in bottlenose dolphins (*Tursiops truncatus*) in Biscayne Bay, Florida. *Environmental Science & Technology*, 41, 7222-7228.
- Liu, J., & Hills, P. (1997). Environmental planning, biodiversity and the development process: The case of Hong Kong's Chinese white dolphins. *Journal of Environmental Management*, 50, 351-367.
- Lockhart, W., Stern, G., Wagemann, R., Hunt, R., Metner, D., DeLaronde, J., et al. (2005). Concentrations of mercury in tissues of beluga whales (*Delphinapterus leucas*) from several communities in the Canadian Arctic from 1981 to 2002. *Science of the Total Environment*, 351-352, 391-412. doi:10.1016/j.scitotenv.2005.01.050
- Lockyer, C., Heide-Jørgensen, M., Jensen, J., Kinze, C., & Sørensen, T. (2001). Age, length and reproductive parameters of harbour porpoises *Phocoena phocoena* from West Greenland. *ICES Journal of Marine Science / Journal du Conseil*, 58, 154-162. doi:10.1006/jmsc.2000.0998
- Lopez, B. D. (2006). Bottlenose dolphin (*Tursiops truncatus*) predation on a marine fin fish farm: Some underwater observations. *Aquatic Mammals*, 32, 305-310.
- Loseto, L., Richard, P., Stern, G., Orr, J., & Ferguson, S. (2006). Segregation of Beaufort Sea beluga whales during the open-water season. *Canadian Journal of Zoology*, 84, 1743-1751. doi:10.1139/Z06-160
- Loseto, L., Stern, G., Connelly, T., Deibel, D., Gemmill, B., Prokopowicz, A., et al. (2009). Summer diet of beluga whales inferred by fatty acid analysis of the eastern Beaufort Sea food web. *Journal of Experimental Marine Biology & Ecology*, 374, 12-18. doi:10.1016/j.jembe.2009.03.015
- Loseto, L., Stern, G., & Ferguson, S. (2008). Size and biomagnification: How habitat selection explains beluga mercury levels. *Environmental Science & Technology*, 42, 3982-3988.
- Lotze, H. K. & Milewski, I. (2004). Two centuries of multiple human impacts and successive changes in a North Atlantic food web. *Ecological Applications*, 14, 1428-1447.
- Luca, M., Andrew, W., Emer, R., Patricia, R., Andrew, R., Jamie, C., et al. (2009). Population structure of short-beaked common dolphins (*Delphinus delphis*) in the North Atlantic Ocean as revealed by mitochondrial and nuclear genetic markers. *Marine Biology*, 156, 821-834. doi:10.1007/s00227-008-1120-y
- Lucke, K., Lepper, P. A., Hoeve, B., Everaarts, E., van Elk, N., & Siebert, U. (2007). Perception of low-frequency acoustic signals by a harbour porpoise (*Phocoena phocoena*) in the presence of simulated offshore wind turbine noise. *Aquatic Mammals*, 33, 55-68.
- Lucy E., R., & Stephen M., D. (2008). Laser photogrammetry to determine dorsal fin size in a population of bottlenose dolphins from Doubtful Sound, New Zealand. *Australian Journal of Zoology*, 56, 239-248.
- Luczkovich, J., Daniel, H., Hutchinson, M., Jenkins, T., Johnson, S., Pullinger, R., et al. (2000). Sounds of sex and death in the sea: Bottlenose dolphin whistles suppress mating choruses of silver perch. *Bioacoustics*, 10, 323-334.

- Lukina, L. (1999). Influence of dolphin-assisted therapy sessions on the functional state of children with psychoneurological symptoms of diseases. *Human Physiology*, 25, 676- 679.
- Lusseau, D. (2003a). Effects of tour boats on the behavior of bottlenose dolphins: Using Markov chains to model anthropogenic impacts. *Conservation Biology*, 17, 1785-1793. doi:10.1111/j.1523-1739.2003.00054.x
- Lusseau, D. (2003b). The emergent properties of a dolphin social network. *Proceedings: Biological Sciences*, 270, 186-188.
- Lusseau, D. (2004). The hidden cost of tourism: Detecting long-term effects of tourism using behavioral information. *Ecology & Society*, 9(1), 2.
- Lusseau, D. (2006a). The short-term behavioral reactions of bottlenose dolphins to interactions with boats in Doubtful Sound, New Zealand. *Marine Mammal Science*, 22, 802-818.
- Lusseau, D. (2006b). Why do dolphins jump? Interpreting the behavioural repertoire of bottlenose dolphins (*Tursiops* sp.) in Doubtful Sound, New Zealand. *Behavioural Processes*, 73, 257-265.
- Lusseau, D. (2007). Evidence for social role in a dolphin social network. *Evolutionary Ecology*, 21, 357-366.
- Lusseau, D., & Conradt, L. (2009). The emergence of unshared consensus decisions in bottlenose dolphins. *Behavioral Ecology and Sociobiology*, 63, 1067-1077. doi:10.1007/s00265-009-0740-7
- Lusseau, D., & Newman, M. E. J. (2004). Identifying the role that animals play in their social networks. *Proceedings: Biological Sciences*, 271, S477-S481.
- Lusseau, D., Schneider, K., Boisseau, O., Haase, P., Slooten, E., & Dawson, S. (2003). The bottlenose dolphin community of Doubtful Sound features a large proportion of long-lasting associations: Can geographic isolation explain this unique trait?. *Behavioral Ecology and Sociobiology*, 54, 396-405. doi:10.1007/s00265-003-0651-y
- Lusseau, D., Whitehead, H., & Gero, S. (2008). Incorporating uncertainty into the study of animal social networks. *Animal Behaviour*, 75, 1809-1815. doi:10.1016/j.anbehav.2007.10.029
- Lusseau, D., Williams, R., Wilson, B., Grellier, K., Barton, T., Hammond, P., et al. (2004). Parallel influence of climate on the behaviour of Pacific killer whales and Atlantic bottlenose dolphins. *Ecology Letters*, 7, 1068-1076. doi:10.1111/j.1461-0248.2004.00669.x
- Lusseau, D., Wilson, B., Hammond, P., Grellier, K., Durban, J., Parsons, K., et al. (2006). Quantifying the influence of sociality on population structure in bottlenose dolphins. *Journal of Animal Ecology*, 75, 14-24. doi:10.1111/j.1365-2656.2005.01013.x
- Lyamin, O., Mukhametov, L., Siegel, J., Nazarenko, E., Polyakova, I., & Shpak, O. (2002). Unihemispheric slow wave sleep and the state of the eyes in a white whale. *Behavioural Brain Research*, 129, 125-129.
- Lyamin, O., Pryaslova, J., Kosenko, P., & Siegel, J. (2007). Behavioral aspects of sleep in bottlenose dolphin mothers and their calves. *Physiology & Behavior*, 92, 725-733. doi:10.1016/j.physbeh.2007.05.064
- Lyamin, O., Pryaslova, J., Lance, V., & Siegel, J. (2005). Animal behaviour: Continuous activity in cetaceans after birth. *Nature*, 435, 1177. doi:10.1038/4351177a
- Lyamin, O., Shpak, O., Nazarenko, E., & Mukhametov, L. (2002). Muscle jerks during behavioral sleep in a beluga whale (*Delphinapterus leucas* L.). *Physiology & Behavior*, 76, 265-270. doi:10.1016/S0031-9384(02)00722-9

- Ma, H., Overstreet, R., Snieszek, J., Solagni, M., & Wayne Coats, D. (2006). Two new species of symbiotic ciliates from the respiratory tract of cetaceans with establishment of the new genus *Planilamina* n. gen. (*Dysterida*, *Kyaroikeidae*). *Journal of Eukaryotic Microbiology*, 53, 407-419. doi:10.1111/j.1550-7408.2006.00124.x
- Madsen, P., Kerr, I., & Payne, R. (2004). Echolocation clicks of two free-ranging, oceanic delphinids with different food preferences: false killer whales, *Pseudorca crassidens*, and Risso's dolphins, *Grampus griseus*. *Journal of Experimental Biology*, 207, 1811-1823.
- Madsen, P., & Wahlberg, M. (2007). Recording and quantification of ultrasonic echolocation clicks from free-ranging toothed whales. *Deep-Sea Research Part I, Oceanographic Research Papers*, 54, 1421-1444. doi:10.1016/j.dsr.2007.04.020
- Maggi, R., Nielsen, O., Ford, S., Breitschwerdt, E., Robinson, J., Lester, S., et al. (2008). Bartonella henselae in captive and hunter-harvested beluga (*Delphinapterus leucas*). *Journal of Wildlife Diseases*, 44, 871-877.
- Maier, R. (1968). Effects of stress upon social distance in dolphins. *Perceptual and Motor Skills*, 27, 862.
- Maldini, D., Mazzuca, L., & Atkinson, S. (2005). Odontocete stranding patterns in the main Hawaiian Islands (1937-2002): How do they compare with live animal surveys? *Pacific Science*, 59, 55-67.
- Mancia, A., Lundqvist, M., Romano, T., Peden-Adams, M., Fair, P., Kindy, M., et al. (2007). A dolphin peripheral blood leukocyte cDNA microarray for studies of immune function and stress reactions. *Developmental and Comparative Immunology*, 31, 520-529.
- Mancia, A., Romano, T., Gefroh, H., Chapman, R., Middleton, D., Warr, G., et al. (2006). The Immunoglobulin G Heavy Chain (IGHG) genes of the Atlantic bottlenose dolphin, *Tursiops truncatus*. *Comparative Biochemistry & Physiology Part B*, 144, 38-46. doi:10.1016/j.cpb.2006.01.014
- Mancia, A., Romano, T., Gefroh, H., Chapman, R., Middleton, D., Warr, G., et al. (2007). Characterization of the immunoglobulin A heavy chain gene of the Atlantic bottlenose dolphin (*Tursiops truncatus*). *Veterinary Immunology and Immunopathology*, 118, 304-309.
- Mancia, A., Warr, G., & Chapman, R. (2008). A transcriptomic analysis of the stress induced by capture-release health assessment studies in wild dolphins (*Tursiops truncatus*). *Molecular Ecology*, 17, 2581-2589. doi:10.1111/j.1365-294X.2008.03784.x
- Manconi, B., Messana, I., Maggiani, F., Olianas, A., Pellegrini, M., Crnjar, R., et al. (2009). Structural and functional characterization of *Delphinus delphis* hemoglobin system. *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 179, 971-983. doi:10.1007/s00360-009-0380-2
- Manger, P., Fuxe, K., Ridgway, S., & Siegel, J. (2004). The distribution and morphological characteristics of catecholaminergic cells in the diencephalon and midbrain of the Bottlenose dolphin (*Tursiops truncatus*). *Brain, Behavior & Evolution*, 64, 42-60. doi:10.1159/000077542
- Manger, P., Ridgway, S., & Siegel, J. (2003). The locus coeruleus complex of the bottlenose dolphin (*Tursiops truncatus*) as revealed by tyrosine hydroxylase immunohistochemistry. *Journal of Sleep Research*, 12, 149-155. doi:10.1046/j.1365-2869.2003.00350.x

- Manger, P., Sum, M., Szymanski, M., Ridgway, S., & Krubitzer, L. (1998). Modular subdivisions of dolphin insular cortex: Does evolutionary history repeat itself?. *Journal of Cognitive Neuroscience*, 10, 153-166. doi:10.1162/089892998562627
- Maniscalco, J. M., Matkin, C. O., Maldini, D., Calkins, D. G., & Atkinson, S. (2007). Assessing killer whale predation on steller sea lions from field observations in Kenai Fjords, Alaska. *Marine Mammal Science*, 23, 306-321. doi:10.1111/j.1748-7692.2007.00103.x
- Mann, J. (2001). Cetacean culture: Definitions and evidence. *Behavioral & Brain Sciences*, 24, 343.
- Mann, J., & Barnett, H. (1999). Lethal tiger shark (*Galeocerdo cuvier*) attack on bottlenose dolphin (*Tursiops* sp.) calf: Defense and reactions by the mother. *Marine Mammal Science*, 15(2), 568-575.
- Mann, J., Connor, R. C., Barre, L. M., Heithaus M. R. (2000). Female reproductive success in bottlenose dolphins (*Tursiops* sp.): Life history, habitat, provisioning, and group-size effects. *Behavioral Ecology*, 11, 210-219.
- Mann, J., Connor, R. C., Tyack, P. L., Whitehead, H. (2000). *Cetacean societies: Field studies of dolphins and whales*. Chicago: The University of Chicago Press.**
- Mann, D., Cook, M., Bauer, G., Fellner, W., & Wells, R. (2005). Auditory evoked potential measurements with cetaceans. *Journal of the Acoustical Society of America*, 117, 2407-2408.
- Mann, J., & Smuts, B. (1998). Natal attraction: Allomaternal care and mother-infant separations in wild bottlenose dolphins. *Animal Behaviour*, 55, 1097-1113. doi:10.1006/anbe.1997.0637
- Mann, J., & Smuts, B. (1999). Behavioral developments in wild bottlenose dolphin newborns (*Tursiops* sp.). *Behaviour*, 136, 529-566. doi:10.1163/156853999501469
- Mann, J., & Watson-Capps, J. (2005). Surviving at sea: Ecological and behavioural predictors of calf mortality in Indian Ocean bottlenose dolphins, *Tursiops* sp. *Animal Behaviour*, 69, 899-909. doi:10.1016/j.anbehav.2004.04.024
- Maresh, J. L., Fish, F. E., Nowacek, D. P., Nowacek, S. M., & Wells, R. S. (2004). High performance turning capabilities during foraging by bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science*, 20(3), 498-509.
- Margulis, S., & Westhus, E. (2008). Evaluation of different observational sampling regimes for use in zoological parks. *Applied Animal Behaviour Science*, 110, 363-376. doi:10.1016/j.applanim.2007.05.001
- Marino, L. (1999). Brain growth in the harbor porpoise and Pacific white-sided dolphin. *Journal of Mammalogy*, 80, 1353.
- Marino, L., & Lilienfeld, S. (2007). Dolphin-assisted therapy: More flawed data and more flawed conclusions. *Anthrozoos*, 20, 239-249. doi:10.2752/089279307X224782
- Marino, L., Rilling, J., Lin, S., & Ridgway, S. (2000). Relative volume of the cerebellum in dolphins and comparison with anthropoid primates. *Brain, Behavior & Evolution*, 56(4), 204. doi:10.1159/000047205
- Marino, L., & Stowe, J. (1997). Lateralized behavior in two captive bottlenose dolphins (*Tursiops truncatus*). *Zoo Biology*, 16, 173-177. doi:10.1002/(SICI)1098-2361(1997)16:2<173::AID-ZOO7>3.0.CO;2-7
- Marino, L., Sudheimer, K., Pabst, D. A., McLellan, W. A., Arshad, S., Naini, G., et al. (2004). Anatomical description of an infant bottlenose dolphin (*Tursiops truncatus*) brain from magnetic resonance images. *Aquatic Mammals*, 30, 315-326.

- Markowitz, T., Harlin, A., & Wuersig, B. (2003). Digital photography improves efficiency of individual dolphin identification. *Marine Mammal Science*, 19(1), 217-223.
- Marques, F., & Buckland, S. (2003). Incorporating Covariates into Standard Line Transect Analyses. *Biometrics*, 59, 924-935.
- Marsh, G., Athanasiadou, M., Athanassiadis, I., Bergman, Å., Endo, T., & Haraguchi, K. (2005). Identification, quantification, and synthesis of a novel dimethoxylated polybrominated biphenyl in marine mammals caught off the coast of Japan. *Environmental Science & Technology*, 39, 8684-8690. doi:10.1021/es051153v
- Marsili, L., Casini, S., Bucalossi, D., Porcelloni, S., Maltese, S., & Fossi, M. (2008). Use of immunofluorescence technique in cultured fibroblasts from Mediterranean cetaceans as new "in vitro" tool to investigate effects of environmental contaminants. *Marine Environmental Research*, 66, 151-153. doi:10.1016/j.marenvres.2008.02.068
- Marsili, L., D'Agostino, A., Bucalossi, D., Malatesta, T., & Fossi, M. (2004). Theoretical models to evaluate hazard due to organochlorine compounds (OCs) in Mediterranean striped dolphin (*Stenella coeruleoalba*). *Chemosphere*, 56, 791-801. doi:10.1016/j.chemosphere.2004.03.014
- Marten, K., & Psarakos, S. (1995). Using self-view television to distinguish between self-examination and social behavior in the bottlenose dolphin (*Tursiops truncatus*). *Consciousness and Cognition: An International Journal*, 4, 205-225. doi:10.1006/ccog.1995.1026
- Marten, K., & Psarakos, S. (1999). Long-term site fidelity and possible long-term associations of wild spinner dolphins (*Stenella longirostris*) seen off Oahu, Hawaii. *Marine Mammal Science*, 15, 1329-1336.
- Marten, K., Herzing, D., Poole, M., & Newman Allman, K. (2001). The acoustic predation hypothesis: Linking underwater observations and recordings during odontocete predation and observing the effects of loud impulsive sounds on fish. *Aquatic Mammals*, 27, 56-66.
- Martin, A., Hall, P., & Richard, P. (2001). Dive behaviour of belugas (*Delphinapterus leucas*) in the shallow waters of Western Hudson Bay. *Arctic*, 54, 276-283.
- Martin, S., Phillips, M., Bauer, E., Moore, P., & Houser, D. (2005). Instrumenting free swimming dolphins echolocating in open water. *Journal of the Acoustical Society of America*, 117, 2301-2307. doi:10.1121/1.1867913
- Martineau, D. (2007). Potential synergism between stress and contaminants in free-ranging cetaceans. *International Journal of Comparative Psychology*, 20, 194-216.
- Martins Silva, J., Jr., Silva, F. J. L., & Sazima, I. (2005). Two presumed interspecific hybrids in the genus *Stenella* (*Delphinidae*) in the tropical West Atlantic. *Aquatic Mammals*, 31, 468-472.
- Mate, B. (1989). Watching whales. *Oregon State Sea Grant*, 53, 2.
- Mate, B., Mesecar, R., & Lagerquist, B. (2007). The evolution of satellite-monitored radio tags for large whales: One laboratory's experience. *Deep-Sea Research Part II, Topical Studies in Oceanography*, 54, 224-247. doi:10.1016/j.dsr2.2006.11.021
- Mate, B., Rossbach, K., Niekirk, S., Wells, R., Irvine, A., Scott, M., & Read, A. (1995). Satellite-monitored movements and dive behavior of a bottlenose dolphin (*Tursiops truncatus*) in Tampa Bay, Florida. *Marine Mammal Science*, 11, 452-463.
- Matkin, C., Barrett-Lennard, L., York, H., Ellifrit, D., & Trites, A. (2007). Ecotypic variation and predatory behavior among killer whales (*Orcinus orca*) off the eastern Aleutian Islands, Alaska. *Fishery Bulletin*, 105, 74-87.

- Matthews, C. (2008). Listening to animals. *Relational Child & Youth Care Practice*, 21, 41-42.
- Matthies, H. (1989). Neurobiological aspects of learning and memory. *Annual Review of Psychology*, 40, 381-404.
- Mattson, M. C., Thomas, J. A., & Aubin, D. S. (2005). Effects of boat activity on the behavior of bottlenose dolphins (*Tursiops truncatus*) in waters surrounding Hilton Head Island, South Carolina. *Aquatic Mammals*, 31, 133-140.
- Maucher, J., Briggs, L., Podmore, C., & Ramsdell, J. (2007). Optimization of blood collection card method/enzyme-linked immunoassay for monitoring exposure of bottlenose dolphin to brevetoxin-producing red tides. *Environmental Science & Technology*, 41, 563-567. doi:10.1021/e0612605
- Maurer, M. (2008). Etude de l'intérêt de l'enfant atteint d'autisme pour le dauphin à partir de deux méthodologies complémentaires. A.N.A.E. *Approche Neuropsychologique des Apprentissages chez l'Enfant*, 20, 143-145.
- May, S. (1998).** *A review of scientific justifications for maintaining cetaceans in captivity. Report for the Whale and Dolphin Conservation Society.* Retrieved from http://www.wdcs.org/submissions_bin/capmayerscjustifications.pdf
- May-Collado, L., Agnarsson, I., & Wartzok, D. (2007). Phylogenetic review of tonal sound production in whales in relation to sociality. *BMC Evolutionary Biology*, 7, 136-156. doi:10.1186/1471-2148-7-136
- May-Collado, L., & Wartzok, D. (2008). A comparison of bottlenose dolphin whistles in the Atlantic Ocean: Factors promoting whistle variation. *Journal of Mammalogy*, 89, 1229-1240.
- Maze, K. S., & Würsig, B. (1999). Bottlenose dolphins of San Luis Pass, Texas: Occurrence patterns, site-fidelity, and habitat use. *Aquatic Mammals*, 25, 91-103.
- Mazzoil, M., McCulloch, S. D., Defran, R. H., & Murdoch, M. E. (2004). Use of digital photography and analysis of dorsal fins for photo-identification of bottlenose dolphins. *Aquatic Mammals*, 30, 209-219.
- Mazzoil, M., McCulloch, S. D., Youngbluth, M. J., Kilpatrick, D. S., Murdoch, M. E., Mase-Guthrie, B., et al. (2008). Radio-tracking and survivorship of two rehabilitated bottlenose dolphins (*Tursiops truncatus*) in the Indian river lagoon, Florida. *Aquatic Mammals*, 34, 54-64.
- Mazzoni, G. (2003). Animals show monitoring, but does monitoring imply awareness?. *Behavioral & Brain Sciences*, 26, 349-350.
- Mazzuca, L., Atkinson, S., Keating, B., & Nitta, E. (1999). Cetacean mass strandings in the Hawaiian Archipelago, 1957-1998. *Aquatic Mammals*, 25, 105-114.
- MacLeod, R. (1994). Terror in the tide. *Natural History*, 103(6), 98.
- MacLeod, R., MacLeod, C. D., Learmonth, J. A., Jepson, P. D., Reid, R. J., Deaville, R., et al. (2007). Mass-dependent predation risk and lethal dolphin-porpoise interactions. *Proceedings of the Royal Society B: Biological Sciences*, 274, 2717-2723.
- McAlpine, D., Kingsley, M., & Daoust, P-Y. (1999). A lactating record-age St. Lawrence beluga (*Delphinapterus leucas*). *Marine Mammal Science*, 15, 854-859.
- McCowan, B. (1995). A new quantitative technique for categorizing whistles using simulated signals and whistles from captive bottlenose dolphins (*Delphinidae, Tursiops truncatus*). *Ethology*, 100, 177-193.
- McCowan, B., & Hanser, S. (1999). Quantitative tools for comparing animal communication systems: Information theory applied to. *Animal Behaviour*, 57, 409.

- McCowan, B., Marino, L., Vance, E., Walke, L., & Reiss, D. (2000). Bubble ring play of bottlenose dolphins (*Tursiops truncatus*): Implications for cognition. *Journal of Comparative Psychology*, 114, 98-106. doi:10.1037/0735-7036.114.1.98
- McCowan, B., & Reiss, D. (1995a). Maternal aggressive contact vocalizations in captive bottlenose dolphins (*Tursiops truncatus*): Wide-band, low-frequency signals during mother/aunt-infant interactions. *Zoo Biology*, 14, 293-309. doi:10.1002/zoo.1430140402
- McCowan, B., & Reiss, D. (1995b). Quantitative comparison of whistle repertoires from captive adult bottlenose dolphins (Delphinidae, *Tursiops truncatus*): A re-evaluation of the signature whistle hypothesis. *Ethology*, 100, 194-209.
- McCowan, B., & Reiss, D. (1995c). Whistle contour development in captive-born infant bottlenose dolphins (*Tursiops truncatus*): Role of learning. *Journal of Comparative Psychology*, 109, 242-260. doi:10.1037/0735-7036.109.3.242
- McCowan, B., & Reiss, D. (1997). Vocal learning in captive bottlenose dolphins: A comparison with humans and nonhuman animals. *Social influences on vocal development* (pp. 178- 207). New York: Cambridge University Press.
- McCowan, B., & Reiss, D. (2001). The fallacy of 'signature whistles' in bottlenose dolphins: A comparative perspective of 'signature information' in animal vocalizations. *Animal Behaviour*, 62, 1151-1162. doi:10.1006/anbe.2001.1846
- McCowan, B., Reiss, D., & Gubbins, C. (1998). Social familiarity influences whistle acoustic structure in adult female bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 24, 27-40.
- McGonigle, M. (2008). Oil and water—18th-Century whale and basking-shark fisheries of Donegal Bay, Ireland. *International Journal of Nautical Archaeology*, 37, 302-312. doi:10.1111/j.1095-9270.2007.00178.x
- McKinney, M., Arukwe, A., De Guise, S., Martineau, D., Béland, P., Dallaire, A., et al. (2004). Characterization and profiling of hepatic cytochromes P450 and phase II xenobiotic-metabolizing enzymes in beluga whales (*Delphinapterus leucas*) from the St. Lawrence River Estuary and the Canadian Arctic. *Aquatic Toxicology*, 69, 35-49.
- McKinney, M., De Guise, S., Martineau, D., Béland, P., Arukwe, A., & Letcher, R. (2006). Biotransformation of polybrominated diphenyl ethers and polychlorinated biphenyls in beluga whale (*Delphinapterus leucas*) and rat mammalian model using an in vitro hepatic microsomal assay. *Aquatic Toxicology*, 77, 87-97. doi:10.1016/j.aquatox.2005.08.016
- McMillan, J. (1992). Tuna, dolphins, and pesticide use: The economics and ecology of GATT. *Journal of Pesticide Reform: A Publication of the Northwest Coalition for Alternatives to Pesticides*, 12(2), 20-21.
- Meagher, E., McLellan, W., Westgate, A., Wells, R., Blum, J., & Pabst, D. (2008). Seasonal patterns of heat loss in wild bottlenose dolphins (*Tursiops truncatus*). *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 178, 529-543. doi:10.1007/s00360-007-0245-5
- Meagher, E., McLellan, W., Westgate, A., Wells, R., Frierson, D., & Pabst, D. (2002). The relationship between heat flow and vacuolation in the dorsal fin of wild bottlenose dolphins *Tursiops truncatus*. *Journal of Experimental Biology*, 205, 3475.
- Melillo, K. E., Dudzinski, K. M., & Cornick, L. A. (2009). Interactions between Atlantic spotted (*Stenella frontalis*) and bottlenose (*Tursiops truncatus*) dolphins off Bimini, The Bahamas, 2003-2007. *Aquatic Mammals*, 35, 281-291.

- Mellink, E., & Orozco-Meyer, A. (2006). Abundance, distribution, and residence of bottlenose dolphins (*Tursiops truncatus*) in the Bahia San Jorge area, northern Gulf of California, Mexico. *Aquatic Mammals*, 32, 133-139.
- Mello, I., & Amundin, M. (2005). Whistle production pre- and post-partum in bottlenose dolphins (*Tursiops truncatus*) in human care. *Aquatic Mammals*, 31, 169-175.
- Mello, I., Nordensten, L., & Amundin, M. (2005). Reactions of three bottlenose dolphin dams with calves to other members of the group in connection with nursing. *Zoo Biology*, 24, 543-555. doi:10.1002/zoo.20071
- Mercado, E., Killebrew, D., Pack, A., Mácha, I., & Herman, L. (2000). Generalization of 'same – different' classification abilities in bottlenosed dolphins. *Behavioural Processes*, 50, 79-94. doi:10.1016/S0376-6357(00)00100-5
- Mercado, E., Murray, S., Uyeyama, R., Pack, A., & Herman, L. (1998). Memory for recent actions in the bottlenosed dolphin (*Tursiops truncatus*): Repetition of arbitrary behaviors using an abstract rule. *Animal Learning & Behavior*, 26, 210-218.
- Mercado, E., Uyeyama, R., Pack, A., & Herman, L. (1999). Memory for action events in the bottlenosed dolphin. *Animal Cognition*, 2, 17-25. doi:10.1007/s100500021
- Meschersky, I., Kholodova, M., & Zvychaynaya, E. (2008). Molecular genetic study of the beluga (*Delphinapterus leucas*: Cetacea, *Monodontidae*) summering in the southern Sea of Okhotsk as compared to North American populations. *Russian Journal of Genetics*, 44, 1105-1110. doi:10.1134/S1022795408090147
- Meyer, W., & Seegers, U. (2004). A preliminary approach to epidermal antimicrobial defense in the *Delphinidae*. *Marine Biology*, 144, 841-844. doi:10.1007/s00227-003-1256-8
- Mignucci-Giannoni, A. A., Montoya-Ospina, R. A., Perez-Zayas, J. J., Rodriguez-Lopez, M. A., & Williams, E. H., Jr. (1999). New records of Fraser's dolphin (*Lagenodelphis hosei*) for the Caribbean. *Aquatic Mammals*, 25, 15-19.
- Mignucci-Giannoni, A. A., Rosario-Delestre, R. J., Alsina-Guerrero, M. M., Falcon-Matos, L., Guzman-Ramirez, L., Williams, E. H., Jr., et al. (2009). Asphyxiation in a bottlenose dolphin (*Tursiops truncatus*) from Puerto Rico due to choking on a black margate (*Anisotremus surinamensis*). *Aquatic Mammals*, 35, 48-54.
- Migura, K. A., & Meadows, D. W. (2002). Short-finned pilot whales (*Globicephala macrorhynchus*) interact with melon-headed whales (*Peponocephala electra*) in Hawaii. *Aquatic Mammals*, 28, 294-297.
- Mihai, M. (1978). Attention and respiratory habituation in captive dolphins. *Revista de Psihologie*, 24, 217-228.
- Mihindukulasuriya, K., Guang, W., St. Leger, J., Nordhausen, R., & Wang, D. (2008). Identification of a novel coronavirus from a Beluga whale by using a panviral microarray. *Journal of Virology*, 82, 5084-5088. doi:10.1128/JVI.02722-07
- Mikaelian, I., Lapointe, J., Labelle, P., Higgins, R., Paradis, M., & Martineau, D. (2001). Dermatophilus-like infection in beluga whales, *Delphinapterus leucas*, from the St. Lawrence estuary. *Veterinary Dermatology*, 12, 59-62.
- Miksis, J., Grund, M., Nowacek, D., Solow, A., Connor, R., & Tyack, P. (2001). Cardiac response to acoustic playback experiments in the captive bottlenose dolphin (*Tursiops truncatus*). *Journal of Comparative Psychology*, 115, 227-232. doi:10.1037/0735-7036.115.3.227
- Miles, J. A., & Herzing, D. L. (2003). Underwater analysis of the behavioural development of free-ranging Atlantic spotted dolphin (*Stenella frontalis*) calves (birth to 4 years of age). *Aquatic Mammals*, 29, 363-377.
- Millera, P., Samarra, F., & Perthuisson, A. (2007). Caller sex and orientation influence spectral characteristics of "two-voice" stereotyped calls produced by free-ranging

- killer whales. *Journal of the Acoustical Society of America*, 121, 3932-3937. doi:10.1121/1.2722056
- Miller, D., Styer, E., Decker, S., & Robeck, T. (2002). Ultrastructure of the spermatozoa from three odontocetes: A killer whale (*Orcinus orca*), a Pacific white-sided dolphin (*Lagenorhynchus albirostris obliquidens*) and a beluga (*Delphinapterus leucas*). *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 31, 158-168. doi:10.1046/j.1439-0264.2002.00385.x
- Miller, P. (2002). Mixed-directionality of killer whale stereotyped calls: A direction of movement cue? *Behavioral Ecology and Sociobiology*, 52, 262-270. doi:10.1007/s00265-002-0508-9
- Miller, P. (2006). Diversity in sound pressure levels and estimated active space of resident killer whale vocalizations. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural & Behavioral Physiology*, 192, 449-459. doi:10.1007/s00359-005-0085-2
- Miller, P., Aoki, K., Rendell, L., & Amano, M. (2008). Stereotypical resting behavior of the sperm whale. *Current Biology*, 18, R21-R23. doi:10.1016/j.cub.2007.11.003
- Miller, P., & Bain, D. (2000). Within-pod variation in the sound production of a pod of killer whales, *Orcinus orca*. *Animal Behaviour*, 60, 617-628. doi:10.1006/anbe.2000.1503
- Miller, P., & Tyack, P. (1998). A small towed beamforming array to identify vocalizing resident killer whales (*Orcinus orca*). *Deep-Sea Research Part II, Topical Studies in Oceanography*, 45, 1389.
- Miller, P., Shapiro, A., Tyack, P., & Solow, A. (2004). Call-type matching in vocal exchanges of free-ranging resident killer whales, *Orcinus orca*. *Animal Behaviour*, 67, 1099-1107. doi:10.1016/j.anbehav.2003.06.017
- Mintzer, V. J., Gannon, D. P., Barros, N. B., & Read, A. J. (2008). Stomach contents of mass-stranded short-finned pilot whales (*Globicephala macrorhynchus*) from North Carolina. *Marine Mammal Science*, 24, 290-302. doi:10.1111/j.1748-7692.2008.00189.x
- Mirimin, L., Coughlan, J., Rogan, E., & Cross, T. (2006). Tetranucleotide microsatellite loci from the striped dolphin (*Stenella coeruleoalba* Meyen, 1833). *Molecular Ecology Notes*, 6, 493-495. doi:10.1111/j.1471-8286.2006.01285.x
- Mitchell, M., Solangi, M., Clemons-Chevis, C., Vanderpool, D., Romagnoli, M., Hoffland, T., et al. (2008). Relationship between plasma iron concentration and gastric pH in captive adult bottlenose dolphins (*Tursiops truncatus*). *American Journal of Veterinary Research*, 69, 900-903.
- Mitchell, R. (1995). Evidence of dolphin self-recognition and the difficulties of interpretation. *Consciousness and Cognition: An International Journal*, 4, 229-234. doi:10.1006/ccog.1995.1029
- Mitchell, R. (2001). On not drawing the line about culture: Inconsistencies in interpretation of nonhuman cultures. *Behavioral & Brain Sciences*, 24, 348.
- Mitchell, R., Yao, P., Sherman, P., & O'Regan, M. (1985). Discriminative responding of a dolphin (*Tursiops truncatus*) to differentially rewarded stimuli. *Journal of Comparative Psychology*, 99, 218-225. doi:10.1037/0735-7036.99.2.218
- Mizroch, S., & Hutchinson, C. (1997). An observed entanglement of *Lagenorhynchus albirostris obliquidens* in the high seas driftnet area in the North Pacific. *Marine Fisheries Review*, 59(3), 41-44.
- Moeller, L., & Beheregaray, L. (2001). Coastal bottlenose dolphins from southeastern Australia are *Tursiops aduncus* according to sequences of the mitochondrial DNA control region. *Marine Mammal Science*, 17, 249-263.

- Møhl, B., Au, W., Pawloski, J., & Nachtigall, P. (1999). Dolphin hearing: Relative sensitivity as a function of point of application of a contact sound source in the jaw and head region. *Journal of the Acoustical Society of America*, 105, 3421-3424. doi:10.1121/1.426959
- Mollenhauer, M., Carter, B., Peden-Adams, M., Bossart, G., & Fair, P. (2009). Gene expression changes in bottlenose dolphin, *Tursiops truncatus*, skin cells following exposure to methylmercury (MeHg) or perfluorooctane sulfonate (PFOS). *Aquatic Toxicology*, 91, 10-18. doi:10.1016/j.aquatox.2008.09.013
- Möller, L., & Beheregaray, L. (2004). Genetic evidence for sex-biased dispersal in resident bottlenose dolphins (*Tursiops aduncus*). *Molecular Ecology*, 13, 1607-1612.
- Möller, L., Beheregaray, L., Allen, S., & Harcourt, R. (2006). Association patterns and kinship in female Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) of southeastern Australia. *Behavioral Ecology and Sociobiology*, 61, 109-117.
- Möller, L., Bilgmann, K., Charlton-Robb, K., & Beheregaray, L. (2008). Multi-gene evidence for a new bottlenose dolphin species in southern Australia. *Molecular Phylogenetics & Evolution*, 49, 674-681. doi:10.1016/j.ympev.2008.08.011
- Mogensen, L., Kinze, C., Werge, T., & Rasmussen, H. (2006). Identification and characterization of a tandem repeat in exon iii of the dopamine receptor D4 (DRD4) gene in cetaceans. *Journal of Heredity*, 97, 279-284. doi:10.1093/jhered/esj024
- Monteiro-Neto, C., Alves-Junior, T. T., Avila, F. J. C., Campos, A. A., Costa, A. F., Silva, C. P. N., et al. (2000). Impact of fisheries on the tucuxi (*Sotalia fluviatilis*) and rough-toothed dolphin (*Steno bredanensis*) populations off Ceara state, northeastern Brazil. *Aquatic Mammals*, 26, 49-56.
- Montie, E., Fair, P., Bossart, G., Mitchum, G., Houde, M., Muir, D., et al. (2008). Cytochrome P4501A1 expression, polychlorinated biphenyls and hydroxylated metabolites, and adipocyte size of bottlenose dolphins from the Southeast United States. *Aquatic Toxicology*, 86, 397-412. doi:10.1016/j.aquatox.2007.12.004
- Montie, E., Hahn, M., Letcher, R., Touhey, K., Reddy, C., & Gebbink, W. (2009). Organohalogen contaminants and metabolites in cerebrospinal fluid and cerebellum gray matter in short-beaked common dolphins and Atlantic white-sided dolphins from the western North Atlantic. *Environmental Pollution*, 157, 2345-2358.
- Mooney, A. (2009). Animal behaviour: Deafening dolphins. *Nature*, 458, 810. doi:10.1038/458810a
- Mooney, T., Nachtigall, P., & Au, W. W. L. (2004). Target strength of a nylon monofilament and an acoustically enhanced gillnet: Predictions of biosonar detection ranges. *Aquatic Mammals*, 30, 220-226.
- Mooney, T., Nachtigall, P., Breese, M., Vlachos, S., & Au, W. (2009). Predicting temporary threshold shifts in a bottlenose dolphin (*Tursiops truncatus*): The effects of noise level and duration. *Journal of the Acoustical Society of America*, 125, 1816-1826. doi:10.1121/1.3068456
- Mooney, T., Nachtigall, P., Castellote, M., Taylor, K., Pacini, A., & Esteban, J. (2008). Hearing pathways and directional sensitivity of the beluga whale, *Delphinapterus leucas*. *Journal of Experimental Marine Biology & Ecology*, 362, 108-116. doi:10.1016/j.jembe.2008.06.004
- Mooney, T., Nachtigall, P., & Yuen, M. (2006). Temporal resolution of the Risso's dolphin, *Grampus griseus*, auditory system. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural & Behavioral Physiology*, 192, 373-380. doi:10.1007/s00359-005-0075-4

- Moore, M., Hayward, B., Lentell, B., Rotstein, D., Garner, M., Bogomolni, A., et al. (2009). Gas bubbles in seals, dolphins, and porpoises entangled and drowned at depth in gillnets. *Veterinary Pathology*, 46, 536-547.
- Moore, M., Weisbrod, A., Shea, D., & Stegeman, J. (2000). Bioaccumulation patterns of polychlorinated biphenyls and chlorinated pesticides in northwest Atlantic pilot whales. *Environmental Toxicology & Chemistry*, 19, 667-677.
- Moore, P. (1997). Cetacean auditory psychophysics. *Bioacoustics*, 8, 61-78.
- Moore, P., Dankiewicz, L., & Houser, D. (2008). Beamwidth control and angular target detection in an echolocating bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 124, 3324-3332. doi:10.1121/1.2980453
- Moore, S. (2000). Variability of cetacean distribution and habitat selection in the Alaskan Arctic, Autumn 1982-91. *Arctic*, 53, 448-460.
- Moore, S., Shelden, K., Litzky, L., Mahoney, B., & Rugh, D. (2000). Beluga habitat associations in Cook Inlet, Alaska. *Marine Fisheries Review*, 62(3), 60-80.
- Moore, S., & Ridgway, S. H. (1995). Whistles produced by common dolphins from the Southern California Bight. *Aquatic Mammals*, 21, 55-63.
- Morell, M., Degollada, E., Alonso, J., Jauniaux, T., & André, M. (2009). Decalcifying odontocete ears following a routine protocol with RDO®. *Journal of Experimental Marine Biology & Ecology*, 376, 55-58. doi:10.1016/j.jembe.2009.05.005
- Mori, C., Morsey, B., Levin, M., Nambiar, P., & De Guise, S. (2006). Immunomodulatory effects of in vitro exposure to organochlorines on t-cell proliferation in marine mammals and mice. *Journal of Toxicology & Environmental Health: Part A*, 69, 283-302. doi:10.1080/15287390500227472
- Morin, P. A., LeDuc, R. G., Robertson, K. M., Hedrick, N. M., Perrin, W. F., Etnier, M., et al. (2006). Genetic analysis of killer whale (*Orcinus orca*) historical bone and tooth samples to identify western U.S. ecotypes. *Marine Mammal Science*, 22, 897-909. doi:10.1111/j.1748-7692.2006.00070.x
- Morisaka, T. (2009). Overview of comparative cognitive studies of dolphins in Japan. *Japanese Psychological Research*, 51, 168-176. doi:10.1111/j.1468-5884.2009.00395.x
- Morisaka, T., & Connor, R. (2007). Predation by killer whales (*Orcinus orca*) and the evolution of whistle loss and narrow-band high frequency clicks in odontocetes. *Journal of Evolutionary Biology*, 20, 1439-1458. doi:10.1111/j.1420-9101.2007.01336.x
- Morisaka, T., Shinohara, M., Nakahara, F., & Akamatsu, T. (2005a). Effects of ambient noise on the whistles of Indo-Pacific bottlenose dolphin populations. *Journal of Mammalogy*, 86, 541-546.
- Morisaka, T., Shinohara, M., Nakahara, F., & Akamatsu, T. (2005b). Geographic variations in the whistles among three Indo-Pacific bottlenose dolphin *Tursiops aduncus* populations in Japan. *Fisheries Science*, 71, 568-576. doi:10.1111/j.1444-2906.2005.01001.x
- Morisaka, T., Shinohara, M., & Taki, M. (2005). Underwater sounds produced by neonatal bottlenose dolphins (*Tursiops truncatus*): II. Potential function. *Aquatic Mammals*, 31, 258-265.
- Mortola, J., & Seguin, J. (2009). End-tidal CO₂ in some aquatic mammals of large size. *Zoology*, 112, 77-85. doi:10.1016/j.zool.2008.06.001
- Morton, A. (2000). Occurrence, photo-identification and prey of Pacific white-sided dolphins (*Lagenorhynchus obliquidens*) in the Broughton archipelago, Canada 1984-1998. *Marine Mammal Science*, 16, 80-93.

- Morton, A., & Symonds, H. (2002). Displacement of *Orcinus orca* by high amplitude sound in British Columbia, Canada. *ICES Journal of Marine Science / Journal du Conseil*, 59, 71. doi:10.1006/jmsc.2001.1136.
- Mossbridge, J., & Thomas, J. (1999). An "acoustic niche" for Antarctic killer whale and leopard seal sounds. *Marine Mammal Science*, 15, 1351-1357.
- Muir, D., Wagemann, R., Grift, N., Norstrom, R., Simon, M., & Lien, J. (1988). Organochlorine chemical and heavy metal contaminants in white-beaked dolphins (*Lagenorhynchus albirostris albirostris*) and pilot whales (*Globicephala melaena*) from the coast of Newfoundland, Canada. *Archives of Environmental Contamination and Toxicology*, 17, 613-629.
- Mukhametov, L., Supin, A., & Polyakova, I. (1977). Interhemispheric asymmetry of the electroencephalographic sleep patterns in dolphins. *Brain Research*, 134, 581-584. doi:10.1016/0006-8993(77)90835-6
- Muller, M., Allen, J., Au, W., & Nachtigall, P. (2008). Time-frequency analysis and modeling of the backscatter of categorized dolphin echolocation clicks for target discrimination. *Journal of the Acoustical Society of America*, 124, 657-666. doi:10.1121/1.2932060
- Muller, M., Au, W., Nachtigall, P., Allen III, J., & Breese, M. (2007). Phantom echo highlight amplitude and temporal difference resolutions of an echolocating dolphin, *Tursiops truncatus*. *Journal of the Acoustical Society of America*, 122, 2255-2262. doi:10.1121/1.2769973
- Muller, M., Battersby, M. Z., Buurman, D., Bossley, M., & Doak, W. (1998). Range and sociability of a solitary bottlenose dolphin *Tursiops truncatus* in New Zealand. *Aquatic Mammals*, 24, 93-104.
- Muller, M., & Bossley, M. (2002). Solitary bottlenose dolphins in comparative perspective. *Aquatic Mammals*, 28, 298-307.
- Munguia-Vega, A., Esquer-Garrigos, Y., Rojas-Bracho, L., Vazquez-Juarez, R., Castro-Prieto, A., & Flores-Ramirez, S. (2007). Genetic drift vs. natural selection in a long-term small isolated population: Major histocompatibility complex class II variation in the Gulf of California endemic porpoise (*Phocoena sinus*). *Molecular Ecology*, 16, 4051-4065. doi:10.1111/j.1365-294X.2007.03319.x.
- Muñoz, P., García-Castrillo, G., López-García, P., González-Cueli, J., De Miguel, M., Marín, C., et al. (2006). Isolation of Brucella species from a live-stranded striped dolphin (*Stenella coeruleoalba*) in Spain. *Veterinary Record: Journal of the British Veterinary Association*, 158, 450-451.
- Muraco, H., Cheng, A., Ravid, N., Arn, D., Hudson, J., & Durrant, B. (2009). Conditioning for conservation: A new approach to detection of luteinizing hormone in a bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 35, 386-393. doi:10.1578/AM.35.3.2009.386
- Murayama, T., Aoki, I., & Ishii, T. (1993). Measurement of the electroencephalogram of the bottlenose dolphin under different light conditions. *Aquatic Mammals*, 19, 171-182.
- Murayama, T., & Tobayama, T. (1997). Preliminary study on stimulus equivalence in beluga (*Delphinapterus leucas*). *Japanese Journal of Animal Psychology*, 47, 79-89.
- Murphy, E. J. (1995). Spatial structure and the Southern ocean ecosystem: Predator-prey linkages in Southern Ocean food webs. *Journal of Animal Ecology*, 64, 333-347.
- Murphy, S., Collet, A., & Rogan, E. (2005). Mating strategy in the male common dolphin (*Delphinus delphis*): What gonadal analysis tells us. *Journal of Mammalogy*, 86, 1247-1258.

- Murphy, S., & Rogan, E. (2006). External morphology of the short-beaked common dolphin, *Delphinus delphis*: Growth, allometric relationships and sexual dimorphism. *Acta Zoologica*, 87, 315-329. doi:10.1111/j.1463-6395.2006.00245.x
- Murray, S. (2006). What's up, doc? *CMAJ: Canadian Medical Association Journal*, 174, 301. doi:10.1503/cmaj.051673
- Murray, S., Mercado, E., & Roitblat, H. (1998). The neural network classification of false killer whale (*Pseudorca crassidens*) vocalizations. *Journal of the Acoustical Society of America*, 104, 3626-3633. doi:10.1121/1.423945
- Nachtigall, P., Au, W. W. L., Pawloski, J. L., Andrews, K., & Oliver, C. W. (2000). Measurements of the low frequency components of active and passive sounds produced by dolphins. *Aquatic Mammals*, 26, 167-174.
- Nachtigall, P., Mooney, T., Taylor, K., Miller, L., Rasmussen, M., Akamatsu, T., et al. (2008). Shipboard measurements of the hearing of the white-beaked dolphin *Lagenorhynchus albirostris albirostris*. *Journal of Experimental Biology*, 211, 642-647. doi:10.1242/jeb.014118
- Nachtigall, P., Pawloski, J., & Au, W. (2003). Temporary threshold shifts and recovery following noise exposure in the Atlantic bottlenosed dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 113, 3425-3437. doi:10.1121/1.1570438
- Nachtigall, P., & Supin, A. (2008). A false killer whale adjusts its hearing when it echolocates. *Journal of Experimental Biology*, 211, 1714-1718. doi:10.1242/jeb.013862
- Nachtigall, P., Supin, A. Y., Pawloski, J., & Au, W. W. (2004). Temporary threshold shifts after noise exposure in the bottlenose dolphin (*Tursiops truncatus*) measured using evoked auditory potentials. *Marine Mammal Science*, 20, 673-687.
- Nachtigall, P., Yuen, M., Mooney, T., & Taylor, K. (2005). Auditory evoked potential measurement methodology for odontocetes and a comparison of measured thresholds with those obtained using psychophysical techniques. *Journal of the Acoustical Society of America*, 117(4), 2407.
- Nachtigall, P., Yuen, M., Mooney, T., & Taylor, K. (2005c). Hearing measurements from a stranded infant Risso's dolphin, *Grampus griseus*. *Journal of Experimental Biology*, 208, 4181-4188. doi:10.1242/jeb.01876
- Nagasaki, F. (1990). The case for scientific whaling. *Nature*, 344, 189.
- Nakahara, F., & Takemura, A. (1997). A survey on the behavior of captive odontocetes in Japan. *Aquatic Mammals*, 23, 135-143.
- Nater, A., Kopps, A., & Krutzen, M. (2009). New polymorphic tetranucleotide microsatellites improve scoring accuracy in the bottlenose dolphin *Tursiops aduncus*. *Molecular Ecology Resources*, 9, 531-534. doi:10.1111/j.1755-0998.2008.02246.x
- Nater, A., Krcotzen, M., & Kopps, A. (2009). New polymorphic tetranucleotide microsatellites improve scoring accuracy in the bottlenose dolphin *Tursiops aduncus*. *Molecular Ecology Resources*, 9, 531-534.
- Nathanson, D. (1998). Long-term effectiveness of dolphin-assisted therapy for children with severe disabilities. *Anthrozoös*, 11, 22-32.
- Nathanson, D. (2007). Reinforcement effectiveness of animatronic and real dolphins. *Anthrozoös*, 20, 181-194. doi:10.2752/175303707X207963
- Natoli, A., Birkun, A., Aguilar, A., Lopez, A., & Hoelzel, A. R. (2005). Habitat structure and the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*). *Proceedings: Biological Sciences*, 272, 1217-1226.

- Natoli, A., Peddemors, V., & Hoelzel, A. (2004). Population structure and speciation in the genus *Tursiops* based on microsatellite and mitochondrial DNA analyses. *Journal of Evolutionary Biology*, 17, 363-375. doi:10.1046/j.1420-9101.2003.00672.x
- Natoli, A., Peddemors, V., & Hoelzel, A. (2008). Population structure of bottlenose dolphins (*Tursiops aduncus*) impacted by bycatch along the east coast of South Africa. *Conservation Genetics*, 9, 627-636.
- Neil, D. (2002). Cooperative fishing interactions between Aboriginal Australians and dolphins in eastern Australia. *Anthrozoös*, 15, 3-18. doi:10.2752/089279302786992694
- Neil, D., & Holmes, B. (2008). Survival of bottlenose dolphin (*Tursiops* sp.) calves at a wild dolphin provisioning program, Tangalooma, Australia. *Anthrozoös*, 21, 57-69. doi:10.2752/089279308X274065
- Nelson, D. L., & Lien, J. (1994). Behaviour patterns of two captive Atlantic white-sided dolphins, *Lagenorhynchus albirostris acutus*. *Aquatic Mammals*, 20, 1-10.
- Neuenhagen, C., García Hartmann, M., & Greven, H. (2007). Histology and morphometrics of testes of the white-sided dolphin (*Lagenorhynchus albirostris acutus*) in bycatch samples from the northeastern Atlantic. *Mammalian Biology*, 72, 283-298. doi:10.1016/j.mambio.2006.10.008
- Neumann, D. R. (2001). The activity budget of free-ranging common dolphins (*Delphinus delphis*) in the northwestern Bay of Plenty, New Zealand. *Aquatic Mammals*, 27, 121-136.
- Neumann, D. R., & Orams, M. B. (2006). Impacts of ecotourism on short-beaked common dolphins (*Delphinus delphis*) in Mercury Bay, New Zealand. *Aquatic Mammals*, 32, 1-9.
- Neumann, D. R., Russell, K., Orams, M. B., Baker, C. S., & Duignan, P. (2002). Identifying sexually mature, male short-beaked common dolphins (*Delphinus delphis*) at sea, based on the presence of a postanal hump. *Aquatic Mammals*, 28, 181-187.
- Newman, E. B. (1950). Hearing. *Annual Review of Psychology*, 1, 49-70.
- Ng, S. L., & Leung, S. (2003). Behavioral response of Indo-Pacific humpback dolphin (*Sousa chinensis*) to vessel traffic. *Marine Environmental Research*, 56, 555-567. doi:10.1016/S0141-1136(03)00041-2
- Nichols, C., Herman, J., Gaggiotti, O. E., Dobney, K. M., Parsons, K., & Hoelzel, A. R. (2007). Genetic isolation of a now extinct population of bottlenose dolphins (*Tursiops truncatus*). *Proceedings of the Royal Society B: Biological Sciences*, 274, 1611-1616.
- Ninomiya, H., & Yoshida, E. (2007). Functional anatomy of the ocular circulatory system: Vascular corrosion casts of the cetacean eye. *Veterinary Ophthalmology*, 10, 231-238.
- Niño-Torres, C., Gallo-Reynoso, J., Galván-Magaña, F., Escobar-Briones, E., & Macko, S. (2006). Isotopic analysis of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$ "a feeding tale" in teeth of the longbeaked common dolphin, *Delphinus capensis*. *Marine Mammal Science*, 22, 831-846.
- Nobbe, G. (1998). Protective sponges. *Wildlife Conservation*, 101(1), 13.
- Noda, K., Aoki, M., Akiyoshi, H., Asaki, H., Ogata, T., Yamauchi, K., et al. (2006). Effect of bovine lactoferrin on the immune responses of captive bottlenosed dolphins (*Tursiops truncatus*) being transported over long distances. *Veterinary Record: Journal of the British Veterinary Association*, 159, 885-888.
- Noël, M., Barrett-Lennard, L., Guinet, C., Dangerfield, N., & Ross, P. (2009). Persistent organic pollutants (POPs) in killer whales (*Orcinus orca*) from the Crozet

- Archipelago, southern Indian Ocean. *Marine Environmental Research*, 68, 196-202. doi:10.1016/j.marenvres.2009.06.009
- Noettestad, L., & Similae, T. (2001). Killer whales attacking schooling fish: Why force herring from deep water to the surface? *Marine Mammal Science*, 17, 343-352.
- Noke, W. D. (2004). The association of echeneids with bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida, USA. *Aquatic Mammals*, 30, 296-298.
- Nollens, H., Green, L., Duke, D., Walsh, M., Chittick, B., Gearhart, S., et al. (2007). Development and validation of monoclonal and polyclonal antibodies for the detection of immunoglobulin G of bottlenose dolphins (*Tursiops truncatus*). *Journal of Veterinary Diagnostic Investigation*, 19, 465-470.
- Nollens, H., Wellehan, J., Saliki, J., Caseltine, S., Jensen, E., Van Bonn, W., et al. (2008). Characterization of a parainfluenza virus isolated from a bottlenose dolphin (*Tursiops truncatus*). *Veterinary Microbiology*, 128, 231-242. doi:10.1016/j.vetmic.2007.10.005
- Nollman, J. (1987). To judge the pain of whales. *Between the Species: A Journal of Ethics*, 3(3), 133-137.
- Nomizo, A., Kudoh, H., & Sakai, T. (2005). Iliocostalis muscles in three mammals (dolphin, goat and human): Their identification, structure and innervation. *Anatomical Science International*, 80, 212-222. doi:10.1111/j.1447-073X.2005.00115.x
- Noonan, M., & Suchak, M. (2005). Long term stability and individual distinctiveness in captive orca vocalizations. *Journal of the Acoustical Society of America*, 117, 2469.
- Nordquist, C., & Hutchins, M. (1985). Killer whale shares food with gulls at the Vancouver Public Aquarium. *Zoo Biology*, 4, 367-374. doi:10.1002/zoo.1430040407
- Noren, S. (2008). Infant carrying behaviour in dolphins: costly parental care in an aquatic environment. *Functional Ecology*, 22, 284-288. doi:10.1111/j.1365-2435.2007.01354.x
- Noren, S., Biedenbach, G., & Edwards, E. (2006). Ontogeny of swim performance and mechanics in bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Biology*, 209, 4724-4731. doi:10.1242/jeb.02566
- Noren, S., Biedenbach, G., Redfern, J., & Edwards, E. (2008). Hitching a ride: The formation locomotion strategy of dolphin calves. *Functional Ecology*, 22, 278-283. doi:10.1111/j.1365-2435.2007.01353.x
- Noren, S., Cuccurullo, V., & Williams, T. (2004). The development of diving bradycardia in bottlenose dolphins (*Tursiops truncatus*). *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 174, 139-147. doi:10.1007/s00360-003-0398-9
- Noren, S., & Edwards, E. F. (2007). Physiological and behavioral development in delphinid calves: Implications for calf separation and mortality due to tuna purse-seine sets. *Marine Mammal Science*, 23, 15-29. doi:10.1111/j.1748-7692.2006.00083.x
- Noren, S., & Wells, R. (2009). Blubber deposition during ontogeny in free-ranging bottlenose dolphins: Balancing disparate roles of insulation and locomotion. *Journal of Mammalogy*, 90, 629-637.
- Norris, S. (2002). Creatures of culture? Making the case for cultural systems in whales and dolphins. *Bioscience*, 52, 9-14.

- Nowacek, D. P. (2002). Sequential foraging behaviour of bottlenose dolphins, *Tursiops truncatus* in Sarasota Bay, FL. *Behaviour*, 139, 1125-1145. doi:10.1163/15685390260437290
- Nowacek, D. P. (2005). Acoustic ecology of foraging bottlenose dolphins (*Tursiops truncatus*), habitat-specific use of three sound types. *Marine Mammal Science*, 21, 587-602.
- Nowacek, D. P., Thorne, L. H., Johnston, D. W., & Tyack, P. L. (2007). Responses of cetaceans to anthropogenic noise. *Mammal Review*, 37, 81-115.**
- Nowacek, S., Wells, R., & Solow, A. (2001). Short-term effects of boat traffic on bottlenose dolphins, *Tursiops truncatus*, in Sarasota Bay, Florida. *Marine Mammal Science*, 17, 673-688.
- Nummela, S., Kosove, J. E., Lancaster, T. E., & Thewissen, J. (2004). Lateral mandibular wall thickness in *tursiops truncatus*: Variation due to sex and age. *Marine Mammal Science*, 20, 491-497.
- O'Brien, J., Steinman, K., & Robeck, T. (2009). Application of sperm sorting and associated reproductive technology for wildlife management and conservation. *Theriogenology*, 71, 98-107. doi:10.1016/j.theriogenology.2008.09.052
- O'Corry-Crowe, G., Suydam, R., Rosenberg, A., Frost, K., & Dizon, A. (1997). Phylogeography, population structure and dispersal patterns of the beluga whale *Delphinapterus leucas* in the western Nearctic revealed by mitochondrial DNA. *Molecular Ecology*, 6, 955-970.
- Oelschläger, H. H. A. (2008). The dolphin brain—A challenge for synthetic neurobiology. *Brain Research Bulletin*, 75, 450-459.
- Oelschläger, H., Haas-Rioth, M., Fung, C., Ridgway, S., & Knauth, M. (2008). Morphology and evolutionary biology of the dolphin (*Delphinus* sp.) brain – MR Imaging and conventional histology. *Brain, Behavior & Evolution*, 71, 68-86. doi:10.1159/000110495
- Ohsumi, S., & Kato, H. (2008). A bottlenose dolphin (*Tursiops truncatus*) with fin-shaped hind appendages. *Marine Mammal Science*, 24, 743-745. doi:10.1111/j.1748-7692.2008.00202.x
- Ognetov, G., & Potelov, V. (1978). Dynamics of visits of white whales to the White Sea. *Soviet Journal of Ecology*, 9(4), 362.
- Olds, M. E. & Fobes, J.L. (1981). The central basis of motivation: Intracranial self-stimulation studies. *Annual Review of Psychology*, 32, 523-574.
- Orams, M. (1994). Tourism and marine wildlife: The wild dolphins of Tangalooma, Australia: A case report. *Anthrozoös*, 7, 195-201.
- Orams, M., & Hill, G. (1998). Controlling the ecotourist in a wild dolphin feeding program: Is education the answer? *Journal of Environmental Education*, 29, 33-38.
- Orams, M., Hill, G., & Baglioni, A.,Jr. (1996). "Pushy" behavior in a wild dolphin feeding program at Tangalooma, Australia. *Marine Mammal Science*, 12, 107-117.
- Orams, M. B. (1995). Development and management of a feeding program for wild bottlenose dolphins at Tangalooma, Australia. *Aquatic Mammals*, 21, 137-147.
- Orós, J., González-Díaz, O., & Monagas, P. (2009). High levels of polychlorinated biphenyls in tissues of Atlantic turtles stranded in the Canary Islands, Spain. *Chemosphere*, 74, 473-478. doi:10.1016/j.chemosphere.2008.08.048
- Orr, J., Aubin, D., Richard, P., & Heide-Joergensen, M. (1998). Recapture of belugas, *Delphinapterus leucas*, tagged in the Canadian arctic. *Marine Mammal Science*, 14, 829-834.

- Orr, J., Martin, A., & Richard, P. (2001). Summer and autumn movements of belugas of the Eastern Beaufort Sea stock. *Arctic*, 54, 223-236.
- Ortiz, G., Feria-Velasco, A., Pacheco-Moisés, F., Rodríguez-Reinoso, S., Cruz-Ramos, J., Rosales-Corral, S., et al. (2009). Scanning electron microscopy of the orbital Harderian gland in the male Atlantic bottlenose dolphin (*Tursiops truncatus*). *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 38, 279-281. doi:10.1111/j.1439-0264.2009.00937.x
- Ortiz, G., Feria-Velasco, A., Tarpley, R., Bitzer-Quintero, O., Rosales-Corral, S., Velázquez Brizuela, I., et al. (2007). The orbital Harderian gland of the male Atlantic bottlenose dolphin (*Tursiops truncatus*): A morphological study. *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 36, 209-214. doi:10.1111/j.1439-0264.2007.00757.x
- Oswald, J., Barlow, J., & Norris, T. (2003). Acoustic identification of nine delphinid species in the Eastern tropical Pacific Ocean. *Marine Mammal Science*, 19, 20-37.
- Oswald, J., Rankin, S., & Barlow, J. (2007). First description of whistles of Pacific Fraser's dolphins *lagenodelphis hosei*. *Bioacoustics*, 16, 99-111.
- Oswald, J., Rankin, S., Barlow, J., & Lammers, M. (2007). A tool for real-time acoustic species identification of delphinid whistles. *Journal of the Acoustical Society of America*, 122, 587-595. doi:10.1121/1.2743157
- Oswald, J., Rankin, S., & Barlow, J. (2008). To whistle or not to whistle? Geographic variation in the whistling behavior of small odontocetes. *Aquatic Mammals*, 34, 288-302.
- Ott, P. H., & Danilewicz, D. (1996). Southward range extension of *Steno bredanensis* in the southwest Atlantic and new records of *Stenella coeruleoalba* for Brazilian waters. *Aquatic Mammals*, 22, 185-189.
- Outridge, P., Hobson, K., & Savelle, J. (2009). Long-term changes of mercury levels in ringed seal (*Phoca hispida*) from Amundsen Gulf, and beluga (*Delphinapterus leucas*) from the Beaufort Sea, western Canadian Arctic. *Science of the Total Environment*, 407, 6044-6051. doi:10.1016/j.scitotenv.2009.08.018
- Overstrom, N. (1983). Association between burst-pulse sounds and aggressive behavior in captive Atlantic bottlenosed dolphins (*Tursiops truncatus*). *Zoo Biology*, 2, 93-103. doi:10.1002/zoo.1430020203
- Oviedo, L., & Silva, N. (2005). Sighting frequency and relative abundance of bottlenose dolphins (*Tursiops truncatus*) along the northeast coast of Margarita Island and Los Frailes Archipelago, Venezuela. *Revista de Biología Tropical*, 53, 595-600.
- Oviedo, L., Silva, N., Bermudez, L., & Odell, D. K. (2005). Distribution of bottlenose dolphins (*Tursiops truncatus*) on the east coast of Isla Margarita and the Los Frailes Archipelago, Venezuela. *Aquatic Mammals*, 31, 442-446.
- Owen, E. C. G., Wells, R. S., & Hofmann, S. (2002). Ranging and association patterns of paired and unpaired adult male Atlantic bottlenose dolphins, *Tursiops truncatus*, in Sarasota, Florida, provide no evidence for alternative male strategies. *Canadian Journal of Zoology*, 80, 2072-2089.
- Oxley, A., Argo, J., & McKay, D. (2005). *Helicobacter* spp. from captive bottlenose dolphins (*Tursiops* spp.) and polar bears (*Ursus maritimus*). *Veterinary Journal*, 170, 377-380.
- Pabst, D., & Rommel, S. (1995). Thermoregulation of the intra-abdominal testes of the bottlenose dolphin (*Tursiops truncatus*). *Journal of Experimental Biology*, 198, 221.
- Pace, D. S. (2000). Fluke-made bubble rings as toys in bottlenose dolphin calves (*Tursiops truncatus*). *Aquatic Mammals*, 26, 57-64.

- Pack, A., & Herman, L. (1995). Sensory integration in the bottlenosed dolphin: Immediate recognition of complex shapes across the senses of echolocation and vision. *Journal of the Acoustical Society of America*, 98, 722-733. doi:10.1121/1.413566
- Pack, A., & Herman, L. (2004). Bottlenosed dolphins (*Tursiops truncatus*) comprehend the referent of both static and dynamic human gazing and pointing in an object-choice task. *Journal of Comparative Psychology*, 118, 160-171. doi:10.1037/0375-7036.1118.2.160
- Pack, A., & Herman, L. (2007). The dolphin's (*Tursiops truncatus*) understanding of human gazing and pointing: Knowing what and where. *Journal of Comparative Psychology*, 121, 34-45. doi:10.1037/0735-7036.121.1.34
- Pack, A., Herman, L., Hoffmann-Kuhnt, M., & Branstetter, B. (2002). The object behind the echo: Dolphins (*Tursiops truncatus*) perceive object shape globally through echolocation. *Behavioural Processes*, 58, 1-26. doi:10.1016/S0376-6357(01)00200-5
- Pack, A., & Herman, L. (2006). Dolphin social cognition and joint attention: Our current understanding. *Aquatic Mammals*, 32, 443-460.
- Paine, R. T. (1980). Food webs: linkage, interaction strength and community infrastructure. *Journal of Animal Ecology*, 49, 667-685.
- Paine, R. T., Ruesink, J. L., Sun, A., Soulanille, E. L., Wonham, M. J., Harley, C. D. G., et al. (1996) Trouble on oiled waters: Lessons from the Exxon Valdez oil spill. *Annual Review of Ecology and Systematics*, 27, 197-235.
- Palacios, D., & Mate, B. (1996). Attack by false killer whales (*Pseudorca crassidens*) on sperm whales (*Physeter macrocephalus*) in the Galapagos Islands. *Marine Mammal Science*, 12, 582-587.
- Pamboris, X. (2004). Lonely Luna's fate. *Wildlife Conservation*, 107(4), 18.
- Pangallo, K. (2008). How does nature deal with persistent pollutants? *Oceanus*, 47(1), 26-27.
- Pangallo, K., & Reddy, C. (2009). Distribution patterns suggest biomagnification of halogenated 1'-Methyl-1-2'-Bipyrroles (MBPs). *Environmental Science & Technology*, 43, 122-127.
- Panigada, S., Zanardelli, M., MacKenzie, M., Donovan, C., Mélin, F., & Hammond, P. (2008). Modelling habitat preferences for fin whales and striped dolphins in the Pelagos Sanctuary (western Mediterranean Sea) with physiographic and remote sensing variables. *Remote Sensing of Environment*, 112, 3400-3412. doi:10.1016/j.rse.2007.11.017
- Parra, G. (2006). Resource partitioning in sympatric delphinids: Space use and habitat preferences of Australian snubfin and Indo-Pacific humpback dolphins. *Journal of Animal Ecology*, 75, 862-874. doi:10.1111/j.1365-2656.2006.01104.x
- Parra, G., Corkeron, P., & Marsh, H. (2006). Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: Implications for conservation. *Biological Conservation*, 129, 167-180. doi:10.1016/j.biocon.2005.10.031
- Parra, G., Corkeron, P. J., & Marsh, H. (2004). The Indo-Pacific humpback dolphin, *Sousa chinensis* (Osbeck, 1765), in Australian waters: A summary of current knowledge. *Aquatic Mammals*, 30, 197-206.
- Parra, G., Schick, R., & Corkeron, P. J. (2006). Spatial distribution and environmental correlates of Australian snubfin and Indo-Pacific humpback dolphins. *Ecography*, 29, 396-406. doi:10.1111/j.2006.0906-7590.04411.x

- Parsons, E. C. M. (1998). The behaviour of Hong Kong's resident cetaceans: The Indo-Pacific hump-back dolphin and the finless porpoise. *Aquatic Mammals*, 24, 91-106.
- Parsons, E. C. M. (2004a). The behavior and ecology of the Indo-Pacific humpback dolphin (*Sousa chinensis*). *Aquatic Mammals*, 30, 38-55.
- Parsons, E. C. M. (2004b). The potential impacts of pollution on humpback dolphins, with a case study on the Hong Kong population. *Aquatic Mammals*, 30, 18-37.
- Parsons, K., Balcomb, K., Ford, J., & Durban, J. (2009). The social dynamics of southern resident killer whales and conservation implications for this endangered population. *Animal Behaviour*, 77, 963-971. doi:10.1016/j.anbehav.2009.01.018
- Parsons, K., Dallas, J., Claridge, D., Durban, J., Balcomb III, K., Thompson, P., et al. (1999). Amplifying dolphin mitochondrial DNA from faecal plumes. *Molecular Ecology*, 8, 1766-1768. doi:10.1046/j.1365-294X.1999.00723-8.x
- Parsons, K., Durban, J. W., & Claridge, D. E. (2003a). Male-male aggression renders bottlenose dolphin (*Tursiops truncatus*) unconscious. *Aquatic Mammals*, 29, 360-362.
- Parsons, K., Durban, J., & Claridge, D. (2003b). Comparing two alternative methods for sampling small cetaceans for molecular analysis. *Marine Mammal Science*, 19, 224-231.
- Parsons, K., Durban, J., Claridge, D., Balcomb, K., Noble, L., & Thompson, P. (2003). Kinship as a basis for alliance formation between male bottlenose dolphins, *Tursiops truncatus*, in the Bahamas. *Animal Behaviour*, 66, 185-194. doi:10.1006/anbe.2003.2186
- Parsons, K., Noble, L., Reid, R., & Thompson, P. (2002). Mitochondrial genetic diversity and population structuring of UK bottlenose dolphins (*Tursiops truncatus*): Is the NE Scotland population demographically and geographically isolated? *Biological Conservation*, 108, 175.
- Paulos, R., Dudzinski, K., & Kuczaj, S. II (2008). The role of touch in select social interactions of Atlantic spotted dolphin (*Stenella frontalis*) and Indo-Pacific bottlenose dolphin (*Tursiops aduncus*). *Journal of Ethology*, 26, 153-164. doi:10.1007/s10164-0070047-y**
- Pearson, H. (2009). Influences on dusky dolphin (*Lagenorhynchus albirostris obscurus*) fission-fusion dynamics in Admiralty Bay, New Zealand. *Behavioral Ecology and Sociobiology*, 63, 1437-1446. doi:10.1007/s00265-009-0821-7
- Peck, A., Pugh, R., Moors, A., Ellisor, M., Porter, B., Becker, P., et al. (2008). Hexabromocyclododecane in white-sided dolphins: Temporal trend and stereoisomer distribution in tissues. *Environmental Science & Technology*, 42, 2650-2655.
- Peddemors, V. M., & Thompson, G. (1994). Beaching behaviour during shallow water feeding by humpback dolphins *Sousa plumbea*. *Aquatic Mammals*, 20, 65-67.
- Peichl, L., Behrmann, G., & Kröger, R. (2001). For whales and seals the ocean is not blue: A visual pigment loss in marine mammals. *The European Journal Of Neuroscience*, 13, 1520-1528.
- Pennisi, E. (2000). Integrating the many aspects of biology. *Science*, 287, 419-421.
- Pepper, R., & Simmons, J. (1973). In-air visual acuity of the bottlenose dolphin. *Experimental Neurology*, 41, 271-276. doi:10.1016/0014-4886(73)90268-9
- Pepperberg, I. (1999). Rethinking syntax: A commentary on E. Kako's 'Elements of syntax in the systems of three language-trained animals.' *Animal Learning & Behavior*, 27, 15-17.

- Perelberg, A., & Schuster, R. (2008). Coordinated breathing in bottlenose dolphins (*Tursiops truncatus*) as cooperation: Integrating proximate and ultimate explanations. *Journal of Comparative Psychology*, 122, 109-120. doi:10.1037/0735-7036.122.2.109
- Perelberg, A., & Schuster, R. (2009). Bottlenose dolphins (*Tursiops truncatus*) prefer to cooperate when petted: Integrating proximate and ultimate explanations ii. *Journal of Comparative Psychology*, 123, 45-55. doi:10.1037/a0013585
- Pérez-Álvarez, M., Alvarez, E., Aguayo-Lobo, A., & Olavarría, C. (2007). Occurrence and distribution of Chilean dolphin (*Cephalorhynchus eutropis*) in coastal waters of central Chile. *New Zealand Journal of Marine & Freshwater Research*, 41, 405-409.
- Perrin, W., Dolar, M., & Robineau, D. (1999). Spinner dolphins (*Stenella longirostris*) of the western Pacific and southeast Asia: Pelagic and shallow-water forms. *Marine Mammal Science*, 15, 1029-1053.
- Perrin, W., & Mesnick, S. (2003). Sexual ecology of the spinner dolphin, *Stenella longirostris*: Geographic variation in mating system. *Marine Mammal Science*, 19, 462-483.
- Perrin, W., Robertson, K. M., van Bree, P. J., & Mead, J. G. (2007). Cranial description and genetic identity of the holotype specimen of *tursiops aduncus* (Ehrenberg, 1832). *Marine Mammal Science*, 23, 343-357. doi:10.1111/j.1748-7692.2007.00119.x
- Pfeiffer, C. J. (1997). Renal cellular and tissue specializations in the bottlenose dolphin (*Tursiops truncatus*) and beluga whale (*Delphinapterus leucas*). *Aquatic Mammals*, 23, 75-84.
- Philips, J., Nachtigall, P., Au, W., Pawloski, J., & Roitblat, H. (2003). Echolocation in the Risso's dolphin, *Grampus griseus*. *Journal of the Acoustical Society of America*, 113, 605-616. doi:10.1121/1.1527964
- Phillips, D. F., & Brugge, J. F. (1985). Progress in neurophysiology of sound localization. *Annual Review of Psychology*, 36, 245-274.
- Piantadosi, C., & Thalmann, E. (2004). Pathology: Whales, sonar and decompression sickness. *Nature*, 428, 1-2. doi:10.1038/nature02527a
- Pichler, F., Dawson, S., Slooten, E., & Baker, C. (1998). Geographic isolation of Hector's Dolphin populations described by mitochondrial DNA sequences. *Conservation Biology*, 12, 676-682. doi:10.1046/j.1523-1739.1998.96390.x
- Pierce, S., Massey, S., Curtis, N., Smith, G. Jr., Olavarría, C., & Maugel, T. (2004). Microscopic, biochemical, and molecular characteristics of the chilean blob and a comparison with the remains of other sea monsters: Nothing but whales. *Biological Bulletin*, 206, 125-133.
- Pillay, P., & Manger, P. (2004). Testing thermogenesis as the basis for the evolution of cetacean sleep phenomenology. *Journal of Sleep Research*, 13, 353-358. doi:10.1111/j.1365-2869.2004.00419.x
- Pilleri, G., & Knuckey, J. (1969). Behaviour patterns of some delphinidae observed in the western Mediterranean. *Zeitschrift für Tierpsychologie*, 26, 48-72.
- Pimper, L., Remis, M., Goodall, R., & Baker, C. (2009). Teeth and bones as sources of dna for genetic diversity and sex identification of Commerson's dolphins (*Cephalorhynchus commersonii*) from Tierra del Fuego, Argentina. *Aquatic Mammals*, 35, 330-333. doi:10.1578/AM.35.3.2009.330
- Pinedo, M. C., Barreto, A. S., Lammardo, M. P., Andrade, A. L. V., & Geracitano, L. (2002). Northernmost records of the spectacled porpoise, Layard's beaked whale,

- Commerson's dolphin, and Peale's dolphin in the southwestern Atlantic Ocean. *Aquatic Mammals*, 28, 32-37.
- Pistorius, P., Taylor, F., Louw, C., Hanise, B., Bester, M., De Wet, C., et al. (2002). Distribution, movement, and estimated population size of killer whales at Marion Island, December 2000. *South African Journal of Wildlife Research*, 32, 86-92.
- Pitman, R. (2003). Good whale hunting. *Natural History*, 112(10), 24-28.
- Pitman, R., Ballance, L., Mesnick, S., & Chivers, S. (2001). Killer whale predation on sperm whales: Observations and implications. *Marine Mammal Science*, 17, 494-507.
- Pitman, R., & Dutton, P. (2004). Killer whale predation on a leatherback turtle in the Northeast Pacific. *Pacific Science*, 58, 497-498.
- Pitman, R., O'Sullivan, S., & Mase, B. (2003). Killer whales (*Orcinus orca*) attack a school of pantropical spotted dolphins (*Stenella attenuata*) in the Gulf of Mexico. *Aquatic Mammals*, 29, 321-324.
- Plön, S., & Bernard, R. (2006). A review of spermatozoan morphology in cetacea with new data for the genus *Kogia*. *Journal of Zoology*, 269, 466-473. doi:10.1111/j.1469-7998.2006.00061.x
- Pond, C. M. (1978). Morphological aspects and the ecological and mechanical consequences of fat deposition in wild vertebrates. *Annual Review of Ecology and Systematics*, 9, 519-570.
- Popov, V. (2005). Bottlenose dolphins audiogram dependence on azimuth: Evoked potential study. *Journal of the Acoustical Society of America*, 117, 2440.
- Popov, V. & Klishin, V. O. (1998). EEG study of hearing in the common dolphin, *Delphinus delphis*. *Aquatic Mammals*, 24, 13-20.
- Popov, V., & Supin, A. (1997). Detection of temporal gaps in noise in dolphins: Evoked potential study. *Journal of the Acoustical Society of America*, 102, 1169-1176. doi:10.1121/1.419935
- Popov, V., & Supin, A. (2007). Analysis of auditory information in the brains of cetaceans. *Neuroscience and Behavioral Physiology*, 37, 285-291. doi:10.1007/s11055-007-0013-8
- Popov, V., & Supin, A. (2009). Comparison of directional selectivity of hearing in a beluga whale and a bottlenose dolphin. *Journal of the Acoustical Society of America*, 126, 1581-1587. doi:10.1121/1.3177273.
- Popov, V., Supin, A., Ding, W., Kexiong, W., Jianqiang, X., & Songhai, L. (2005). Evoked potential audiogram of the Yangtze finless porpoise *Neophocaena phocaenoides asiaeorientalis* (L.). *Journal of the Acoustical Society of America*, 117, 2728-2731. doi:10.1121/1.1880712
- Popov, V., Supin, A., & Klishin, V. (2001). Auditory brainstem response recovery in the dolphin as revealed by double sound pulses of different frequencies. *Journal of the Acoustical Society of America*, 110, 2227-2233. doi:10.1121/1.1404382
- Popov, V., Supin, A., Klishin, V., & Bulgakova, T. (2006). Monaural and binaural hearing directivity in the bottlenose dolphin: Evoked-potential study. *Journal of the Acoustical Society of America*, 119, 636-644. doi:10.1121/1.2141093
- Popov, V., Supin, A., Klishin, V., Tarakanov, M., & Pletenko, M. (2008). Evidence for double acoustic windows in the dolphin, *Tursiops truncatus*. *Journal of the Acoustical Society of America*, 123, 552-560. doi:10.1121/1.2816564
- Popov, V., Supin, A. Y., Pletenko, M. G., Tarakanov, M. B., Klishin, V. O., Bulgakova, T. N., et al. (2007). Audiogram variability in normal bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 33, 24-33.

- Prahl, S., Ketten, D., & Siebert, U. (2008). Examinations of ears in harbor porpoises (*phocoena phocoena*) from the North and Baltic seas. *Bioacoustics*, 17, 85-87.
- Preen, A. (2004). Distribution, abundance and conservation status of dugongs and dolphins in the southern and western Arabian Gulf. *Biological Conservation*, 118, 205. doi:10.1016/j.biocon.2003.08.014
- Proust, J. (2003). Does metacognition necessarily involve metarepresentation? *Behavioral & Brain Sciences*, 26, 352.
- Psarakos, S., Herzing, D. L., & Marten, K. (2003). Mixed-species associations between Pantropical spotted dolphins (*Stenella attenuata*) and Hawaiian spinner dolphins (*Stenella longirostris*) off Oahu, Hawaii. *Aquatic Mammals*, 29, 390-395.
- Pulster, E., & Maruya, K. (2008). Geographic specificity of Aroclor 1268 in bottlenose dolphins (*Tursiops truncatus*) frequenting the Turtle/Brunswick River Estuary, Georgia (USA). *Science of the Total Environment*, 393, 367-375. doi:10.1016/j.scitotenv.2007.12.031
- Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., & Ridoux, V. (2007). Food and feeding ecology of the common dolphin (*Delphinus delphis*) in the oceanic northeast Atlantic and comparison with its diet in neritic areas. *Marine Mammal Science*, 23(1), 30-47. doi:10.1111/j.1748-7692.2006.00088.x
- Pyle, P., Schramm, M., Keiper, C., & Anderson, S. (1999). Predation on a white shark (*Carcharodon carcharias*) by a killer whale (*Orcinus orca*) and a possible case of competitive displacement. *Marine Mammal Science*, 15, 563-568.
- Pyper, W. (2002). Gene sleuths get behind the whale. *Ecos*, 111, 7.
- Quérouil, S., Silva, M., Cascão, I., Magalhães, S., Seabra, M., Machete, M., et al. (2008). Why do dolphins form mixed-species associations in the Azores? *Ethology*, 114, 1183-1194. doi:10.1111/j.1439-0310.2008.01570.x
- Quick, N., & Janik, V. (2008). Whistle rates of wild bottlenose dolphins (*Tursiops truncatus*): Influences of group size and behavior. *Journal of Comparative Psychology*, 122, 305-311. doi:10.1037/0735-7036.122.3.305
- Quintana-Rizzo, E., Mann, D., & Wells, R. (2006). Estimated communication range of social sounds used by bottlenose dolphins (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 120, 1671-1683. doi:10.1121/1.2226559
- Quintana-Rizzo, E., & Wells, R. (2001). Resighting and association patterns of bottlenose' dolphins (*Tursiops truncatus*) in the Cedar Keys, Florida: Insights into social organization. *Canadian Journal of Zoology*, 79, 447-457. doi:10.1139/cjz-79-3-447
- Raga, J., Banyard, A., Domingo, M., Corteyn, M., Van Bressem, M., Fernández, M., et al. (2008). Dolphin morbillivirus epizootic resurgence, Mediterranean Sea. *Emerging Infectious Diseases*, 14, 471-473.
- Ralston, J., & Herman, L. (1995). Perception and generalization of frequency contours by a bottlenose dolphin (*Tursiops truncatus*). *Journal of Comparative Psychology*, 109, 268-277. doi:10.1037/0735-7036.109.3.268
- Rankin, S., & Barlow, J. (2005). Variation in acoustic behavior of delphinids in the Pacific Ocean based on school size and species composition. *Journal of the Acoustical Society of America*, 117, 2469.
- Rasmussen, M., Lammers, M., Beedholm, K., & Miller, L. (2006). Source levels and harmonic content of whistles in white-beaked dolphins (*Lagenorhynchus albirostris albirostris*). *Journal of the Acoustical Society of America*, 120, 510-517. doi:10.1121/1.2202865

- Rasmussen, M., & Miller, L. A. (2002). Whistles and clicks from white-beaked dolphins, *Lagenorhynchus albirostris albirostris*, recorded in Faxafloia Bay, Iceland. *Aquatic Mammals*, 28, 78-89.
- Rasmussen, M., Wahlberg, M., & Miller, L. (2004). Estimated transmission beam pattern of clicks recorded from free-ranging white-beaked dolphins (*Lagenorhynchus albirostris albirostris*). *Journal of the Acoustical Society of America*, 116, 1826-1831. doi:10.1121/1.1775274
- Rayment, W., Fresne, S., Webster, T., Brogden, S., Dawson, S., & Slooten, E. (2009). Kernel density estimates of alongshore home range of Hector's dolphins at Banks Peninsula, New Zealand. *Marine Mammal Science*, 25, 537-556.
- Rayment, W., & Webster, T. (2009). Observations of Hector's dolphins (*Cephalorhynchus hectori*) associating with inshore fishing trawlers at Banks Peninsula, New Zealand. *New Zealand Journal of Marine & Freshwater Research*, 43, 911-916.
- Rayne, S., Ikonomou, M., Ross, P., Ellis, G., & Barrett-Lennard, L. (2004). PBDEs, PBBs, and PCNs in three communities of free-ranging killer whales (*Orcinus orca*) from the Northeastern Pacific Ocean. *Environmental Science & Technology*, 38, 4293-4299.
- Razafindrakoto, Y., Andrianarivelo, N., & Rosenbaum, H. C. (2004). Sightings, catches, and other records of Indo-Pacific humpback dolphins in the coastal waters of Madagascar. *Aquatic Mammals*, 30, 103-110.
- Read, A., Urian, K., Wilson, B., & Waples, D. (2003). Abundance of bottlenose dolphins in the bays, sounds, and estuaries of North Carolina. *Marine Mammal Science*, 19, 59-73.
- Reeves, R., Leatherwood, S., & Baird, R. (2009). Evidence of a possible decline since 1989 in False Killer Whales (*Pseudorca crassidens*) around the Main Hawaiian Islands. *Pacific Science*, 63, 253-261.
- Rehn, N., Teichert, S., & Thomsen, F. (2007). Structural and temporal emission patterns of variable pulsed calls in free-ranging killer whales (*Orcinus orca*). *Behaviour*, 144, 307-329.
- Rehtanz, M., Bossart, G., Doescher, B., Rector, A., Van Ranst, M., Fair, P., et al. (2009). Bottlenose dolphin (*Tursiops truncatus*) papillomaviruses: Vaccine antigen candidates and screening test development. *Veterinary Microbiology*, 133, 43-53. doi:10.1016/j.vetmic.2008.06.017
- Reich, S., Jimenez, B., Marsili, L., Hernandez, L. M., Schurig, V., & Gonzalez, M. J. (1999). Congener specific determination and enantiomeric ratios of chiral polychlorinated biphenyls in striped dolphins (*Stenella coeruleoalba*) from the Mediterranean Sea. *Environmental Science & Technology*, 33, 1787-1793.
- Reid, K., Mann, J., Weiner, J., & Hecker, N. (1995). Infant development in two aquarium bottlenose dolphins. *Zoo Biology*, 14, 135-147. doi:10.1002/zoo.1430140207
- Reif, J., Fair, P., Adams, J., Joseph, B., Kilpatrick, D., Sanchez, R., et al. (2008). Evaluation and comparison of the health status of Atlantic bottlenose dolphins from the Indian River Lagoon, Florida, and Charleston, South Carolina. *Journal of the American Veterinary Medical Association*, 233, 299-307.
- Reif, J., Mazzoil, M., McCulloch, S., Varela, R., Goldstein, J., Fair, P., et al. (2006). Lobomycosis in Atlantic bottlenose dolphins from the Indian River Lagoon, Florida. *Journal of the American Veterinary Medical Association*, 228, 104-108.
- Reiss, D., & Marino, L. (1995). Self-view television as a test of self-awareness: Only in the eye of the beholder? *Consciousness and Cognition: An International Journal*, 4, 235-238. doi:10.1006/ccog.1995.1030

- Reiss, D., & Marino, L. (2001). Mirror self-recognition in the bottlenose dolphin: A case of cognitive convergence. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 5937.
- Reiss, D., & McCowan, B. (1993). Spontaneous vocal mimicry and production by bottlenose dolphins (*Tursiops truncatus*): Evidence for vocal learning. *Journal of Comparative Psychology*, 107, 301-312. doi:10.1037/0735-7036.107.3.301
- Reiss, D., McCowan, B., & Marino, L. (1997). Communicative and other cognitive characteristics of bottlenose dolphins. *Trends in Cognitive Sciences*, 1, 140-145. doi:10.1016/S1364-6613(97)01046-2
- Remington, N., Stevens, R., Wells, R., Holm, A., Dhungana, S., Taboy, C., et al. (2007). Genetic diversity of coastal bottlenose dolphins revealed by structurally and functionally diverse hemoglobins. *Gene*, 398, 123-131. doi:10.1016/j.gene.2007.02.050
- Rendell, L., & Whitehead, H. (2001). Culture in whales and dolphins. *Behavioral & Brain Sciences*, 24, 309-324.**
- Renner, M., & Bell, K. (2008). A white killer whale in the Central Aleutians. *Arctic*, 61, 102-104.
- Resendes, A., Almeria, S., Dubey, J., Obon, E., Juan-Salles, C., Degollada, E., et al. (2002). Disseminated toxoplasmosis in a Mediterranean pregnant Risso's dolphin (*Grampus griseus*) with transplacental fetal infection. *Journal of Parasitology*, 88, 1029-1032.
- Rettew, D. (2009). New findings in developmental research with animals. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48, 96. doi:10.1097/CHI.0b013e3181930689
- Reyes, L. M. (2006). Cetaceans of central Patagonia, Argentina. *Aquatic Mammals*, 32, 20-30.
- Richard, C. C. (2007). Dolphin social intelligence: Complex alliance relationships in bottlenose dolphins and a consideration of selective environments for extreme brain size evolution in mammals. *Philosophical Transactions: Biological Sciences*, 362, 587-602.
- Richards, D., Wolz, J., & Herman, L. (1984). Vocal mimicry of computer-generated sounds and vocal labeling of objects by a bottlenosed dolphin, *Tursiops truncatus*. *Journal of Comparative Psychology*, 98, 10-28. doi:10.1037/0735-7036.98.1.10
- Richardson, M., & Oelschläger, H. (2002). Time, pattern, and heterochrony: A study of hyperphalangy in the dolphin embryo flipper. *Evolution & Development*, 4, 435-444. doi:10.1046/j.1525-142X.2002.02032.x
- Richardson, W., & Würsig, B. (1997). Influences of man-made noise and other human actions on cetacean behaviour. *Marine and Freshwater Behaviour and Physiology*, 29, 183-209. doi:10.1080/10236249709379006
- Ridgway, S. (2002). Asymmetry and symmetry in brain waves from dolphin left and right hemispheres: Some observations after anesthesia, during quiescent hanging behavior, and during visual obstruction. *Brain, Behavior and Evolution*, 60, 265-274. doi:10.1159/000067192
- Ridgway, S., & Carder, D. (1997). Hearing deficits measured in some *Tursiops truncatus*, and discovery of a deaf/mute dolphin. *Journal of the Acoustical Society of America*, 101, 590-594. doi:10.1121/1.418122
- Ridgway, S., Carder, D., Finneran, J., Keogh, M., Kamolnick, T., Todd, M., et al. (2006). Dolphin continuous auditory vigilance for five days. *Journal of Experimental Biology*, 209, 3621-3628. doi:10.1242/jeb.02405

- Ridgway, S., Carder, D., Kamolnick, T., Smith, R., Schlundt, C., & Elsberry, W. (2001). Hearing and whistling in the deep sea: Depth influences whistle spectra but does not attenuate. *Journal of Experimental Biology*, 204, 3829-3841.
- Ridgway, S., & Harrison, R. (1999). *Handbook of marine mammals Vol. 6*. London: Academic Press.**
- Ridgway, S., Houser, D., Finneran, J., Carder, D., Keogh, M., Van Bonn, W., et al. (2006). Functional imaging of dolphin brain metabolism and blood flow. *Journal of Experimental Biology*, 209, 2902-2910. doi:10.1242/jcb.02348
- Ridgway, S., Keogh, M., Carder, D., Finneran, J., Kamolnick, T., Todd, M., et al. (2009). Dolphins maintain cognitive performance during 72 to 120 hours of continuous auditory vigilance. *Journal of Experimental Biology*, 212, 1519-1527. doi:10.1242/jeb.027896
- Ridoux, V., Guinet, C., Liret, C., Creton, P., Steenstrup, R., & Beauplet, G. (1997). A video sonar as a new tool to study marine mammals in the wild: Measurements of dolphin swimming speed. *Marine Mammal Science*, 13, 196-206.
- Riesch, R., Ford, J., & Thomsen, F. (2006). Stability and group specificity of stereotyped whistles in resident killer whales, *Orcinus orca*, off British Columbia. *Animal Behaviour*, 71, 79-91.
- Riesch, R., Ford, J., & Thomsen, F. (2008). Whistle sequences in wild killer whales (*o*). *Journal of the Acoustical Society of America*, 124, 1822-1829. doi:10.1121/1.2956467.
- Rimskaya-Korsakova, L. (1998). Modeling of total evoked potentials of the dolphin brainstem responses to tonal stimuli. *Sensory Systems*, 12, 285-291.
- Rimskaya-Korsakova, L., & Dubrovsky, N. (1998). Involvement of two sound conduction pathways to the cochlea in forming spatial sound images in dolphins. *Sensory Systems*, 12, 369-375.
- Rimskaya-Korsakova, L., & Dubrovsky, N. (2006). Dolphins' echolocation strategy of target identification: Is it determined by the peripheral auditory encoding? *Acoustical Physics*, 52, 446-454. doi:10.1134/S1063771006040129
- Ritter, F. (2002). Behavioural observations of rough-toothed dolphins (*Steno bredanensis*) off La Gomera, Canary Islands (1995-2000), with special reference to their interactions with humans. *Aquatic Mammals*, 28, 46-59.
- Ritter, F. (2007). Behavioral responses of rough-toothed dolphins to a dead newborn calf. *Marine Mammal Science*, 23, 429-433.
- Rivamonte, L. A. (2009). Bottlenose dolphin (*Tursiops truncatus*) double-slit pupil asymmetries enhance vision. *Aquatic Mammals*, 35, 269-280.
- Robeck, T., & Monfort, S. (2006). Characterization of male killer whale (*Orcinus orca*) sexual maturation and reproductive seasonality. *Theriogenology*, 66, 242-250. doi:10.1016/j.theriogenology.2005.11.007
- Robineau, D., & Vely, M. (1994). *Stenella clymene* (Cetacea, Delphinidae) from the coast of West Africa. *Journal of Mammalogy*, 75, 766.
- Roch, M., Soldevilla, M., Burtenshaw, J., Henderson, E., & Hildebrand, J. (2007). Gaussian mixture model classification of odontocetes in the Southern California Bight and the Gulf of California. *Journal of the Acoustical Society of America*, 121, 1737-1748. doi:10.1121/1.2400663
- Rocklin, D., Santoni, M., Culoli, J., Tomasini, J., Pelletier, D., & Mouillot, D. (2009). Changes in the catch composition of artisanal fisheries attributable to dolphin depredation in a Mediterranean marine reserve. *ICES Journal of Marine Science / Journal du Conseil*, 66, 699-707.

- Rodriguez-Lopez, M. A., & Mignucci-Giannoni, A. A. (1999). A stranded pygmy killer whale (*Feresa attenuata*) in Puerto Rico. *Aquatic Mammals*, 25, 119-121.
- Rogers, C. A., Brunnick, B. J., Herzing, D. L., & Baldwin, J. D. (2004). The social structure of bottlenose dolphins, *Tursiops truncatus*, in the Bahamas. *Marine Mammal Science*, 20, 688-708.
- Rohan J. C. C., Rowe, L., Dawson, S., & Slooten, E. (2008). Abundance and demography of bottlenose dolphins in Dusky Sound, New Zealand, inferred from dorsal fin photographs. *New Zealand Journal of Marine & Freshwater Research*, 42, 439-449.
- Rohr, J., & Fish, F. (2004). Strouhal numbers and optimization of swimming by odontocete cetaceans. *Journal of Experimental Biology*, 207, 1633-1642. doi:10.1242/jeb.00948
- Rohr, J., Fish, F., & Gilpatrick, J Jr.. (2002). Maximum swim speeds of captive and free-ranging delphinids: Critical analysis of extraordinary performance. *Marine Mammal Science*, 18, 1-19.
- Rohr, J., Latz, M., Fallon, S., Nauen, J., & Hendricks, E. (1998). Experimental approaches towards interpreting dolphin-stimulated bioluminescence. *Journal of Experimental Biology*, 201, 1447-1460.
- Roitblat, H., Penner, R., & Nachtigall, P. (1990). Matching-to-sample by an echolocating dolphin (*Tursiops truncatus*). *Journal of Experimental Psychology: Animal Behavior Processes*, 16, 85-95. 2116, doi:10.1037/0097-7403.16.1.85
- Roman, J. (2001). Against the tide. *Wildlife Conservation*, 104(3), 12.
- Romanenko, E. (2004). Some results of studying the acoustics of dolphins. *Acoustical Physics*, 50, 343-349. doi:10.1134/1.1739503
- Rooney, A., & Meritt, D. (1999). Microsatellite diversity in captive bottlenose dolphins (*Tursiops truncatus*). *Journal of Heredity*, 90, 228-233.
- Rosas, F. C. W., Monteiro-Filho, E. L. A., Marigo, J., Santos, R. A., Andrade, A. L. V., Rautenberg, M., et al. (2002). The striped dolphin, *Stenella coeruleoalba* (Cetacea:Delphinidae), on the coast of São Paulo State, southeastern Brazil. *Aquatic Mammals*, 28, 60-66.
- Rose, N. A., Parsons, E. C. M., & Garinato, R. (2009). *The case against marine mammals in captivity*. (4th ed). The Humane Society of the United States and the World Society for the Protection of Animals.**
- Rosel, P., Forgetta, V., & Dewar, K. (2005). Isolation and characterization of twelve polymorphic microsatellite markers in bottlenose dolphins (*Tursiops truncatus*). *Molecular Ecology Notes*, 5, 830-833. doi:10.1111/j.1471-8286.2005.01078.x
- Rosel, P., Hansen, L., & Hohn, A. (2009). Restricted dispersal in a continuously distributed marine species: Common bottlenose dolphins *Tursiops truncatus* in coastal waters of the western North Atlantic. *Molecular Ecology*, 18, 5030-5045. doi:10.1111/j.1365-294X.2009.04413.x
- Rosing-Asvid, A., Heide-Jørgensen, M., & Richard, P. (1998). Dive patterns of belugas (*Delphinapterus leucas*) in waters near eastern Devon Island. *Arctic*, 51, 17-26.
- Ross, P. (2006). Fireproof killer whales (*Orcinus orca*): Flame-retardant chemicals and the conservation imperative in the charismatic icon of British Columbia, Canada. *Canadian Journal of Fisheries & Aquatic Sciences*, 63, 224-234. doi:10.1139/F05-244
- Rossbach, K. A. (1999). Cooperative feeding among bottlenose dolphins (*Tursiops truncatus*) near Grand Bahama Island, Bahamas. *Aquatic Mammals*, 25, 163-167.

- Rossi-Santos, M., Baracho, C., Cipolotti, S., & Marcovaldi, E. (2008). Cetacean sightings near south Georgia islands, south Atlantic Ocean. *Polar Biology*, 31, 63-68. doi:10.1007/s00300-007-0333-8
- Rotstein, D., Burdett, L., McLellan, W., Schwacke, L., Rowles, T., Terio, K., et al. (2009). Lobomycosis in offshore bottlenose dolphins (*Tursiops truncatus*), North Carolina. *Emerging Infectious Diseases*, 15, 588-590. doi:10.3201/eid1504.081358
- Rotstein, D., Harms, C., Lovewell, G., & Hohn, A. (2007). Gastric leiomyoma in a free-living Atlantic bottlenosed dolphin (*Tursiops truncatus*). *Veterinary Record: Journal of the British Veterinary Association*, 160, 130-131.
- Rowe, L. E., & Dawson, S. M. (2009). Determining the sex of bottlenose dolphins from doubtful sound using dorsal fin photographs. *Marine Mammal Science*, 25, 19-34. doi:10.1111/j.1748-7692.2008.00235.x
- Ruelas-Inzunza, J., Horvat, M., Perez-Cortes, H., & Paez-Osuna, F. (2003). Methylmercury and total mercury distribution of tissues of gray whales (*Eschrichtius robustus*) and spinner dolphins (*Stenella longirostris*) stranded along the lower Gulf of California, Mexico. *Ciencias Marinas*, 29, 1-8.
- Rugh, D., Shelden, K., & Mahoney, B. (2000). Distribution of belugas, in Cook Inlet, during June/July 1993-2000. *Marine Fisheries Review*, 62(3), 6-21.
- Ruiz, C., Nollens, H., Venn-Watson, S., Green, L., Wells, R., Walsh, M., et al. (2009). Baseline circulating immunoglobulin G levels in managed collection and free-ranging bottlenose dolphins (*Tursiops truncatus*). *Developmental and Comparative Immunology*, 33, 449-455.
- Ruiz, C., Sánchez, A., & Maldonado, F. (2009). Social and individual behavior of a group of bottlenose dolphins (*Tursiops truncatus*) in open and closed facilities. *Veterinaria Mexico*, 40, 381-387.
- Russell, J., Simonoff, J., & Nightengale, J. (1997). Nursing behaviors of beluga calves (*Delphinapterus leucas*) born in captivity. *Zoo Biology*, 16, 247-262. doi:10.1002/(SICI)1098-2361(1997)16:3<247::AID-ZOO5>3.0.CO;2-A
- Rutenko, A., & Vishnyakov, A. (2006). Time sequences of sonar signals generated by a beluga whale when locating underwater objects. *Acoustical Physics*, 52, 314-323. doi:10.1134/S1063771006030122
- Ryabov, V., & Zaslavskii, G. (1998). Interference resistance of the Dolphin echolocator. *Sensory Systems*, 12, 151-156.
- Ryabov, V., & Zaslavskii, G. (1999). Targeting differentiation in conditions of jamming reflections and monaural hearing in a bottlenose dolphin. *Sensory Systems*, 13, 325- 333.
- Saayman, G., Tayler, C., & Bower, D. (1973). Diurnal activity cycles in captive and free-ranging Indian Ocean bottlenose dolphins (*Tursiops aduncus* Ehrenburg). *Behaviour*, 44(3), 212-233. doi:10.1163/156853973X00409
- Sakai, M., Hishii, T., Takeda, S., & Kohshima, S. (2006a). Flipper rubbing behaviors in wild bottlenose dolphins (*Tursiops aduncus*). *Marine Mammal Science*, 22, 966-978.
- Sakai, M., Hishii, T., Takeda, S., & Kohshima, S. (2006b). Laterality of flipper rubbing behaviour in wild bottlenose dolphins (*Tursiops aduncus*): Caused by asymmetry of eye use? *Behavioural Brain Research*, 170, 204-210.
- Samuels, A., & Gifford, T. (1997). A quantitative assessment of dominance relations among bottlenose dolphins. *Marine Mammal Science*, 13, 70-99.

- Samuels, A., & Spradlin, T. (1995). Quantitative behavioral study of bottlenose dolphins in swim-with-dolphin programs in the united states. *Marine Mammal Science*, 11, 520-544.
- Samuels, A., & Tyack, P. L. (2000). Flukeprints. In J. Mann, R. C. Connor, P. L. Tyack, & H. Whitehead (Eds.), *Cetacean societies: Field studies of dolphins and whales* (pp. 9-44). Chicago: The University of Chicago Press.**
- Samuels, A., & Worthy, G. (2004). Variability in fatty acid composition of bottlenose dolphin (*Tursiops truncatus*) blubber as a function of body site, season, and reproductive state. *Canadian Journal of Zoology*, 82, 1933-1942. doi:10.1139/Z05-001
- Santos, L. (1995). When dolphins go bad. *Sciences*, 35(5), 47.
- dos Santos, M. E., Louro, S., Couchinho, M., & Brito, C. (2005). Whistles of bottlenose dolphins (*Tursiops truncatus*) in the Sado Estuary, Portugal: Characteristics, production rates, and long-term contour stability. *Aquatic Mammals*, 31, 453-462.
- Sapr'gkyn, V. (1977). The invariability of stimuli recognition by dolphins. *Fiziologicheskii Zhurnal SSSR im I. M. Sechenova*, 63, 1083-1087.
- Sargeant, B., & Mann, J. (2009). Developmental evidence for foraging traditions in wild bottlenose dolphins. *Animal Behaviour*, 78, 715-721. doi:10.1016/j.anbehav.2009.05.037
- Sargeant, B., Mann, J., Berggren, P., & Krützen, M. (2005). Specialization and development of beach hunting, a rare foraging behavior, by wild bottlenose dolphins (*Tursiops* sp.). *Canadian Journal of Zoology*, 83, 1400-1410.
- Sargeant, B., Wirsing, A., Heithaus, M., & Mann, J. (2007). Can environmental heterogeneity explain individual foraging variation in wild bottlenose dolphins (*Tursiops* sp.)? *Behavioral Ecology and Sociobiology*, 61, 679-688. doi:10.1007/s00265-006-0296-8
- Sarko, D., Marino, L., & Reiss, D. (2002). A bottlenose dolphin's (*Tursiops truncatus*) responses to its mirror image: Further analysis. *International Journal of Comparative Psychology*, 15, 69-76.
- Sassu, R., & Cozzi, B. (2007). The external and middle ear of the striped dolphin *Stenella coeruleoalba* (Meyen 1833). *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C*, 36, 197-201. doi:10.1111/j.1439 0264.2006.00750.x
- Saulitis, E., Matkin, C., Barrett-Lennard, L., Heise, K., & Ellis, G. (2000). Foraging strategies of sympatric killer whale (*Orcinus orca*) populations in Prince William Sound, Alaska. *Marine Mammal Science*, 16, 94-109.
- Saulitis, E., Matkin, C., & Fay, F. (2005). Vocal repertoire and acoustic behavior of the isolated AT1 killer whale subpopulation in southern Alaska. *Canadian Journal of Zoology*, 83, 1015-1029. doi:10.1139/Z05-089.
- Sayigh, L., Esch, H., Wells, R., & Janik, V. (2007). Facts about signature whistles of bottlenose dolphins, *Tursiops truncatus*. *Animal Behaviour*, 74, 1631-1642. doi:10.1016/j.anbehav.2007.02.018
- Sayigh, L., Tyack, P., Wells, R., & Scott, M. (1990). Signature whistles of free-ranging bottlenose dolphins *Tursiops truncatus*: Stability and mother-offspring comparisons. *Behavioral Ecology and Sociobiology*, 26, 247-260. doi:10.1007/BF00178318
- Sayigh, L., Tyack, P., Wells, R., & Scott, M. (1995). Sex difference in signature whistle production of free-ranging bottlenose dolphins, *Tursiops truncatus*. *Behavioral Ecology and Sociobiology*, 36, 171-177. doi:10.1007/BF00177793

- Sayigh, L., Tyack, P., Wells, R., Solow, A., Scott, M., & Irvine, A. (1999). Individual recognition in wild bottlenose dolphins: A field test using playback experiments. *Animal Behaviour*, 57, 41-50. doi:10.1006/anbe.1998.0961
- Scano, P., Cesare Marincola, F., Locci, E., & Lai, A. (2006). ^1H and ^{13}C NMR studies of melon and head blubber of the striped dolphin (*Stenella coeruleoalba*). *Lipids*, 41, 1039-1048.
- Scano, P., Maxia, C., Maggiani, F., Crnjar, R., Lai, A., & Sirigu, P. (2005). A histological and NMR study of the melon of the striped dolphin (*Stenella coeruleoalba*). *Chemistry & Physics of Lipids*, 134, 21-28. doi:10.1016/j.chemphyslip.2004.10.001
- Schaefer, A. M., Reif, J. S., Goldstein, J. D., Ryan, C. N., Fair, P. A., & Bossart, G. D. (2009). Serological evidence of exposure to selected viral, bacterial, and protozoal pathogens in free-ranging atlantic bottlenose dolphins (*Tursiops truncatus*) from the Indian River Lagoon, Florida, and Charleston, South Carolina. *Aquatic Mammals*, 35, 163-170.
- Scheel, D., Matkin, C., & Saulitis, E. (2001). Distribution of killer whale pods in Prince William Sound, Alaska 1984-1996. *Marine Mammal Science*, 17, 555-569.
- Scheer, M., Hofmann, B., & Behr, I. (2004). Ethogram of selected behaviors initiated by free-ranging short-finned pilot whales (*Globicephala macrorhynchus*) and directed to human swimmers during open water encounters. *Anthrozoös*, 17, 244-258. doi:10.2752/089279304785643267
- Schiavini, A., Pedraza, S., Crespo, E., Gonzalez, R., & Dans, S. (1999). Abundance of dusky dolphins (*Lagenorhynchus obscurus*) off north and central Patagonia, Argentina, in spring and a comparison with incidental catch in fisheries. *Marine Mammal Science*, 15, 828-840.
- Schlundt, C., Dear, R., Green, L., Houser, D., & Finneran, J. (2007). Simultaneously measured behavioral and electrophysiological hearing thresholds in a bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 122, 615-622. doi:10.1121/1.2737982
- Schlundt, C., Finneran, J., Carder, D., & Ridgway, S. (2000). Temporary shift in masked hearing thresholds of bottlenose dolphins, *Tursiops truncatus*, and white whales, *Delphinapterus leucas*, after exposure to intense tones. *Journal of the Acoustical Society of America*, 107, 3496-3508. doi:10.1121/1.429420
- Schmidt, K. (1994). Scientists count a rising tide of whales in the seas. *Science*, 263, 25-26.
- Schneider, D. (1999). Shedding light on dolphins. *American Scientist*, 87, 216.
- Schofield, T. D., Early, G., Wenzel, F. W., Matassa, K., Perry, C., Beekman, G., et al. (2008). Rehabilitation and homing behavior of a satellite-tracked harbor porpoise (*Phocoena phocoena*). *Aquatic Mammals*, 34, 1-8.
- Schrope, M. (2002). Whale deaths caused by US Navy's sonar. *Nature*, 415, 106-108.
- Schuller, D., Kadko, D., & Smith, C. (2004). Use of $^{210}\text{Pb}/^{226}\text{Ra}$ disequilibria in the dating of deep-sea whale falls. *Earth & Planetary Science Letters*, 218, 277-289. doi:10.1016/S0012-821X(03)00690-3
- Schusterman, R. J., Thomas, J. A., & Wood, F. G. (1986). *Dolphin cognition and behavior: A comparative approach*. Hillsdale, NJ: Lawrence Erlbaum Associates.**
- Schwacke, L., Hall, A., Townsend, F., Wells, R., Hansen, L., Hohn, A., et al. (2009). Hematologic and serum biochemical reference intervals for free-ranging common bottlenose dolphins (*Tursiops truncatus*) and variation in the distributions of clinicopathologic values related to geographic sampling site. *American Journal of Veterinary Research*, 70, 973-985.

- Schwacke, L., Voit, E., Hansen, L., Wells, R., Mitchum, G., Hohn, A., & Fair, P. A. (2002). Probabilistic risk assessment of reproductive effects of polychlorinated diphenyls on bottlenose dolphins (*Tursiops truncatus*) from the Southeast United States coast. *Environmental Toxicology & Chemistry*, 21, 2752-2764.
- Schwartzkopff, J. (1977). Auditory communication in lower animals: Role of auditory physiology. *Annual Review of Psychology*, 28, 61-84.
- Scott, E., Mann, J., Watson-Capps, J., Sargeant, B., & Connor, R. (2005). Aggression in bottlenose dolphins: Evidence for sexual coercion, male-male competition, and female tolerance through analysis of tooth-rake marks and behaviour. *Behaviour*, 142, 21-44. doi:10.1163/1568539053627712
- Scott, M., & Chivers, S. (2009). Movements and diving behavior of pelagic spotted dolphins. *Marine Mammal Science*, 25, 137-160.
- Secchi, E. R., & Siciliano, S. (1995). Comments on the southern range of the spinner dolphin (*Stenella longirostris*) in the western South Atlantic. *Aquatic Mammals*, 21, 105-108.
- Secchi, E. R., & Vaske Jr., T. (1998). Killer whale (*Orcinus orca*) sightings and depredation on tuna and swordfish longline catches in southern Brazil. *Aquatic Mammals*, 24, 117-122.
- Segura, I., Rocha-Olivares, A., Flores-Ramírez, S., & Rojas-Bracho, L. (2006). Conservation implications of the genetic and ecological distinction of *Tursiops truncatus* ecotypes in the Gulf of California. *Biological Conservation*, 133, 336-346. doi:10.1016/j.biocon.2006.06.017.
- Seibel, B., & Dierssen, H. (2003). Cascading trophic impacts of reduced biomass in the Ross Sea, Antarctica: Just the tip of the iceberg? *Biological Bulletin*, 205, 93.
- Sekiguchi, K. (1995). Occurrence, behavior and feeding habits of harbor porpoises (*Phocoena phocoena*) at Pajaro Dunes, Monterey Bay, California. *Aquatic Mammals*, 21, 91-103.
- Sekiguchi, Y., Arai, K., & Kohshima, S. (2006a). Sleep behaviour: Activity and sleep in dolphins. *Nature*, 441, E10-E11. doi:10.1038/nature04899
- Sekiguchi, Y., Arai, K., & Kohshima, S. (2006b). Sleep behaviour: Sleep in continuously active dolphins. *Nature*, 441, E9-E10. doi:10.1038/nature04898
- Sekiguchi, Y., & Kohshima, S. (2003). Resting behaviors of captive bottlenose dolphins (*Tursiops truncatus*). *Physiology & Behavior*, 79, 643-653. doi:10.1016/S0031-9384(03)00119-7
- Servais, V. (1999). Some comments on context embodiment in zootherapy: The case of the Autidolfin project. *Anthrozoös*, 12, 5-15. doi:10.2752/089279399787000417
- Sezaki, K., Hirosaki, Y., Watabe, S., & Hashimoto, K. (1984). Electrophoretic characters of the hybrids between two dolphins *Tursiops truncatus* and *Grampus griseus*. *Bulletin of the Japanese Society of Scientific Fisheries*, 50, 1771-1776.
- Shaffer, S., Costa, D., Williams, T., & Ridgway, S. (1997). Diving and swimming performance of white whales, *Delphinapterus leucas*: A assessment of plasma.. *Journal of Experimental Biology*, 200, 3091-3099.
- Shaffer, S., & Costa, P. (2006). A database for the study of marine mammal behavior: Gap analysis, data standardization, and future directions. UC Santa Cruz: Retrieved from: <http://www.escholarship.org/uc/item/044838bg>**
- Shakespeare, M. (2006). Rescue and rehab. *Wildlife Conservation*, 109(6), 56-57.
- Shane, S. (1995). Behavior patterns of pilot whales and Risso's dolphins off Santa Catalina Island, California. *Aquatic Mammals*, 21, 195-197.
- Shane, S., Tepley, L., & Costello, L. (1993). Life-threatening contact between a woman and a pilot whale captured on film. *Marine Mammal Science*, 9, 331-336.

- Shane, S., Wells, R., & Würsig, B. (1986). Ecology, behavior and social organization of the bottlenose dolphin: A review. *Marine Mammal Science*, 2, 34-63.
- Shelden, K., Rugh, D., Mahoney, B., & Dahlheim, M. (2003). Killer whale predation on belugas in Cook Inlet, Alaska: Implications for a depleted population. *Marine Mammal Science*, 19, 529-544.
- Shirai, K., & Sakai, T. (1997). Haematological findings in captive dolphins and whales. *Australian Veterinary Journal*, 75, 512-514.
- Shirakihara, K., Shirakihara, M., & Yamamoto, Y. (2007). Distribution and abundance of finless porpoise in the Inland Sea of Japan. *Marine Biology*, 150, 1025-1032. doi:10.1007/s00227-006-0363-8
- Shirakihara, M., Seki, K., Takemura, A., Shirakihara, K., Yoshida, H., & Yamazaki, T. (2008). Food habits of finless porpoises *neophocaena phocaenoides* in Western Kyushu, Japan. *Journal of Mammalogy*, 89, 1248-1256.
- Shirakihara, M., Yoshida, H., & Shirakihara, K. (2003). Indo-Pacific bottlenose dolphins *Tursiops aduncus* in Amakusa, western Kyushu, Japan. *Fisheries Science*, 69, 654-656. doi:10.1046/j.1444-2906.2003.00669.x
- Shoham-Frider, E., Amiel, S., Roditi-Elasar, M., & Kress, N. (2002). Risso's dolphin (*Grampus griseus*) stranding on the coast of Israel (eastern Mediterranean). Autopsy results and trace metal concentrations. *Science of the Total Environment*, 295, 157-166.
- Shoham-Frider, E., Kress, N., Wynne, D., Scheinin, A., Roditi-Elsar, M., & Kerem, D. (2009). Persistent organochlorine pollutants and heavy metals in tissues of common bottlenose dolphin (*Tursiops truncatus*) from the Levantine Basin of the Eastern Mediterranean. *Chemosphere*, 77, 621-627. doi:10.1016/j.chemosphere.2009.08.048
- Shpak, O. V., Lyamin, O. I., Manger, P. R., Siegel, J., & Mukhametov, L. M. (2009). States of rest and activity in the Commerson's dolphin *Cephalorhynchus commersonii*. *Journal of Evolutionary Biochemistry & Physiology*, 45, 111-119.
- Shulman, S. (1990). Dolphin excused military service. *Nature*, 348(6298), 189.
- Shyan, M., & Herman, L. (1987). Determinants of recognition of gestural signs in an artificial language by Atlantic bottle-nosed dolphins (*Tursiops truncatus*) and humans (*Homo sapiens*). *Journal of Comparative Psychology*, 101, 112-125. doi:10.1037/0735-7036.101.2.112
- Shyan, M., Merritt, D., Kohlmeier, N., Barton, K., & Tenge, J. (2002). Effects of pool size on free-choice selections by Atlantic bottlenosed dolphins at one zoo facility. *Journal of Applied Animal Welfare Science*, 5, 215-225. doi:10.1207/S15327604JAWS0503_05
- Siebert, U., Prenger-Berninghoff, E., & Weiss, R. (2009). Regional differences in bacterial flora in harbour porpoises from the North Atlantic: Environmental effects? *Journal of Applied Microbiology*, 106, 329-337. doi:10.1111/j.1365-2672.2008.04006.x
- Siebenalen, J., & Caldwell, D. (1956). Cooperation among adult dolphins. *Journal of Mammalogy*, 37, 126-128.
- Silber, G. K., & Fertl, D. (1995). Intentional beaching by bottlenose dolphins (*Tursiops truncatus*) in the Colorado River Delta, Mexico. *Aquatic Mammals*, 21, 183-186.
- Silva, F., & da Silva Jr., J. (2009). Circadian and seasonal rhythms in the behavior of spinner dolphins (*Stenella longirostris*). *Marine Mammal Science*, 25, 176-186.
- Silva Jr., J. M., Pandolfo, L. J., & Sazima, I. (2004). Vomiting behavior of the spinner dolphin (*Stenella longirostris*) and squid meals. *Aquatic Mammals*, 30, 271-274.

- Silva, M., Prieto, R., Magalhães, S., Seabra, M., Santos, R., & Hammond, P. (2008). Ranging patterns of bottlenose dolphins living in oceanic waters: implications for population structure. *Marine Biology*, 156, 179-192. doi:10.1007/s00227-008-1075-z
- Simard, P., & Gowans, S. (2004). Two calves in echelon: An alloparental association in Atlantic white-sided dolphins (*Lagenorhynchus albirostris acutus*)? *Aquatic Mammals*, 30, 330-334.
- Simard, P., & Gowans, S. (2008). Group movements of white-beaked dolphins (*Lagenorhynchus albirostris albirostris*) near Halifax, Canada. *Aquatic Mammals*, 34, 331-337.
- Simila, T. (1997). Sonar observations of killer whales (*Orcinus orca*) feeding on herring schools. *Aquatic Mammals*, 23, 119-126.
- Simmonds, M. (2006). Into the brains of whales. *Applied Animal Behaviour Science*, 100, 103-116. doi:10.1016/j.applanim.2006.04.015
- Simon, M., Hanson, M., Murrey, L., Tougaard, J., & Ugarte, F. (2009). From captivity to the wild and back: An attempt to release Keiko the killer whale. *Marine Mammal Science*, 25, 693-705. doi:10.1111/j.1748-7692.2009.00287.x
- Simon, M., McGregor, P., & Ugarte, F. (2007). The relationship between the acoustic behaviour and surface activity of killer whales (*Orcinus orca*) that feed on herring (*Clupea harengus*). *Acta Ethologica*, 10, 47-53. doi:10.1007/s10211-007-0029-7
- Simon, M., Ugarte, F., Wahlberg, M., & Miller, L. (2006). Icelandic killer whales *Orcinus orca* use a pulsed call suitable for manipulating the schooling behaviour of Herring *Clupea harengus*. *Bioacoustics*, 16, 57-74.
- Simon, M., Wahlberg, M., Ugarte, F., & Miller, L. (2005). Acoustic characteristics of underwater tail slaps used by Norwegian and Icelandic killer whales (*Orcinus orca*) to debilitate herring (*Clupea harengus*). *Journal of Experimental Biology*, 208, 2459-2466. doi:10.1242/jeb.01619
- Sinclair, E., Moore, S., Friday, N., Zeppelin, T., & Waite, J. (2005). Do patterns of Steller sea lion (*Eumetopias jubatus*) diet, population trend and cetacean occurrence reflect oceanographic domains from the Alaska Peninsula to the central Aleutian Islands? *Fisheries Oceanography*, 14, 223-242. doi:10.1111/j.1365-2419.2005.00375.x
- Sleno, G., Kingsley, M., & Gosselin, S. (2001). Movements and dive behaviour of belugas in Northern Quebec. *Arctic*, 54, 262-275.
- Slooten, E., Fletcher, D., & Taylor, B. (2000). Accounting for uncertainty in risk assessment: Case study of Hector's dolphin mortality due to gillnet entanglement. *Conservation Biology*, 14, 1264-1270. doi:10.1046/j.1523-1739.2000.00099-411.x
- Slooten, E., Rayment, W., & Dawson, S. (2006). Offshore distribution of Hector's dolphins at Banks Peninsula, New Zealand: Is the Banks Peninsula Marine Mammal sanctuary large enough? *New Zealand Journal of Marine & Freshwater Research*, 40, 333-343.
- Smith, B. (2007). Fish tales. *Wildlife Conservation*, 110(1), 26-31.
- Smith, B. (2009). Witness to extinction: How we failed to save the Yangtze river dolphin. *Marine Mammal Science*, 25, 987-991.**
- Smith, H., Samuels, A., & Bradley, S. (2008). Reducing risky interactions between tourists and free-ranging dolphins (*Tursiops* sp.) in an artificial feeding program at Monkey Mia, Western Australia. *Tourism Management*, 29, 994-1001. doi:10.1016/j.tourman.2008.01.001
- Smith, J., Schull, J., Strote, J., McGee, K., Egnor, R., & Erb, L. (1995). The uncertain response in the bottlenosed dolphin (*Tursiops truncatus*). *Journal of*

Experimental Psychology: General, 124, 391-408. doi:10.1037/0096-3445.124.4.391

- Smith, T., Dietz, R., Heide-Jorgensen, M., Orr, J., & Richard, P. (2001). Summer and autumn movements and habitat use by belugas in the Canadian high Arctic and adjacent areas. *Arctic*, 54, 207-222.
- Smith, T., & Mooney, M. (1994). Shape of scapular fossae in freshwater and marine dolphins. *Journal of Mammalogy*, 75, 515.
- Smith, T., Siniff, D., Reichle, R., & Stone, S. (1981). Coordinated behavior of killer whales, *Orcinus orca*, hunting a crabeater seal, *Lobodon carcinophagus*. *Canadian Journal of Zoology*, 59, 1185-1189.
- Smolker, R., Mann, J., & Smuts, B. (1993). Use of signature whistles during separations and reunions by wild bottlenose dolphin mothers and infants. *Behavioral Ecology and Sociobiology*, 33, 393-402. doi:10.1007/BF00170254
- Smolker, R., & Pepper, J. (1999). Whistle convergence among allied male bottlenose dolphins (*Delphinidae*, *Tursiops* sp.). *Ethology*, 105, 595-617. doi:10.1046/j.1439-0310.1999.00441.x
- Smolker, R., Richards, A., Connor, R., Mann, J., & Berggren, P. (1997). Sponge carrying by dolphins (*Delphinidae*, *Tursiops* sp.): A foraging specialization involving tool use? *Ethology*, 103, 454-465.
- Smolker, R., Richards, A., Connor, R., & Pepper, J. (1992). Sex differences in patterns of association among Indian Ocean bottlenose dolphins. *Behaviour*, 123, 38-69. doi:10.1163/156853992X00101
- Snowdon, C. T. (1983). Ethology, comparative psychology, and animal behavior. *Annual Review of Psychology*, 34, 63-94.
- Sobel, N., Supin, A., & Myslobodsky, M. (1994). Rotational swimming tendencies in the dolphin (*Tursiops truncatus*). *Behavioural Brain Research*, 65, 41-45. doi:10.1016/0166-4328(94)90071-X
- Soldevilla, M., Henderson, E., Campbell, G., Wiggins, S., Hildebrand, J., & Roch, M. (2008). Classification of Risso's and Pacific white-sided dolphins using spectral properties of echolocation clicks. *Journal of the Acoustical Society of America*, 124, 609-624. doi:10.1121/1.2932059
- Solntseva, G. N. (2002). Early embryogenesis of the vestibular apparatus in mammals with different ecologies. *Aquatic Mammals*, 28, 159-169.
- Songhai, L., Akamatsu, T., Ding, W., Kexiong, W., Shouyue, D., Xiujiang, Z., et al. (2008). Indirect evidence of boat avoidance behavior of Yangtze finless porpoises. *Bioacoustics*, 17, 174-176.
- Songhai, L., Ding, W., Kexiong, W., Akamatsu, T., Zhiqiang, M., & Jiabo, H. (2007). Echolocation click sounds from wild inshore finless porpoise (*Neophocaena phocaenoides sunameri*) with comparisons to the sonar of riverine *N. p. asiaticorientalis*. *Journal of the Acoustical Society of America*, 121, 3938-3946. doi:10.1121/1.2721658
- Songhai, L., Ding, W., Kexiong, W., Jiangiang, X., & Tomorari, A. (2007). The ontogeny of echolocation in a Yangtze finless porpoise (*Neophocaena phocaenoides asiaticorientalis*). *Journal of the Acoustical Society of America*, 122, 715-718. doi:10.1121/1.2747203
- Songhai, L., Kexiong, W., Ding, W., & Akamatsu, T. (2005). Origin of the double- and multi pulse structure of echolocation signals in Yangtze finless porpoise (*Neophocaena phocaenoides asiaticorientalis*). *Journal of the Acoustical Society of America*, 118, 3934-3940. doi:10.1121/1.2126919

- Songhai, L., Wang, K., Wang, D., Shouyue, D., & Tomonari, A. (2008). Simultaneous production of low- and high-frequency sounds by neonatal finless porpoises. *Journal of the Acoustical Society of America*, 124, 716-718. doi:10.1121/1.2945152
- Sorensen, K., Ridgway, S., & Venn-Watson, S. (2008). Trace and non-trace elements in blood cells of bottlenose dolphins (*Tursiops truncatus*): Variations with values from liver function indicators. *Journal of Wildlife Diseases*, 44, 304-317.
- Spanier, E., Goffman, O., Kerem, D., & Lavalli, K. (2000). Injury of an Indian Ocean bottlenose dolphin (*Tursiops aduncus*) in the Red Sea by a stingray spine. *Aquatic Mammals*, 26, 196-201.
- Spear, N. E., Miller, J. S., & Jagielo, J. A. (1990). Animal memory and learning. *Annual Review of Psychology*, 41, 169-211.
- Speckman, S., & Piatt, J. (2000). Historic and current use of lower Cook Inlet, Alaska, by belugas. *Marine Fisheries Review*, 62(3), 22-27.
- Spitz, J., Richard, E., Meynier, L., Pusineri, C., & Ridoux, V. (2006). Dietary plasticity of the oceanic striped dolphin, *Stenella coeruleoalba*, in the neritic waters of the Bay of Biscay. *Journal of Sea Research*, 55, 309-320. doi:10.1016/j.seares.2006.02.001
- Spitz, J., Rousseau, Y., & Ridoux, V. (2006). Diet overlap between harbour porpoise and bottlenose dolphin: An argument in favour of interference competition for food? *Estuarine Coastal & Shelf Science*, 70, 259-270. doi:10.1016/j.ecss.2006.04.020
- Spong, P., & White, D. (1971). Visual acuity and discrimination learning in the dolphin (*Lagenorhynchus albirostris obliquidens*). *Experimental Neurology*, 31, 431-436. doi:10.1016/0014-4886(71)90245-7
- Springer, A., Estes, J., Van Vliet, G., Williams, T., Doak, D., Danner, E., et al. (2003). Sequential megafaunal collapse in the North Pacific Ocean: An ongoing legacy of industrial whaling? *Proceedings of the National Academy of Sciences of the United States of America*, 100, 12223-12228. doi:10.1073/pnas.1635156100
- Springer, S. (1957). Some observations of the behavior of schools of fishes in the Gulf of Mexico and adjacent waters. *Ecology*, 38, 166-171.
- St. Aubin, D., & Geraci, J. (1992). Thyroid hormone balance in Beluga whales, *Delphinapterus leucas*: dynamics after capture and influence of thyrotropin. *Canadian Journal of Veterinary Research*, 56, 1-5.
- Stafne, G., & Manger, P. (2004, October). Predominance of clockwise swimming during rest in Southern Hemisphere dolphins. *Physiology & Behavior*, 82, 919-926. doi:10.1016/j.physbeh.2004.08.001
- Starodubtsev, Y. (1993). The formation of complex skill in black sea dolphins under free choice conditions. *Neuroscience and Behavioral Physiology*, 23, 210-217. doi:10.1007/BF01182919
- Stavros, H., Bossart, G., Hulsey, T., & Fair, P. (2007). Trace element concentrations in skin of free-ranging bottlenose dolphins (*Tursiops truncatus*) from the southeast Atlantic coast. *Science of the Total Environment*, 388, 300-315.
- Stefanie K., G., Richard C., C., Robert K., E., & Frank, C. (2005). A division of labour with role specialization in group-hunting bottlenose dolphins (*Tursiops truncatus*) off Cedar Key, Florida. *Proceedings: Biological Sciences*, 272, 135-140.
- Steiner, A., & Bossjey, M. (2008). Some reproductive parameters of an estuarine population of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*). *Aquatic Mammals*, 34, 84-92.
- Stern, G., Macdonald, C., Armstrong, D., Dunn, B., Fuchs, C., Harwood, L., et al. (2005). Spatial trends and factors affecting variation of organochlorine contaminants

- levels in Canadian Arctic beluga (*Delphinapterus leucas*). *Science of the Total Environment*, 351-352, 344-368. doi:10.1016/j.scitotenv.2004.10.033.
- Stockin, K. A., Binedell, V., Wiseman, N., Brunton, D. H., & Orams, M. B. (2009). Behavior of free-ranging common dolphins (*Delphinus sp.*) in the Hauraki Gulf, New Zealand. *Marine Mammal Science*, 25, 283-301. doi:10.1111/j.1748-7692.2008.00262.x
- Stockin, K. A., & Burgess, E. A. (2005). Opportunistic feeding of an adult humpback whale (*Megaptera novaeangliae*) migrating along the coast of southeastern Queensland, Australia. *Aquatic Mammals*, 31, 120-123.
- Stockin, K. A., Law, R., Duignan, P., Jones, G., Porter, L., Mirimin, L., et al. (2007). Trace elements, PCBs and organochlorine pesticides in New Zealand common dolphins (*Delphinus sp.*). *Science of the Total Environment*, 387, 333-345. doi:10.1016/j.scitotenv.2007.05.016
- Stockin, K. A., & Visser, I. N. (2005). Anomalously pigmented common dolphins (*Delphinus sp.*) off northern New Zealand. *Aquatic Mammals*, 31, 43-51.
- Stolen, M., Odell, D., & Barros, N. (2002). Growth of bottlenose dolphins (*Tursiops truncatus*) from the Indian River lagoon system, Florida, U.S.A. *Marine Mammal Science*, 18, 348-357.
- Storelli, M., & Marcotrigiano, G. (2003). Levels and congener pattern of polychlorinated biphenyls in the blubber of the Mediterranean bottlenose dolphins *Tursiops truncatus*. *Environment International*, 28, 559-566.
- Streever, B. (2002). Science and emotion, on ice: The role of science on Alaska's North slope. *Bioscience*, 52, 179.
- Stewart, J., Blakely, J., Karpowicz, P., Kalanxhi, E., Thatcher, B., & Martin, B. (2004). Unusually weak oxygen binding, physical properties, partial sequence, autoxidation rate and a potential phosphorylation site of beluga whale (*Delphinapterus leucas*) myoglobin. *Comparative Biochemistry & Physiology Part B*, 137, 401. doi:10.1016/j.cbpc.2004.01.007
- Stewart, R., Campana, S., Jones, C., & Stewart, B. (2006). Bomb radiocarbon dating calibrates beluga (*Delphinapterus leucas*) age estimates. *Canadian Journal of Zoology*, 84, 1840-1852. doi:10.1139/Z06-182
- Sudo, R., Uranishi, M., Kawaminami, T., Ihara, M., Iizuka, S., Ueda, M., et al. (2008). Sighting survey of cetaceans in the Tsugaru Strait, Japan. *Fisheries Science*, 74, 211-213. doi:10.1111/j.1444-2906.2007.01513.x
- Sukhoruchenko, M. (2004). Auditory detection of paired pulses by a dolphin in the presence of a pulse jam. *Acoustical Physics*, 50, 485-492. doi:10.1134/1.1776228
- Sukhoruchenko, M. (2008). Behavioral studies of the auditory discrimination of paired pulses with identical pulse spacings by a dolphin. *Acoustical Physics*, 54, 874-879. doi:10.1134/S1063771008060183
- Sullivan, S., & Mullin, K. (1997). Killer whales (*Orcinus orca*) in the Northern Gulf of Mexico. *Marine Mammal Science*, 13, 141-147.
- Summers, A. (2001). Spring-loaded. *Natural History*, 110(10), 76.
- Supin, A. (1994). Auditory mechanisms in dolphins. *Sensory Systems*, 8, 250-254.
- Supin, A., Nachtigall, P., Au, W., & Breese, M. (2004). The interaction of outgoing echolocation pulses and echoes in the false killer whale's auditory system: Evoked-potential study. *Journal of the Acoustical Society of America*, 115, 3218-3225. doi:10.1121/1.1707088
- Supin, A., Nachtigall, P., Au, W., & Breese, M. (2005). Invariance of evoked-potential echo-responses to target strength and distance in an echolocating false killer

- whale. *Journal of the Acoustical Society of America*, 117, 3928-3935. doi:10.1121/1.1914150
- Supin, A., Nachtigall, P., & Breese, M. (2006). Source-to-sensation level ratio of transmitted biosonar pulses in an echolocating false killer whale. *Journal of the Acoustical Society of America*, 120, 518-526. doi:10.1121/1.2202862
- Supin, A., Nachtigall, P., & Breese, M. (2007). Evoked-potential recovery during double click stimulation in a whale: A possibility of biosonar automatic gain control. *Journal of the Acoustical Society of America*, 121, 618-625. doi:10.1121/1.2382343
- Supin, A., Nachtigall, P., & Breese, M. (2008). Hearing sensitivity during target presence and absence while a whale echolocates. *Journal of the Acoustical Society of America*, 123, 534-541. doi:10.1121/1.2812593
- Supin, A., Nachtigall, P., & Breese, M. (2009). Forward-masking based gain control in odontocete biosonar: An evoked-potential study. *Journal of the Acoustical Society of America*, 125, 2432-2442. doi:10.1121/1.3089589
- Supin, A., Nachtigall, P., Pawloski, J., & Au, W. (2003). Evoked potential recording during echolocation in a false killer whale *Pseudorca crassidens*. *Journal of the Acoustical Society of America*, 113, 2408-2411. doi:10.1121/1.1561497
- Supin, A. Y., & Popov, V. V. (2007). Improved techniques of evoked-potential audiometry in odontocetes. *Aquatic Mammals*, 33, 14-23.
- Sutaria, D., & Jefferson, T. A. (2004). Records of Indo-Pacific humpback dolphins (*Sousa chinensis*, Osbeck, 1765) along the coasts of India and Sri Lanka: An overview. *Aquatic Mammals*, 30, 125-136.
- Suttle, C. (2007). Marine viruses — major players in the global ecosystem. *Nature Reviews Microbiology*, 5, 801-812. doi:10.1038/nrmicro1750
- Suydam, R., Lowry, L., Frost, K., O'Corry-Crowe, G., & Pikok, J. (2001). Satellite tracking of Eastern Chukchi Sea beluga whales into the Arctic Ocean. *Arctic*, 54, 237-243.
- Suzuki, M., Endo, N., Nakano, Y., Kato, H., Kishiro, T., & Asahina, K. (2008). Localization of aquaporin-2, renal morphology and urine composition in the bottlenose dolphin and the Baird's beaked whale. *Journal of Comparative Physiology B: Biochemical, Systemic, & Environmental Physiology*, 178, 149-156. doi:10.1007/s00360-007-0204-1
- Suzuki, M., Kashiwabara, T., Koie, H., Suzuki, C., Hirako, K., & Saito, S. (2008). Usage of high-performance mattresses for transport of Indo-Pacific bottlenose dolphin. *Zoo Biology*, 27, 331-340.
- Suzuki, M., Uchida, S., Ueda, K., Tobayama, T., Katsumata, E., Yoshioka, M., et al. (2003). Diurnal and annual changes in serum cortisol concentrations in Indo-Pacific bottlenose dolphins *Tursiops aduncus* and killer whales *Orcinus orca*. *General & Comparative Endocrinology*, 132, 427-433. doi:10.1016/S0016-6480(03)00100-X
- Svane, I. (2005). Occurrence of dolphins and seabirds and their consumption of by-catch during prawn trawling in Spencer Gulf, South Australia. *Fisheries Research*, 76, 317-327. doi:10.1016/j.fishres.2005.07.012
- Szymanski, M., Bain, D., Kiehl, K., Pennington, S., Wong, S., & Henry, K. (1999). Killer whale (*Orcinus orca*) hearing: Auditory brainstem response and behavioral audiograms. *Journal of the Acoustical Society of America*, 106, 1134-1141. doi:10.1121/1.427121
- Szymanski, M., Supin, A., Bain, D., & Henry, K. (1998). Killer whale (*Orcinus orca*) auditory evoked potentials to rhythmic clicks. *Marine Mammal Science*, 14, 676-691.

- Takahashi, Y., Ohwada, S., Watanabe, K., Ropert-Coudert, Y., Zenitani, R., Naito, Y., et al. (2006). Does elastin contribute to the persistence of corpora albicantia in the ovary of the common dolphin (*Delphinus delphis*). *Marine Mammal Science*, 22, 819-830.
- Tamaki, N., Morisaka, T., & Taki, M. (2006). Does body contact contribute towards repairing relationships? The association between flipper-rubbing and aggressive behavior in captive bottlenose dolphins. *Behavioural Processes*, 73, 209-215. doi:10.1016/j.beproc.2006.05.010
- Tarakanov, M., Pletenko, M., Popov, V., & Supin, A. (2005). Audiogram variability in normal bottlenose dolphins. *Journal of the Acoustical Society of America*, 117, 2441.
- Tarakanov, M., Pletenko, M., & Supin, A. (1998). Spectral-temporal sound signal pattern discrimination by the auditory system in dolphins. *Sensory Systems*, 12, 377-385.
- Taubenberger, J., & Tsai, M. (1996). Two morbilliviruses implicated in bottlenose dolphin epizootics. *Emerging Infectious Diseases*, 2, 213-216.
- Tayler, C., & Saayman, G. (1973). Imitative behaviour by Indian Ocean bottlenose dolphins (*Tursiops aduncus*) in captivity. *Behaviour*, 44, 286-298. doi:10.1163/156853973X00436
- Taylor, B., Brotheridge, R., Jessup, D., & Stott, J. (2002). Measurement of serum immunoglobulin concentration in killer whales and sea otters by radial immunodiffusion. *Veterinary Immunology and Immunopathology*, 89, 187-196.
- Taylor, G., Nudds, R., & Thomas, A. (2003). Flying and swimming animals cruise at a Strouhal number tuned for high power efficiency. *Nature*, 425, 707-711. doi:10.1038/nature02000
- Taylor, K. A., Nachtigall, P. E., Mooney, T. A., Supin, A. Y., & Yuen, M. M. L. (2007). A portable system for the evaluation of the auditory capabilities of marine mammals. *Aquatic Mammals*, 33, 93-99.
- Taylor, M. (1986). Stunning whales and deaf squids. *Nature*, 323, 298-299. doi:10.1038/323298a0
- Teilmann, J., Miller, L. A., Kirketerp, T., Kastelein, R. A., Madsen, P. T., Nielsen, B. K., et al. (2002). Characteristics of echolocation signals used by a harbour porpoise (*Phocoena phocoena*) in a target detection experiment. *Aquatic Mammals*, 28, 275-284.
- Ternes, A. (1986). Picnic à la dauphine. *Natural History*, 95(4), 70.
- Tezanos-Pinto, G., Scott Baker, C., Russell, K., Martien, K., Baird, R., Hutt, A., et al. (2009). A worldwide perspective on the population structure and genetic diversity of bottlenose dolphins (*Tursiops truncatus*) in New Zealand. *Journal of Heredity*, 100, 11-24. doi:10.1093/jhered/esn039
- Thaheld, F. (2004). A method to explore the possibility of nonlocal correlations between brain electrical activities of two spatially separated animal subjects. *Biosystems*, 73, 205-216. doi:10.1016/j.biosystems.2004.01.001
- Thayer, V., Read, A., Friedlaender, A., Colby, D., Hohn, A., McLellan, W., et al. (2003). Reproductive seasonality of western Atlantic bottlenose dolphins off North Carolina, U.S.A. *Marine Mammal Science*, 19, 617-629.
- Thomas, C. (2009). Sonar doesn't appear to deafen dolphins. *ScienceNOW*, April 8, 2009. Retrieved from <http://news.sciencemag.org/sciencenow/2009/04/08-04.html>.
- Thomas, J., Kastelein, R., & Awbrey, F. (1990). Behavior and blood catecholamines of captive belugas during playbacks of noise from an oil drilling platform. *Zoo Biology*, 9, 393- 402. doi:10.1002/zoo.1430090507

- Thomas, J. A., & Kastelein, R. (1990). *Sensory abilities of cetaceans*. New York: Plenum.**
- Thomas, R., Fristrup, K., & Tyack, P. (2002). Linking the sounds of dolphins to their locations and behavior using video and multichannel acoustic recordings. *Journal of the Acoustical Society of America*, 112, 1692-1701. doi:10.1121/1.1494805
- Thompson, R., & Herman, L. (1977). Memory for lists of sounds by the bottle-nosed dolphin: Convergence of memory processes with humans? *Science*, 195, 501-503. doi:10.1126/science.835012
- Thompson, R., & Herman, L. (1981). Auditory delayed discriminations by the dolphin: Nonequivalence with delayed-matching performance. *Animal Learning & Behavior*, 9, 9-15.
- Thomsen, F., Franck, D., & Ford, J. (2001). Characteristics of whistles from the acoustic repertoire of resident killer whales (*Orcinus orca*) off Vancouver Island, British Columbia. *Journal of the Acoustical Society of America*, 109, 1240-1246. doi:10.1121/1.1349537
- Thornton, A. (1994). Environmental threats to whales. *AWI Quarterly*, 43(1), 7.
- Tibbs, R., Elghetany, M., Tran, L., van Bonn, W., Romano, T., & Cowan, D. (2005). Characterization of the coagulation system in healthy dolphins: The coagulation factors, natural anticoagulants, and fibrinolytic products. *Comparative Clinical Pathology*, 14, 95-98. doi:10.1007/s00580-005-0567-1
- Timberlake, W. (1993) Animal behavior: a continuing synthesis. *Annual Review of Psychology*, 44, 675-706.
- Tolley, K., Read, A., Wells, R. S., Urian, K. W., Scott, M. D., Irvine, A. B., et al. (1995). Sexual dimorphism in wild bottlenose dolphins (*Tursiops truncatus*) from Sarasota, Florida. *Journal of Mammalogy*, 76, 1190-1198.
- Tomonari, A., Ding, W., Kexiong, W., & Yasuhiko, N. (2005). Biosonar behaviour of free ranging porpoises. *Proceedings of the Royal Society B: Biological Sciences*, 272, 797-801.
- Tomy, G., & Muir, D. (2000). Levels of C10-C13 Polychloro-n-Alkanes in marine mammals from the Arctic and the St. Lawrence. *Environmental Science & Technology*, 34, 1615-1620.
- Tornero, V., Borrell, A., Aguilar, A., Forcada, J., & Lockyer, C. (2006). Organochlorine contaminant and retinoid levels in blubber of common dolphins (*Delphinus delphis*) off northwestern Spain. *Environmental Pollution*, 140, 312-321.
- Tornero, V., Borrell, A., Forcada, J., Pubill, E., & Aguilar, A. (2004). Retinoid and lipid patterns in the blubber of common dolphins (*Delphinus delphis*): Implications for monitoring vitamin A status. *Comparative Biochemistry & Physiology Part B*, 137, 391. doi:10.1016/j.cbpc.2004.01.001
- Tosh, C. A., de Bruyn, P., & Bester, M. N. (2008). Preliminary analysis of the social structure of killer whales, *Orcinus orca*, at subantarctic Marion Island. *Marine Mammal Science*, 24, 929-940. doi:10.1111/j.1748-7692.2008.00233.x
- Tremel, D. P., Thomas, J. A., Ramirez, K. T., Dye, G. S., Bachman, W. A., Orban, A. N., et al. (1998). Underwater hearing sensitivity of a Pacific white-sided dolphin, *Lagenorhynchus albirostris obliquidens*. *Aquatic Mammals*, 24, 63-69.
- Triossi, F., Pace, D. S., Terranova, M. L., & Renzi, P. (1998). The development of suckling behavior in two captive-born calves of bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 24, 75-83.
- Trites, A., Deecke, V., Gregr, E., Ford, J., & Olesiuk, P. (2007). Killer whales, whaling, and sequential megafaunal collapse in the North Pacific: A comparative analysis

- of the dynamics of marine mammals in Alaska and British Columbia following commercial whaling. *Marine Mammal Science*, 23, 751-765.
- Tromans, A. (2003). Echolocation: Volume control. *Nature*, 423, 815.
- Trone, M., Kuczaj, S., & Solangi, M. (2005). Does participation in dolphin-human interaction programs affect bottlenose dolphin behaviour? *Applied Animal Behaviour Science*, 93, 363-374. doi:10.1016/j.applanim.2005.01.003
- Tryland, M., Kleivane, L., Alfredsson, A., Kjeld, M., Arnason, A., Stuen, S., et al. (1999). Evidence of Brucella infection in marine mammals in the North Atlantic Ocean. *Veterinary Record: Journal of the British Veterinary Association*, 144, 588-592.
- Tryland, M., Thoresen, S., Kovacs, K., & Lydersen, C. (2006). Serum chemistry of free-ranging white whales (*Delphinapterus leucas*) in Svalbard. *Veterinary Clinical Pathology*, 35, 199-203.
- Tschudin, A. (2001). 'Mindreading' mammals? Attribution of belief tasks with dolphins. *Animal Welfare*, 10, S119-S127.
- Tschudin, A., Call, J., Dunbar, R., Harris, G., & van der Elst, C. (2001). Comprehension of signs by dolphins (*Tursiops truncatus*). *Journal of Comparative Psychology*, 115, 100-105. doi:10.1037/0735-7036.115.1.100
- Tsikliras, A., Koutrakis, E., Kallianiotis, A., & Economidis, P. (2004). Massive leaping of Sardinella Aurita Valenciennes, 1847 under probable risk of predation in northern Aegean sea (Greece). *Marine and Freshwater Behaviour and Physiology*, 37, 31-34. doi:10.1080/1023624042000199971
- Tsur, I., Yakobson, B., Elad, D., Moffett, D., & Kennedy, S. (1997). Morbillivirus infection in a bottlenose dolphin from the Mediterranean Sea. *European Journal of Veterinary Pathology: Official Journal of the European Society of Veterinary Pathology*, 3, 83-85.
- Tsutsui, S., Tanaka, M., Miyazaki, N., & Furuya, T. (2001). Pacific white-sided dolphins (*Lagenorhynchus albirostris obliquidens*) with anomalous colour patterns in Volcano Bay, Hokkaido, Japan. *Aquatic Mammals*, 27, 172-182.
- Tuerk, K., Kucklick, J., Becker, P., Stapleton, H., & Baker, J. (2005). Persistent organic pollutants in two dolphin species with focus on toxaphene and polybrominated. *Environmental Science & Technology*, 39, 692-698.
- Turl, C. (1993). Low-frequency sound detection by a bottlenose dolphin. *Journal of the Acoustical Society of America*, 94, 3006-3008. doi:10.1121/1.407333
- Turl, C., Penner, R., & Au, W. (1987). Comparison of target detection capabilities of the beluga and bottlenose dolphin. *Journal of the Acoustical Society of America*, 82, 1487-1491. doi:10.1121/1.395192
- Turner, J., Clark, L. S., Haubold, E. A., Worthy, G. A. J., & Cowan, D. F. (2006). Organ weights and growth profiles in bottlenose dolphins (*Tursiops truncatus*) from the northwestern Gulf of Mexico. *Aquatic Mammals*, 32, 46-57.
- Turner, J., & Worthy, G. (2003). Skull morphometry of bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Mexico. *Journal of Mammalogy*, 84, 665-673.
- Turvey, S. (2008). Witness to extinction: How we failed to save the Yangtze river dolphin. Oxford, UK: Oxford University Press.**
- Tyack, P. (1981). Why do whales sing? *The Sciences*, 2, 22-25.
- Tyack, P. (1986). Whistle repertoires of two bottlenosed dolphins, *Tursiops truncatus*: Mimicry of signature whistles? *Behavioral Ecology and Sociobiology*, 18, 251-257. doi:10.1007/BF00300001
- Tyack, P. (1991). If you need me, whistle. *Natural History*, 100(8), 60-61.
- Tyack, P. (1997). Development and social functions of signature whistles in bottlenose dolphins *Tursiops truncatus*. *Bioacoustics*, 8, 21-46.

- Tyack, P. (2000). Dolphins whistle a signature tune. *Science*, 289, 1310.
- Tyack, P. (2001). Cetacean culture: Humans of the sea? *Behavioral & Brain Sciences*, 24, 358.
- Tyack, P. (2008). Convergence of calls as animals form social bonds, active compensation for noisy communication channels, and the evolution of vocal learning in mammals. *Journal of Comparative Psychology*, 122, 319-331. doi:10.1037/a0013087
- Tyack, P., & Recchia, C. (1991). A datalogger to identify vocalizing dolphins. *Journal of the Acoustical Society of America*, 90, 1668-1671. doi:10.1121/1.401908
- Tynan, C., Ainley, D., Barth, J., Cowles, T., Pierce, S., & Spear, L. (2005). Cetacean distributions relative to ocean processes in the northern California Current System. *Deep-Sea Research Part II, Topical Studies in Oceanography*, 52, 145-167. doi:10.1016/j.dsr2.2004.09.024
- Tyson, R., Nowacek, D., & Miller, P. (2007). Nonlinear phenomena in the vocalizations of North Atlantic right whales (*Eubalaena glacialis*) and killer whales (*Orcinus orca*). *Journal of the Acoustical Society of America*, 122, 1365-1373. doi:10.1121/1.2756263
- Uemura, Y., Asakuma, S., Nakamura, T., Arai, I., Taki, M., & Urashima, T. (2005). Occurrence of a unique sialyl tetrasaccharide in colostrum of a bottlenose dolphin (*Tursiops truncatus*). *BBA - General Subjects*, 1725, 290-297. doi:10.1016/j.bbagen.2005.05.011
- Ulrich, Ridgway, & Ulrich. (1999). Enrofloxacin serum bioactivity in bottlenose dolphins, *Tursiops truncatus*, following oral administration of 5 mg/kg in whole fish. *Journal of Veterinary Pharmacology and Therapeutics*, 22, 170-173. doi:10.1046/j.1365-2885.1999.00199.x
- Urian, K., & Duffield, D. (1996). Seasonality of reproduction in bottlenose dolphins, *Tursiops truncatus*. *Journal of Mammalogy*, 77, 394.
- Urian, K., Hofmann, S., Wells, R. S., & Read, A. J. (2009). Fine-scale population structure of bottlenose dolphins (*Tursiops truncatus*) in Tampa Bay, Florida. *Marine Mammal Science*, 25, 619-638. doi:10.1111/j.1748-7692.2009.00284.x
- Valsecchi, E., & Amos, W. (1996). Microsatellite markers for the study of cetacean populations. *Molecular Ecology*, 5, 151-156.
- Valsecchi, E., Amos, W., Raga, J. A., Podesta M., Sherwin, W. (2004). The effects of inbreeding on mortality during a morbillivirus outbreak in the Mediterranean striped dolphin (*Stenella coeruleoalba*). *Animal Conservation*, 7, 139-146.
- van Bergen, Y. (2005). Chubby calves combat the cold. *Journal of Experimental Biology*, 208, 3.
- Van Opzeeland, I. C., Corkeron, P. J., Leyssen, T., Simila, T., & Van Parijs, S. M. (2005). Acoustic behaviour of Norwegian killer whales, *Orcinus orca*, during carousel and seiner foraging on spring-spawning herring. *Aquatic Mammals*, 31, 110-119.
- Van Parijs, S., & Corkeron, P. (2001). Vocalizations and behaviour of Pacific Humpback dolphins, *Sousa chinensis*. *Ethology*, 107, 701-716. doi:10.1046/j.1439-0310.2001.00714.x
- Van Parijs, S., Leyssen, T., & Similä, T. (2004). Sounds produced by Norwegian killer whales, *Orcinus orca*, during capture. *Journal of the Acoustical Society of America*, 116, 557-560. doi:10.1121/1.1763954
- Van Parijs, S., Lydersen, C., & Kovacs, K. (2003). Sounds produced by individual white whales, *Delphinapterus leucas*, from Svalbard during capture (L). *Journal of the Acoustical Society of America*, 113, 57-60. doi:10.1121/1.1528931

- Van Parijs, S., Smith, J., & Corkeron, P. (2002). Using calls to estimate the abundance of inshore dolphins: A case study with Pacific humpback dolphins, *Sousa chinensis*. *Journal of Applied Ecology*, 39, 853-864.
- van Tienderen, P. H. (1995). Life cycle trade-offs in matric population models. *Ecology*, 76, 2482-2489.
- van Waerebeek, K., & Read, A. (1994). Reproduction of dusky dolphins, *Lagenorhynchus obscurus*, from coastal Peru. *Journal of Mammalogy*, 75, 1054-1062.
- van Waerebeek, K., Reyes, J. C., & Alfaro, J. (1993). Helminth parasites and phoronts of dusky dolphins *Lagenorhynchus albirostris obscurus* (Gray, 1828) from Peru. *Aquatic Mammals*, 19, 159-169.
- Varela, R., Schwacke, L., Fair, P., & Bossart, G. (2006). Effects of duration of capture and sample handling on critical care blood analytes in free-ranging bottlenose dolphins. *Journal of the American Veterinary Medical Association*, 229, 1955-1961.
- Vaughan, K., Del Crew, J., Hermanson, G., Wloch, M., Riffenburgh, R., Smith, C., et al. (2007). A DNA vaccine against dolphin morbillivirus is immunogenic in bottlenose dolphins. *Veterinary Immunology & Immunopathology*, 120, 260-266. doi:10.1016/j.vetimm.2007.06.036
- Vaughn, R., Shelton, D., Timm, L., Watson, L., & Würsig, B. (2007). Dusky dolphin (*Lagenorhynchus albirostris obscurus*) feeding tactics and multi-species associations. *New Zealand Journal of Marine & Freshwater Research*, 41, 391-400.
- Vaughn, R., Würsig, B., Shelton, D., Timm, L., & Watson, L. (2008). Dusky dolphins influence prey accessibility for seabirds in Admiralty Bay, New Zealand. *Journal of Mammalogy*, 89, 1051-1058.
- Vecchione, A., Peden-Adams, M. M., Romano, T. A., & Fair, P. A. (2008). Recent cytokine findings and implications toward health assessment of the bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 34, 93-101.
- Venn-Watson, S., Jensen, E., & Ridgway, S. (2007). Effects of age and sex on clinicopathologic reference ranges in a healthy managed Atlantic bottlenose dolphin population. *Journal of the American Veterinary Medical Association*, 231, 596-601.
- Venn-Watson, S., Jensen, E., & Smith, C. (2008). Assessment of increased serum aminotransferases in a managed Atlantic bottlenose dolphin (*Tursiops truncatus*) population. *Journal of Wildlife Diseases*, 44, 318-330.
- Venn-Watson, S., & Ridgway, S. (2007). Big brains and blood glucose: Common ground for diabetes mellitus in humans and healthy dolphins. *Comparative Medicine*, 57, 390-395.
- Venn-Watson, S., Rivera, R., Smith, C., Saliki, J., Caseltine, S., Leger, J., et al. (2008). Exposure to novel parainfluenza virus and clinical relevance in 2 bottlenose dolphin (*Tursiops truncatus*) populations. *Emerging Infectious Diseases*, 14, 397-405.
- Verfuß, U., Miller, L., & Schnitzler, H. (2005). Spatial orientation in echolocating harbor porpoises (*Phocoena phocoena*). *Journal of Experimental Biology*, 208, 3385-3394. doi:10.1242/jeb.01786
- Vermeulen, E., & Cammareri, A. (2009). Residency patterns, abundance, and social composition of bottlenose dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina. *Aquatic Mammals*, 35, 378-385. doi:10.1578/AM.35.3.2009.378.

- Vetter, W., Stoll, E., Garson, M., Fahey, S., Gaus, C., & Müller, J. (2002). Spong halogenated natural products found at parts-per-million levels in marine mammals. *Environmental Toxicology & Chemistry*, 21, 2014-2019. doi:10.1897/1551-5028(2002)021<2014:SHNPFA>2.0.CO;2
- Viaud-Martinez, K., Brownell, R., Komnenou, A., & Bohonak, A. (2008). Genetic isolation and morphological divergence of Black Sea bottlenose dolphins. *Biological Conservation*, 141, 1600-1611. doi:10.1016/j.biocon.2008.04.004
- Viddi, F. A., & Lescrauwaet, A.-K. (2005). Insights on habitat selection and behavioural patterns of Peale's dolphins (*Lagenorhynchus albirostris australis*) in the Strait of Magellan, southern Chile. *Aquatic Mammals*, 31, 176-183.
- Villano, M. (2007). Eavesdropping on whales' mealtime conversation. *Oceanus*, 45(3), 20-25.
- Viricel, A., Strand, A., Rosel, P., Ridoux, V., & Garcia, P. (2008). Insights on common dolphin (*Delphinus delphis*) social organization from genetic analysis of a mass-stranded pod. *Behavioral Ecology and Sociobiology*, 63, 173-185. doi:10.1007/s00265-008-0648-7
- Visser, I. N. (1998). Prolific body scars and collapsing dorsal fins on killer whales (*Orcinus orca*) in New Zealand waters. *Aquatic Mammals*, 24, 71-81.
- Visser, I. N. (2000). Killer whale (*Orcinus orca*) interactions with longline fisheries in New Zealand waters. *Aquatic Mammals*, 26, 241-252.
- Visser, I. N. (2005). First observations of feeding on thresher (*Alopias vulpinus*) and hammerhead (*Sphyrna zygaena*) sharks by killer whales (*Orcinus orca*) specialising on Elasmobranch prey. *Aquatic Mammals*, 31, 83-88.
- Visser, I. N., Berghan, J., van Meurs, R., & Fertl, D. (2000). Killer whale (*Orcinus orca*) predation on a shortfin mako shark (*Isurus oxyrinchus*) in New Zealand waters. *Aquatic Mammals*, 26, 229-231.
- Visser, I. N., & Fertl, D. (2000). Stranding, resighting, and boat strike of a killer whale (*Orcinus orca*) off New Zealand. *Aquatic Mammals*, 26, 232-240.
- Visser, I., & Maekelaeinen, P. (2000). Variation in eye-patch shape of killer whales (*Orcinus orca*) in New Zealand Waters. *Marine Mammal Science*, 16, 459-469.
- Vogel, G. (1998). DNA suggests cultural traits affect whales' evolution. *Science*, 282, 1616.
- von Fersen, L., & Delius, J. (2000). Acquired equivalences between auditory stimuli in dolphins (*Tursiops truncatus*). *Animal Cognition*, 3, 79-83. doi:10.1007/s100710000063
- von Fersen, L., Schall, U., & Güntürkün, O. (2000). Visual lateralization of pattern discrimination in the bottlenose dolphin (*Tursiops truncatus*). *Behavioural Brain Research*, 107, 177-181. doi:10.1016/S0166-4328(99)00142-4
- Voronin, L., & Kozarovitskii, L. (1969). Some features of higher nervous activity in the Black Sea dolphin of the aphelin species (*Tursiops truncatus*). *Zhurnal Vysshei Nervnoi Deyatel'nosti*, 19(1), 233-242.
- Wafo, E., Sarrazin, L., Diana, C., Dhermain, F., Schembri, T., Lagadec, V., et al. (2005). Accumulation and distribution of organochlorines (PCBs and DDTs) in various organs of *Stenella coeruleoalba* and a *Tursiops truncatus* from Mediterranean littoral environment (France). *Science of the Total Environment*, 348, 115-127. doi:10.1016/j.scitotenv.2004.12.078
- Wahlberg, M., Bejder, L., & Madsen, P. (2008). Noise levels and masking potential of small whale-watching and research vessels around two delphinid species. *Bioacoustics*, 17, 166-168.

- Waite, J., Friday, N., & Moore, S. (2002). Killer whale (*Orcinus orca*) distribution and abundance in the central and southeastern Bering Sea, July 1999 and June 2000. *Marine Mammal Science*, 18, 779-786.
- Wang, D. (2009). Population status, threats and conservation of the Yangtze finless porpoise. *Chinese Science Bulletin*, 54, 3473-3484. doi:10.1007/s11434-009-0522-7
- Wang, J., & Chou, L. (2000). Differences in the external morphology of two sympatric species of bottlenose dolphins (genus *Tursiops*) in the water of China. *Journal of Mammalogy*, 81, 1157-1166.
- Wang, J., Chou, L., & White, B. (1999). Mitochondrial DNA analysis of sympatric morphotypes of bottlenose dolphins (genus: *Tursiops*) in Chinese waters. *Molecular Ecology*, 8, 1603-1612. doi:10.1046/j.1365-294X.1999.00741.x
- Wang, J., Hung, S. K., & Yang, S.-C. (2004). Records of Indo-Pacific humpback dolphins, *Sousa chinensis* (Osbeck, 1765), from the waters of western Taiwan. *Aquatic Mammals*, 30, 189-196.
- Wang, J., Frasier, T., Yang, S., & White, B. (2008). Detecting recent speciation events: the case of the finless porpoise (genus *Neophocaena*). *Heredity*, 101, 145-155. doi:10.1038/hdy.2008.40
- Waples, K., & Gales, N. (2002). Evaluating and minimising social stress in the care of captive bottlenose dolphins (*Tursiops aduncus*). *Zoo Biology*, 21, 5-26. doi:10.1002/zoo.10004
- Ward, E., Holmes, E., & Balcomb, K. (2009). Quantifying the effects of prey abundance on killer whale reproduction. *Journal of Applied Ecology*, 46, 632-640. doi:10.1111/j.1365-2664.2009.01647.x
- Ward, E., Parsons, K., Holmes, E., Balcomb, I., & Ford, J. (2009). The role of menopause and reproductive senescence in a long-lived social mammal. *Frontiers in Zoology*, 61, 10. doi:10.1186/1742-9994-6-4
- Warren, J. M. (1965). The comparative psychology of learning. *Annual Review of Psychology*, 16, 95-118.
- Warren-Sinith, A. B., & Dunn, W. L. (2006). Epimeletic behaviour toward a seriously injured juvenile bottlenose dolphin (*Tursiops* sp.) in Port Phillip, Victoria, Australia. *Aquatic Mammals*, 32, 357-362.
- Watanabe, N., Hatano, J., Asahina, K., Iwasaki, T., & Hayakawa, S. (2007). Molecular cloning and histological localization of LH-like substances in a bottlenose dolphin (*Tursiops truncatus*) placenta. *Comparative Biochemistry & Physiology Part A: Molecular & Integrative Physiology*, 146, 105-118. doi:10.1016/j.cbpa.2006.09.011
- Watson, A., Bahr, R. J., & Alexander, J. W. (2004). Thoracolumbar kyphoscoliosis and compression fracture of a thoracic vertebra in a captive bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 30, 275-278.
- Watson, A., Bahr, R. J., & Matheson, T. (2008). Metacarpo-phalangeal anomalies in bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals*, 34, 65-70.
- Watson, A., & Gee, L. E. (2005). Laryngeal displacement and asphyxiation by a beheaded sheepshead (*Archosargus probatocephalus*) in a bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 31, 447-452.
- Watson-Capps, J., & Mann, J. (2005). The effects of aquaculture on bottlenose dolphin (*Tursiops* sp.) ranging in Shark Bay, Western Australia. *Biological Conservation*, 124, 519-526. doi:10.1016/j.biocon.2005.03.001
- Watts, P., & Draper, B. (1988). Beluga whale ecology: A concern of freshwater biologists? *Freshwater Biology*, 20, 211-213.

- Watwood, S., Owen, E., Tyack, P., & Wells, R. (2005). Signature whistle use by temporarily restrained and free-swimming bottlenose dolphins, *Tursiops truncatus*. *Animal Behaviour*, 69, 1373-1386. doi:10.1016/j.anbehav.2004.08.019
- Watwood, S., Tyack, P., & Wells, R. (2004). Whistle sharing in paired male bottlenose dolphins, *Tursiops truncatus*. *Behavioral Ecology and Sociobiology*, 55, 531-543. doi:10.1007/s00265-003-0724-y
- Wazura, K., Strong, J., Glenn, C., & Bush, A. (1986). Helminths of the beluga whale (*Delphinapterus leucas*) from the Mackenzie River Delta, Northwest Territories. *Journal of Wildlife Diseases*, 22, 440-442.
- Weaver, A. (2003). Conflict and reconciliation in captive bottlenose dolphins, *tursiops truncatus*. *Marine Mammal Science*, 19, 836-846.
- Webb, N., & Drummond, P. (2001). The effect of swimming with dolphins on human well-being and anxiety. *Anthrozoös*, 14, 81-85.
- Weber, P., Howle, L., Murray, M., & Fish, F. (2009). Lift and drag performance of odontocete cetacean flippers. *Journal of Experimental Biology*, 212, 2149-2158. doi:10.1242/jeb.029868
- Webster, T., Dawson, S., & Slooten, E. (2009). Evidence of sex segregation in Hector's dolphin (*Cephalorhynchus hectori*). *Aquatic Mammals*, 35, 212-219. doi:10.1578/AM.35.2.2009.212
- Wedekin, L. L., Freitas, A., Engel, M. H., & Sazima, I. (2004). Rough-toothed dolphins (*Steno bredanensis*) catch diskfishes while interacting with humpback whales (*Megaptera novaeangliae*) off Abrolhos Bank breeding ground, Southwest Atlantic. *Aquatic Mammals*, 30, 327-329.
- Wehle, D., & Coleman, F. (1983). Plastics at sea. *Natural History*, 92(2), 20.
- Weihs, D. (2002). Dynamics of dolphin porpoising revisited. *Integrative & Comparative Biology*, 42, 1071-1078.
- Weihs, D., Fish, F. E., & Nicastro, A. J. (2007). Mechanics of remora removal by dolphin spinning. *Marine Mammal Science*, 23, 707-714. doi:10.1111/j.1748-7692.2007.00131.x
- Weilgart, L. (2007). A brief review of known effects of noise on marine mammals. *International Journal of Comparative Psychology*, 20, 159-168.
- Weinrich, M., Belt, C., & Morin, D. (2001). Behavior and ecology of the Atlantic white-sided dolphin (*Lagenorhynchus acutus*) in coastal New England waters. *Marine Mammal Science*, 17, 231-248.
- Weir, C. (2006). First confirmed records of Clymene dolphin, *Stenella clymene* (Gray, 1850), from Angola and Congo, South-East Atlantic Ocean. *African Zoology*, 41, 297-300.
- Weir, C. (2008a). Overt responses of humpback whales (*Megaptera novaeangliae*), sperm whales (*Physeter macrocephalus*), and Atlantic spotted dolphins (*Stenella frontalis*) to seismic exploration off Angola. *Aquatic Mammals*, 34, 71-83.
- Weir, C. (2008b). Short-finned pilot whales (*Globicephala macrorhynchus*) respond to an airgun ramp-up procedure off Gabon. *Aquatic Mammals*, 34, 349-354.
- Weir, C. (2009). Distribution, behaviour and photo-identification of Atlantic humpback dolphins *Sousa teuszii* off Flamingos, Angola. *African Journal of Marine Science*, 31, 319-331. doi:10.2989/AJMS.2009.31.3.5.993
- Weir, C., Debrah, J., Ofori-Danson, P., Pierpoint, C., & Van Waerebeek, K. (2008). Records of Fraser's dolphin, *Lagenodelphis hosei fraser* 1956 from the Gulf of Guinea and Angola. *African Journal of Marine Science*, 30, 241-246. doi:10.2989/AJMS.2008.30.2.4.554

- Weir, J., Duprey, N., & Würsig, B. (2008). Dusky dolphin (*Lagenorhynchus albirostris obscurus*) subgroup distribution: Are shallow waters a refuge for nursery groups? *Canadian Journal of Zoology*, 86, 1225-1234. doi:10.1139/Z08-101
- Weiβa, B., Ladich, F., Spong, P., & Symonds, H. (2006). Vocal behavior of resident killer whale matrilines with newborn calves: The role of family signatures. *Journal of the Acoustical Society of America*, 119, 627-635. doi:10.1121/1.2130934
- Weiba, B., Symonds, H., Spong, P., & Ladich, F. (2007). Intra- and intergroup vocal behavior in resident killer whales, *Orcinus orca*. *Journal of the Acoustical Society of America*, 122, 3710-3716. doi:10.1121/1.2799907
- Weiss, J. (2006). Foraging habitats and associated preferential foraging specializations of bottlenose dolphin (*Tursiops truncatus*) mother-calf pairs. *Aquatic Mammals*, 32, 10-19.
- Weller, D. W., Cockcroft, V. G., Wuersig, B., Lynn, S. K., & Fertl, D. (1997). Behavioral responses of bottlenose dolphins to remote biopsy sampling and observations of surgical biopsy wound healing. *Aquatic Mammals*, 23, 49-58.
- Wells, R. (1991). Bringing up baby. *Natural History*, 100(8), 56.
- Wells, R. (2009). Learning from nature: Bottlenose dolphin care and husbandry. *Zoo biology*, 28, 635-651.**
- Wells, R., Bassos-Hull, K., & Norris, K. (1998). Experimental return to the wild of two bottlenose dolphins. *Marine Mammal Science*, 14, 51-71.
- Wells, R., Manire, C. A., Byrd, L., Smith, D. R., Gannon, J. G., Fauquier, D., & Mullin, K. D. (2009). Movements and dive patterns of a rehabilitated Risso's dolphin, *Grampus griseus*, in the Gulf of Mexico and Atlantic Ocean. *Marine Mammal Science*, 25, 420-429. doi:10.1111/j.1748-7692.2008.00251.x
- Wells, R., Rhinehart, H., Cunningham, P., Whaley, J., Baran, M., Koberna, C., & Costa, D. (1999). Long distance offshore movements of bottlenose dolphins. *Marine Mammal Science*, 15, 1098-1114.
- Wells, R., Sutton, G., DeLynn, R., Scott, M., Socha, V., Barros, N., et al. (2008). Consequences of injuries on survival and reproduction of common bottlenose dolphins (*Tursiops truncatus*) along the west coast of Florida. *Marine Mammal Science*, 24, 774-794.
- Wells, R., Sweeney, J., Townsend, F., Balmer, B., Hohn, A., et al. (2007). Annual, seasonal and individual variation in hematology and clinical blood chemistry profiles in bottlenose dolphins (*Tursiops truncatus*) from Sarasota Bay, Florida. *Comparative Biochemistry & Physiology Part A: Molecular & Integrative Physiology*, 148, 266-277. doi:10.1016/j.cbpa.2007.04.017
- Wells, R., Tornero, V., Borrell, A., Aguilar, A., Rowles, T., Rhinehart, H., et al. (2005). Integrating life-history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Science of the Total Environment*, 349, 106-119. doi:10.1016/j.scitotenv.2005.01.010
- Welsh, L., & Herzing, D. (2008). Preferential association among kin exhibited in a population of Atlantic spotted dolphins (*Stenella frontalis*). *International Journal of Comparative Psychology*, 21, 1-11.
- West, K., Oftedal, O., Carpenter, J., Krames, B., Campbell, M., & Sweeney, J. (2007). Effect of lactation stage and concurrent pregnancy on milk composition in the bottlenose dolphin. *Journal of Zoology*, 273, 148-160. doi:10.1111/j.1469-7998.2007.00309.x

- Westgate, A. (2007). Geographic variation in cranial morphology of short-beaked common dolphins (*Delphinus delphis*) from the North Atlantic. *Journal of Mammalogy*, 88, 678-688.
- Westgate, A., McLellan, W., Wells, R., Scott, M., Meagher, E., & Pabst, D. (2007). A new device to remotely measure heat flux and skin temperature from free-swimming dolphins. *Journal of Experimental Marine Biology & Ecology*, 346, 45-59. doi:10.1016/j.jembe.2007.02.014
- Westgate, A., & Read, A. (2007). Reproduction in short-beaked common dolphins (*Delphinus delphis*) from the western North Atlantic. *Marine Biology*, 150, 1011-1024. doi:10.1007/s00227-006-0394-1
- Wever, E. G. (1951). Hearing. *Annual Review of Psychology*, 2, 65-78.
- White, D., Spong, P., Cameron, N., & Bradford, J. (1971). Visual discrimination learning in the killer whale (*Orcinus orca*). *Behavior Research Methods & Instrumentation*, 3, 187-188.
- Whitehead, H. (1985). Why whales leap. *Scientific American*, 252, 84-93.
- Whitehead, H. (1997). Analysing animal social structure. *Animal Behaviour*, 53, 1053-1067. doi:10.1006/anbe.1996.0358
- Whitehead, H. (1998). Cultural selection and genetic diversity in matrilineal whales. *Science*, 282, 1708.
- Whitehead, H. (2009). Estimating abundance from one-dimensional passive acoustic surveys. *Journal of Wildlife Management*, 73, 1000-1009. doi:10.2193/2008-303
- Whitehead, H., Worm, B., & McGill, B. (2008). Diversity of deep-water cetaceans in relation to temperature: Implications for ocean warming. *Ecology Letters*, 11, 1198-1207.
- Whiten, A. (2001). Imitation and cultural transmission in apes and cetaceans. *Behavioral & Brain Sciences*, 24, 359.
- Wiley, D., Early, G., Mayo, C. A., & Moore, M. J. (2001). Rescue and release of mass stranded cetaceans from beaches on Cape Cod, Massachusetts, USA; 1990-1999: A review of some response actions. *Aquatic Mammals*, 27, 162-171.
- Wiley, D., Moller, J., Pace, R., & Carlson, C. (2008). Effectiveness of voluntary conservation agreements: Case study of endangered whales and commercial whale watching. *Conservation Biology the Journal of the Society for Conservation Biology*, 22, 450-457.
- Wilke, M., Bossley, M., & Doak, W. (2005). Managing human interactions with solitary dolphins. *Aquatic Mammals*, 31, 427-433.
- Williams, A., Petersen, S., Goren, M., & Watkins, B. (2009). Sightings of killer whales *Orcinus orca* from longline vessels in South African waters, and consideration of the regional conservation status. *African Journal of Marine Science*, 31, 81-86. doi:10.2989/AJMS.2009.31.1.7.778
- Williams, N. (2006). A whale's tale. *Current Biology*, 16, R110. doi:10.1016/j.cub.2006.02.016
- Williams, R., & Ashe, E. (2007). Killer whale evasive tactics vary with boat number. *Journal of Zoology*, 272, 390-397. doi:10.1111/j.1469-7998.2006.00280
- Williams, R., Hall, A., & Winship, A. (2008). Potential limits to anthropogenic mortality of small cetaceans in coastal waters of British Columbia. *Canadian Journal of Fisheries & Aquatic Sciences*, 65, 1867-1878. doi:10.1139/F08-098
- Williams, R., Hammond, P., & Lusseau, D. (2009). The role of social aggregations and protected areas in killer whale conservation: The mixed blessing of critical habitat. *Biological Conservation*, 142, 709-719.

- Williams, R., Lusseau, D., & Hammond, P. (2006). Estimating relative energetic costs of human disturbance to killer whales (*Orcinus orca*). *Biological Conservation*, 133, 301-311. doi:10.1016/j.biocon.2006.06.010
- Williams, R., & Noren, D. P. (2009). Swimming speed, respiration rate, and estimated cost of transport in adult killer whales. *Marine Mammal Science*, 25, 327-350. doi:10.1111/j.1748-7692.2008.00255.x
- Williams, R., & Thomas, L. (2009). Cost-effective abundance estimation of rare animals: Testing performance of small-boat surveys for killer whales in British Columbia. *Biological Conservation*, 142, 1542-1547.
- Williams, T., Davis, R., Fuiman, L., Francis, J., Le Boeuf, B., Horning, M., et al. (2000). Sink or swim: Strategies for cost-efficient diving by marine mammals. *Science*, 288, 133-137.
- Williams, T., & Friedl, W. (1992). Travel at low energetic cost by swimming and wave-riding bottlenose dolphins. *Nature*, 355(6363), 821-823.
- Williams, T., & Haun, J. (1999). The diving physiology of bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Biology*, 202, 2739.
- Williams, T., Estes, J. A., Doak, D. F., Springer, A. M. (2004). Killer appetites: assessing the role of predators in ecological communities. *Ecology*, 85, 3373-3384.
- Wilson, B., Reid, R. J., Grellier K., Thompson, P. M., & Hammond, P. S. (2004). Considering the temporal when managing the spatial: A population range expansion impacts protected areas-based management for bottlenose dolphins. *Animal Conservation*, 7, 331-338.
- Williamson, C. (2008). Dolphin assisted therapy: Can swimming with dolphins be a suitable treatment? *Developmental Medicine & Child Neurology*, 50, 477-477.
- Willis, P., & Dill, L. (2007). Mate guarding in male Dall's porpoises (*Phocoenoides dalli*). *Ethology*, 113, 587-597. doi:10.1111/j.1439-0310.2007.01347.x
- Wilson, B., Thompson, P., & Hammond, P. (1997). Habitat use by bottlenose dolphins: Seasonal distributions and stratified movement patterns in the Moray Firth, Scotland. *Journal of Applied Ecology*, 34, 1365-1374.
- Wilson, C., & Tisdell, C. (2003). Conservation and economic benefits of wildlife-based marine tourism: Sea turtles and whales as case studies. *Human Dimensions of Wildlife*, 8, 49-58. doi:10.1080/10871200390180145
- Wilson, J., Cooke, S., Moore, M., Martineau, D., Mikaelian, I., Metner, D., et al. (2005). Systemic effects of arctic pollutants in beluga whales indicated by cyp1a1 expression. *Environmental Health Perspectives*, 113, 1594. doi:10.1289/ehp.7664
- Wilson, R. P., Grant, W. S., Duffy, D. C. (1986). Recording devices on free-ranging marine mammals: Does measurement affect foraging performance? *Ecology*, 67, 1091-1093.
- Wintle, B. (2007). Adaptive management, population modeling and uncertainty analysis for assessing the impacts of noise on cetacean populations. *International Journal of Comparative Psychology*, 20, 237-249.
- Wirsing, A., Heithaus, M., Frid, A., & Dill, L. (2008). Seascapes of fear: Evaluating sublethal predator effects experienced and generated by marine mammals *Marine Mammal Science*, 24, 1-15.
- Wiszniewski, J., Allen, S., & Möller, L. (2009). Social cohesion in a hierarchically structured embayment population of Indo-Pacific bottlenose dolphins. *Animal Behaviour*, 77, 1449-1457. doi:10.1016/j.anbehav.2009.02.025
- Wolkers, H., Bavel, B., Derocher, A., Wiig, Ø., Kovacs, K., Lydersen, C., et al. (2004). Congener-specific accumulation and food chain transfer of polybrominated

- diphenyl ethers in two Arctic food chains. *Environmental Science & Technology*, 38, 1667-1674.
- Wolkers, H., Corkeron, P., Van Parijs, S., Simila, & Van Bavel, B. (2007). Accumulation and transfer of contaminants in killer whales (*Orcinus orca*) from Norway: Indications for contaminant metabolism. *Environmental Toxicology & Chemistry*, 26, 1582-1590.
- Wolkomir, J., & Wolkomir, R. (1996). Marine explorer. *Wildlife Conservation*, 99(4), 36.
- Wood, C. (1999). A period of human contact by two free-ranging bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science*, 15, 863-871.
- Wright, A., Aguilar Soto, N., Baldwin, A., Bateson, M., Beale, C., Clark, C., et al. (2007). Do marine mammals experience stress related to anthropogenic noise? *International Journal of Comparative Psychology*, 20, 274-316.
- Würsig, B. (1989). Cetaceans. *Science*, 244, 1550-1557. doi:10.1126/science.2662403
- Würsig, B. (2007). Whalewatcher: A global guide to watching whales, dolphins and porpoises in the wild. *Quarterly Review of Biology*, 82, 66.
- Würsig, B., & Würsig, M. (1979). Day and night of the dolphin. *Natural History*, 88(3), 60-67.
- Würsig, B., Lynn, S. K., Jefferson, T. A., & Mullin, K. D. (1998). Behaviour of cetaceans in the northern Gulf of Mexico relative to survey ships and aircraft. *Aquatic Mammals*, 24, 41-50.
- Xitco, M., Gory, J., & Kuczaj, S. (2001). Spontaneous pointing by bottlenose dolphins (*Tursiops truncatus*). *Animal Cognition*, 4, 115-123. doi:10.1007/s100710100107
- Xitco, M., Gory, J., & Kuczaj, S. (2004). Dolphin pointing is linked to the attentional behavior of a receiver. *Animal Cognition*, 7, 231-238. doi:10.1007/s10071-004-0217-z
- Xitco, M., & Roitblat, H. (1996). Object recognition through eavesdropping: Passive echolocation in bottlenose dolphins. *Animal Learning & Behavior*, 24, 355-365.
- Yaman, S., von Fersen, L., Dehnhardt, G., & Güntürkün, O. (2003). Visual lateralization in the bottlenose dolphin (*Tursiops truncatus*): Evidence for a population asymmetry? *Behavioural Brain Research*, 142, 109-114. doi:10.1016/S0166-4328(02)00385-6
- Yamazaki, T., Oda, S., & Shirakihara, M. (2008). Stomach contents of an Indo-Pacific bottlenose dolphin stranded in Amakusa, western Kyushu, Japan. *Fisheries Science*, 74, 1195-1197. doi:10.1111/j.1444-2906.2008.01640.x
- Yang, W., Chou, L., Jepson, P., Brownell, R. Jr., Cowan, D., P-H., C., et al. (2008). Unusual cetacean mortality event in Taiwan, possibly linked to naval activities. *Veterinary Record: Journal of the British Veterinary Association*, 162, 184-186.
- Yazdi, P. (2002). A possible hybrid between the dusky dolphin (*Lagenorhynchus albirostris obscurus*) and the southern right whale dolphin (*Lissodelphis peronii*). *Aquatic Mammals*, 28, 211-217.
- Yazdi, P., & Kilian, A. (1999). Energy expenditure of swimming bottlenose dolphins (*Tursiops truncatus*). *Marine Biology*, 134, 601-607.
- Yeates, L., & Houser, D. (2008). Thermal tolerance in bottlenose dolphins (*Tursiops truncatus*). *Journal of Experimental Biology*, 211, 3249-3257. doi:10.1242/jeb.020610
- Yodzis, P. (1994). Predator-prey theory and management of multispecies fisheries. *Ecological Applications*, 4, 51-58.
- Yodzis, P. (1998). Local trophodynamics and the interaction of marine mammals and fisheries in the Benguela ecosystem. *Journal of Animal Ecology*, 67, 635-658.

- Yu, J., Sun, Y., & Xia, Z. (2009). The Rescue, Rehabilitation, and Release of a Stranded Finless Porpoise (*Neophocaena phocaenoides sunameri*) at Bohai Bay of China. *Aquatic Mammals*, 35, 220-225.
- Yuen, H. (1961). Bow wave riding of dolphins. *Science*, 134, 1011-1012. doi:10.1126/science.134.3484.1011
- Yuen, M., Nachtigall, P., Breese, M., & Vlachos, S. (2007). The perception of complex tones by a false killer whale (*Pseudorca crassidens*). *Journal of the Acoustical Society of America*, 121, 1768-1774. doi:10.1121/1.2436640
- Yuen, Q., Ying, M., Brook, F., & Kinoshita, R. (2009). Reliability of 2D ultrasound measurements of testis size in dolphins taken under voluntary behavior. *Ultrasound in Medicine & Biology*, 35, 1005-1009. doi:10.1016/j.ultrasmedbio.2008.12.005
- Yurk, H., Barrett-Lennard, L., Ford, J., & Matkin, C. (2002). Cultural transmission within maternal lineages: Vocal clans in resident killer whales in southern Alaska. *Animal Behaviour*, 63, 1103-1119. doi:10.1006/anbe.2002.3012
- Zagzebski, K. A., Gulland, F. M. D., Haulena, M., Lander, M. E., Greig, D. J., Gage, L. J., et al. (2006). Twenty-five years of rehabilitation of odontocetes stranded in central and northern California, 1977 to 2002. *Aquatic Mammals*, 32, 334-345.
- Zaitseva, K., & Korolev, V. (1994). Resolving abilities of the echolocation system of the bottlenosed dolphins in discrimination of moving target velocity. *Sensory Systems*, 8, 75-79.
- Zaitseva, K., & Korolev, V. (1996). Studies of the ability of dolphin sonar to detect the acceleration of a moving target. *Sensory Systems*, 10, 192-195.
- Zaitseva, K., Korolev, V., & Akhi, A. (2008). Recognition of classes of noise-like signals by dolphins *Tursiops truncatus*. *Journal of Evolutionary Biochemistry & Physiology*, 44, 230-236.
- Zappulli, V., Mazzariol, S., Cavicchioli, L., Petterino, C., Bargelloni, L., & Castagnaro, M. (2005). Fatal necrotizing fasciitis and myositis in a captive common bottlenose dolphin (*Tursiops truncatus*) associated with *Streptococcus agalactiae*. *Journal of Veterinary Diagnostic Investigation*, 17, 617-622.
- Zegers, B., Mets, A., van Bommel, R., Minkenberg, C., Hamers, T., Kamstra, J., et al. (2005). Levels of hexabromocyclododecane in harbor porpoises and common dolphins from Western European Seas, with evidence for stereoisomer-specific biotransformation by cytochrome P450. *Environmental Science & Technology*, 39, 2095-2100.
- Zeppel, H. (2008). Education and conservation benefits of marine wildlife tours: Developing free-choice learning experiences. *Journal of Environmental Education*, 39, 3-17. doi:10.3200/JOEE.39.3.3-18
- Zerbini, A., Waite, J., Durban, J., LeDuc, R., Dahlheim, M., & Wade, P. (2007). Estimating abundance of killer whales in the nearshore waters of the Gulf of Alaska and Aleutian Islands using line-transect sampling. *Marine Biology*, 150, 1033-1045. doi:10.1007/s00227-006-0347-8
- Zerbini, A. N., & Santos, M. C. D. O. (1997). First record of the pygmy killer whale *Feresa attenuata* (Gray, 1874) for the Brazilian coast. *Aquatic Mammals*, 23, 105-109.
- Zhukovskii, Y., Zaitseva, K., Krivchenko, A., & Korolev, V. (2006). Mechanisms by which dolphins identify target movement characteristics and their technical use in hydrolocation and radiolocation. *Neuroscience and Behavioral Physiology*, 36, 801-809. doi:10.1007/s11055-006-0090-0

- Zollett, E., & Read, A. (2006). Depredation of catch by bottlenose dolphins (*Tursiops truncatus*) in the Florida king mackerel (*Scomberomorus cavalla*) troll fishery. *Fishery Bulletin*, 104, 343-349.
- Zolman, E. (2002). Residence patterns of bottlenose dolphins (*Tursiops truncatus*) in the Stono river estuary, Charleston County, South Carolina, U.S.A. *Marine Mammal Science*, 18, 879-892.
- Zornetzer, H., & Duffield, D. (2003). Captive-born bottlenose dolphin x common dolphin (*Tursiops truncatus* x *Delphinus capensis*) intergeneric hybrids. *Canadian Journal of Zoology*, 81, 1755-1762. doi:10.1139/Z03-150