

Lawrence Berkeley National Laboratory

Recent Work

Title

Investigation of interactions of U(VI) with bacteria by laser spectroscopic methods

Permalink

<https://escholarship.org/uc/item/1pz1210q>

Author

Nitsche, H.

Publication Date

2001-05-03

LBNL-47658 Abs.

This abstract was prepared for an oral presentation at the 8th International Conference on “Chemistry and Migration Behaviour of Actinides and Fission Products in the Geosphere (Migration 01)” to be held in Bregenz, Austria, September 16- 21, 2001.

prepared: February 28, 2001

Abstract Form Migration'01

Speaker's name: Dr. Roger Knopp

Institute: Lawrence Berkeley National Laboratory, The Glenn T. Seaborg Center

Address: 1 Cyclotron Road, MS 70A-1150, Berkeley, CA, 94720, USA

Phone number: +510 486 7709

Fax number: +510 486 7444

E-mail address: rknopp@lbl.gov

Investigation of Interactions of U(VI) with Bacteria by Laser Spectroscopic Methods

R. Knopp¹, P.J. Panak¹, L.A. Wray¹, N. Renninger², J.D. Keasling², H. Nitsche^{1,3}

¹) Lawrence Berkeley National Laboratory, The Glenn T. Seaborg Center, Berkeley, CA 94720, USA

²) University of California at Berkeley, Department of Chemical Engineering, Berkeley, CA 94720, USA

³) University of California at Berkeley, Department of Chemistry, Berkeley, CA 94720-1460, USA

Bacteria are omnipresent in natural environments. They may play a role in the immobilization or transportation of actinides in aquifers. For example, the gram-positive *Bacillus sphaericus* was found to take up relatively large amounts of hexavalent uranium and plutonium [1]. This study by Panak et al. also indicated that the uptake occurs extracellularly via phosphate-containing cell structures like teichoic acids. We investigated in detail the nature of the functional groups involved in this uranium binding process by employing time-resolved laser-induced fluorescence spectroscopy (TRLFS) and Raman spectroscopy. In addition to living cells, we also investigated spores, intact dead cells and decomposed cells. By comparing fluorescence spectra of U(VI) interacting with living cells, spores, and intact dead cells to U(VI) adenosine monophosphate (AMP), we found very similar spectra for all these complex species. The AMP resembles very well the structure of teichoic acids with a phosphate group directly bound to an organic rest. U(VI), in contact with decomposed cells, shows a very different fluorescence spectrum than U(VI)-AMP. The spectrum is nearly identical to the spectrum of U(VI) that was precipitated with NaH₂PO₄ at the same pH than for the bacterial studies. Raman spectra revealed that the decomposition of the bacteria leads to enzymatic production of H₂PO₄⁻. This increases the immobilization of U(VI) compared to the immobilization by living cells. In addition to sorption and decomposition, the natural bacterial metabolism can express complexing agents (phosphates or organic chelators). We investigated this with gram-negative *Pseudomonas aeruginosa* that were genetically engineered to over express phosphate. The kinetics and the mechanisms of the interaction of U(VI) with the released phosphate were studied by spectroscopy. We observed quantitative precipitation of U(VI) phosphate. In contrast, *P. aeruginosa*, not induced to express phosphate, does not change the fluorescence of U(VI) and therefore, it can be concluded that no significant interaction of natural metabolic byproducts of these bacteria with U(VI) occurs under the given conditions.

[1] Panak, P.; Raff J., S.; Selenska-Pobell, S.; Geipel, G.; Bernhard, G.; Nitsche, H., *Radiochim. Acta* 88, 71-76, 2000.

Preference:

Session No.: B4

Oral: X

=====
Acknowledgements: This work is supported by the U.S. Department of Energy under Contract No. DE-AC03-76SF00098