

Lawrence Berkeley National Laboratory

Recent Work

Title

ASSOCIATED PRODUCTION IN PION-NUCLEON COLLISIONS AND CHARGE INDEPENDENCE

Permalink

<https://escholarship.org/uc/item/1q02w83n>

Author

Barshay, Saul.

Publication Date

1957-06-28

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-3834

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

ASSOCIATED PRODUCTION IN PION-NUCLEON COLLISIONS
AND CHARGE INDEPENDENCE

Saul Barchay

June 28, 1957

Printed for the U. S. Atomic Energy Commission

ASSOCIATED PRODUCTION IN PION-NUCLEON COLLISIONS AND CHARGE INDEPENDENCE

Saul Barshay

Radiation Laboratory
University of California
Berkeley, California

June 28, 1957

Using the method of Low,¹ we have derived the equation for the Heisenberg scattering amplitude describing the processes $\pi + N \rightarrow \Lambda + K$ and $\pi + N \rightarrow \Sigma + K$. In the "one-meson" approximation¹ with pair intermediate states omitted we obtain a linear integral equation for this amplitude. The inhomogeneous term is the Born approximation, and the kernel of the uncrossed one-meson term is the pion-nucleon scattering amplitude. The crossed one-meson term involves the pion-hyperon scattering amplitude and may be treated with the inhomogeneous term. A partial-wave study of the Born approximation has been made, using the Hamiltonian suggested recently by Gell-Mann² with the assumption of spin-parity 0^- for the K mesons and $(1/2)^+$ for the Λ and Σ hyperons. S, P, and D waves have been included to first order in an expansion in powers of the ratio of the intermediate-state baryon kinetic energy to its mass. At 1 Bev pion bombarding energy we find that a coupling of the form

$$g (\bar{\Lambda} N K - \bar{\Sigma}_a N \tau_a K + \text{hermitian conjugate})$$

yields the strong backward preference of the Λ hyperons in the center-of-mass system that has been observed experimentally.^{3,4} Also in accord with

¹F. E. Low, Phys. Rev. 97, 1392 (1955).

²M. Gell-Mann, A Model for the Strong Couplings Phys. Rev. (to be published).

³Budde, Chretien, Qeitner, Samios, Schwartz, and Steinberger, Phys. Rev. 103, 1827 (1956).

⁴D. Glaser, private communication.