

UCSF

UC San Francisco Previously Published Works

Title

HIV and risk behaviors among visitors of inmates at the Great Tehran Prison, Iran, 2018.

Permalink

<https://escholarship.org/uc/item/1q78t4dt>

Journal

The Open AIDS Journal, 14(1)

ISSN

1874-6136

Authors

Najafi, Zeinab
Shekarbeigi, Saeede
Koochak, Hamid Emadi
et al.

Publication Date

2020

DOI

10.2174/1874613602014010035

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Published in final edited form as:

Open AIDS J. 2020 ; 14(1): 35–40. doi:10.2174/1874613602014010035.

HIV and risk behaviors among visitors of inmates at the Great Tehran Prison, Iran, 2018

Zeinab Najafi¹, Saeede Shekarbeigi¹, Hamid Emadi Koochak¹, Behnam Farhoudi², SeyedAhmad SeyedAlinaghi¹, Ali Mirzazadeh³

¹Iranian Research Center for HIV/AIDS (IRCHA), Iranian Institute for Reduction of High-Risk Behavior, Tehran, Iran

²Clinical Research Development Center, Amir-Almomenin Hospital, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran

³University of California, University of San Francisco, San Francisco, United States

Abstract

Objectives: The objective of the present study was to determine the prevalence of HIV and risk behaviors among the inmates' visitors at the Great Tehran Prison in 2018. This population is more likely to be infected with HIV due to their relationship with inmates. However, they have not been prioritized to the other populations in comparison to inmates who are usually prioritized over others.

Method: Using a convenient sampling method, we recruited inmates' visitors at the Great Tehran Prison in 2018. An anonymous questionnaire collected information on the participant's demographic characteristics, relationship with inmate, imprisonment history, drug-related risk behaviors, and HIV-related risk behaviors. Rapid HIV blood test detected HIV positive people. To examine the association between risk behaviors and HIV infection, we used a multivariate logistic regression model to calculate the adjusted Odds Ratio (OR) (adjusted Odds ratio of 23.140, 95% CI OD: 1.994–268.5).¹

¹. It should be noted that according to the latest Spectrum projections, Iran's HIV population in 2018 stood at 60,760. For more information see: <https://www.unaids.org/en/regionscountries/countries/islamicrepublicofiran>

Correspondence to: SeyedAhmadSeyedAlinaghi, **Address:** Iranian Research Center for HIV/AIDS, Imam Khomeini Hospital, Keshavarz Blvd., Tehran, Iran, **Tel/Fax:** +98 (21) 66947984, s_a_alinaghi@yahoo.com.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The Ethical committee of Tehran University of Medical Sciences, Iran, approved the study protocol under reference no. 8923101017.

HUMAN AND ANIMAL RIGHTS

No animals were used in this research. All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013

CONSENT FOR PUBLICATION

The participants were recruited using a convenience sampling method and interviewed after collecting informed written consent.

AVAILABILITY OF DATA & MATERIALS

The data supporting the findings of the article are available in the Iranian Research Center for HIV/AIDS at <https://research.tums.ac.ir/>, reference number 8923101017.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

Results: 1131 visitors were asked to participate in the study; out of which 1120 consented (99%). Out of the 1120 participants, 55.2% were female, and the majority (72.4%) aged older than 35 years. Of those who had a sexual partner, 50% did not use a condom in the last 6 months. Overall, HIV prevalence was 0.4% in the inmate visitors, but it was significantly higher among those with a history of drug injection ($p=0.003$).

Conclusion: Our findings indicated that overall the HIV prevalence is low among inmates' visitors; Subgroups with a history of injection have high HIV prevalence which needs to be prioritized for HIV screening and treatment.

Keywords

HIV/AIDS; Inmate Visitors; Prison; Iran

1. Introduction

There are about 220,000 inmates in Iran [1]. The prevalence of HIV among inmates is higher than the general population [2-3] which is attributed to the low socioeconomic condition, lack of knowledge about HIV and its transmission routes, drug injection, high-risk sexual behaviors, and tattoos in inmates [4-5] [6]. Prisons and other temporary residential facilities (such as camps and shelters) have reported having a higher rate of HIV transmission [7, 8].

Imprisonment not only increases the risk of HIV transmission among the incarcerated populations but also it is likely the risk to be higher among other members of society through unsafe sexual or drug practices [9]. Given the short average time of being in prison (about 6 months) and the risk of acquiring HIV infection within the prison and then transmitting to others upon release, most of the HIV programs have included inmates population as a priority population [9]. One study documented that 50% of Hispanic inmates had sex within the first 12 hours after release from prison [10]. The majority of men in relationships had unprotected intercourse with their primary partners within days of paroling from prison [11]. These findings indicated a high potential risk of HIV transmission to inmates' sexual partners. According to a study in Iran, turnover of prisoners back to society is high and while prisoners are not separated in prison based on the history of drug use, they might transmit the HIV within the prison which in turn can amplify the possibility of transmitting infection outside of the prison to inmates' sexual partners or people in their drug use circle [4].

Inmates who used to inject or use drugs still have fair access to drugs inside the prisons [12], and of course, upon release, they most likely continue to drug use, share injection kits, and even overdose [13]. This puts their family members who also use drugs at risk for HIV infection.

Besides the potential higher risk for HIV infection among inmates' visitors, this group has not been studied very well. As the rate of HIV infection among inmates is higher than in society, this study was designed to investigate the prevalence of HIV infection among inmates' visitors whom may be exposed to HIV infection more than other people. Moreover,

investigation of related risk factors was another objective of the present study which helps to reduce the HIV infection risk among this population.

2. Methods

In this cross-sectional study, we measured the demographic characteristics, HIV prevalence, and HIV-related risk behaviors in 1131 visitors from which 1120 accepted to participate of male inmates at the Great Tehran Prison. The participants were recruited using a convenience sampling method and interviewed after collecting informed written consent.

The questionnaire included sections on demographic characteristics, the relationship withinmates, imprisonment history, drug injection-related risk behaviors, and HIV-related risk behaviors. Counseling was also provided regarding HIV transmission and prevention measures and the introduction of HIV testing and counseling services. This was done by two trained researchers in the prison's visiting room. Those who provided consent for HIV testing were tested by a rapid HIV blood test kit, and if the test results were reactive, they were retested on a second rapid test. Those who were reactive for both rapid tests were considered to be HIV positive.

HIV point prevalence was calculated as the number of HIV-positive individuals divided by the number of test results overall and stratified by demographic characteristics, incarceration history, and drug-use practices. Variables associated with HIV in bivariate analysis at the $p < 0.10$ level were included as potential independent predictors. The final model retained those variables associated with infection at the $p < 0.05$ level.

The data was analyzed using the SPSS software version 22. Variables associated with HIV infection in bivariate analysis at the $p < 0.10$ levels were included as potential independent predictors. The final model retained those variables associated with HIV infection at the $p < 0.05$ level.

Ethical considerations

The protocol of the study was reviewed and approved by the Institutional Review Board (IRB) of Tehran University of Medical Sciences. All the participants were informed and familiarized with the study objectives and methods. The participants were assured about their information anonymity and confidentiality. Informed written consent was obtained from all the participants prior to the study.

3. Results

Out of the 1120 participants, majority were female (55.2%), aged older than 35 years (72.4%), married (83.1%), and first degree family member (i.e. spouse, daughter, sister, mother) (Table 1). The most prevalent high-risk behaviors among participants were unprotected sex (47.9%), history of addiction (17.6%), marriage to a person with at least one risk factor (16.4%), having multiple partners (14.2%), history of incarceration (11.2%) (Fig. 1).

Overall, 4 (0.4%) were HIV positive. HIV prevalence was significantly higher among those with a history of drug injection than others ($p = 0.003$). In multivariate analysis, a history of

injection remained independently associated with HIV infection (adjusted Odds ratio of 23,140, 95% CI OD: 1,994-268,5), $p = 0.003$).

4. Discussion

To the best of our knowledge, this study was the first study to assess the HIV prevalence and risk behaviors of inmates' visitors in Iran. Our findings indicated that overall the HIV prevalence is low among inmates' visitors; however, those with a history of injection have the highest HIV prevalence which needs to be prioritized for HIV screening and treatment.

Unsafe sex is high among partners of inmates. Female partners of male inmates reported having good knowledge about HIV and its transmission routes; however, they did not use condoms in sex with their partners when they were released from the prison as they thought they did not have sex or other risk behaviors during incarceration. This is not true as the majority of male inmates had a history of injection and 20% had a second partner [14]. We also found that half of the inmates' visitors did not use condoms during the last 6 months and about three percent reported a history of drug injection.

Among women visiting their incarcerated partners at several prisons in California, the HIV prevalence was 1.23%; Factors associated with HIV prevalence were age, incarceration period, history of drug injection, history of needle sharing, and tattooing [15]. While HIV prevalence was lower in our study, (i.e. 0.4%), we also found that history of drug injection is the main predictor for HIV. Similar to findings in our study, other concerning risk factors include lack of practical knowledge and positive attitudes toward HIV prevention practices and high frequency of unprotected sex [16].

Male inmates have a high frequency of risk behaviors (unsafe sex or injection), before and after incarceration. Unprotected sex was reported by 46% of the prisoners for the time prior to incarceration, and by 18% at one month and by 24% at three months post-release [17].

Partners of male inmates are at risk for other mental health problems like anxiety and depression which may lead to lower self-esteem and also misjudgment about risk behaviors and risk perception [15]. Having multiple partnerships among partners of male inmates and higher risk rates of STDs and HIV infection were reported in the past [18]. Women with multiple partnerships reported more history of STDs treatment than those who were in a one-one relationship (26% vs. 11%) [19]. This also explains the slightly higher HIV prevalence among our study population.

5. LIMITATION:

Our study had three main limitations. First, the number of HIV positive cases was low in our study and so we did not have the power to fully assess the risk factors of HIV infection. Second, the collected risk behaviors were self-reported hence they were likely to be under-reported. And last, we did not study the HIV and risk behaviors of inmates and their visitors together and so were not able to assess the proportion and predictors of unsafe relationships.

CONCLUSION:

Despite the limitations, we found the HIV prevalence to be low among the inmates' visitors. Overall Subgroups with a history of injection had high HIV prevalence which needs to be prioritized for HIV screening and treatment. Being in a prison for a while provides a window opportunity for prevention interventions, family support, and behavioral change in both inmates and their families. However, right now our best option for intervention is to continue our programs of distributing syringes inside and outside of prison. HIV programs in Iran need to use this opportunity to further improve their response to drug-related HIV epidemics in the country.

Acknowledgements

The authors express thanks to all staff in the Great Tehran Prison for their help. This study was supported by Tehran University of Medical Sciences (grant no.8923101017). For this paper, we also received support from the University of California, San Francisco's International Traineeships in AIDS Prevention Studies (ITAPS), U.S. NIMH, R25MH064712.

References

- [1]. Walmsley R. World Prison Population List. http://www.prisonstudies.org/sites/default/files/resources/downloads/wppl_12.pdf; 2018.
- [2]. Nunn A, Cornwall A, Fu J, Bazerman L, Loewenthal H, Beckwith C. Linking HIV-positive jail inmates to treatment, care, and social services after release: results from a qualitative assessment of the COMPASS program. *Journal of Urban Health*. 2010;87(6):954–68. [PubMed: 21046470]
- [3]. Hosseini M, SEYED ANS, KHEYR AP, ESMAEILI JGR, Shirzad H, Karami N, et al. Prevalence and correlates of co-infection with human immunodeficiency virus and hepatitis C virus in male injection drug users in Iran. 2010;13(4): 318–323.
- [4]. Navadeh S, Mirzazadeh A, Gouya MM, Farnia M, Alasvand R, Haghdoost A-A. HIV prevalence and related risk behaviours among prisoners in Iran: results of the national biobehavioural survey, 2009. *Sex Transm Infect*. 2013;89(Suppl 3):iii33–iii6. [PubMed: 23986417]
- [5]. Kheirandish P, Seyedalinalghi S, Hosseini M, Jahani M, Shirzad H, Foroughi M, et al. Prevalence and correlates of HIV infection among male injection drug users in detention in Tehran, Iran. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2010;53(2):273–5. [PubMed: 20104123]
- [6]. Catalan-Soares BC, Almeida RTP, Carneiro-Proietti ABF. Prevalence of HIV-1/2, HTLV-I/II, hepatitis B virus (HBV), hepatitis C virus (HCV), *Treponema pallidum* and *Trypanosoma cruzi* among prison inmates at Manhuaçu, Minas Gerais State, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical*. 2000;33(1):27–30.
- [7]. Altice FL, Mostashari F, Selwyn PA, Checko PJ, Singh R, Tanguay S, et al. Predictors of HIV infection among newly sentenced male prisoners. *Journal of acquired immune deficiency syndromes and human retrovirology: official publication of the International Retrovirology Association*. 1998;18(5):444–53.
- [8]. Allwright S, Bradley F, Long J, Barry J, Thornton L, Parry JV. Prevalence of antibodies to hepatitis B, hepatitis C, and HIV and risk factors in Irish prisoners: results of a national cross sectional survey. *Bmj*. 2000;321(7253):78–82. [PubMed: 10884256]
- [9]. Macalino GE, Vlahov D, Sanford-Colby S, Patel S, Sabin K, Salas C, et al. Prevalence and incidence of HIV, hepatitis B virus, and hepatitis C virus infections among males in Rhode Island prisons. *American journal of public health*. 2004;94(7):1218–23. [PubMed: 15226146]
- [10]. Gomez C, Marin B, Morales T. Freedom and HIV prevention: Challenges facing Latino inmates leaving prison. San Francisco: University of California, San Francisco, Center for AIDS ...; 1996.

- [11]. Grinstead OA, Zack B, Faigeles B, Grossman N, Blea L. Reducing postrelease HIV risk among male prison inmates: A peer-led intervention. *Criminal Justice and Behavior*. 1999;26(4):453–65.
- [12]. Simpler AH, Langhinrichsen-Rohling J. Substance use in prison: how much occurs and is it associated with psychopathology?. *Addiction Research & Theory*. 2005 10 1;13(5):503–11.
- [13]. Seaman S, Brettle R, Gore SM. Mortality from overdose among injecting drug users recently released from prison: database linkage study. *Bmj*. 1998;316(7129):426–8. [PubMed: 9492665]
- [14]. Grinstead OA, Zack B, Faigeles B. Collaborative research to prevent HIV among male prison inmates and their female partners. *Health Education & Behavior*. 1999;26(2):225–38. [PubMed: 10097966]
- [15]. Comfort M, Grinstead OA, Faigeles B, Zack B. Reducing HIV risk among women visiting their incarcerated male partners. *Criminal Justice and Behavior*. 2000;27(1):57–71.
- [16]. Grinstead O, Comfort M, McCartney K, Koester K, Neilands T. Bringing it home: Design and implementation of an HIV/STD intervention for women visiting incarcerated men. *AIDS Education & Prevention*. 2008;20(4):285–300. [PubMed: 18673062]
- [17]. Grinstead OA, Faigeles B, Comfort M, Seal D, Nealey-Moore J, Belcher L, et al. HIV, STD, and hepatitis risk to primary female partners of men being released from prison. *Women & health*. 2005;41(2):63–80.
- [18]. Margolis AD, MacGowan RJ, Grinstead O, Sosman J, Kashif I, Flanigan TP, et al. Unprotected sex with multiple partners: implications for HIV prevention among young men with a history of incarceration. *Sexually transmitted diseases*. 2006;33(3):175–80. [PubMed: 16505732]
- [19]. Wagstaff DA, Kelly JA, Perry MJ, Sikkema KJ, Solomon LJ, Heckman TG, et al. Multiple partners, risky partners and HIV risk among low-income urban women. *Family Planning Perspectives*. 1995:241–5. [PubMed: 8666088]




Figure 1.
The trend of HIV prevalence (%) among prisoners in Iran between 1999 and 2018².

².SeyedAlinaghi SA, Taj L, Mazaheri E, Ahsani-Nasab S, Abedinzadeh N, Mcfarland W, Mohraz M, Mirzazadeh A. HIV in Iran: Epidemiology, Response, Challenges and Way Forward. *AIDS. In Press*

Table1.

Demographic characteristics and HIV prevalence of people visiting incarcerated men, Great Tehran Prison, Iran, 2018

Variable	N* (%)	HIV prevalence, N (%)
Gender		
Male	618 (55.2)	2 (0.4)
Female	502 (44.8)	2 (0.3)
Age group		
26-35	309 (27.6)	1 (0.3)
36-45	290 (25.9)	2 (0.7)
46-59	295 (26.3)	1 (0.3)
>60	95 (8.5)	0 (0.0)
Marital status		
Single	73 (12.6)	3 (4.1)
Married	482 (83.1)	0 (0.0)
Divorced	20 (3.4)	1 (5.0)
Widowed	5 (0.9)	0 (0.0)
Education level		
Illiterate	97 (16.7)	1 (1.0)
Primary school	90 (15.5)	0 (0.0)
Did not complete high school	135 (23.3)	1 (0.3)
Completed high school	175 (30.2)	0 (0.0)
More than high school	83 (14.3)	0 (0.0)
Who you are visiting today?		
Spouse	285 (25.4)	1 (0.3)
Son	347 (31.0)	0 (0.0)
Brother	319 (28.5)	0 (0.0)
Father	47 (4.2)	1 (2.1)
Grandchild	7 (0.6)	0 (0.0)
Friend	59 (5.3)	2 (3.4)
Other relatives	40 (3.6)	0 (0.0)
Client	16 (1.4)	0 (0.0)

* Subgroups do not always add up to total due to missing data

Table2.

Statistical measures for the age of participants

Variable	Mean	Std. Deviation	Range
age	40.31	12.851	67

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Behaviors and HIV prevalence among people visiting incarcerated men, Great Tehran Prison, Iran, 2018

Variables	N (%)	HIV prevalence, N (%)
Reported history of incarceration *	125(11.2)	2 (1.6)
Reported history of addiction	197(17.6)	2 (1.0)
Reported history of injection **	27 (2.4)	2 (7.4)
Reported ever sharing needles or equipment	8 (0.7)	0(0.0)
Reported ever having unprotected sex	537(47.9)	3 (0.5)
Reported having multiple partners ***	159(14.2)	2 (1.2)
Reported history of male-male sex	13 (1.2)	0(0.0)
Receiving blood and blood products	3 (0.3)	0(0.0)
Reported history of STD	82 (7.3)	1 (1.2)
Married to a person with at least one risk factor ^l	184(16.4)	1 (0.5)
Reported ever having unprotected sex with HIV ⁺ partner	1 (0.1)	0(0.0)
Reported having occupational or non-occupational exposure	1 (0.1)	0(0.0)
History of tattooing	(2) 22	0 (0.0)
Condom use during the last 6 months		
Always	19 (6.3)	0 (0.0)
Sometimes	77 (25.7)	0 (0.0)
Rarely	54 (18)	0 (0.0)
Never	150 (50)	2 (1.3)
Routes of administration in drug users		
Smoking	122 (10.9)	1 (0.8)
Oral consumption	26 (2.3)	0(0.0)
Injection	3 (0.3)	0(0.0)
Smoking/ Oral	6 (0.5)	0(0.0)
Oral/ Injection	1 (0.1)	0(0.0)
All	3 (0.3)	0(0.0)

* P=0.06

** P= 0.003

*** P=0.10

^l Drug injection,unprotected sexual contact