Lawrence Berkeley National Laboratory

Recent Work

Title

VECTOR-TO-VECTOR POLARIZATION TRANSFER IN DEUTERON-PROTON ELASTIC SCATTERING

Permalink

<https://escholarship.org/uc/item/1q81d328>

Authors

Rad, P.N.
Birchall, J.
Conzett, H.E.
et al.

Publication Date

1974-09-01

VECTOR-TO-VECTOR POLARIZATION TRANSFER IN
DEUTERON-PROTON ELASTIC SCATTERING

F. N. Rad, J. Birchall, H. E. Conzett and R. Roy

September, 1974

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

VECTOR-TO-VECTOR POLARIZATION TRANSFER IN DEUTERON-PROTON ELASTIC SCATTERING*

F. N. Rad, J. Birchall[†], H. E. Conzett and R. Roy[‡]Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

September 1974

Abstract:

An angular distribution of the polarization transfer coefficient $K_Y^{Y'}(\theta)$ in d-p elastic scattering has been measured at $E_d = 45.4$ MeV. Our results are in very good agreement with the prediction of a recent three-body calculation.

We report in this letter on the first significant comparison between experimental and theoretically calculated vector-to-vector polarization transfer coefficients in nucleon-deuteron elastic scattering.

Considerable theoretical progress has been made recently in fitting the nucleon analyzing-powers measured in N-d elastic scattering at energies below 50 MeV.¹⁻³ These three-nucleon calculations, based on the Faddeev equations with separable nucleon-nucleon interactions, have also provided predictions of the deuteron vector and tensor polarizations and, more recently, of polarization transfer coefficients⁴ at energies up to $E_N = 23$ MeV. Subsequent determinations of the deuteron analyzing-powers were in good agreement with the calculations.⁵ These studies showed that the P-wave part of the N-N interaction was chiefly responsible for the vector analyzing power, whereas, in addition, the two-body $^3S_1 - ^3D_1$ tensor force was required in order to provide agreement with the observed tensor analyzing powers.

As a further test of the theoretical predictions, we report on measurements of the polarization of the recoil protons in d-p elastic scattering with incident 45.4 MeV vector polarized deuterons.⁶ These ${}^1\text{H}(\vec{d}, \vec{p}){}^2\text{H}$ measurements provide determinations of the vector-to-vector polarization transfer coefficient, $K_Y^{Y'}(\theta)$, defined by the equation⁷

$$p_{Y'} = (A_{Y'} + \frac{3}{2} p_Y K_Y^{Y'}) / (1 + \frac{3}{2} p_Y A_{Y'}), \quad (1)$$

where p_Y is the polarization of the purely vector polarized incident deuteron beam, $p_{Y'}$ is the observed polarization of the outgoing (recoil) proton, and A_Y and $A_{Y'}$ are, respectively, the deuteron vector analyzing power and the proton analyzing power in d-p elastic scattering. The Madison convention⁸ is followed in defining the cartesian forms of the polarization and analyzing power, and the y and y' axes are taken along $\vec{k}_d \times \vec{k}_p$. From time-reversal invariance, the same coefficient describes the transfer of vector polarization in the inverse ${}^2\text{H}(\vec{p}, \vec{d}){}^1\text{H}$ scattering process at the equivalent proton energy of 22.7 MeV.⁷ Thus, our measurements can be compared directly with Pieper's⁴ calculated $K_Y^{Y'}(\theta)$ at that energy.

Previous measurements of vector-to-tensor polarization transfer coefficients in p-d scattering were made at two angles at several energies between 5 and 9 MeV.⁹ The values were consistent with zero within a typical error of ± 0.05 . Pieper's later three-body calculation⁴ gave predicted values of these coefficients to be less than 0.1 for energies below 10 MeV, in essential agreement with the data. Of particular interest was his prediction that the vector-to-vector transfer coefficients differed more substantially from zero. A single measured value of $K_X^{X'}(\theta)$ at 9 MeV¹⁰ was two standard deviations from the calculated value, but it was clear that more data were

required for a realistic comparison with the predictions. His calculation also predicted substantial values of $K_y^{y'}(\theta)$ at $E_p = 22.7$ MeV.

Our experimental arrangement is shown in Fig. 1. The polarized deuteron beam from the Berkeley 88-inch cyclotron passed through a high pressure liquid-nitrogen-cooled hydrogen gas target, positioned in a 36-inch diameter scattering chamber. The entrance and exit windows of the gas target were Havar foils of 25 μ m in thickness. The areal density of the gas was 12.8 mg/cm² at the operating pressure of 12.4 atm, and the mean energy in the gas was 45.4 MeV. Two silicon polarimeters, described in detail elsewhere,¹¹ were used to determine the polarization of the scattered protons. They were positioned at equal scattering angles to the left and right of the beam axis. The angular resolution, defined by tantalum collimators, was 0.8° (FWHM). Two monitor counters were placed left and right of the beam axis at a scattering angle of $\theta \simeq 22^\circ$ and azimuthal angles near 70° and 110°. A helium gas target cell and a pair of ΔE -E counter telescopes at equal left and right scattering angles were positioned in a smaller scattering chamber downstream from the main scattering chamber, and this provided continuous monitoring of the beam polarization. The vector polarization of the beam, with zero tensor component, was near $0.81 \pm .03$ of the maximum possible value $p_y = 2/3$. The analyzing power of d-⁴He scattering at this energy had been measured previously.¹² For each determination of $K_y^{y'}(\theta)$, data were taken with the spin vector of the beam orientated alternately up and down with respect to the scattering plane. This mode of operation along with the left-right polarimeters provided several internal consistency checks. The relative yield of elastic doubly-scattered protons into the LL detector of the silicon polarimeter is given by:

$$LL^{\pm} = (1 \mp 3/2 p_y A_y) (1 + p_y A_2), \quad (2)$$

where the (+) and (-) superscripts correspond to the incident beam spin orientation of up and down, respectively, and A_2 is the silicon proton analyzing power in the second scattering.¹³ Inserting p_y from Eq. (1),

$$LL^{\pm} = 1 \mp 3/2 p_y A_y + A_2 A_y, \pm 3/2 A_2 p_y K_y^{y'} \quad (3)$$

From this and similar expressions for the relative yields $LR^{\pm}$, $RL^{\pm}$, and $RR^{\pm}$ one obtains¹⁴

$$3/2 p_y A_2 K_y^{y'} = \frac{(LL^+ - LL^-) - (LR^+ - LR^-)}{LL^+ + LL^- + LR^+ + LR^-} \quad (4a)$$

$$= \frac{(RR^- - RR^+) - (RL^- - RL^+)}{RR^- + RR^+ + RL^- + RL^+} \quad (4b)$$

$$= \frac{(LL^+ - RR^+) - (LR^+ - RL^+)}{LL^+ + RR^+ + LR^+ + RL^+} \quad (4c)$$

$$= \frac{(RR^- - LL^-) - (RL^- - LR^-)}{RR^- + LL^- + RL^- + LR^-} \quad (4d)$$

First order systematic errors due to instrumental asymmetries and beam misalignments were eliminated by averaging the values of $K_y^{y'}(\theta)$ from Eqs. (4a) and (4b), and similarly from Eqs. (4c) and (4d). The two average values obtained in this way agreed to within ± 0.006 at all angles. The effective proton analyzing power A_2 of the silicon polarimeter was determined with polarized protons elastically scattered from ^4He and ^{12}C , for which $K_y^{y'}(\theta) = 1.0$. With p_y known Eq. (4) then results in a value for A_2 . Details of the silicon polarimeter and of the calibration procedures and results are reported elsewhere.¹¹

Our experimental results are shown in Fig. 2, where the errors indicated are purely statistical. The solid curve is Pieper's calculation; and it is seen that the agreement with experiment is excellent. Since this constitutes the first significant comparison between experimental and calculated polarization-transfer coefficients in N-d elastic scattering, this agreement represents yet another substantial success of the three-body calculations, particularly in view of the fact that there has been no adjustment of the two-body input parameters in order to fit these data.

Despite the excellent agreement between these experimental and predicted values of $K_Y^{y'}(\theta)$, it recently has become clear that further calculations are necessary to address unresolved questions in N-d scattering. Pieper compared his perturbative calculation⁴ with Doleschall's exact calculation² for the same input N-N interactions, and he found that there were significant differences between the two calculations of the nucleon polarization near 23 MeV. In his most recent calculation, Doleschall³ improved the P-wave and the $^3S_1 - ^3D_1$ tensor interactions to give better agreement with the N-N phase-shift parameters. This resulted in improved fits to the nucleon and deuteron¹⁵ vector analyzing-power data. An appropriate question, then, is the extent to which such improved N-N interactions would change the calculated values of $K_Y^{y'}(\theta)$. As has been suggested,^{15,16} an important program now is to specifically examine the sensitivities of the various analyzing-powers and polarization-transfer coefficients to the details of the input two-body interactions.

We are grateful to R. M. Larimer for her assistance during the course of the experiment.

FOOTNOTES AND REFERENCES

* Work performed under the auspices of the U. S. Atomic Energy Commission.

† Present Address: Physics Department, University of Basel, Switzerland

‡ National Research Council of Canada Post Doctoral Fellow.

1. S. C. Pieper, Nucl. Phys. A193, 529 (1973).
2. P. Doleschall, Nucl. Phys. A201, 264 (1973).
3. P. Doleschall, Nucl. Phys. A220, 491 (1974).
4. S. C. Pieper, Phys. Rev. C 8, 1702 (1973).
5. J. S. C. McKee, H. E. Conzett, R. M. Larimer, and Ch. Leemann, Phys. Rev. Letters 29, 1613 (1972); A. Fiore, J. Arvieux, Ngugen Van Sen, G. Perrin, F. Merchez, J. C. Gondrand, C. Perrin, J. L. Durand, and R. Darves - Blanc, Phys. Rev. C 8, 2019 (1973).
6. Preliminary results were reported at the International Conference on Few Body Problems in Nuclear and Particle Physics, Laval University, Quebec, Canada, August 27-31, 1974.
7. G. G. Ohlsen, Rep. Prog. Phys. 35, 717 (1972).
8. Polarization Phenomena in Nuclear Reactions, edited by H. H. Barschall and W. Haeberli (Univ. of Wisconsin Press, Madison, 1971), p. xxv.
9. G. G. Ohlsen, C. K. Mitchell, and G. C. Salzman, Few Particle Problems in the Nuclear Interaction, edited by I. Slaus, S. A. Moszkowski, R. P. Haddock, and W. T. H. van Oers (North-Holland, Amsterdam, 1973), p. 460.
10. G. G. Ohlsen, C. K. Mitchell, G. C. Salzman, J. L. Gammel, D. C. Dodder, and K. Witte, Bull. Am. Phys. Soc. 17, 922 (1972).
11. J. Birchall, H. E. Conzett, W. Dahme, J. Arvieux, F. N. Rad, R. Roy, and R. M. Larimer, Nuclear Instruments and Methods, in press.

12. Ch. Leeman, H. E. Conzett, W. Dahme, J. A. Macdonald, and J. P. Meulders, Bull. Amer. Phys. Soc. 17, 562 (1972), and to be published.
13. For the spin-up orientation, the negative sign in the first term of Eq. (2), and correspondingly in the denominator of Eq. (1), results from the fact that the incident deuteron is scattered to the right, so p_y is opposite to $\vec{k}_i \times \vec{k}_f$.
14. Equation (4b) is analagous to an expression for $p_{y2} D$ which appeared in work of P. Schnabel, Thesis, University of Paris, 1971 (unpublished). There D is the depolarization parameter in proton elastic scattering from nuclei.
15. F. N. Rad, J. Birchall, H. E. Conzett, S. Chintalapudi, and R. Roy, Bull. Amer. Phys. Soc. 19, 476 (1974). Submitted to Phys. Rev. Letters.
16. H. E. Conzett, Proc. Int'l. Conf. on Few Body Problems in Nuclear and Particle Physics, Laval University, Quebec, Canada, August 27-31 1974, in press.

FIGURE CAPTIONS

Fig. 1. Schematic diagram of the experimental arrangement, showing the left and right silicon polarimeters. The detectors labelled L and R are the left and right silicon analyzers, respectively. LL, LR, RL, and RR are the silicon side detectors of the polarimeters.

Fig. 2. The vector-to-vector polarization transfer coefficient, $K_Y^{y'}(\theta)$, in d-p elastic scattering at $E_d = 45.4$ MeV. The solid curve is the calculated result from Ref. 4.

XBL7410-4413

Fig. 2

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720