Lawrence Berkeley National Laboratory

Recent Work

Title

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

Permalink

<https://escholarship.org/uc/item/1qh3d9ph>

Author

Bradley, Dan F.

Publication Date

1955-12-21

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

Dan F. Bradley

December 21, 1955

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

Contents

Abstract	3
Introduction	4
Experimental Methods	6
Results and Discussion	
Chromatography	7
Stationary States	8
Transients	23
Effect of Atmosphere	27
Acknowledgment	27
References	28

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

Dan F. Bradley

Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

December 21, 1955

ABSTRACT

Transient changes in the P^{32} activity of PPA, PGA, OP, GP, ATP, ADP, and RDP when the illumination is terminated and initiated have been measured in Scenedesmus. The signs of the transients, their order of appearance, and the number of maxima and minima are consistent with a model involving production of ATP from OP and ADP by the oxidation of photochemically produced $TPNH_2$, the reduction of PGA with the photo-produced ATP and $TPNH_2$, the formation of GP from the resulting triose phosphate, the phosphorylation of RMP formed from the GP mediated by photoproduced ATP, and the production of PPA from PGA. The effect of CO_2 upon the stationary state and transients of PGA and RDP is consistent with the carboxylation of RDP to form PGA.

PHOSPHATE TRANSIENTS IN PHOTOSYNTHESIS

Dan F. Bradley*

Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

December 21, 1955

INTRODUCTION

When a photosynthesizing plant is suddenly placed in the dark, the production of the primary photoproducts and consumption of primary photoreactants cease, resulting in a transient decrease in the concentrations of the products and increase in the concentrations of the reactants. Numerous experiments have recently been designed to determine which plant metabolites are produced and which are consumed by the photochemical reaction by observing which ones decrease and increase when illumination is suddenly terminated.

O. Kandler¹ observed that the TCA-soluble inorganic phosphate content of Chlorella pyrenoidosa fell in the 30 seconds after the beginning of illumination by about 20%, whereas in the first 90 seconds after darkening it rose about 30%, although there was no essential difference between the stationary states in the light and in the dark. These observations could be interpreted in terms of a photochemically driven consumption of orthophosphate (OP)_i[†] presumably in the formation of ATP.

M. Calvin and P. Massini² studied transients in Scenedesmus obliquus, using C¹⁴ as a tracer and paper chromatography to isolate and identify the products. They found: "When illumination is interrupted there appears a

* Present address: Section of Physical Chemistry, National Institute of Mental Health, Bethesda 14, Maryland.

† The following abbreviations are used herein: LDT, light-to-dark transients; DLT, dark-to-light transient; OP, orthophosphate; PGA, phosphoglyceric acid; GP, glucose phosphates; PPA, phosphopyruvic acid; ATP, adenosine triphosphate; ADP, adenosine diphosphate; RDP, ribulose diphosphate; RMP, ribulose monophosphate; TPN and TPNH₂, oxidized and reduced triphosphopyridine nucleotide.

sudden great increase in the concentration of PGA followed by a slow decrease after two minutes, and an almost complete depletion of the diphosphate area. Analysis of the monophosphate area showed that the amount of sedoheptulose phosphate decreased also. The concentration of malic acid decreases as well. The rate of labeling of glutamic acid is increased greatly after a short induction period; citric acid, which contains little activity during the whole light period, shows a sudden increase in the dark, followed by a slow decrease, the labeling of sucrose continues at the same rate as in the light for about two minutes, after which it is stopped almost completely. ... In the second experiment the diphosphate area, which in the first experiment contained almost the same number of counts as PGA during the light, had only about 15% of it. This value dropped to 5% in the dark. The PGA showed a hardly significant rise in the dark during the first two minutes, but again a slow decrease after five minutes. ... In the light following the dark the diphosphates, PGA, and malic acid increase again."

B. L. Strehler,³ using the firefly luminescent system as an assay tool, has observed the influence of light and other environmental factors on ATP content of Chlorella pyrenoidosa. He found that "After a period of anaerobiosis the ATP level rises upon illumination or admission of air to higher levels (2 to 10 times dark levels). The ATP levels with respect to time show a maximum at approximately one minute; thereafter a decrease." On the other hand, "Illumination causes a decrease in the ATP after a period of dark aerobiosis, and the subsequent dark period is characterized by a rapid increase in the ATP level. This effect is observable in the presence or absence of CO₂, indicating that the ATP-utilizing reactions are largely prior to or independent of carboxylation reactions." Strehler concludes "that ATP is both generated and used by light-driven reactions."

A. T. Wilson⁴ has observed transients in photosynthesizing Scenedesmus obliquus when the CO₂ tension is suddenly reduced. He found that PGA and FMP decrease and the RDP area, RMP, FDP, GDP, TP, and glycolic acid increase in the first 50 seconds of lowered CO₂ tension. As PGA is the first to decrease and RDP the first to increase, Wilson concluded that the CO₂-fixing reaction involved the formation of two PGA molecules from RDP and CO₂. The other transient changes are secondary, resulting from the effect of the primary transients in PGA and RDP on reactions further from the carboxylation step, and hence appear at later times.

In order to determine which of the numerous chemicals described above experiencing light-dark transients are primary photochemicals, it would therefore seem necessary to determine the order in time of these transients. Comparison of the data of Kandler, Calvin and Massini, and Strehler obtained under different experimental conditions and with different algae seem insufficient for this purpose, so we have measured light-dark transients in Scenedesmus obliquus using P^{32} as a tracer and paper chromatography to isolate and identify the products.

EXPERIMENTAL METHODS

Twenty-four-hour sterile Scenedesmus obliquus cultures were grown from 10% inoculum in an atmosphere of 4% CO_2 -air with an incident light intensity of 1.8×10^4 ergs/cm²/sec at 25°C. The harvested cells were centrifuged and decanted three times from their culture medium to remove traces of OP- P^{31} , which interfered with subsequent measurements. Suspensions of the cells in distilled water were made up so as to exhibit an optical density of the ethanol-extracted pigments at the 6600 Å red maximum of chlorophyll-a of 8 OD units, corresponding to 9×10^{-5} M chlorophyll-a in suspension and approximately 10 mm^3 packed cells per ml suspension. The suspension was transferred to a lucite reaction vessel 3/8 in. thick which was cooled by a running-tap-water jacket. A tube inserted in the solution bubbled the appropriate gas mixture at several hundred bubbles per min, stirring the suspension. Illumination was provided by a bank of seven GE 6-watt white fluorescent lights at an intensity of 4×10^4 ergs/cm²/sec. The intensity in the vicinity of the reaction vessel when the lights were off was less than 40 ergs/cm²/sec.

The suspension was pretreated with the desired gas mixture and lighting conditions for 15 to 30 min. Sufficient isotopically pure OP- P^{32} in physiological saline (0.9%) was then added to the suspension so as to give a final activity of 3 µc/ml (approximately 0.01% NaCl) in the suspension. After 56 min of exposure to OP- P^{32} , $1 \pm .02$ ml of the suspension was removed by pipet and expelled into 4 ml of simmering absolute ethanol. The sampling,

which required about 3 sec between the time the cells were withdrawn from the reaction vessel and contacted the ethanol, was repeated at 58 and 60 min. At 60-1/4 min the illumination was stopped--or started--and samples were withdrawn 1/4, 1/2, 3/4, 1, 1-1/2, 2, 3, 4, and 5 min thereafter and treated with simmering ethanol. Each of the twelve samples was then brought to a boil over a steam bath, and allowed to extract at room temperature for 1 to 2 hr. The cell residues were spun off and resuspended in 25% v/v ethanol-water, brought to a boil, and allowed to extract for 1 to 2 hr. The 80% and 25% extracts were combined and diluted to 8 ml with ethanol. Aliquots of 0.5 ml were spotted on Whatman No. 4 paper for one-dimensional chromatography with the Buchanan modification of the Hanes-Isherwood solvent system: t-butanol--water--picric acid, 80v/20v/2w.^{5, 6}

The chromatograms were run in dry or water-saturated boxes, depending on which separations were desired, and the dried papers exposed to duPont X-ray film for several days. The P^{32} activity in each of the 6 or 7 main spots was counted with a mica-window GM tube through the developed X-ray film; 2-mm-thick lead shields were used to isolate the desired spots from the neighboring spots. A 1.1-mm soft glass plate calibrated to give $8.7 \pm .2\%$ transmission of the spot activity was used whenever the count was greater than 5000 counts per min. Counts were taken for 2 min, or at least 1000 counts for each spot. Whenever any count indicated the existence of a maximum or minimum in the transient curves, the spot was recounted and the average of the two counts reported. Each spot is reported as the percentage of the total P^{32} in the original suspension, measured by counting an appropriate aliquot of the suspension spotted on the developed chromatogram.

RESULTS AND DISCUSSION

Chromatography

Six principal P^{32} spots were observed which accounted for roughly three-fourths of the total P^{32} , the rest of the activity remaining associated with the cell hulks or appearing in several minor spots of sugar phosphates

and phospholipids on the chromatogram. Cochromatography showed OP, PGA, and GP spots to be quite free of contaminants. PPA was identified by its R_F value and assumed to be pure because of its good separation from other spots. While the ADP and ATP spots contain these compounds as indicated by cochromatography, two-dimensional chromatography showed the presence of other minor contaminants of undetermined nature. In experiments carried out in CO_2 -free atmosphere the activity in the RDP spot, identified by its R_F value, was high, and in several instances it was not possible to obtain good resolution of the GP and RDP spots; in these cases the sum of GP and RDP is reported. Table I lists the ratios of the R_F 's of each spot to the R_F of orthophosphate to indicate what type of contaminants might be expected in each spot. Columns 1 and 2 also demonstrate the effect of saturating the box with water vapor before chromatography.

The results of ten experiments are tabulated in Tables II through XI. Seven experiments involved light-to-dark transients (LDT) and three dark-to-light transients (DLT). Experiments 1557, 1559, and 1564 were identical as to experimental conditions but were carried out with different harvests of Scenedesmus. The same is true of 1556 and 1566 as well as of 1562 and 1565. Experiment 1565 was carried out with the suspension remaining after Experiment 1564, after the light had been off for 36 minutes. Similarly Experiment 1567 followed 1566 after 56 minutes of darkness. In order to better visualize these transients the data are represented as curves in Fig. 1 through 10, drawn so that each curve possesses the minimum curvature while passing within 0.2% of each experimental point.

Stationary States

The one-hour P^{32} incorporation was generally sufficiently long so that the activity in each compound was nearly time-invariant, or at most varied only slowly with respect to the LDT and DLT. As the turnover time of most compounds in Scenedesmus in photosynthesis is of the order of several minutes we expect that the distribution of P^{32} corresponds nearly to the relative concentrations of the compounds. The averages of the three pretransient points for experiments in Tables II-XI are presented in Table XII.

Table I

Ratios of R_F of chromatogram spots to R_F of orthophosphate.
(Solvent: picric acid-t-butanol-water.)

Spot	P^{32} Activity		Known Spots			Hanes-Isherwood		Wilson P^{32} Activity
	Dry Box	H ₂ O- Satd. Box	Paper 1	Paper 2	Paper 3	Whatman 1	Whatman 4	
Adenosine triphosphate	0.06	0.16	0	0.14	0.08	0.07	0.08	
Adenosine diphosphate	0.19	0.34	0.15	0.21	0.28	0.24	0.26	
Glucose phosphate	0.42	0.54	0.44	0.44		0.41	0.44	0.40
Phosphoglyceric acid	0.90	0.93	0.89			0.90		0.89
Phosphopyruvic acid	1.19	1.12						1.20
Uridine triphosphate			0.06	0.13	0.08			
Uridine diphosphate			0.26	0.28	0.28			
Uridine monophosphate			0.60					
Adenosine monophosphate						0.64		
Diphosphopyridine nucleotide			0.03	0.12				
Fructose-6-phosphate					0.66	0.59		0.61
Mannose monophosphate								0.52
Ribose-5-phosphate								0.58
Sedoheptulose monophosphate								0.52
Fructose diphosphate			0.66			0.59	0.62	0.58
Ribulose diphosphate								0.58
Glucose diphosphate								0.58
Phosphoglycer-aldehyde			0.60		0.44			
Phosphoglycolic acid								1.06
Dihydroxyacetone phosphate			0.70		0.76			

Table II

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1564.
Atmosphere 4% CO_2 , 96% air.

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D	5D
PGA	18.6	18.9	20.2	20.1	20.7	20.7	19.6	17.8	17.1	17.1	16.4	15.7
PPA	3.2	3.3	3.1	3.5	3.7	3.7	3.7	3.7	3.3	3.5	3.4	3.1
GP	3.9	4.2	4.4	4.0	3.7	3.3	3.0	2.7	2.7	2.9	3.1	3.4
OP	11.0	12.8	10.9	17.4	12.2	12.9	11.8	12.9	15.4	15.0	13.4	11.1
ATP	11.2	10.5	9.9	8.2	9.8	9.8	9.3	10.0	8.7	10.1	10.1	11.1
ADP	16.6	16.9	17.7	14.1	16.8	18.7	18.7	19.5	20.2	19.2	18.7	18.9
Totals	64.5	66.6	66.2	67.3	66.9	69.1	66.1	66.6	67.9	67.8	65.1	63.3

Table III

Percent of total P^{32} found in principal spots on chromatogram of dark-to-light transients extracted from Scenedesmus cells. Experiment 1565
(carried out with suspension remaining after Experiment 1564). Atmosphere 4% CO_2 , 96% air

Substance isolated	Conditions: Time (in minutes) in light or dark											
	36D	38D	40D	1/4L	1/2L	3/4L	1L	1-1/2L	2L	3L	4L	5L
PGA	16.9	17.4	16.6	16.5	18.0	18.2	17.7	20.0	19.8	19.5	21.1	22.3
PPA	3.1	3.1	3.0	2.6	2.4	2.3	2.0	1.9	1.9	1.9	2.0	2.1
GP	2.8	2.9	3.0	3.7	3.7	3.7	3.6	4.0	3.8	3.9	3.9	3.8
OP	10.3	9.6	10.6	9.0	7.3	7.3	6.6	9.1	9.2	8.5	8.5	7.5
ATP	10.6	10.1	10.5	10.8	10.8	9.5	9.7	10.7	9.9	9.3	9.7	9.2
ADP	14.6	14.9	15.5	13.7	13.7	14.0	14.5	13.3	13.0	13.0	14.4	14.0
Totals	58.3	58.0	59.2	56.3	55.9	55.0	54.1	59.0	57.6	56.1	59.6	58.9

Table IV

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1557
(same as 1564 with different harvest of Scenedesmus).
Atmosphere 4% CO_2 , 96% air.

Substance isolated	Conditions: Time (in minutes) in light or dark										
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D
PGA	15.5	16.1	15.3	17.7	18.2	18.0	17.5	17.7	15.3	14.1	14.7
PPA	1.5	1.6	1.6	2.1	2.1	2.1	2.1	2.2	2.2	2.2	2.1
GP	3.9	3.8	3.6	3.3	3.0	2.6	2.8	2.5	2.7	2.4	1.8
OP	37.4	34.2	32.2	37.3	35.2	31.0	36.2	35.2	35.1	38.6	34.7
ATP	6.4	6.7	6.9	6.4	6.0	(5.9)	5.8	6.7	6.4	5.7	(5.7)
ADP	10.5	11.2	11.8	9.4	8.5		12.0	9.7	11.2	9.6	
Totals	75.2	73.6	71.4	76.2	73.0		76.4	74.0	72.9	72.6	

Table V

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1559
(same as 1564 and 1557 but with different harvest of Scenedesmus.)
Atmosphere 4% CO_2 , 96% air.

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D	5D
PGA	20.9	21.1	22.4	23.5	23.5	23.6	23.2	20.8	20.8	18.4	19.4	18.2
PPA	1.7	1.6	1.7	2.3	2.3	2.6	2.5	2.3	2.4	2.5	2.7	2.8
GP	3.9	4.9	4.6	3.7	3.8	3.8	3.4	3.0	2.9	2.8	2.8	2.6
OP	28.0	33.0	32.5	34.0	30.2	27.5	33.2	33.9	31.3	34.6	34.0	29.7
ATP	4.1	4.1	4.3	4.2	4.3	4.1	4.3	3.9	4.2	4.0	4.1	4.0
ADP	7.2	8.1	7.3	5.7	7.7	7.2	9.0	8.3	9.5	9.5	9.5	9.7
Totals	65.8	72.8	72.8	73.4	71.8	68.8	75.6	71.5	71.1	71.8	72.5	67.0

Table VI

Percent of total P^{32} found in principal spots on chromatogram of dark-to-light transients extracted from Scenedesmus cells. Experiment 1562
(same as 1565 but with different harvest of Scenedesmus).
Atmosphere 4% CO_2 , 96% air.

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56D	58D	60D	1/4L	1/2L	3/4L	1L	1-1/2L	2L	3L	4L	5L
PGA	5.9	6.2	6.5	5.9	8.2	9.0	9.3	8.9	9.3	9.8	10.2	10.4
PPA	1.3	1.6	1.5	1.3	1.3	1.2	1.0	0.9	0.9	0.9	1.1	1.1
GP	2.3	2.1	2.2	2.8	2.7	2.8	2.9	2.8	3.1	3.5	3.3	3.5
OP	67.0	61.7	63.5	62.4	56.9	56.5	56.0	54.0	53.8	53.0	52.0	52.5
ATP	5.2	5.0	5.1	5.8	6.2	5.2	5.5	6.1	5.3	6.6		
ADP	8.9	8.0	9.2	12.3	10.0	8.7	9.6	10.2	7.3	10.6		
Totals	90.6	84.6	88.0	90.5	85.3	83.4	84.3	82.9	79.7	84.4		

Table VII

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1568.
Atmosphere 95.4% N_2 , 4.6% CO_2 .

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D	5D
PGA	17.2	17.8	17.8	18.2	19.2	17.7	18.2	15.1	13.5	13.2	12.5	11.7
PPA	1.9	1.8	1.9	2.2	2.4	2.3	2.3	2.3	2.2	2.2	2.2	2.2
GMP	4.4	4.7	4.7	4.4	4.3	3.7	3.8	3.7	3.7	4.1	4.0	4.2
OP	30.7	29.7	28.1	29.4	26.3	24.7	24.8	26.3	23.1	29.4	29.0	31.2
ATP	9.8	9.0	9.7	8.6	9.1	9.9	9.5	9.0	8.7	7.1	7.7	7.6
ADP	12.8	13.8	14.5	12.1	14.4	14.8	15.4	16.5	17.0	12.0	12.1	14.0
Totals	76.8	76.8	76.7	74.9	75.7	73.1	74.0	72.9	68.2	68.0	67.5	70.9

Table VIII

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1566.
Atmosphere 100% N_2 .

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D	5D
PGA	4.8	4.4	4.8	5.1	5.9	7.2	7.2	7.4	7.7	8.4	7.9	8.1
PPA	1.1	1.1	1.0	1.2	1.3	1.4	1.5	1.5	1.5	1.7	1.9	2.0
GP	2.8	2.5	2.9	3.8	4.1	3.9	4.6	4.1	3.3	3.3	3.3	3.2
OP	17.8	17.0	16.3	19.0	18.0	16.8	16.9	16.3	17.7	17.1	20.9	23.2
ATP	4.0	4.0	4.5	3.4	3.6	3.3	3.7	3.7	3.7	4.2	3.5	3.6
ADP	13.6	12.8	11.7	9.7	10.1	11.1	9.5	10.6	12.6	11.3	11.9	11.8
RDP	23.4	22.6	21.8	20.8	19.1	20.4	19.2	17.8	20.1	18.0	16.0	15.1
Totals	67.5	64.4	63.0	63.0	62.1	64.1	62.6	61.4	66.6	64.0	65.4	67.0

Table IX

Percent of total P^{32} found in principal spots on chromatogram of dark-to-light transients extracted from Scenedesmus cells. Experiment 1567,
Atmosphere N_2 .

Substance isolated	Conditions: Time (in minutes) in light or dark											
	46D	48D	50D	1/4L	1/2L	3/4L	1L	1-1/2L	2L	3L	4L	5L
PGA	17.5	7.1	7.5	3.4	3.0	2.9	3.2	3.3	2.9	3.2	3.3	3.6
PPA	2.3	2.6	2.4	1.5	1.0	1.0	1.0	0.9	0.8	0.8	1.2	1.2
OP	30.4	30.6	29.1	27.0	23.2	24.8	21.3	20.9	20.2	16.4	16.4	12.1
GP+RDP	4.4	3.9	4.0	7.2	9.6	10.6	10.3	11.1	10.3	11.6	12.6	12.3
ATP+ADP	22.7	23.2	22.8	27.6	29.1	27.6	24.9	26.2	24.4	23.8	23.2	23.2
Totals	67.3	67.4	65.8	66.7	65.9	66.9	69.7	62.4	58.6	55.8	56.7	52.4

Table X

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1556
(same as 1566 but with different harvest of Scenedesmus).
Atmosphere 100% N_2 .

Substance isolated	Conditions: Time (in minutes) in light or dark										
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D
PGA	5.1	5.9	5.2	6.0	6.8	8.0	7.0	7.3	7.6	7.9	8.1
PPA	0.9	1.1	1.0	1.3	1.4	1.6	1.5	1.6	1.7	1.9	2.2
GMP	4.5	3.8	3.4	5.8	6.6	5.9	6.6	5.8	5.4	5.3	5.0
OP	14.7	14.1	12.8	15.1	13.1	13.8	13.8	12.2	13.6	15.2	18.7
ATP	4.0	4.5	4.6	4.6	4.2	4.2	4.2	4.5	4.4	4.0	4.1
ADP	8.2	9.2	10.1	7.4	7.4	7.5	7.4	8.2	8.5	9.8	9.1
RDP	15.3	15.0	15.8	12.1	13.6	17.1	14.8	15.0	13.4	12.8	
Totals	52.7	53.6	52.9	52.3	53.1	58.1	55.3	54.6	54.6	56.9	

Table XI

Percent of total P^{32} found in principal spots on chromatogram of light-to-dark transients extracted from Scenedesmus cells. Experiment 1570.
Atmosphere air.

Substance isolated	Conditions: Time (in minutes) in light or dark											
	56L	58L	60L	1/4D	1/2D	3/4D	1D	1-1/2D	2D	3D	4D	5D
PGA	6.9	7.1	6.6	8.2	10.2	12.6	14.6	12.1	11.9	16.3	14.7	15.0
PPA	1.7	1.9	1.8	1.8	2.2	2.9	3.1	2.8	3.1	3.9	4.2	3.9
OP	39.8	37.6	37.6	38.6	36.2	38.0	33.5	31.4	31.0	32.6	33.1	31.4
GP+RDP	9.8	10.7	9.0	8.2	8.6	6.6	5.4	5.1	4.5	4.0	3.5	3.8
ATP	6.4	6.3	8.1	7.5	7.0	6.6	6.4	9.8	9.1	7.9	7.6	6.3
ADP	8.4	10.1	10.0	7.3	8.7	9.8	9.5	10.7	12.6	10.0	10.2	10.0
Totals	73.2	73.7	73.1	71.6	72.9	76.5	72.5	71.9	72.2	74.7	73.3	70.4

Figs. 1-10. Curves showing percent of P^{32} in the principal transients appearing in Scenedesmus cells during the first few minutes of transition from light to dark or from dark to light.

In each graph the experiment number and the composition of the atmosphere are indicated at the top; the time (in minutes) preceding and following the turning off or turning on of the light appear to left and right of the axis of ordinates.

MU-10759

MU-10760

MU-10761

Table XII

Stationary levels of phosphates, as averages of percent of total activity found at pretransient times in Scenedesmus cells (see Tables II through XI).

Light								Dark		
Spot	Experiment Number and Atmosphere							1567 N ₂	1562 Air-CO ₂	1565 Air-CO ₂
	1556 N ₂	1566 N ₂	1570 Air	1568 N ₂ -CO ₂	1557 Air-CO ₂	1559 Air-CO ₂	1564 Air-CO ₂			
PGA	5.4	4.7	6.9	17.6	15.6	21.5	19.2	7.4	6.2	17.0
PPA	1.0	1.1	1.8	1.9	1.6	1.7	3.2	2.4	1.5	3.1
GP	3.9	2.7	*	4.6	3.7	4.5	4.2	4.1**	2.2	2.9
OP	13.9	17.0	38.3	29.5	34.6	31.2	11.8	30.0	64.0	10.2
ATP	4.4	4.2	6.9	9.5	6.7	4.2	10.5	22.9	5.1	10.4
ADP	9.2	12.7	9.5	13.7	11.2	7.5	17.1		8.7	15.0
RDP	15.4	22.6	9.9		negli- gible					

* Included with RDP

** All sugar phosphates

The OP activity varies widely, from a minimum of about 10% to a high of 64%. Such wide variation is to be expected because the OP activity represents both intracellular OP and the OP that has not yet been incorporated from the medium. The rate of OP-P³² incorporation is sensitive to light and the extracellular OP-P³¹ content; e.g., the OP = 64% is the result of slower incorporation in the dark; when the culture medium rather than distilled water is used for the suspension the extracellular OP-P³¹ is so high that there is no detectable organic P³² in one-hour light incorporation. We expect that the 10% values correspond to the true intracellular OP content, in good agreement with the data of Kamen and Spiegelan⁷ and Emerson, Stauffer, and Umbreit.⁸

The RDP level is negligible in the presence of CO₂ but is the most active spot obtained in a CO₂-free atmosphere. On the other hand, the PGA level is three to four times as great with CO₂ as without. Both these facts are consistent with the assumption that the carboxylation reaction of photosynthesis is the addition of CO₂ to RDP to form 2 PGA's. Emerson, Stauffer, and Umbreit⁸ first observed that the "resistant" phosphate ester is several times as high in the light with CO₂ as without--the major stationary-state difference they were able to observe--but owing to peculiarities in their analytical data they refrained from referring to this resistant phosphate as PGA. Wintermans and Tjia⁹ found that a labile phosphate is formed in the light in the absence but not in the presence of CO₂, which is probably RDP and related to the observations above.

When the photosynthetic rate is high in the presence of CO₂, there is little change from light to dark in the activity distribution (from Experiment 1564 to Experiment 1565). The relatively small difference in light-dark stationary states plagued investigators in the past who hoped to find in larger differences clues as to the nature of the photosynthetic process. Goodman, Bradley, and Calvin¹⁰ found only small differences in P³² distribution between light followed by 15 minutes of dark and between dark and 15 minutes of light in Scenedesmus. Kandler¹ found no essential differences in stationary-state light and dark OP level. Strehler³ found that the stationary-state ATP levels were approximately equal in light and dark. Gest and Kemen¹¹ found "no significant or consistent differences between light and dark samples with respect to P³¹ content of the fractions examined."

When the photosynthetic rate is slowed by a nitrogen atmosphere, there are considerable light-dark stationary-state differences, as indicated by

Experiments 1566 and 1567. The drop of RDP from 23% to a negligible activity, the rise of OP from 17% to 30%, and the smaller rise of other spots contrast with the experiment in the presence of CO₂, and suggests a photochemically driven consumption of OP, formation of RDP, and consumption of PGA. The terms "consumption" and "formation" are somewhat misleading since, if a stationary state is reached in photosynthesis each intermediate is both "formed" and "consumed" at the same rate by light-driven reactions. Therefore, "consumption" as we use it here means that the utilization of the compound is closer in the metabolic network to the light reactions than the formation of the compound.

Transients

Tables XIII and XIV give the signs of the initial transients together with the estimated times required to reach the first maximum (or minimum). Comparable values are included from the experiments of Strehler,³ Kandler,¹ and Calvin and Massini.² When no maximum or minimum is reached within 5 minutes no time is given. Our sign of DLT for PGA is not the same as reported by Calvin and Massini, since the minimum is not deep and is rapidly reached, so that at their earliest experimental point (1/2 minute) the sign of DLT has changed from - to +. There is no such reason apparent for the difference in sign of the LDT of ATP between our results and those of Strehler.

These signs may be compared with what would be expected on the basis of the models of Bassham *et al.*,¹² Arnon, Allen, and Whatley,¹³ Vishniac and Ochoa,¹⁴ and others:

When the formation of TPNH₂ (R₁) is stopped by darkness, the reduction of PGA ceases (R₃), causing a buildup in PGA level, i. e., LDT of PGA is positive.

Table XIII

Sign of Transient upon Darkening and Time Required for the First Maximum or Minimum (Minutes)											
Substance produced	Experiment										
	1556	1566	1570	1568	1557	1559	1564	Kandler	Strehler	Calvin and Massini	Theory
N ₂	+	+	+	+	+	+	+			+	
O ₂			+		+	+	+	+	+	+	
CO ₂				+	+	+	+	+	+	+	
PGA	+	+	+	+	+	+	+			+	+
	3/4		1	1/2	1/2	1/2	1/2			1/2	
PPA	+	+	+	+	+	+	+				+
				1/2			1				
GP	+	+		-	-	-	-				-
	3/4	1		1			1-3/4				
OP	+	+	+	+	+	+	+	+			+ or -
	1/4	1/4	1/4	1/4	1/4	1/4	1/4	1-1/2			
ATP	0	-	+	-	-	-	-		+		- or +
		1/2	1/4	1/4	3/4	3/4	3/4		3		
ADP	-	-	-	-	-	-	-				+ or -
	1/2	1/4	1/4	1/4	1/4	1/4	1/4				
RDP	-	-	-							-	-
	1/4	1/2								1/2	

Table XIV

Sign of Transient upon Illumination and Time Required for First Maximum or Minimum (Minutes)							
Substance produced	Experiment						
	1567	1562	1565	Kandler	Calvin and Massini	Strehler	Theory
N ₂	+	+	+		+	+	
O ₂		+	+	+	+		
CO ₂		+	+	+	+		
PGA	- 1-1/4	- 1/4	- 1/4		+		-
PPA	-	- 2	- 2-1/2				-
GP		+	+				+
OP	-	- 4	- 1	- 1/2			+ or -
ATP	+ 1/2	+ 1/2	+ 1/4			+	- or +
ADP		+ 1/2	- 1/4				+ or -
RDP					+		+

As PGA increases part will be converted to extra PPA (R_7), making its LDT also positive. As the production of TP decreases (R_3 is stopped) the GP and RDP, which are in quasi equilibrium with it, drop; i.e., negative LDT. The situation is more complicated with ATP, ADP, and OP, since they are produced and consumed by reactions just one step from the photochemical reaction (R_2 , R_3). If step (R_2) is the first to feel the effect of the stopping of (R_1), then an increase in OP and ADP is expected with a decrease in ATP. If (R_3) is the first to feel the (R_1) stopping, then the opposite signs would be predicted. If (R_2) and (R_3) both felt the effects simultaneously no transients in OP, ADP, or ATP would occur, or at least the transients would be small and would change sign rapidly. As seen in Table XIII, all signs with the exception of ADP (or

OP and ATP) are in agreement with the theoretical predictions. A single major exception appears in the positive sign of LDT with GP in a N_2 atmosphere, which might be interpreted in terms of a requirement of photoproduced ATP in the transformation of GP to RDP.

On the basis of similar reasoning the sign predictions for the DLT are given in Table XIV; they are in complete agreement with the experimental results.

Further predictions as to the relative times required to reach the first maximum or minimum can be made on the basis of the model if we assume that the compounds closest to the photochemical process have the shortest times. Since it appears from the signs that (R_2) is closer to (R_1) than (R_3) (temporally) we would predict the following order of increasing times,

ATP, OP, ADP $\longrightarrow$ PGA $\longrightarrow$ PPA, GP, RDP,

which is consistent with the observed order, with the possible exception that we would expect the RDP time to be longer than observed. This may indicate that photochemically produced ATP is used to phosphorylate RMP to RDP, making RDP closer to the R_1 than would appear from the model. In the DLT the PGA reaches its minimum surprisingly rapidly, again suggesting that activity is rapidly incorporated into PGA via "photosynthetic phosphorylation" of RMP to RDP mediated by ATP.

The long time required for the OP minimum in DLT may be interpreted in terms of its initial increased utilization by the R_2 reaction and by the failure of the sluggish (R_3) reaction to keep pace, so that PGA also builds up continuously after its first dip and OP fails to reach a minimum within a short time. The ATP does not build up during this period, presumably because of its use in RMP phosphorylation.

A third method of prediction based on the model depends on the assumption that those compounds closest to the photochemical act and involved in the largest numbers of reactions experience the largest number of maxima and minima in a given transient time period. We would expect, therefore, that the order of decreasing numbers of maxima and minima in the first five-minute period would be OP, ADP, ATP $\longrightarrow$ PGA $\longrightarrow$ PPA, TP, RDP, which is in good agreement with the order actually observed in Fig. 1 through 10, again with the exception of RDP, which has more than the expected number of nodes, suggesting its photodrive formation from RMP.

Effect of Atmosphere

Perhaps the most interesting effect of atmospheric condition is the CO_2 effect on the PGA transients: the initial transients in PGA are smaller in the presence of CO_2 than in its absence. In the LDT the initial positive PGA transient results from the carboxylation of residual RDP, which is very much smaller in the presence of CO_2 than in its absence. Calvin and Massini² noticed that the PGA LDT was very small in their "second experiment," where the RDP level was low for an unknown reason. In the DLT the initial transient arises from the initiation of PGA reduction while the second arises from PGA production from RDP. In the presence of CO_2 this formation can occur rapidly, so that the first minimum occurs quite soon, followed by a positive DLT. In the absence of CO_2 the formation from RDP is inhibited so that the initial DLT of PGA continues for several minutes and reaches a very deep minimum.

ACKNOWLEDGEMENT

I wish to express my appreciation to Professor Melvin Calvin for his advice, interest, and encouragement in the course of this work.

This work was done under the auspices of the U. S. Atomic Energy Commission.

REFERENCES

1. O. Kandler, Z. Naturforsch. 5b, 423 (1950).
2. M. Calvin and P. Massini, *Experientia* 8, 445 (1952).
3. B. L. Strehler, *Arch. Biochem. Biophys.* 43, 67 (1953)
4. A. T. Wilson, Thesis, University of California Radiation Laboratory Report No. UCRL-2589, May 1954.
5. C. S. Hanes and F. A. Isherwood, *Nature* 164, 1107 (1949).
6. J. G. Buchanan, private communication.
7. M. D. Kamen and S. Spiegelman, *Cold Spring Harbor Symposia Quant. Biol.* 13, 151 (1948).
8. J. F. Stauffer and W. W. Umbreit, *Am. J. Botany* 31, 107 (1944).
9. J. F. G. M. Winternans and J. E. Tjia, *Koninkl. Ned. Akad. Wetenschap. Proc.* 55C, 34, (1952).
10. M. Goodman, D. F. Bradley, and M. Calvin, *J. Am. Chem. Soc.* 75, 1962 (1953).
11. H. Gest and M. D. Kamen, *J. Biol. Chem.* 176, 299 (1948).
12. J. A. Bassham et al., *J. Am. Chem. Soc.* 76, 1760 (1954).
13. D. I. Arnon, M. B. Allen, and F. R. Whatley, *Nature* 174, 394 (1954).
14. W. Vishniac and S. J. Ochoa, *Biol. Chem.* 198, 501 (1952).