Lawrence Berkeley National Laboratory
LBL Publications

Title
LEPTON BRANCHING FRACTIONS

Permalink
https://escholarship.org/uc/item/1qz658vf

Author
Aihara, H.

Publication Date
1984-06-01

eScholarship.org Powered by the California Digital Library
University of California

LBL-18014
Preprint

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

Physics Division


RECEIVED
LAWRENCE
BERKELEY LABORATORY
JUL 23 1984
LIBRARY AND
DOCUMENTS SECTION

Submitted to Physical Review Letters

τ LEPTON BRANCHING FRACTIONS

PEP-4 TPC Collaboration

June 1984



TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 6782.

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

LBL-18014

LBL-18014

τ Lepton Branching Fractions

H. Aihara, M. Alston-Garnjost, D.H. Badtke, J.A. Bakken, A.Barbaro-Galtieri, A.V.Barnes, B.A.Barnett, B.J.Blumenfeld, A.D.Bross, C.D.Buchanan, O.Chamberlain, J.Chiba, C-Y.Chien, A.R. Clark, A. Cordier, O.I. Dahl, C.T. Day, K.A. Derby, P.H.Eberhard, D.L.Fancher, H.Fujii, T.Fujii, B.Gabioud, J.W.Gary, W.Gorn, N.J.Hadley, J.M.Hauptman, W.Hofmann, J.E. Huth, J. Hylen, H. Iwasaki, T. Kamae, R.W. Kenney, L.T.Kerth, R.I.Koda, R.R.Kofler, K.K.Kwong, J.G.Layter, C.S.Lindsey, S.C.Loken, X-Q.Lu, G.R.Lynch, L.Madansky, R.J.Madaras, R.M.Majka, P.S.Martin, K.Maruyama, J.N.Marx, J.A.J.Matthews, S.O.Melnikoff, W.Moses, P.Nemethy, D.R.Nygren, P.J.Oddone, D.A.Park, A.Pevsner, M.Pripstein, P.R.Robrish, M.T.Ronan, R.R.Ross, F.R.Rouse, R.R.Sauerwein, G.Shapiro, M.D.Shapiro, B.C.Shen, W.E.Slater, M.L.Stevenson, D.H. Stork, H.K. Ticho, N. Toge, R.F. van Daalen Wetters, G.J.VanDalen, R.van Tyen, M.R.Wayne, W.A.Wenzel, H.Yamamoto, M.Yamauchi, W-M.Zhang

Lawrence Berkeley Laboratory,
University of California, Berkeley, CA 94720,
University of California, Los Angeles, CA 90024,
University of California, Riverside, CA92521,
Johns Hopkins University, Baltimore, MD 21218,
University of Massachusetts, Amherst, MA 01003,
University of Tokyo, Tokyo 113, JAPAN.

June 1984

Submitted to Physical Review Letters

Abstract

From a study of τ decays using the TPC detector at PEP we have measured the one-, three- and five-charged-particle inclusive branching fractions of the τ to be (85.2±1.7)%, (14.8±1.7)% and <0.3%, respectively. Using particle identification by dE/dx in the TPC we have searched for charged K mesons in three-prong τ decays, and have set an upper limit on B($\tau^{\pm} \rightarrow K^{\pm}$ + 2 charged + neutrals) of 0.6%.

PACS numbers: 14.60.Jj, 13.35.+s


Tau leptons produced in e^+e^- annihilation are useful tools for testing various aspects of the standard model of electroweak interactions. In particular, since the τ is the only lepton massive enough to decay into hadrons, one can use the decays of the τ to learn about the couplings of the hadronic weak current to leptons. One characteristic feature of these τ decays is the small multiplicity of charged particles. We report here on the measurement of this multiplicity. Using K meson identification by dE/dx, we also set limits on the strangeness content of three-prong τ decays, which can give information on the Cabibbo-suppressed part of the weak current.

The results presented in this Letter are based on data collected with the PEP-4 Time Projection Chamber (TPC) detector at the Stanford Linear Accelerator Center e^+e^- storage ring PEP. The data correspond to an integrated luminosity of 77 pb^{-1} at a center-of-mass energy of 29 Gev.

The TPC is the central tracking chamber of the PEP-4 detector and has been described in detail elsewhere.[1,2] Charged particles are identified in the TPC through a simultaneous measurement of momentum and dE/dx energy loss. Ionization created by particles passing through the TPC drifts in axial electric and magnetic fields to arrays of proportional wires at the endcaps. The pulse heights from the wires provide up to 183 samples of ionization per track. The average dE/dx energy loss of the particle is defined as

the mean of the smallest 65% of the individual samples. The dE/dx resolution is 3.7%. Three-dimensional track coordinates are derived from signals induced on fifteen rows of segmented cathode pads beneath the sense wires, and are used to measure the momentum (p) of the particle with a resolution given by $(dp/p)^2=(0.06)^2+(0.035p)^2$, with p in Gev/c.

To measure the branching fractions of the τ decaying into one, three and five charged particles (B_1, B_3 and B_5) $\tau^+\tau^-$ events of the "1+3" and "1+5" topologies are selected:



For the selection of these event topologies, charged tracks with momenta greater than 300 MeV/c and angles greater than 30 degrees from the beam direction are accepted as "good" if i) the extrapolated orbit passes the nominal interaction point within 10 cm in the beam direction and within 5 cm in the plane perpendicular to the beams, and ii) the track is not identified as part of a geometrically reconstructed e^+e^- pair from photon conversions in the 0.2 r.l. of material before the TPC. An event is selected as a $\tau^+\tau^-$ candidate if it has 4 or 6 good tracks which satisfy the following criteria:

1) The total charge is zero.

2) At least 3 good tracks are each > 140 degrees from one of the other good tracks (called the "isolated" track).

3) The invariant mass of all the good tracks except the isolated one is < 2 Gev/c^2 (assuming pion rest masses).

4) The good tracks (except the isolated one) are not likely to be electrons, as determined by dE/dx measurements in the TPC.

In addition, for the event as a whole we require that

5) The scalar sum of the total visible charged momenta is a) > 4.5 Gev/c, and b) < 24.0 Gev/c.

6) The angle between the highest momentum track and any other good track is < 178 degrees.

7) The total number of reconstructed tracks is < 10.

Criteria 1 and 2 select the τ-pair topology. Background from multihadron events is reduced by the invariant mass criterion (3), the momentum sum criterion (5b) and the total multiplicity criterion (7). Radiative Bhabha and μ-pair events are reduced by the dE/dx criterion

(4) and the acolinearity criterion (6). Background from two-photon events is reduced by the topological criterion (2) and the momentum sum criterion (5a). Unreconstructed photon conversion pairs are reduced by the dE/dx criterion (4). Residual beam-gas events are eliminated by the topological criterion (2).

With these selection criteria we find 669 τ-pair candidates in the "1+3" topology, and 4 candidates in the "1+5" topology. No candidates were found in a "1+7" topology. Seven events in the "1+3" sample are eliminated as radiative Bhabhas since two of the four tracks in each event shower in the barrel electromagnetic calorimeter.[3] Two events are rejected as radiative μ-pairs because two of the four tracks traverse the full thickness of the muon system.[4] This leaves a "1+3" candidate sample of 660 events.

The detection efficiencies corresponding to the selection criteria are determined by modeling $\tau^+\tau^-$ production and decay using the α^3 QED cross-section[5] and a statistical model[6] adjusted to agree with published branching fractions.[7] The efficiency for detecting $\tau^+\tau^-$ events in the "1+3" ("1+5") topology is 25% (10%). The most important limitations on the total efficiency are the probabilities of 40% (20%) to have 4 (6) good tracks of zero total charge in the detector fiducial volume, 80% (75%) for the 140 degree topological criterion, and 90% (75%) for the

invariant mass cut. The trigger efficiency is estimated to be 96% by comparing several independent triggers. Radiative corrections increase the lowest-order total cross section by a factor of 1.32. These include vacuum polarization, vertex corrections and bremsstrahlung with photon energies up to the kinematic maximum.

A Monte Carlo simulation is used to calculate the contamination of the final event sample by multihadron, two-photon and radiative Bhabha events and by other τ-pair topologies. Based on the measured luminosity we estimate that the total background in the "1+3" ("1+5") event sample is 52.0±5.7 (5.7±1.7) events. The most important contributions are 20 (2.4) events from multihadron events, 13 (1.1) from $e^+e^-\to\tau^+\tau^-\to$"1+1" topology with unreconstructed photon conversion pairs and 9 (1.1) from $e^+e^-\to e^+e^-\tau^+\tau^-$.

We observe four τ-pair candidates in the "1+5" topology, while the estimated background is 5.7±1.7 events. Thus our signal is consistent with zero. Taking into account the statistical uncertainty in the observed number of events and the background, and the systematic uncertainty in the luminosity and detection efficiency, we calculate an upper limit on the τ five-charged-particle inclusive branching fraction

$$B_5 < 0.3\% \qquad (90\%\ \mathrm{CL}).$$

Our result is compatible with the limit $B_5<0.5\%$ from the MARK II collaboration[8] at PEP, but is in poor agreement with the measurement $B_5=(1.0\pm0.4)\%$ from the CELLO collaboration[9] at PETRA.

The number of events in the "1+3" topology is proportional to the product B_1B_3. Since we found that B_5 is consistent with zero, we assume that $B_1=1-B_3$. Combining the measured number of "1+3" events above background (608.0±26.3) with the luminosity, detection efficiency and theoretical cross section, we find that the τ three-charged-particle inclusive branching fraction is

$$B_3 = (14.8 \pm 0.9 \pm 1.5)\%.$$

The first error is statistical. The second error reflects the systematic uncertainty in the luminosity (7%), trigger efficiency (3%) and detection efficiency (4%). Our result agrees with recent values from MARK II[8] and CELLO[9]. However, earlier experiments (see Ref. 7 for a compilation) give values for B_3 about twice as large.

The 660 events in the "1+3" topology sample are used to measure the charged K meson content of three-prong τ decays. To reduce the background to the K signal, only tracks in three-prong decays having the same sign charge as the parent τ are considered. For Cabibbo-suppressed decays like $\tau^-\to K^-\pi^+\pi^-\nu_\tau$ the K always has the same sign as the τ because a τ^-, for example, decays via a W^-. This W^- couples to a

strange (s) quark, which then gives a K^-. For non-Cabibbo-suppressed decays like $\tau^- \to K^- K^+ \pi^- \nu_\tau$, the two kaons are of opposite charge as they come from a $s\bar{s}$ quark pair. Thus one kaon always has the same sign as the parent τ. Of the two same sign tracks from three-prong τ decays the one with the higher momentum is chosen, because the heavier particle (ie. the K) usually has a greater momentum. The resulting 660 tracks are then each subjected to the following cuts before the dE/dx analysis is used to identify kaons:

1) The track must not be near the edges of the endcap detectors, where the wire gain is lower and less well known.

2) The track must have at least 80 samples of ionization to insure a good dE/dx measurement in the TPC.

3) The track momentum must be in the range 3.3-10.0 Gev/c, above the region where the K and π dE/dx values overlap, and below the region where momentum smearing is a problem.

These cuts result in a sample of 224 tracks. The number of kaons in the sample is determined by a maximum-likelihood fit to the dE/dx distribution of these tracks, as described in detail in Ref. 1. The fit gives 6.5±3.6 kaons.

The K selection efficiency corresponding to the above criteria is determined using a Monte Carlo simulation of τ-pair production[5] and decay via $\tau^- \to K^- \pi^+ \pi^- \nu_\tau$ and $K^- K^+ \pi^- \nu_\tau$. The efficiency is 35% for each channel and is not significantly different for similar decay modes including neutral pions.

The background in the K sample from kaons in multihadron events is estimated using the LUND Monte Carlo to be 1.0±0.8 kaons. The background-subtracted signal of 5.5±3.7 kaons corresponds to a branching fraction for $\tau^\pm \to K^\pm$ + 2 charged + neutrals of (0.3±0.2±0.1)%, where the first error is statistical and the second error reflects the systematic uncertainty in the dE/dx fit, luminosity, trigger efficiency and K selection efficiency. Since the statistical significance of this result is not strong, we prefer to quote an upper limit

$$B(\tau^\pm \to K^\pm + 2 \text{ charged} + \text{neutrals}) < 0.6\% \quad (90\% \text{ CL}).$$

This is the first limit on the charged kaon content of three-prong τ decays. Measurements[10] of the charged kaon content of one-prong τ decays give values 2-3 times larger than this limit.

By using dE/dx to identify kaons and pions individually, in contrast with the statistical separation used above, we have searched for specific groups of charged particles that could contribute to the ($K^\pm$+2 charged) system

in three-prong τ decays, such as $K^-\pi^+\pi^-$, $\overline{K^{*0}}(892)\pi^-$, $K^-K^+\pi^-$, $K^{*0}(892)K^-$, Q(1280) and Q(1400). We find one candidate for $\tau^- \to K^{*0}(892)K^-$ + neutrals. Using a Monte Carlo simulation we estimate the background for this reaction to be about 0.05 events from $\tau^- \to \pi^-\pi^+\pi^-$ + neutrals and about 0.1 events from multihadron events. One event corresponds to a branching fraction for $\tau^- \to K^{*0}(892)K^-$ + neutrals of 0.2%. However, our conclusion is that we do not find a statistically significant signal for this channel or any of the other five individual channels, and that the upper limits we can set on their branching fractions are not more significant than our result for $B(\tau^\pm \to K^\pm$ + 2 charged + neutrals).

Our upper limit on the charged kaon content of three-prong τ decays can be compared with the calculations of Tsai.[11] In particular, he finds that the branching fraction for $\tau \to Q\nu_\tau$, which can contribute to the ($K^\pm$ + 2 charged) system in three-prong τ decays, is given by

$$B(\tau \to Q\nu_\tau) = (M_{K^*}/M_Q)^2 (\tan^2\theta_c) B(\tau \to \rho\nu_\tau)(F_Q/F_\rho)$$

where $F_i = [1-(M_i/M_\tau)^2]^2[1+2(M_i/M_\tau)^2]$ and θ_c is the Cabibbo angle. Using $\tan^2\theta_c = 0.056$ and $B(\tau \to \rho\nu_\tau) = 22.1\%$,[7] one finds that $B(\tau \to Q\nu_\tau)$ is about 0.2-0.3%. This prediction for $B(\tau \to Q\nu_\tau)$ can be regarded as an approximate value for the sum of $B(\tau \to Q(1280)\nu_\tau)$ and $B(\tau \to Q(1400)\nu_\tau)$, and is consistent with our upper limit on the charged kaon content of three-prong τ

decays.

In summary, we have measured the one-, three- and five-charged-particle inclusive branching fractions of the τ, and have set an upper limit on the charged K meson content of three-prong τ decays.

We acknowledge the efforts of the PEP staff, and of the engineers, programmers and technicians of the collaborating institutions who made this work possible. This work was supported by the Department of Energy under contract numbers DE-AC03-76SF00098, DE-AM03-76SF00034, and DE-AC02-76ER03330, the National Science Foundation, and the Joint Japan-U.S. Collaboration in High Energy Physics.

References

1. PEP-4 TPC Collaboration, H.Aihara et al., Phys.Rev.Lett. 52, 577 (1984).

2. PEP-4 TPC Collaboration, H.Aihara et al., IEEE Trans. Nucl.Sci. NS30, 63,76,162 (1983).

3. PEP-4 TPC Collaboration, H.Aihara et al., Nucl.Instr.and Meth. 217, 259 (1983), and IEEE Trans.Nucl.Sci. NS30, 117 (1983).

4. PEP-4 TPC Collaboration, H.Aihara et al., IEEE Trans. Nucl.Sci. NS30, 67 (1983).

5. F.A.Berends and R.Kleiss, Nucl.Phys. B177, 237 (1981).

6. T.Sjöstrand, Comput.Phys.Commun. 27, 243 (1982).

7. Particle Data Group, C.G.Wohl et al., Rev.Mod.Phys. 56, No. 2, Part II (1984)

8. MARK II Collaboration, C.A.Blocker et al., Phys.Rev.Lett. 49, 1369 (1982).

9. CELLO Collaboration, H.J.Behrend et al., Phys.Lett. 114B, 282 (1982).

10. MARK II Collaboration, J.M.Dorfan et al., Phys.Rev.Lett. 46, 215 (1981); MARK II Collaboration, C.A.Blocker et al., Phys.Rev.Lett. 48, 1586 (1982); DELCO Collaboration, G.B.Mills et al., Phys.Rev.Lett. 52, 1944 (1984).

11. Y.S. Tsai, Phys.Rev. D4, 2821 (1971).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720