

Lawrence Berkeley National Laboratory

Recent Work

Title

TWO-PHOTON DECAY AND LIFETIME OF THE $22s_{1/2}$ STATE OF HYDROGEN-LIKE ARGON

Permalink

<https://escholarship.org/uc/item/1rg7t83z>

Authors

Schmieder, Robert W.
Marrus, Richard.

Publication Date

1970-09-01

C 23

TWO-PHOTON DECAY AND LIFETIME OF THE
 $2^2s_{1/2}$ STATE OF HYDROGEN-LIKE ARGON

RECEIVED
LAWRENCE
RADIATION LABORATORY

NOV 2 1970

Robert W. Schmieder and Richard Marrus

LIBRARY AND
DOCUMENTS SECTION

September 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

C 23

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

TWO-PHOTON DECAY AND LIFETIME OF THE $2^2s_{1/2}$ STATE OF HYDROGEN-LIKE ARGON*

Robert W. Schmieder and Richard Marrus

Lawrence Radiation Laboratory
and Department of Physics
University of California
Berkeley, California 94720

September 1970

Abstract:

The lifetime of the $2^2s_{1/2}$ state of the hydrogen-like atom Ar XVIII has been measured by direct observation of spontaneous two-photon decay in a beam-foil time-of-flight experiment. Identification of the two-photon mode was made using photon counting techniques to observe the single-photon continuum, and the peak resulting from summing the energies of photon pairs detected in coincidence. The measured lifetime, $\tau(2^2s_{1/2}) = 3.54(25) \times 10^{-9}$ sec (95% confidence) is in excellent agreement with theoretical predictions.

The decay of the $2^2s_{1/2}$ state of hydrogen-like atoms has been a subject of long-standing theoretical interest. It was first shown by Breit and Teller¹ in 1940 that this state should decay to the $1^2s_{1/2}$ ground state primarily by the simultaneous emission of two photons. In this process, which we designate $2E1$, the emission spectrum is a continuum, and the sum of the energies of the two photons equals the $1s$ - $2s$ energy separation. Spitzer and Greenstein² invoked this mechanism to explain the continuous spectrum of planetary nebulae, and using non-relativistic theory, they accurately computed the predicted spectrum

and lifetime of metastable hydrogen. Later, Shapiro and Breit³ obtained the decay rate for a hydrogen-like atom of atomic number Z :

$$A_{2E1} (2^2s_{1/2} - 1^2s_{1/2}) = 8.226 Z^6 \text{ sec}^{-1}, \quad (1)$$

which agrees with Spitzer and Greenstein for $Z = 1$. Recently, a closed form expression for the spectral distribution has been obtained⁴ and several authors⁵ have theoretically treated the two-photon decay mode in the 2^1S_0 and 2^3S_1 states of the helium isoelectronic sequence.

All of these calculations are non-relativistic, and consequently are accurate only for small Z . Relativistic effects make it possible for the $2^2s_{1/2}$ state to decay to the $1^2s_{1/2}$ state by magnetic dipole (M1) radiation, a process that vanishes in the non-relativistic approximation. This mechanism also was first noted by Breit and Teller, and recent calculations by Drake⁶ and by Schwartz⁷ reveal that to relative accuracy $1/Z$, the M1 rate is

$$A_{M1} (2^2s_{1/2} - 1^2s_{1/2}) = 2.50 \times 10^{-6} Z^{10} \text{ sec}^{-1}, \quad (2)$$

and hence for $Z = 18$ contribute about 4% of the total decay probability of the $2^2s_{1/2}$ state.

Experimentally, the two photon decay mode was first observed in He II by Lipeles, Novick, and Tolk,⁸ who reported detection of coincidences and angular distribution measurements consistent with the $1 + \cos^2\theta$ prediction. Rough spectral measurements made with broad-band crystal filters also have been reported.⁹ A continuous spectrum observed in a plasma¹⁰ has been attributed to the two-photon decay of the 2^1S_0 state of helium-like atom Ne IX. The relativistic magnetic dipole mode has been observed recently in the solar corona and

in the laboratory.^{12,13} Two recent measurements of the 2^1S_0 lifetime in He I, which presumably decay by two-photon emission, have been reported,¹⁵ but there has so far been no measurement of the lifetime of the $2^2s_{1/2}$ state of any hydrogen-like atom.

In this letter we report the direct observation of the two-photon decay mode in the hydrogen-like atom Ar XVIII using coincident photon counting techniques and the measurement of the lifetime of the $2^2s_{1/2}$ state, using the beam-foil time-of-flight method. The result is

$$\tau(2^2s_{1/2}) = 3.54(25) \times 10^{-9} \text{ sec} \quad , \quad (3)$$

where the error indicates 95% confidence.

The apparatus used in this measurement has been described in previous communications,¹²⁻¹⁴ and only a brief summary is given here. Ions of ^{40}Ar in the +14 charge state are accelerated in the Berkeley HILAC to an energy of 412 MeV ($\beta = v/c = 0.148$) and passed through a thin foil, from which they emerge distributed among the +16 (helium-like), +17 (hydrogen-like), and +18 (fully stripped) charge states. A significant fraction of the +16 and +17 ions may emerge highly excited, but they undergo fast radiative or non-radiative de-excitation to the ground or metastable states. The (forbidden) radiative decay of the metastable states is detected in flight a few tens of cm downstream of the foil by a pair of Si(Li) solid state x-ray detectors placed symmetrically perpendicular to the beam and about 2 cm from it. The photons were detected both singly and in coincidence, using standard high-rate coincidence circuitry with a resolving time $2\tau \sim 1 \mu\text{sec}$. In the singles mode, the energy E_1, E_2 of every detected photon in each detector was recorded. In the coincidence mode, the

detection of two photons in separate detectors within a time interval $|T_1 - T_2| \lesssim 5 \mu\text{sec}$ was defined as a "coincidence," and resulted in the storage of E1, E2, E1 + E2, and T1 - T2. With a typical beam current of 1 namp, single counting rates were typically $10^2 - 10^3 \text{ sec}^{-1}$, and coincidence rates were $0.1 - 1 \text{ sec}^{-1}$.

Since the lifetimes and spectra of the two-photon decays from 2^1S_0 and $2^2S_{1/2}$ states are similar, it was necessary to discriminate against the helium-like atoms in favor of the hydrogen-like atoms of interest here. To this end, a two-foil technique was used. A thick ($> 100 \mu\text{g}/\text{cm}^2$) beryllium foil capable of producing near charge equilibrium in the beam was placed ahead of the steering magnet (at position A in Fig. 1 of Ref. 13). The steering magnet was then set so that only fully-stripped ions were passed into our apparatus. The beam was then passed through an extremely thin ($< 10 \mu\text{g}/\text{cm}^2$) carbon foil, much thinner than is necessary to produce charge equilibrium. Since the capture of two or more electrons in this thin foil is a less likely process than the capture of one electron, the resulting beam has a substantially higher ratio of +17/+16 than with the near charge-equilibrated beam.

A typical singles spectrum taken in this way, using a foil-detector separation of 25 cm, is shown in Fig. 1. It consists of a broad continuum between the detector threshold and $E \sim 3 \text{ keV}$, in general agreement with the predictions.^{2,4} The observed spectrum also includes the composite effects of absolute detector efficiency (the step at 1.84 keV is due to absorption in the detector at the K-edge of silicon), detector resolution ($\cong 0.2 \text{ keV}$), doppler broadening due to the large acceptance angle of the detectors, low energy x-rays, electronic noise, high energy background, and contributions from the decays of

any helium-like ions that might be present. Although the correction of the data for all these effects is not yet complete, these data clearly verify the qualitative nature of the two-photon spectrum.

The purity of the beam as mostly hydrogen-like atoms can be assessed in several ways: (1) by comparing spectra like Fig. 1 with similar spectra taken with a nearly charge-equilibrated beam (measured to be 25% (+16), 50% (+17), 25% (+18)); (2) by observing the spectrum at very large foil-detector separations where only the magnetic dipole line $^{13} 2^3S_1 - 1^1S_0$ in Ar XVII is present; (3) by measuring the sum energy of coincident photons (= 3.10 keV for Ar XVII, 3.30 keV for Ar XVIII). Using these methods, we estimate that the contribution of helium-like atoms to the two photon spectrum for the conditions of Fig. 1 is less than a few percent.

To further verify the two-photon nature of the observed continuum, the detectors were operated in the coincidence mode, so that only coincident events (defined above) were stored. Figure 2a is a plot of the number of events versus the time delay $T_1 - T_2$ between the two photons. The zero in this plot was generated by introducing a fixed delay in one detector; it was calibrated using a pulse generator to simulate a true coincidence. The peak in the time spectrum of Fig. 2a is strong evidence that real coincident events are being observed.

The spectra observed as coincidences are shown in Figs. 2b and 2c. These spectra represent only the true coincidences appearing under the peak of the time spectrum in Fig. 2a. The contribution to that peak by accidental coincidences was removed by subtracting the events occurring away from the peak (presumably all accidentals), suitably normalized. The main difference between the singles spectra in Figs. 1 and 2b is that the coincidence mode symmetrically discriminates

against the ends of the continuum, thus peaking it more strongly at the center. The spectrum of the sum energy $E_1 + E_2$ observed as true coincidences is shown in Fig. 2c. Except for residual noise due to accidentals which have been removed, the spectrum is a single strong peak with a width roughly equal to the system resolution, indicating that this peak represents a single line. That this line falls at 3.3 keV is also strong evidence of the two-photon mode in Ar XVIII and not Ar XVII. The fact that the singles spectrum observed in singles mode (Fig. 1) is the same as that observed in coincidence mode (Fig. 2b), and the association of the latter (via Figs. 2a, 2c) with the $2E_1$ decay mode in Ar XVIII, permits measuring the lifetime in the singles mode.

The decay of the $2^2s_{1/2}$ state was observed by varying the separation between the foil and the detectors. A set of spectra like Fig. 1 was taken for various separations (normalized to a fixed amount of integrated beam current), and the total number of counts in the interval $0.75 \leq E \leq 2.5$ keV was obtained. A plot of the normalized count rate versus distance is shown in Fig. 3. At large distances the count rate levels off due to background and this is subtracted off to leave the pure $2^2s_{1/2}$ decay. From the measured decay length and known beam velocity, the mean $(1/e)$ lifetime was determined. After a 1% correction for the relativistic time dilation, the proper mean lifetime was found to be

$$\tau(2^2s_{1/2}) = 3.54(25) \times 10^{-9} \text{ sec.}$$

This result is in excellent agreement with the theoretical predictions. The non-relativistic formula (1) predicts the value $\tau(2^2s_{1/2}) = 3.57 \times 10^{-9}$ sec. If we combine this result with formula (2) for the relativistic magnetic dipole decay using

$$\frac{1}{\tau} = A_{2E1} + A_{M1} \quad , \quad (4)$$

the result $\tau(2^2s_{1/2}) = 3.47 \times 10^{-9}$ sec is obtained. Thus, our experimental results are in good agreement with the non-relativistic calculation.

We wish to again express our gratitude to A. Ghiorso, R. M. Diamond, F. Goulding, F. Grobelch, E. Lampo, J. Walton, D. MacDonald, and W. Davis for their encouragement and assistance in this work. G. W. F. Drake and C. Schwartz kindly informed us of their work prior to publication.

FOOTNOTES AND REFERENCES

* Work performed under the auspices of the U. S. Atomic Energy Commission.


1. G. Breit and E. Teller, *Astrophys. J.* 91, 215 (1940).
2. L. Spitzer and J. L. Greenstein, *Astrophys. J.* 114, 407 (1951).
3. J. Shapiro and G. Breit, *Phys. Rev.* 113, 179 (1959).
4. B. A. Zon and L. P. Rapoport, *JETP Letters* 7, 52 (1968).
5. G. A. Victor and A. Dalgarno, *Phys. Rev. Letters* 18, 1105 (1967); G. W. F. Drake and A. Dalgarno, *Astrophys. J.* 152, L121 (1968); G. W. F. Drake, G. A. Victor, and A. Dalgarno, *Phys. Rev.* 180, 25 (1969); O. Bely, *J. Phys. B* 1, 718 (1968); O. Bely and P. Faucher, *Astron. and Astrophys.* 1, 37 (1969).
6. G. W. F. Drake, University of Windsor (to be published).
7. C. Schwartz, University of California, (private communication).
8. M. Lipeles, R. Novick, and N. Tolk, *Phys. Rev. Letters* 15, 690 (1965).
9. C. J. Artura, N. Tolk, and R. Novick, *Astrophys. J.* 157, L181 (1969).
10. R. C. Elton, L. J. Palumbo, and H. R. Griem, *Phys. Rev. Letters* 20, 783 (1968).
11. A. H. Gabriel and C. Jordan, *Nature* 221, 947 (1969); W. M. Neupert and M. Swartz, *Astrophys. J.* 160, L189 (1970).
12. R. Marrus and R. W. Schmieder, *Phys. Letters* 32A, 431 (1970).
13. R. W. Schmieder and R. Marrus, (to be published).
14. R. Marrus and R. W. Schmieder, (to be published).
15. A. S. Pearl, *Phys. Rev. Letters* 24, 703 (1970); R. S. Van Dyke, Jr., C. E. Johnson, and H. A. Shugart, University of California (to be published).

FIGURE CAPTIONS

Fig. 1. Typical energy spectrum observed in singles mode. This uncorrected spectrum is predominantly the two-photon decay of metastable Ar XVIII. The edge at 1.84 keV is an instrumental effect (see text). The detector resolution is indicated.

Fig. 2. Spectra observed in coincidence mode: (a) Time difference between coincident photon pairs. (b) Single photon energy spectrum, including only events which participated in a true coincidence. (c) Sum of energies of photons participating in a true coincidence.

Fig. 3. Decay curve obtained by recording the singles count rate versus foil-detector separation. At large distances the observed counts level off to a constant value, which was subtracted from each point to give the two-photon counts. The error bars indicate the statistical error.



XBL 709-6649

Fig. 3

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720