

UC Merced

Proceedings of the Annual Meeting of the Cognitive Science Society

Title

Hierarchical temporal organization of speech in children and adolescents whostutter

Permalink

<https://escholarship.org/uc/item/1tg2p07g>

Journal

Proceedings of the Annual Meeting of the Cognitive Science Society, 42(0)

Authors

Franke, Mona

Kello, Christopher

Falk, Simone

Publication Date

2020

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Hierarchical temporal organization of speech in children and adolescents who stutter

Mona Franke

Ludwig-Maximilians-University, Munich, Germany

Christopher Kello

University of California Merced, Merced, California, United States

Simone Falk

Universit  de Montral, Montreal, Quebec, Canada

Abstract

With 10 to 20 sounds per second, fluent speech requires extremely skilled motor coordination. Therefore, young speakers with an immature or malfunctioning speech production system may be particularly challenged by the temporal aspects of fluent speech. In the present study, we examine nested temporal bout structure (Abney et al., 2014) to investigate how young speakers (children 9-12; adolescents 13-17 years old) who do and do not stutter might differ in their temporal organization of speech during reading. Allan Factor analyses show that nested clustering of peak amplitudes at short time-scales (≤ 300 ms) differs between children and adolescents, pointing to developmental differences in the temporal organization of syllabic structure. Greater nested clustering at longer timescales (≥ 300 ms – 10 s) was characteristic of stuttering, particularly in adolescents whose stutter risks to persist into adulthood. We discuss these findings in light of theories of stuttering and the acquisition of fluent speech