Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

PRODUCTION OF CHARMONIUM STATES BY RESONANT $p\bar{p}$ ANNIHILATION

Permalink

<https://escholarship.org/uc/item/1tj3414b>

Author

Cahn, Robert N.

Publication Date

1980-02-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division

Submitted to Physics Letters B

PRODUCTION OF CHARMONIUM STATES BY RESONANT
 $p\bar{p}$ ANNIHILATION

Robert N. Cahn and Mahiko Suzuki

February 1980

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 21 1980

LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*

LBL-10505 C.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

PRODUCTION OF CHARMONIUM STATES BY RESONANT $p\bar{p}$ ANNIHILATION*

Robert N. Cahn[†]

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

and

Mahiko Suzuki

Department of Physics and Lawrence Berkeley Laboratory

University of California
Berkeley, California, 94720

ABSTRACT

Cross sections are estimated for the production of charmonium states in $p\bar{p}$ annihilation on resonance. For processes like $p\bar{p} \rightarrow \eta_c \rightarrow \gamma\gamma$ the cross sections are about 1 nb. Non-resonant contributions are estimated and found to be much smaller. Similar rates are found for $p\bar{p} \rightarrow \eta_c' \rightarrow \gamma\gamma$ and $p\bar{p} \rightarrow {}^1P_1 \rightarrow \eta_c \gamma \rightarrow \gamma\gamma\gamma$. These may thus furnish a unique means of observing these charmonium states. Upsilononic states might be observable in colliding ring $p\bar{p}$ experiments, but the cross sections are likely to be in the picobarn range.

* This work was supported by the U.S. Department of Energy under Contract W-7405-ENG-48 and by the National Science Foundation under Contract PHY-77-23512.

[†] A. P. Sloan Foundation Fellow.

Any resonance which can be produced in e^+e^- annihilation, can, in principle be produced as well by $p\bar{p}$ annihilation. In addition, states with J^{PC} differing from that of the photon, 1^{--} , may also be produced in $p\bar{p}$ annihilation. Resonant production of the charmonium states $1^1S_0(\eta_c)$, $2^1S_0(\eta_c')$, and 1^1P_1 would be an attractive possibility with $\bar{p}$ beams of about 4-5 GeV. Of course the coupling of these states to the $p\bar{p}$ system is likely to be feeble and the large background of purely conventional hadronic events poses a serious problem.

Here we pursue the suggestion¹ that the hadronic background can be overcome by looking for specific few body final states such as $p\bar{p} \rightarrow \eta_c \rightarrow \gamma\gamma$. It is straightforward to estimate the cross section for this and similar processes. They are in the nanobarn range, as we shall show below. The more difficult task is to estimate the non-resonant sources of the same final state. This we shall do using experimental data. Both vector meson dominance² and Regge treatments indicate that nonresonant cross sections are small. Our conclusion is that the non-resonant sources are probably not too serious a problem.

Certainly there are substantial difficulties besides the ones we evaluate here. The resonant cross sections are down by $10^7 - 10^8$ from the total hadronic cross section. Most of this background can be eliminated by vetoing events with charged particles present and by excluding events where there are any particles, charged or neutral, near the forward or backward directions. The resonant processes, of course, yield events with isotropic or nearly isotropic distributions in the center of mass, unlike the typical hadronic

events which peak in the forward and backward directions. The severe kinematical constraints satisfied by the events we seek could be used to reduce the background.

The cross section for production of a spin J -resonance in $p\bar{p}$ annihilation is given at the peak ($\sqrt{s} = m_{\text{res}}$) by

$$\sigma_{\text{res}} = \frac{4\pi}{s - 4m_p^2} (2J + 1) B_i, \quad (1)$$

where B_i is the branching ratio of the resonance into $p\bar{p}$. In practice, $\bar{p}$ beams are not monoenergetic and a correction must be made for this. The spread in beam energy produces a spread in the center of mass energy, ΔW . If ΔW is much bigger than the resonance width, Γ_t , the effective cross section is

$$\sigma_{\text{eff}} = \sigma_{\text{res}} \frac{\Gamma_t}{\Delta W}. \quad (2)$$

We estimate the branching ratio of the charmonium states in $p\bar{p}$ using the data for $\psi \rightarrow p\bar{p}$. The measured branching³ ratio is $(0.21 \pm 0.02)\%$. However, the total width for the ψ includes decays through a single virtual photon which account for about 30% of the width⁴. Since these decays are absent for the particles of interest here, we use a branching ratio of about 3×10^{-3} . In this way we estimate

$$\begin{aligned} \sigma_{\text{res}}(p\bar{p} \rightarrow \eta_c) &\approx 3\mu\text{b}, \\ \sigma_{\text{res}}(p\bar{p} \rightarrow \eta_c') &\approx 1.8\mu\text{b} \\ \sigma_{\text{res}}(p\bar{p} \rightarrow {}^1P_1(c\bar{c})) &\approx 5.4\mu\text{b} \end{aligned} \quad (3)$$

We have taken the values⁵

$$\begin{aligned} m(\eta_c) &= 2.98 \text{ GeV}, \\ m(\eta_c') &\approx 3.50 \text{ GeV}, \\ m({}^1P_1) &\approx 3.50 \text{ GeV}. \end{aligned} \quad (4)$$

The production cross sections must now be multiplied by the branching ratios to the final states of interest. For the η_c , there are indications⁶ that the total width is about 20 MeV, while the $\gamma\gamma$ decay is expected⁷ to have a partial width of about 10keV. On this basis, we assume

$$\begin{aligned} B(\eta_c \rightarrow \gamma\gamma) &= 0.5 \times 10^{-3}, \\ B(\eta_c' \rightarrow \gamma\gamma) &= 0.5 \times 10^{-3}. \end{aligned} \quad (5)$$

Combining these rates with those in Eq. (3), we have the estimates

$$\begin{aligned} \sigma_{\text{res}}(p\bar{p} \rightarrow \eta_c \rightarrow \gamma\gamma) &\approx 1.5 \text{ nb}, \\ \sigma_{\text{res}}(p\bar{p} \rightarrow \eta_c' \rightarrow \gamma\gamma) &\approx 0.9 \text{ nb}. \end{aligned} \quad (6)$$

The competing non-resonant processes will be estimated later.

The 1P_1 resonance has the signature ${}^1P_1 \rightarrow \eta_c \rightarrow \gamma\gamma\gamma$. The total width of 1P_1 is expected to be roughly in the same range as those of the ψ and ψ' since it decays through three gluons, or a little wider because of a logarithmic threshold enhancement⁸.

The branching ratio, $B(^1P_1 \rightarrow \gamma n_c)$ can be estimated fairly reliably from a comparison with the measured branching ratio⁹ $B(^3P_1 \rightarrow \psi \gamma) = (23.4 \pm 0.8)\%$. Using the theoretical relation, $\Gamma(^1P_1) = (5/6) \Gamma(^3P_1)$, we find

$$B(^1P_1 \rightarrow \gamma n_c) \approx \frac{6}{5} B(^3P_1 \rightarrow \psi \gamma) \left(\frac{p(n_c)}{p(\psi)} \right)^3 \approx 28\%. \quad (7)$$

Together with Eqs. (3) and (5), this yields

$$\sigma_{\text{res}}(^1P_1 \rightarrow \gamma n_c \rightarrow \gamma \gamma \gamma) \approx 0.8 \text{ nb}. \quad (8)$$

It must be borne in mind that the small total width of 1P_1 may result in a much smaller effective peak cross section.

If the proton were structureless, like the electron, we could calculate the non-resonant process $p\bar{p} \rightarrow \gamma \gamma$ using quantum electrodynamics. With just a Dirac coupling, the result is that the total cross section at $s = m(n_c)^2$ is about 30 nb. This is clearly an overestimate since we have ignored the tendency of the proton to emit additional pions when it undergoes a large momentum transfer. Indeed, it is well known that $p\bar{p}$ annihilation at $s \approx 10 \text{ GeV}^2$ results in final states with only two particles very infrequently.

This can be made quantitative using the vector dominance model.¹⁰ A bubble chamber experiment with 5.7 GeV incident antiprotons found an upper limit, at 90% confidence level, of $10 \mu\text{b}$ for the reaction $p\bar{p} \rightarrow \rho^0 \rho^0$. This cross section would lead to cross section for $p\bar{p} \rightarrow \gamma \gamma$ of

$$\begin{aligned} \sigma(p\bar{p} \rightarrow \gamma \gamma) &= \sigma(p\bar{p} \rightarrow \rho^0 \rho^0) \cdot \alpha^2 \left(\frac{4\pi}{g_\rho} \right)^2 \\ &< 10 \mu\text{b} \times \alpha^2 \times \frac{1}{2.3^2} = 0.1 \text{ nb} \end{aligned} \quad (9)$$

at $s \approx 12 \text{ GeV}^2$. Not only is this much smaller than the cross sections estimated in Eq. (6), but furthermore, the angular distribution is likely to be peaked in the forward and backward directions, unlike the distribution from the resonant process.

As a second means of estimating the non-resonant cross section, we use a very crude Regge model. The scattering $p\bar{p} \rightarrow \gamma \gamma$ is presumed to occur through N or Δ Regge exchange. This same exchange controls backward πp scattering and backward pion-photoproduction. Using these two processes, factorization and Δ -dominance we can predict the cross section for $\gamma p \rightarrow p \gamma$:

$$\frac{d\sigma}{du}(\gamma p \rightarrow p \gamma) = 3 \frac{\left[\frac{d\sigma}{du}(\gamma p \rightarrow p \pi^0) \right]^2}{\frac{d\sigma}{du}(\pi^- p \rightarrow p \pi^-)}. \quad (10)$$

The factor of three arises from the Clebsch-Gordon relation $g(\Delta^+ \pi^0 p) = \sqrt{\frac{2}{3}} g(\Delta^{++} p)$ together with a correction for the presence of two photon polarizations. The minor phase space correction has been ignored. The relation follows if the process is dominated by Δ Regge exchange. Using the data of Refs. 11 and 12, we estimate that for $P_{\text{LAB}} = 6 \text{ GeV}$

$$\frac{d\sigma}{du}(\gamma p \rightarrow p \gamma) \approx 0.08 \times e^{-2.2|u|} \text{ nb GeV}^{-2} \quad (11)$$

where u is measured in GeV^2 . By crossing symmetry (again neglecting a small phase space correction),

$$\frac{d\sigma}{dt}(\bar{p}p \rightarrow \gamma\gamma) \approx 0.08 e^{-2.2|t|_{\text{nb}}} \text{GeV}^{-2}. \quad (12)$$

A similar analysis can be made with $\pi^+p \rightarrow p\pi^+$ scattering, with similar results. Integrating Eq. (12) we find

$$\sigma(\bar{p}p \rightarrow \gamma\gamma) \approx 0.04 \text{ nb}. \quad (13)$$

This estimate is appropriate for $s \approx 12 \text{ GeV}^2$. To estimate the cross section at the η_c where $s \approx 9 \text{ GeV}^2$ we must know the energy dependence of the cross section. The Regge form, $\sigma \sim s^{2\alpha-2}$, with $\alpha = 0.2$, gives a fairly small correction. In fact the data suggest a rather more dramatic s^{-3} dependence. To be conservative we use this estimate. We have then

$$\sigma(\bar{p}p \rightarrow \gamma\gamma) \approx 0.1 \text{ nb} \quad (14)$$

at $s = 9 \text{ GeV}^2$. This crude estimate is in reasonable agreement with the vector dominance estimate. Although neither ought to be trusted to more than a factor of two, they are reassuringly small, especially since most of the cross section will be peaked in the forward and backward directions.

The background for $\bar{p}p \rightarrow {}^1P_1 \rightarrow \gamma\gamma\gamma$ comes from several sources. The purely electromagnetic process analogous to $e^+e^- \rightarrow \gamma\gamma\gamma$ would yield about 0.2 nb if the proton form factors were ignored. In

fact, the form factors will suppress the cross section considerably.

A second source of $\gamma\gamma\gamma$ final states is the two-body process $\bar{p}p \rightarrow \gamma\pi^0$, and its analogues, $\bar{p}p \rightarrow \gamma\eta$ and $\bar{p}p \rightarrow \gamma\eta'$. These cross sections can also be estimated using Regge analysis. Again we use the measured cross section¹¹ for $\gamma p \rightarrow p\pi^0$. Using crossing and adding together the forward and backward π^0 production we have

$$\begin{aligned} \frac{d\sigma}{dt}(\bar{p}p \rightarrow \pi^0\gamma) + \frac{d\sigma}{dt}(\bar{p}p \rightarrow \gamma\pi^0) &= 2 \frac{d\sigma}{du}(\gamma p \rightarrow p\pi^0) \\ &\approx 15 \times e^{-2.8|t|_{\text{nb}}} \text{GeV}^{-2} \end{aligned}$$

at $p_{\text{LAB}} = 6 \text{ GeV}$, and

$$\sigma(\bar{p}p \rightarrow \pi^0\gamma) \approx 5 \text{ nb}, \text{ at } \sqrt{s} = 3.5 \text{ GeV}. \quad (14)$$

This is consistent with $\sigma(\bar{p}p \rightarrow \pi^0\gamma) < 80 \text{ nb}$ at $p_{\text{LAB}} = 6 \text{ GeV}$, the limit obtained using vector dominance and the existing experimental upper limit¹³ on $\sigma(\bar{p}p \rightarrow \pi^0\rho^0)$. The η and η' have weaker couplings to $\bar{p}p$ and we estimate¹⁴ at $\sqrt{s} = 3.5 \text{ GeV}$

$$\sigma_{\text{tot}}(\bar{p}p \rightarrow \eta\gamma) \times B(\eta \rightarrow \gamma\gamma) \approx 0.07 \text{ nb}, \quad (15)$$

$$\sigma_{\text{tot}}(\bar{p}p \rightarrow \eta'\gamma) \times B(\eta' \rightarrow \gamma\gamma) \approx 0.05 \text{ nb}.$$

These events would have to be separated from the true ${}^1P_1 \rightarrow \eta_c \rightarrow \gamma\gamma\gamma$ signal by kinematic constraints. Once again, eliminating events with forward and backward production would suppress the background considerably.

An alternative to searching for neutral final states is to use the clear leptonic signature of ψ decay. For example, using Eq. (1) and the known leptonic branching ratios, we have

$$\begin{aligned}\sigma(p\bar{p} \rightarrow \psi \rightarrow \mu^+ \mu^-) &\approx 0.36\mu\text{b}, \\ \sigma(p\bar{p} \rightarrow \psi' \rightarrow \mu^+ \mu^-) &\approx 0.03\mu\text{b}.\end{aligned}\tag{16}$$

The competing backgrounds are negligible. It should be possible, as well, to observe $p\bar{p} \rightarrow \chi \rightarrow \gamma\psi \rightarrow \gamma \mu^+ \mu^-$. Estimated cross sections are given in Table I.

We conclude with some speculations on the possibility of observing upsilonic states using $p\bar{p}$ colliding rings. The branching ratios of the resonances into $p\bar{p}$ might well be lower by a factor of ten than those assumed for the charmonium states. In addition, the unitarity limit, Eq. (1), is lower by a factor of ten as well. Even if the branching ratio were 0.5×10^{-3} , the cross sections of interest would be about 10^{-35} cm^2 . While this would give a few events per hour given a luminosity of $10^{32} \text{ cm}^{-2} \text{ sec}^{-1}$, it is not clear that this is really practicable. It may be feasible to explore the 3P_1 and 3P_2 states of $(b\bar{b})$ through $\gamma\ell\bar{\ell}$ final states, since the analogous $(c\bar{c})$ states are predicted to have relatively large production cross sections.

In summary, we find that the resonant production cross sections for charmonium states in $p\bar{p}$ annihilation are large enough (in the nanobarn range) to permit observation, even for some states which have not been seen in e^+e^- annihilation. The non-resonant cross sections which compete are estimated to be smaller than the resonant cross sections. Estimates for the production of $(b\bar{b})$ states are less certain, but the cross-sections are considerably smaller, perhaps in the picobarn range. Even in the charmonium

case, extremely careful experiments will be required to find the events of interest which are only a few parts in 10^8 of the total cross section.

We would like to thank our colleagues M. Alston-Garnjost, R.W. Kenny, and T.P. Pun for bringing this problem to our attention and for their explanations of the attendant experimental problems. In addition, we would like to thank J. D. Jackson, H. Steiner, and F. Gilman for their suggestions.

REFERENCES

1. T. P. Pun, private communication.
2. We thank J. D. Jackson for reminding us of this tried and true procedure.
3. Particle Data Group, Review of Particle Properties, Phys. Lett. 75B (1978) 1.
4. Cited in particle Data Group compilation, Ref. 3. Included are the decays into e^+e^- , $\mu^+\mu^-$, and an additional amount equal to about twice the partial width into e^+e^- , corresponding to second order electromagnetic decays into hadrons.
5. The η_c mass is that found by the Crystal Ball Experiment, E. Bloom, Invited talk at 9th International Symposium on Lepton and Photon Interactions at High Energy, Batavia, IL, August 23-29, 1979.
6. A preliminary result of the Crystal Ball Group, given by T. Burnett at the Irvine Conference, November 30, 1979.
7. T. Appelquist and H. D. Politzer, Phys. Rev. D12, (1975) 1404.
8. R. Barbieri, R. Gratto, and E. Remiddi, Phys. Lett. 61B, (1976) 465.
9. Cited in Particle Data Group compilation, Ref. 3.
10. A. Accensi, et al., Physics Letters 20, (1966) 557.
11. D. Tompkins et al., Phys. Rev. Lett. 23 (1969) 725. V. Barger and P. Weiler, Phys. Letters 30B, (1969) 105.
12. A. Ashmore et al., Phys. Rev. Lett. 19, (1967) 460 and Phys. Rev. Lett. 21, (1968) 387; J. Orear et al., Phys. Rev. Lett. 21, (1968) 389; V. Barger and D. Cline, Phys. Rev. Lett. 21, (1968) 392.

11

TABLE I

Estimated peak resonant production cross sections for formation of charmonium states in $p\bar{p}$ annihilation. The background processes are non-resonant cross sections with the same final state as the resonant process. A dash in the final column indicates that the cross section is negligible.

Signature	Resonant Cross Section (nb)	Non-resonant process	Non-resonant Cross Section (nb)
$p\bar{p} \rightarrow \eta_c \rightarrow \gamma\gamma$	1.5	$p\bar{p} \rightarrow \gamma\gamma$	0.1
$p\bar{p} \rightarrow \eta_c' \rightarrow \gamma\gamma$	0.9	$p\bar{p} \rightarrow \gamma\gamma$	0.04
$p\bar{p} \rightarrow {}^1P_1 \rightarrow \gamma\eta_c \rightarrow \gamma\gamma\gamma$	0.8	$p\bar{p} \rightarrow \gamma\pi^0 \rightarrow \gamma\gamma\gamma$	5
		$\rightarrow \gamma\eta \rightarrow \gamma\gamma\gamma$	0.07
		$\rightarrow \gamma\eta' \rightarrow \gamma\gamma\gamma$	0.05
$p\bar{p} \rightarrow \psi \rightarrow \mu^+\mu^-$	400	$p\bar{p} \rightarrow \mu^+\mu^-$	-
$p\bar{p} \rightarrow \psi' \rightarrow \mu^+\mu^-$	30	$p\bar{p} \rightarrow \mu^+\mu^-$	-
$p\bar{p} \rightarrow X_0 \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	5	$p\bar{p} \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	-
$p\bar{p} \rightarrow X_1 \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	90	$p\bar{p} \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	-
$p\bar{p} \rightarrow X_2 \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	100	$p\bar{p} \rightarrow \gamma\psi \rightarrow \gamma\mu^+\mu^-$	-

13. T. Ferbel et al., Phys. Rev. 173, (1968) 1307.
14. Only N exchange contributes to the processes. SU(3) symmetry gives the relation between the Regge residues,

$$\beta^2(\bar{p}p\eta) = \frac{1}{3} \left(\frac{D-3F}{D+F} \right)^2 \beta^2(\bar{p}p\pi^0),$$

where the D and F are the usual octet pseudoscalar meson baryon couplings.