

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

Targeting IGF2BP3 Enhances Anti-Leukemic Effects of Menin-MLL Inhibition in MLL-AF4 Leukemia

### Permalink

<https://escholarship.org/uc/item/1tj7v7p2>

### Journal

Blood, 142(Supplement 1)

### ISSN

0006-4971

### Authors

Lin, Tasha L  
Ritter, Alexander J  
Jaiswal, Amit K  
[et al.](#)

### Publication Date

2023-11-28

### DOI

10.1182/blood-2023-174813

Peer reviewed

# Targeting IGF2BP3 enhances antileukemic effects of menin-MLL inhibition in MLL-AF4 leukemia

Tasha L. Lin,<sup>1</sup> Amit K. Jaiswal,<sup>2</sup> Alexander J. Ritter,<sup>3</sup> Jenna Reppas,<sup>2</sup> Tiffany M. Tran,<sup>2,4</sup> Zachary T. Neeb,<sup>3</sup> Sol Katzman,<sup>5</sup> Michelle L. Thaxton,<sup>2</sup> Amanda Cohen,<sup>2</sup> Jeremy R. Sanford,<sup>3,5</sup> and Dinesh S. Rao<sup>2,6,7</sup>

<sup>1</sup>Division of Hematology and Oncology, Department of Medicine and <sup>2</sup>Department of Pathology and Laboratory Medicine, University of California, Los Angeles, Los Angeles, CA; <sup>3</sup>Department of Molecular, Cell and Developmental Biology and Center for Molecular Biology of RNA, University of California Santa Cruz, Santa Cruz, CA; <sup>4</sup>Department of Medicine, University of California, Los Angeles, Los Angeles, CA; <sup>5</sup>Center for Biomolecular Science & Engineering, University of California Santa Cruz, Santa Cruz, CA; <sup>6</sup>Jonsson Comprehensive Cancer Center and <sup>7</sup>Broad Stem Cell Research Center, University of California, Los Angeles, Los Angeles, CA

## Key Points

- Knockdown of IGF2BP3 sensitizes MLL-AF4 leukemia to menin-MLL inhibition.
- These data support the role of IGF2BP3 as an oncogenic amplifier in MLL-AF4 leukemogenesis and its therapeutic potential.

RNA-binding proteins (RBPs) are emerging as a novel class of therapeutic targets in cancer, including in leukemia, given their important role in posttranscriptional gene regulation, and have the unexplored potential to be combined with existing therapies. The RBP insulin-like growth factor 2 messenger RNA-binding protein 3 (IGF2BP3) has been found to be a critical regulator of MLL-AF4 leukemogenesis and represents a promising therapeutic target. Here, we study the combined effects of targeting IGF2BP3 and menin-MLL interaction in MLL-AF4-driven leukemia in vitro and in vivo, using genetic inhibition with CRISPR-Cas9-mediated deletion of *Igf2bp3* and pharmacologic inhibition of the menin-MLL interaction with multiple commercially available inhibitors. Depletion of *Igf2bp3* sensitized MLL-AF4 leukemia to the effects of menin-MLL inhibition on cell growth and leukemic initiating cells in vitro. Mechanistically, we found that both *Igf2bp3* depletion and menin-MLL inhibition led to increased differentiation in vitro and in vivo, seen in functional readouts and by gene expression analyses. IGF2BP3 knockdown had a greater effect on increasing survival and attenuating disease than pharmacologic menin-MLL inhibition with small molecule MI-503 alone and showed enhanced antileukemic effects in combination. Our work shows that IGF2BP3 is an oncogenic amplifier of MLL-AF4-mediated leukemogenesis and a potent therapeutic target, providing a paradigm for targeting leukemia at both the transcriptional and posttranscriptional level.

## Introduction

Leukemias driven by translocations in the mixed-lineage leukemia 1 gene (MLL1/KMT2A) represent a subgroup of particular interest because of their unique clinical and biological characteristics and poor prognosis. Chromosomal rearrangements in the MLL1 gene result in fusion proteins, with several partners that are part of the superelongation complex, leading to aberrant gene expression.<sup>1-3</sup> Comprehensive genomic studies of patient-derived MLL-rearranged leukemia cells show that these leukemias have very few genetic alterations,<sup>4,5</sup> suggesting that epigenetic and transcriptional

Submitted 5 July 2023; accepted 22 November 2023; prepublished online on *Blood Advances* First Edition 4 December 2023; final version published online 10 January 2024. <https://doi.org/10.1182/bloodadvances.2023011132>.

RNA-sequencing data are shared via deposit to the NCBI Short Read Archive repository (accession number PRJNA918326).

Plasmids, vectors, and other renewable resources included in the article are available on request from the corresponding author, Dinesh S. Rao ([drao@mednet.ucla.edu](mailto:drao@mednet.ucla.edu)).

The full-text version of this article contains a data supplement.

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

dysregulation driven by the fusion proteins are largely responsible for leukemogenesis. This is further borne out by the demonstration of the sufficiency of MLL fusion protein overexpression, for example, MLL-AF9 and MLL-Af4, to drive leukemia in both murine and human hematopoietic stem and progenitor cells (HSPCs).<sup>6-8</sup> MLL-leukemogenesis requires the interaction of the N-terminal end of the fusion protein with menin, in addition to its interactions with downstream chromatin modifiers and the superelongation complex.<sup>9-11</sup> Menin is a molecular adapter protein that in turn interacts with epigenetic regulators and cell signaling molecules, driving massive dysregulation of gene expression.

In addition to epigenetic and transcriptional mechanisms, the importance of posttranscriptional regulation by factors such as microRNAs and RNA-binding proteins (RBPs) has been increasingly appreciated in leukemogenesis. Our group implicated the oncofetal RBP insulin-like growth factor 2 messenger RNA (mRNA)-binding protein 3 (IGF2BP3) as a critical regulator of MLL-AF4-mediated leukemogenesis.<sup>12,13</sup> We found that MLL-AF4 transcriptionally induced *Igf2bp3* and that IGF2BP3 regulates important leukemogenic transcripts, amplifying the aberrant gene expression initiated by MLL-AF4.<sup>13</sup> Furthermore, IGF2BP3 is required for the efficient initiation of MLL-Af4-driven leukemia and the function of leukemic initiating cells (LICs), thereby establishing a critical role of IGF2BP3 in the pathogenesis of MLL-driven leukemia and its potential as a promising therapeutic target.

Existing strategies for treating MLL-rearranged leukemia include chemotherapy, small molecule inhibitors, and CD19-directed therapies. Novel therapeutic strategies for MLL-rearranged leukemia have been directed toward targeting the menin-MLL interaction as well as various epigenetic factors.<sup>14,15</sup> Multiple potent small molecule inhibitors that have been developed to target the menin-MLL1 interaction, including MI-503, MI-463, MI-3454, and VTP-50469, have demonstrated impressive single-agent activity in preclinical studies, with derivatives entering phase 1 to 2 clinical trials.<sup>16-20</sup> Nonetheless, combinatorial strategies remain a cornerstone of cancer-directed therapy to combat issues with dose-limiting toxicities and drug resistance from single-agent therapy. Therefore, we sought to explore a novel therapeutic strategy of combinatorial targeting of leukemia at the transcriptional and posttranscriptional level. Our approach combined the pharmacologic inhibition of the menin-MLL interaction with the genetic inhibition of *IGF2BP3*.

Here, we tested the combined effects of IGF2BP3 knockdown via CRISPR/Cas9-mediated deletion and menin-MLL inhibition using commercially available menin-MLL inhibitors in human B-cell acute lymphoblastic leukemia (B-ALL) cell lines and a murine model for MLL-Af4-driven leukemia.<sup>6</sup> Depletion of IGF2BP3 sensitized MLL-AF4 leukemia to the effects of menin-MLL inhibition on cell growth and LICs. Mechanistically, we found that both knockdown of IGF2BP3 and menin-MLL inhibition led to increased differentiation of MLL-Af4 leukemia, with a more dramatic effect seen when both were combined. Concordantly, gene expression analyses demonstrated an upregulation of differentiation genes with MI-503 treatment and IGF2BP3 knockdown. Lastly, we found decreased leukemic engraftment and significantly increased survival with IGF2BP3 knockdown in comparison to MI-503 treatment and enhanced antileukemic effects with the combination. Together, our work supports the idea that IGF2BP3 is an oncogenic amplifier of MLL-AF4-mediated leukemogenesis, and IGF2BP3 represents an

exciting therapeutic target. Our data also support a novel combinatorial therapeutic strategy of targeting leukemia at both the transcriptional and posttranscriptional levels.

## Methods

### Cell culture

RS4;11 (American Type Culture Collection (ATCC) CRL-1873), NALM6 (ATCC CRL-3273), and SEM (DMZ-ACC 546) were cultured as previously described.<sup>21</sup> Immortalized MLL-Af4-transformed HSPCs derived from mouse bone marrow (BM; MLL-Af4 lineage-negative [ $\text{Lin}^-$ ] cells) were cultured in Iscove modified Dulbecco medium with 15% fetal bovine serum, supplemented with murine 100 ng/mL stem cell factor, 4 ng/mL murine interleukin-6, 50 ng/mL human Fms-like tyrosine kinase 3 ligand (FLT3L), and 50 ng/mL mouse thrombopoietin (mTPO).

### Plasmids and viral transduction

The MSCV-MLL-FLAG-Af4 plasmid was generously provided by Michael Thirman (University of Chicago) through a material transfer agreement.<sup>21</sup> Single-guide RNAs (sgRNAs) against mouse *Igf2bp3* and nontargeting (NT) guides were cloned into an inhouse MSCV-hu6-sgRNA-EFS-mCherry vector.<sup>21</sup> Single-guide RNAs against human *IGF2BP3* and nontargeting guides were cloned into pLKO.sgRNA.EFS.tRFP (Addgene 57823). Sequences are available in supplemental Table 1. Generation of retroviral and lentiviral supernatants and viral transduction were performed according to standard procedures.

### CRISPR/Cas9-mediated deletion of IGF2BP3 in cell lines

SEM, RS4;11, and NALM6 were depleted for IGF2BP3 using lentiviral delivery of CRISPR/Cas9 components in a 2-vector system as previously described.<sup>21</sup> MLL-Af4  $\text{Lin}^-$  cells were initially isolated from Cas9-green fluorescent protein (Cas9-GFP) mouse BM and immortalized after transduction with MLL-Af4 retrovirus, as previously described.<sup>21</sup> Cells then underwent a second transduction with retroviral supernatant containing sgRNA against *Igf2bp3* (I3sg2 and I3sg3) or nontargeting (NT-1 and NT-2) and sorted on GFP and mCherry positivity.

### Molecular biology assays

mRNA extracts and cell lysates were prepared, and western blot and reverse transcription quantitative polymerase chain reaction (qRT-PCR) were performed as previously described.<sup>12</sup> Reverse transcription quantitative polymerase chain reaction (RT-qPCR) primer sequences are listed in supplemental Table 1. Antibodies used were IGF2BP3 anti-rabbit polyclonal (RNP009, Medical and Biological Laboratories) and  $\beta$ -actin anti-mouse monoclonal (A5441, Sigma-Aldrich).

### In vitro assays for cell viability and apoptosis, with drug treatment

Cells were treated with drug or 0.1% dimethyl sulfoxide (DMSO) control, using concentrations and time periods specified in figure legends. The following menin-MLL inhibitors were used: MI-503 (SML2520, Sigma-Aldrich), MI-463 (HY-19809, MedChemExpress), and MI-538 (HY-19810, MedChemExpress). Cell viability

assays were performed using a luminescent assay based on adenosine triphosphate quantitation (CellTiterGlo, Promega). To measure apoptosis, cells were analyzed for annexin V by flow cytometry (BV421, BD Biosciences, BDB563873) and caspase-3/7 activity by a luminescent-based assay (Caspase-Glo 3/7 Assay, Promega).

### Methylcellulose-based colony-forming unit assays

The assay was performed by seeding drug-treated MLL-Af4 Cas9 Lin<sup>-</sup> cells into MethoCult colony-forming media (STEMCELL Technologies, M3434) at seeding densities of 250 to 2500. Cells were cultured in MethoCult media for 10 to 12 days and counted for total colony number and morphologic subtypes.

### Flow cytometry

Isolation of single-cell suspensions and staining with fluorochrome-conjugated antibodies were performed per standard procedures.<sup>12</sup> Antibodies are provided in supplemental Table 2. Flow cytometry was performed on a BD LSRII and analyzed using FlowJo software.

### RNA sequencing

Paired-end RNA sequencing was performed on an Illumina Nova-Seq S4 (UC Davis Genome Center). RNA-sequencing reads were mapped to the mouse genome assembly mm18 using STAR 2.5.3 and Bowtie 2.<sup>30,31</sup> Differentially expressed genes were identified using DESeq2.<sup>25</sup> Multiple testing correction was performed using the Benjamini-Hochberg method. Significant differentially expressed genes have adjusted *P* value  $\leq .05$  and absolute log<sub>2</sub>FC  $\geq 1$ . Enrichment analyses were completed with Metascape.<sup>29</sup> Overlap between differentially expressed genes and publicly available gene sets for MLL-AF4 chromatin immunoprecipitation (CHIP) and IGF2BP3 cross-link immunoprecipitation (CLIP) in CD11b<sup>+</sup> and Lin<sup>-</sup> leukemic cells was evaluated using Fisher exact test. Raw data have been deposited to the NCBI Short Read Archive (accession number PRJNA918326).

### Mice and BM transplantation

C57BL/6J, B6.SJL-*Ptprca*<sup>a</sup> *Pepc*<sup>b</sup>/BoyJ (B6 CD45.1), and B6J.129(Cg)-Gt(ROSA)26Sor<sup>tm1.1(CAG-cas9\*,EGFP)Fezh/J</sup> (Cas9-GFP BL/6J) mice were obtained from The Jackson Laboratory. MLL-Af4 Lin<sup>-</sup> cells were transplanted into 8- to 10-week-old CD45.1 female recipients (or C57BL/6J in pilot experiments). Recipients were conditioned with busulfan 30 mg/kg intraperitoneally twice, three days and two days prior to transplantation, followed by retro-orbital injection of 2x10<sup>5</sup> Lin<sup>-</sup> cells with 10<sup>5</sup> CD45.1 carrier marrow cells (with the exception of the experiment in supplemental Figure 4, which used a dose of 3x10<sup>5</sup> Lin<sup>-</sup> cells). This study received approval from the Chancellor's Animal Research Committee at UCLA, the responsible body for animal use approvals at UCLA. No human participants were involved in the research.

### Histopathology

Fixation, sectioning, and analysis were performed as previously described (DSR).<sup>32</sup>

### Statistics

Data shown represent mean  $\pm$  standard deviation for continuous numerical data. Two-tailed Student *t* tests or 1-way analysis of variance followed by Bonferroni multiple comparisons test were

performed using GraphPad Prism software and conducted as described in the figure legends. Survival analyses were performed using the Kaplan-Meier method with comparisons made using log-rank tests, followed by Bonferroni correction for multiple comparisons.

## Results


### IGF2BP3 knockdown increases sensitivity of MLL-r leukemia cells to menin-MLL inhibition in vitro

We assessed the effects of combining menin-MLL inhibition with IGF2BP3 knockdown in human B-ALL cell lines and murine immortalized HSPCs transformed with MLL-Af4 (referred herein as MLL-Af4 Lin<sup>-</sup>) as a means of inhibiting MLL-AF4 leukemogenesis at the transcriptional and posttranscriptional level. To knock down IGF2BP3 in human B-ALL cell lines, we used a 2-vector lentiviral system as previously described to deliver Cas9 and sgRNA targeting *IGF2BP3* (I3-sg2) or NT in 2 MLL-AF4-expressing human B-ALL cell lines, RS4;11 and SEM, and an additional human B-ALL cell line without MLL-AF4 translocation, NALM6. To deplete IGF2BP3 in MLL-Af4 Lin<sup>-</sup> cells, MLL-Af4 Cas9-GFP Lin<sup>-</sup> cells were transduced with a retroviral vector containing sgRNA targeting *Igf2bp3* (I3-sg2 or I3-sg3) as previously described<sup>13,21</sup> (Figure 1A; supplemental Figures 1B and 2A). MLL-Af4 Lin<sup>-</sup> cells treated with menin-MLL inhibitor, MI-503, for 4 days showed a dose-dependent decrease in IGF2BP3 expression, consistent with our prior findings that MLL-Af4 drives *Igf2bp3* expression<sup>13</sup> (Figure 1B).

IGF2BP3 knockdown resulted in dramatically enhanced menin-MLL inhibition of cell growth in SEM and RS4;11 treated with MI-503, MI-463, and MI-538 for 4 days in luminescence-based cell viability assays (Figure 1C-D). In contrast, there was no difference in sensitivity to the effects of menin-MLL inhibition on cell growth in NALM6, a human B-ALL cell line without MLL-AF4 translocation, with IGF2BP3 knockdown (Figure 1E). This synergistic effect of IGF2BP3 knockdown and menin-MLL inhibition on cell growth was also seen in our murine MLL-Af4 Lin<sup>-</sup> cells across multiple menin-MLL inhibitors and 2 different guides targeting *Igf2bp3* (Figure 1F; supplemental Figure 2D). MI-503 did not cause a significant increase in apoptosis of MLL-Af4 NT Lin<sup>-</sup> cells by annexin V staining or caspase 3/7 activity, except at the highest concentration of 1.0  $\mu$ M, similar to the limited effect on cell growth (Figure 1F-I). MI-503 had a greater effect on apoptosis in MLL-Af4 Lin<sup>-</sup> I3KO cells by increased caspase 3/7 activity and annexin V positivity (Figure 1G-J). Together, these findings highlight the sensitization of MLL-Af4 leukemia cells to menin-MLL inhibition with IGF2BP3 knockdown, with enhanced inhibition of cell growth and increased apoptosis.

### IGF2BP3 knockdown enhances effect of menin-MLL inhibition on LIC and differentiation

MLL-AF4 inhibition has been shown to decrease LICs and increase differentiation.<sup>7,22</sup> To characterize the combined effects of IGF2BP3 knockdown and menin-MLL inhibition on LICs in vitro, we seeded MI-503-treated MLL-Af4 I3KO and NT cells into end point colony formation assays and performed flow cytometric analyses, including measurements of c-Kit expression, an LIC marker that has been reported in MLL model systems.<sup>6,13,18,23,24</sup> IGF2BP3 knockdown more significantly reduced the total colony number than MI-503 treatment (Figure 2A). MI-503-treated MLL-Af4 Lin<sup>-</sup> I3KO cells showed further reduction in total colony formation,



**Figure 1. IGF2BP3 knockdown increases sensitivity of MLL-r leukemia cells to menin-MLL inhibition.** (A) Western blot analysis showing IGF2BP3 knockdown in RS4;11, NALM6, and SEM cell lines and MLL-Af4 Lin<sup>-</sup> cells (using I3KO sgRNA targeting human and mouse genes and NT guide). (B) Western blot analysis showing IGF2BP3 expression in MI-503 treated MLL-Af4 Lin<sup>-</sup> cells depleted (I3KO) or nondepleted for IGF2BP3 (NT). Cells were treated with MI-503 (0.2 μM, 1.0 μM, or DMSO control) for 4 days. (C-E) Dose-response curves from cell viability assays, using CellTiterGlo, of human B-ALL cell lines, SEM, RS4;11, and NALM6, depleted (I3KO) vs nondepleted (NT) for IGF2BP3 treated with menin-MLL inhibitors (MI-503, MI-463, and MI-538) for 4 days. Viability has been normalized to DMSO control-treated cells not depleted for IGF2BP3 (NT DMSO); mean ± standard deviation (SD); n = 6. (F) Dose-response curves from cell viability assays, using CellTiterGlo, of MLL-Af4 Lin<sup>-</sup> cells depleted for IGF2BP3 (I3KO) vs nondepleted (NT) treated with menin-MLL inhibitors for 4 days. Viability has been normalized to DMSO control-treated cells not depleted for IGF2BP3 (NT DMSO); mean ± SD; n = 6. (G) Increased caspase 3/7 activity of MI-503 treated MLL-Af4 Lin<sup>-</sup> cells depleted for IGF2BP3 (I3KO) vs nondepleted (NT). Cells treated with MI-503 for 4 days at various concentrations

compared with IGF2BP3 knockdown alone, suggesting that the combined inhibitory effect on LICs is at least additive and is dose dependent (Figure 2A-C). Both MI-503 treatment and IGF2BP3 knockdown in MLL-Af4 Lin<sup>-</sup> cells showed a shift toward more differentiated colony morphologies in colony formation assays, decreased c-Kit expression, increased expression of maturation markers by flow cytometry, and morphologic changes consistent with increased differentiation. Decreased granulocyte-macrophage colony-forming unit (CFU-GM) and increased granulocyte colony-forming unit (CFU-G) and macrophage colony-forming unit (CFU-M) progenitor colonies were seen with MI-503 and in I3KO cells, with the maximal effect seen in the combination (Figure 2D-F). By flow cytometry, we found a marked reduction in the number of Lin<sup>-</sup> cells with I3KO, consistent with increased differentiation (Figure 2G,J-M). We also saw a decrease in c-Kit<sup>+</sup> cells, more profoundly with I3KO than with MI-503 treatment, and with the maximal effect seen in combination (Figure 2H). Furthermore, expression of CD16/32, a marker for granulocyte-macrophage differentiation, was increased with MI-503 treatment (Figure 2O,Q), whereas expression of CD34, a marker that has been used to identify and enrich LICs, was decreased with I3KO (Figure 2N,P). This pattern of increased differentiation with MI-503 treatment and IGF2BP3 depletion was seen morphologically in cytospin preparations (supplemental Figure 3). Altogether, these findings confirm that both IGF2BP3 and menin inhibition affect the number and differentiation state of LICs and that combined inhibition appears to have an additive effect in our in vitro assay readouts.

### IGF2BP3 knockdown in MLL-Af4 Lin<sup>-</sup> cells affects the global and MLL-AF4 transcriptome and leads to the upregulation of genes involved in differentiation

To gain insight into the phenotypic findings described, we next examined gene expression from cells with combined menin-MLL inhibition and IGF2BP3 knockdown. We performed RNA-sequencing analysis in MLL-Af4 Lin<sup>-</sup> cells with the following treatment conditions: (1) MLL-Af4 Lin<sup>-</sup> cells depleted for IGF2BP3 (I3KO) and treated with MI-503 (at 0.2 and 1.0 μM) or DMSO control and (2) MLL-Af4 Lin<sup>-</sup> cells nondepleted for IGF2BP3 (NT) and treated with MI-503 (at 0.2 and 1.0 μM) or DMSO control. Comparison of DMSO-treated MLL-Af4 Lin<sup>-</sup> I3KO vs NT cells showed dramatic changes in the global transcriptome with 4346 upregulated and 1720 downregulated genes by differential expression analysis with DESeq2<sup>25,26</sup> (Figure 3A; supplemental Figure 4A). In contrast, fewer genes are differentially expressed in MLL-Af4 Lin<sup>-</sup> NT with MI-503 than with DMSO (Figure 3A; supplemental Figure 4A). Remarkably, there is a highly significant overlap between genes that are differentially expressed with IGF2BP3 knockdown and MI-503 treatment (Hypergeometric test;  $P = 0$ ). Seventy-eight percent of genes that are upregulated with MI-503 treatment are also upregulated with IGF2BP3 knockdown, and 34% of the genes that are downregulated with MI-503 treatment are downregulated with IGF2BP3 knockdown (Figure 3B).

Next, we looked at the overlap between differentially expressed genes with MI-503 and I3KO with MLL-AF4 ChIP targets and

IGF2BP3 CLIP targets identified in CD11b<sup>+</sup> and Lin<sup>-</sup> MLL-Af4 leukemic cells.<sup>13,27,28</sup> We found significant overlap of differentially expressed genes with MI-503 treatment with IGF2BP3 CLIP targets (by Fisher exact test; Figure 3D), suggesting that IGF2BP3 directly regulates genes that are affected by menin-MLL inhibition. Conversely, there was an overlap seen between differentially expressed genes with IGF2BP3 knockdown and MLL-AF4 ChIP targets identified in SEM and RS4;11, human MLL-AF4 leukemia cell lines (Figure 3D). These overlap patterns highlight the interaction between the MLL-AF4 and IGF2BP3 transcriptomes and suggest a mechanism for the synergistic inhibition of leukemia by targeting both MLL-AF4 and IGF2BP3.

In comparisons of upregulated genes in NT and I3KO cells, we identified a significant enrichment by Metascape analysis<sup>29</sup> in pathways involved in cell differentiation and activation, particularly in leukocytes: neutrophil degranulation, regulation of cell-cell adhesion, cell activation, myeloid leukocyte differentiation, and leukocyte migration (Figure 3E). Although fewer genes were differentially expressed with MI-503 treatment, partial pathway enrichment overlap was observed (eg, neutrophil degranulation), but additional distinct pathways were affected by MI-503 treatment (eg, oxidative stress and redox pathways and some metabolic pathways; Figure 3F). Pathway enrichment analyses of genes differentially expressed with MI-503 treatment in I3KO cells also revealed pathways important in hematopoietic differentiation (eg, leukocyte migration and neutrophil degranulation; Figure 3C; supplemental Figure 4D). Pathway enrichment analyses of downregulated genes in NT vs I3KO, NT with and without MI-503 treatment, and I3KO with and without MI-503 treatment revealed a variety of pathways involved in signaling, stem cell homeostasis, and immune-related pathways (supplemental Figure 4B-D).

Given the phenotypic alterations to a more differentiated state and the strong enrichment for hematopoietic differentiation pathways among upregulated genes, we next looked to validate our findings using individual qRT-PCR for these genes of interest. As anticipated, we found a decrease in the expression of *Igf2bp3* and its known target mRNAs, *Myc* and *Hoxa9*, in NT cells treated with MI-503 (Figure 4A-C). Next, we focused on the enriched differentially upregulated genes involved in leukocyte differentiation, *Fcnb*, *Prg2*, *Mmrn1*, *Elane*, *Ets1*, *Pram1*, *Cebpd*, and *Cebpe*. A significant increase was seen in all of these genes with *Igf2bp3* depletion (Figure 4D-K). Fold change was significantly higher with I3KO than with MI-503 treatment (Figure 4D-K). For genes *Fcnb*, *Prg2*, *Mmrn1*, and *Pram1*, MI-503 treatment led to a significant increase in expression in MLL-Af4 Lin<sup>-</sup> I3KO, whereas a nonsignificant increase was seen with the other genes of interest, *Elane*, *Ets1*, *Cebpd*, and *Cebpe*, suggesting a saturating effect of *Igf2bp3* depletion on upregulation of these genes. Together, our analyses demonstrate that similar, but not identical, impacts on gene expression are engendered by genetic inhibition of IGF2BP3 and menin-MLL inhibition with MI-503. We propose that these impacts on gene expression underlie the phenotypic effects seen upon the coinhibition of transcriptional and posttranscriptional gene expression regulation.

**Figure 1 (continued)** for dose response. Caspase 3/7 activity measured using Caspase-Glo 3/7 and normalized to the activity of DMSO-treated control; mean ± SD, n = 3. (H) Increased annexin V positivity in MLL-Af4 Lin<sup>-</sup> I3KO cells (vs NT) and with MI-503 treatment (vs DMSO control); mean ± SD; n = 3 (1-way analysis of variance [ANOVA] with Bonferroni multiple comparisons test; \*\* $P < .01$ ; \*\*\*\* $P < .0001$ ). (I-J) Histograms from representative samples for annexin V staining, analyzed by flow cytometry. kDa, kilodalton.




Figure 2.

## IGF2BP3 knockdown and MI-503 treatment decrease engraftment of MLL-Af4 leukemia cells in vivo

To further characterize our findings in vivo, we developed and validated a congenic CD45.1/CD45.2 transplantation model for the murine MLL-Af4 leukemia, using the earlier-described cellular systems (MLL-Af4 Lin<sup>-</sup> NT and I3KO) derived from murine BM. In timed experiments terminated at 8 to 10 weeks, I3KO demonstrated a clear reduction in engraftment (assessed using mCherry<sup>+</sup> and GFP<sup>+</sup> cells; supplemental Figure 5B), gross leukemia or histologic preleukemia (supplemental Figure 5C-D), and spleen weights (supplemental Figure 5E). At the cellular level, I3KO led to reductions in cell counts, CD11b<sup>+</sup> cells, and engraftment in the spleen (supplemental Figure 5F-G) and BM (supplemental Figure 5I-K). Next, we transplanted either NT or I3KO Lin<sup>-</sup> cells (derived from CD45.2<sup>+</sup> mice) into CD45.1<sup>+</sup> mice after 5 days of treatment with MI-503 in vitro (Figure 5A). Six weeks after transplantation, evaluation of peripheral blood showed that both treatment with MI-503 and IGF2BP3 depletion led to decreased CD45.2<sup>+</sup> cells in the peripheral blood, reflective of decreased leukemic engraftment (Figure 5B). In end point experiments at 8.5 weeks, IGF2BP3 depletion led to a significantly decreased proportion of mice with gross leukemia at necropsy, decreased BM counts, and decreased CD11b<sup>+</sup> count and CD45.2<sup>+</sup> percentage in the spleen and BM (Figure 5C,D-I). MI-503 treatment resulted in a modest decrease in the proportion of mice with gross leukemia, total cellular and CD11b<sup>+</sup> counts, and CD45.2<sup>+</sup> percentage in the spleen and BM (Figure 5D).

In addition to the aforementioned end point experiments, we evaluated mice for overall survival and leukemia-free survival in a separate experiment. MI-503 treatment showed a modest effect with a delay in leukemia progression but without improvement in overall survival in mice that received transplantation with MLL-Af4 leukemia (NT; comparison of MI-503 vs DMSO; Figure 6A-B). In contrast, IGF2BP3 depletion significantly increased overall and leukemia-free survival in mice that received transplantation with MLL-Af4 leukemia (NT DMSO vs I3KO DMSO; Figure 6A-B). This increase in overall and leukemia-free survival was also seen in mice that received transplantation with MI-503-treated MLL-Af4 leukemia cells (comparison NT MI-503 vs I3KO MI-503). Interestingly, histopathologic analysis showed morphologic changes consistent with increased differentiation in both the BM and spleens of mice that received transplantation with MI-503-treated or IGF2BP3-depleted MLL-Af4 Lin<sup>-</sup> cells, with the cells appearing most differentiated in mice that received transplantation with MI-503 treated I3KO cells (Figure 6C; supplemental Figure 6A). Furthermore, in BM samples from these mice, we saw an upregulation in

granulocytic differentiation-related genes, *Cebpe*, *Elane*, *Fcnb*, *Prg1*, *Mmrr1*, and *Ets1*, previously identified in our evaluation of the Lin<sup>-</sup> cells in vitro (Figure 6D-F; supplemental Figure 6C-G). Together, the data confirm that MI-503 and I3KO both have impacts on leukemogenesis, with an additive effect on leukemic engraftment and cellular differentiation.

## Discussion

In this study, we tested a novel combinatorial strategy of targeting leukemia at the transcriptional and posttranscriptional level using commercially available inhibitors of menin-MLL and genetic inhibition of IGF2BP3, an RBP critical to MLL-Af4-mediated leukemogenesis.<sup>13</sup> We found that IGF2BP3 knockdown enhanced the therapeutic effect of pharmacologic menin-MLL inhibition in MLL-Af4-driven leukemia in our in vitro and in vivo studies. Both IGF2BP3 knockdown and the menin-MLL inhibition negatively affect the number and function of LICs, as shown by functional readouts in end point colony formation assays and decreased c-Kit<sup>+</sup> cells by flow cytometry. Furthermore, detailed evaluation of colony morphologies, flow cytometry, histopathology, and RNA-sequencing data show a consistent shift toward increased differentiation with IGF2BP3 knockdown and menin-MLL inhibition, highlighting a mechanism of their combined antileukemic effects.

RBPs are proving to be viable therapeutic targets in cancer. Small molecule inhibitors have been developed against RBPs, including Musashi-2, HuR, LIN28B, MBNL1, and IGF2BP2, and studied as single agents but have not been combined with therapies directed toward upstream transcriptional regulators.<sup>33-38</sup> Here, we have shown a potent effect of combining genetic inhibition of the RBP IGF2BP3 and pharmacologic inhibition of the menin-MLL interaction, resulting in decreased cell growth in vitro and leukemic burden in vivo. These results highlight an important interaction between MLL and IGF2BP3 in the gene regulation of MLL-Af4-driven leukemia and support the combined targeting of menin-MLL and IGF2BP3 for the treatment of MLL-rearranged leukemia. Our finding demonstrates the feasibility of a novel treatment paradigm, concomitantly targeting transcriptional and posttranscriptional gene regulation.

Regarding underlying mechanisms of IGF2BP3 function in leukemogenesis, we have previously shown that deletion of *Igf2bp3* led to decreased number and function of MLL-Af4 LICs.<sup>13</sup> Borkin et al showed that treatment with menin-MLL inhibitors, MI-503 and MI-563, led to decreased c-Kit expression in MLL-r leukemia cells.<sup>18</sup> In this study, we demonstrated a decrease in the number of Lin<sup>-</sup> and c-Kit<sup>+</sup> cells with IGF2BP3 knockdown and MI-503 treatment, consistent with prior observations showing decreased

**Figure 2. Combined IGF2BP3 knockdown and menin inhibition increases differentiation of MLL-Af4 leukemia.** (A) Total colony numbers of MI-503-treated MLL-Af4 Lin<sup>-</sup> cells, depleted (I3KO) or nondepleted (NT) for IGF2BP3. MLL-Af4 Lin<sup>-</sup> NT and I3KO cells were treated with MI-503 0.5 μM for 4 days and seeded in methylcellulose colony formation assays at various initial seeding densities and cultured for 10 days. (B,C) Total colony number was reduced with both I3KO and MI-503 treatment at 0.5 μM (B) and 0.5 μM (C) in methylcellulose colony formation assays at initial seeding density of 2500. (D) Proportion of CFU-GM colonies are decreased with both I3KO and MI-503 (at 0.5 μM). (E-F) Proportion of CFU-G and -M colonies are increased with both I3KO and MI-503 (at 0.5 μM). (G-I) Lin<sup>-</sup>, c-kit<sup>+</sup>, and Lin<sup>-</sup>c-kit<sup>+</sup> cells are decreased with both I3KO and MI-503. MLL-Af4 Lin<sup>-</sup> NT and I3KO cells were treated with MI-503 0.5 mM for 7 days and then analyzed by flow cytometric immunophenotyping; mean ± SD, n = 3; 1-way ANOVA with Bonferroni multiple comparison's test (\*P < .05; \*\*P < .01; \*\*\*P < .001; \*\*\*\*P < .0001). (J-M) Corresponding representative flow cytometry plots of MLL-Af4 Lin<sup>-</sup> NT and I3KO cells treated with MI-503 (or DMSO control), showing expression of c-kit and Lin markers. (N) Decreased CD34 expression by mean fluorescence intensity with I3KO; mean ± SD; n = 3; student t test, \*P < .05. (O) Decreased CD16/32 expression by mean fluorescence intensity with MI-503 treatment; mean ± SD; n = 3; Student t test, \*\*P < .01. (P-Q) Corresponding representative flow cytometry plots of MLL-Af4 Lin<sup>-</sup> NT and I3KO cells treated with MI-503 (or DMSO control), showing expression of CD16/32 and CD34. CFU-GM, granulocyte-macrophage colony-forming units; CFU-G, granulocyte colony-forming units; CFU-M, macrophage colony-forming units; MFI, mean fluorescence intensity; prop, proportion.






Figure 3.

CD11b<sup>+</sup>c-kit<sup>+</sup> cells in I3KO/MLL-Af4 mice and decreased c-Kit<sup>+</sup> expression in MI-503-treated MLL-AF9-transformed BM<sup>13,18</sup> and the prior characterization of menin-MLL inhibitors in promoting differentiation of MLL-rearranged leukemia.<sup>18,19</sup> We observed a decrease in total colony numbers and a shift toward more differentiated colony morphologies in end point colony formation assays, consistent with the prior characterization of these menin-MLL inhibitors. Notably, IGF2BP3 knockdown enhanced the effect of MI-503 in these readouts of LIC function and number, suggesting that concomitant inhibition of IGF2BP3 and menin-MLL additively target the leukemic stem cell, a known critical driver of refractory and relapsed disease.<sup>39-42</sup>

Furthermore, we tested the combined inhibition of IGF2BP3 and the menin-MLL interaction on the ability to initiate MLL-Af4 leukemia in vivo. We demonstrated that transformed MLL-Af4 HSPCs can efficiently engraft and initiate leukemia in syngeneic mice, after limited time in culture and CRISPR/Cas9-mediated deletion of *Igf2bp3*. Treatment of these MLL-Af4-transformed HSPCs with menin-MLL inhibitor in vitro and CRISPR/Cas9-mediated deletion of *Igf2bp3* effectively reduced leukemic engraftment in vivo, showing that IGF2BP3 and menin-MLL inhibition reduce the number and function of LICs and is consistent with the results from our in vitro colony formation assays. Hence, this represents a single integrated system that allows for both in vitro and in vivo analyses of leukemogenesis.

Our RNA-sequencing data in drug-treated and knockout cells provide some mechanistic insight into the enhanced antileukemic effects seen with combinatorial inhibition. We demonstrated that both menin-MLL inhibition and IGF2BP3 upregulated genes involved in leukocyte/granulocyte differentiation, with a stronger effect seen with *Igf2bp3* depletion than pharmacologic menin-MLL inhibition alone and an additive effect in combination with menin-MLL inhibition. These results help provide a molecular explanation for the increased differentiation seen with menin-MLL inhibitor treatment, previously reported in studies of MI-503, MI-463, and VTP-50469.<sup>16,18,19</sup> Furthermore, the pattern of upregulation of genes involved in granulocyte differentiation, seen with MI-503 treatment alone, IGF2BP3 knockdown alone, and the combination, mirror functional and phenotypic outputs of antileukemic effects on cell growth, differentiation, and LIC number and function.


We demonstrated that menin-MLL inhibition reduces IGF2BP3 expression and the expression of known IGF2BP3 targets that are important in leukemogenesis, such as *Hoxa9* and *Myc*, and that the addition of IGF2BP3 knockdown further reduced the expression of these important oncogenic transcripts. This effect supports our understanding of IGF2BP3 as an oncogenic amplifier of

MLL-Af4-mediated leukemogenesis. Additionally, this study provides possible targets of interest that can be studied to further understand the interaction between MLL-Af4 and IGF2BP3. For example, we identified *Elane* as a differentially upregulated gene with MI-503 treatment and IGF2BP3 knockdown, which is interestingly also a direct target of IGF2BP3, in MLL-Af4 Lin<sup>-</sup> cells.<sup>13</sup> Additional studies using functional genomic analyses to knock out direct targets of IGF2BP3 would help define the contribution of these transcripts to MLL-Af4 leukemogenesis and elucidate the mechanisms by which IGF2BP3 contributes to leukemogenesis. Our model system of MLL-Af4 leukemia using Cas9-expressing MLL-Af4-transformed HSPCs lends itself readily to these functional genomic analyses for accelerated screening of important functional targets, using CRISPR sgRNA libraries. The myeloid lineage of the MLL-Af4 leukemia generated in this system in vitro and in vivo is a potential limitation to the mouse model in modeling human disease. However, human MLL-AF4 leukemia is marked by lineage infidelity, particularly after treatment.<sup>43,44</sup> Furthermore, Lin et al demonstrated important similarities between the phenotypic and molecular aspects of MLL-Af4-induced pro-B-ALL and acute myeloid leukemia in mice.<sup>6</sup>


The molecular mechanisms of IGF2BP3 as an RBP remain an area of active investigation. Prior work from our group and others has shown the ability of IGF2BP3 and other RBPs to affect RNA splicing and RNA stability through interaction with microRNAs, the RNA-induced silencing complex, as potential molecular mechanisms underlying the role and function of RBP in oncogenesis.<sup>12,13,45-48</sup> Whether these functions, particularly the proposed role in mRNA splicing, interact with the aberrant transcriptional regulation mediated by menin-MLL fusion proteins remains unknown. In this regard, it is interesting to note that METTL3, an RNA N<sup>6</sup> methyl adenosine (m<sup>6</sup>A) methyltransferase, binds to promoters of genes whose mRNA is modified.<sup>49</sup> Given the overlap between MLL- and IGF2BP3-regulated gene expression, it is interesting to speculate that a similar mechanism connecting transcriptional and posttranscriptional regulation may exist.

Despite the strong evidence for IGF2BP3 in leukemogenesis, there may be alternative models to consider. A recent publication has suggested that IGF2BP3, among several other RBPs, may be important in negatively regulating the stability of the human MLL-AF4 transcript.<sup>50</sup> Although some of the findings in this study are in agreement with ours, there are significant differences in other observations, perhaps because of significantly different models and readouts and differential induction of IGF2BP3 by chimeric fusion proteins that were tested. Nonetheless, this work does bring up new possibilities that IGF2BP3 may directly regulate MLL-AF4: a possibility that requires further work to fully explore.


**Figure 3. Increased upregulation of genes involved in differentiation with IGF2BP3 knockdown and menin-MLL inhibition in MLL-Af4 leukemia.** (A) Volcano plots of differentially expressed genes with IGF2BP3 knockdown (top) and with MI-503 treatment (left) in MLL-Af4 Lin<sup>-</sup> NT cells, using DESeq analysis on RNA-sequencing samples from MLL-Af4 Lin<sup>-</sup> NT or I3KO cells treated with MI-503 1.0 μM or DMSO vehicle. (B) Venn diagram of shared differentially expressed genes with IGF2BP3 knockdown and MI-503 treatment in MLL-Af4 Lin<sup>-</sup> NT cells. (C) Volcano plots of differentially expressed genes with IGF2BP3 knockdown (top) and with MI-503 treatment (left) in MLL-Af4 Lin<sup>-</sup> I3KO cells, using DESeq analysis on RNA-sequencing samples from MLL-Af4 Lin<sup>-</sup> NT or I3KO cells treated with MI-503 1.0 μM or DMSO vehicle. (D) Overlap between differentially expressed genes with MI-503 in MLL-Af4 Lin<sup>-</sup> NT or I3KO cells and with IGF2BP3 knockdown and known MLL-AF4 ChIP targets from previously published data sets in SEM and RS4;11 and known IGF2BP3 eCLIP targets from previously published data sets of MLL-Af4 CD11b<sup>+</sup> and MLL-Af4 Lin<sup>-</sup> cells. *P* values calculated using Fisher exact test. (E) Pathway enrichment for upregulated genes with IGF2BP3 knockdown using Metascape analysis webtool on MLL-Af4 Lin<sup>-</sup> IGF2BP3 DESeq data set with an adjusted *P* < .05 cutoff. (F) Pathway enrichment for upregulated genes with MI-503 treatment using Metascape analysis webtool on MLL-Af4 Lin<sup>-</sup> NT MI-503 DESeq data set with an adjusted *P* < .05 cutoff. FC, fold change.



**Figure 4. Increased upregulation of genes involved in differentiation with IGF2BP3 knockdown and menin-MLL inhibition in MLL-Af4 leukemia, validated by qRT-PCR.** (A-C) MI-503 treatment at 1.0  $\mu$ M leads to downregulation of *Igf2bp3* and known IGF2BP3 targets, *Myc* and *Hoxa9*, in MLL-Af4 Lin<sup>-</sup> cells. (D-K) Upregulation of genes involved in differentiation with IGF2BP3 knockdown and MI-503 treatment. mRNA expression was measured by qRT-PCR in MLL-Af4 Lin<sup>-</sup> cells depleted (I3KO) or nondepleted (NT) for IGF2BP3 and treated with MI-503 1.0  $\mu$ M or DMSO control. Expression shown as fold change from NT DMSO (mean  $\pm$  SD; n = 2; 1-way ANOVA with Bonferroni multiple comparisons test, \* $P$  < .05; \*\* $P$  < .01; \*\*\* $P$  < .001; \*\*\*\* $P$  < .0001).



**Figure 5. Combinatorial inhibition of menin-MLL and IGF2BP3 decreases leukemic engraftment and burden and increases survival in vivo.** (A) Schematic representation of BM transplantation of MI-503–treated MLL-Af4 Lin<sup>−</sup> cells, depleted (I3KO) or nondepleted (NT) for IGF2BP3. Cells treated with MI-503 0.5  $\mu$ M (or DMSO control) for 5 days in vitro were transplanted into CD45.1 recipients after busulfan conditioning. (B) Decreased peripheral blood engraftment of leukemic cells by CD45.2<sup>+</sup> percentage with MI-503 treatment and I3KO at D42 (mean  $\pm$  SD; n = 8 mice per group; 1-way ANOVA with Bonferroni multiple comparison's test, \*  $P < .05$ ). (C) Decreased



**Figure 6. Combinatorial inhibition of menin-MLL and IGF2BP3 increases survival in vivo and is accompanied by concordant changes in gene expression.** (A-B) Increased overall survival and leukemia survival with IGF2BP3 knockdown and MI-503 treatment in NT groups (n = 8 mice per group; Kaplan-Meier with log-rank test, \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ ). Comparisons made vs NT DMSO. Follow-up to 15 weeks with terminal sac of remaining mice. (C) Photomicrographs of Wright-stained BM smears from mice as above; original magnification,  $\times 1000$ ; scale bar, 10  $\mu\text{m}$ . (D-F) Expression of genes of interest in the BM of mice that received transplantation with MI-503–treated MLL-Af4 Lin<sup>-</sup> NT or I3KO cells was measured by qRT-PCR. MLL-Af4 Lin<sup>-</sup> cells depleted (I3KO) or nondepleted (NT) for IGF2BP3 were treated with MI-503 0.5 mM (MI-503) or carrier control (DMSO) for 5 days in vitro before transplantation. Mice were euthanized at 8.5 weeks at first signs of the first mouse developing terminal leukemia. Gene expression data shown from selected mice in each group. Shown as fold change from NT DMSO (mean  $\pm$  SD; n = 2; 1-way ANOVA with Bonferroni multiple comparisons test, \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ ; \*\*\*\* $P < .0001$ ).

**Figure 5 (continued)** proportion of mice with leukemia with MI-503 treatment and IGF2BP3 knockdown, (8 mice per group; Fisher exact test; \* $P < .05$ ; \*\* $P < .01$ ). Mice were all evaluated at necropsy at 8.5 weeks after the first mouse developed signs of terminal leukemia. (D-I) Decreased leukemic burden seen with IGF2BP3 knockdown and MI-503 treatment in NT groups, based on total counts, CD11b<sup>+</sup> counts, and CD45.2<sup>+</sup> percentage by flow cytometry in the spleen (D-F) and BM (G-I) (mean  $\pm$  SD; n = 7-8 mice per group; 1-way ANOVA with Bonferroni multiple comparison's test, \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ ; \*\*\*\* $P < .0001$ ). Mice were all evaluated at necropsy at 8.5 weeks after the first mouse developed signs of terminal leukemia.

Our data have demonstrated that IGF2BP3 is a high value therapeutic target as we look toward methods of inhibiting IGF2BP3 in combination with existing experimental therapeutics, including menin-MLL inhibitors. One limitation of our study is that *Igf2bp3* deletion occurred before the functional readouts that we pursued. Therefore, the role of IGF2BP3 in leukemia maintenance remains an important and unanswered question. Future studies, using a conditional knockout system and/or protein degradation systems, will help us further assess the therapeutic potential of targeting IGF2BP3. Our initial attempts at developing an inducible knockdown using cyclization recombinase-locus of cross-over, P1 (Cre-LoxP) system resulted in a significantly leaky system that demonstrated the anti-leukemic effects of IGF2BP3 deletion, but deletion of *Igf2bp3* occurred concurrent with leukemia induction (supplemental Figure 7).<sup>51</sup> Nonetheless, in this study, we have shown that targeting IGF2BP3 had potent antileukemic effects against MLL-Af4 leukemia in vitro and in vivo and that this results in additional antileukemic effects with menin-MLL inhibitors. Our studies confirm a role of IGF2BP3 as an oncogenic amplifier of MLL-AF4-driven leukemia and suggest a promising and novel combinatorial approach to targeting leukemia at the transcriptional and posttranscriptional level.

## Acknowledgments

The authors thank members of the Rao Lab for helpful discussions regarding the research.

This work was supported by R01CA264986 from the National Institutes of Health (D.S.R. and J.R.S.) and R03CA251854 (D.S.R.); Hematology Training Grant (T32HL066992), Tumor Immunology Training Grant (NIH T32CA009120), a training grant from the California Institute of Regenerative Medicine, the Broad

Stem Cell Research Center, and the UCLA Specialty Training and Advanced Research Program (T.L.L.); and the Tumor Cell Biology Training Grant (NIH T32CA009056) (T.M.T.). Flow cytometry was performed in the UCLA Jonsson Comprehensive Cancer Center and Center for AIDS Research Flow Cytometry Core Facility that is supported by National Institutes of Health awards AI-28697, and award number P30CA016042, the UCLA Jonsson Comprehensive Cancer Center, the UCLA AIDS Institute, and the David Geffen School of Medicine at UCLA.

## Authorship

Contribution: T.L.L., A.K.J., A.J.R., J.R., T.M.T., Z.T.N., S.K., M.L.T., and A.C. performed experiments; T.L.L., A.K.J., A.J.R., S.K., and D.S.R. analyzed results and made the figures; T.L.L., J.R.S., and D.S.R. designed the research and wrote the manuscript; and T.L.L., A.K.J., A.J.R., J.R., T.M.T., Z.T.N., S.K., M.L.T., A.C., J.R.S., and D.S.R. reviewed and edited the manuscript.

Conflict-of-interest disclosure: D.S.R. serves on advisory boards and has consulted for AbbVie, Inc, a pharmaceutical company that markets drugs for acute leukemia. The remaining authors declare no competing financial interests.

ORCID profiles: A.J.R., 0000-0002-1998-7357; Z.T.N., 0000-0001-8930-3574; J.R.S., 0000-0002-1131-1627; D.S.R., 0000-0002-0794-9337.

Correspondence: Dinesh S. Rao, Department of Pathology and Laboratory Medicine, David Geffen School of Medicine at UCLA, 650 Charles E Young Dr, 12-272 Factor, Los Angeles, CA 90095; email: [drao@mednet.ucla.edu](mailto:drao@mednet.ucla.edu).

## References

1. Luo Z, Lin C, Shilatifard A. The super elongation complex (SEC) family in transcriptional control. *Nat Rev Mol Cell Biol*. 2012;13(9):543-547.
2. Cucinotta CE, Arndt KM. SnapShot: transcription elongation. *Cell*. 2016;166(4):1058-1058.e1.
3. Collins CT, Hess JL. Deregulation of the HOXA9/MEIS1 axis in acute leukemia. *Curr Opin Hematol*. 2016;23(4):354-361.
4. Mullighan CG, Goorha S, Radtke I, et al. Genome-wide analysis of genetic alterations in acute lymphoblastic leukaemia. *Nature*. 2007;446(7137):758-764.
5. Radtke I, Mullighan CG, Ishii M, et al. Genomic analysis reveals few genetic alterations in pediatric acute myeloid leukemia. *Proc Natl Acad Sci U S A*. 2009;106(31):12944-12949.
6. Lin S, Luo RT, Ptasinska A, et al. Instructive role of MLL-fusion proteins revealed by a model of t(4;11) pro-B acute lymphoblastic leukemia. *Cancer Cell*. 2016;30(5):737-749.
7. Krivtsov AV, Twomey D, Feng Z, et al. Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature*. 2006;442(7104):818-822.
8. Barabé F, Gil L, Celton M, et al. Modeling human MLL-AF9 translocated acute myeloid leukemia from single donors reveals RET as a potential therapeutic target. *Leukemia*. 2017;31(5):1166-1176.
9. Yokoyama A, Somerville TC, Smith KS, Rozenblatt-Rosen O, Meyerson M, Cleary ML. The menin tumor suppressor protein is an essential oncogenic cofactor for MLL-associated leukemogenesis. *Cell*. 2005;123(2):207-218.
10. Chen YX, Yan J, Keeshan K, et al. The tumor suppressor menin regulates hematopoiesis and myeloid transformation by influencing Hox gene expression. *Proc Natl Acad Sci U S A*. 2006;103(4):1018-1023.
11. Caslini C, Yang Z, El-Osta M, Milne TA, Slany RK, Hess JL. Interaction of MLL amino terminal sequences with menin is required for transformation. *Cancer Res*. 2007;67(15):7275-7283.
12. Palanichamy JK, Tran TM, Howard JM, et al. RNA-binding protein IGF2BP3 targeting of oncogenic transcripts promotes hematopoietic progenitor proliferation. *J Clin Invest*. 2016;126(4):1495-1511.
13. Tran TM, Philipp J, Bassi JS, et al. The RNA-binding protein IGF2BP3 is critical for MLL-AF4-mediated leukemogenesis. *Leukemia*. 2022;36(1):68-79.

14. Chan AKN, Chen CW. Rewiring the epigenetic networks in MLL-rearranged leukemias: epigenetic dysregulation and pharmacological interventions. *Front Cell Dev Biol.* 2019;7:81.
15. Bernt KM, Armstrong SA. Targeting epigenetic programs in MLL-rearranged leukemias. *Hematology Am Soc Hematol Educ Program.* 2011;2011:354-360.
16. He S, Malik B, Borkin D, et al. Menin-MLL inhibitors block oncogenic transformation by MLL-fusion proteins in a fusion partner-independent manner. *Leukemia.* 2016;30(2):508-513.
17. Kühn MWM, Song E, Feng Z, et al. Targeting chromatin regulators inhibits leukemogenic gene expression in NPM1 mutant leukemia. *Cancer Discov.* 2016;6(10):1166-1181.
18. Borkin D, He S, Miao H, et al. Pharmacologic inhibition of the Menin-MLL interaction blocks progression of MLL leukemia in vivo. *Cancer Cell.* 2015;27(4):589-602.
19. Krivtsov AV, Evans K, Gadrey JY, et al. A Menin-MLL inhibitor induces specific chromatin changes and eradicates disease in models of MLL-rearranged leukemia. *Cancer Cell.* 2019;36(6):660-673.e11.
20. Klossowski S, Miao H, Kempinska K, et al. Menin inhibitor MI-3454 induces remission in MLL1-rearranged and NPM1-mutated models of leukemia. *J Clin Invest.* 2020;130(2):981-997.
21. Jaiswal AK, Truong H, Tran TM, et al. Focused CRISPR-Cas9 genetic screening reveals USO1 as a vulnerability in B-cell acute lymphoblastic leukemia. *Sci Rep.* 2021;11(1):13158.
22. Cozzio A, Passequé E, Ayton PM, Karsunky H, Cleary ML, Weissman IL. Similar MLL-associated leukemias arising from self-renewing stem cells and short-lived myeloid progenitors. *Genes Dev.* 2003;17(24):3029-3035.
23. Somerville TC, Cleary ML. Identification and characterization of leukemia stem cells in murine MLL-AF9 acute myeloid leukemia. *Cancer Cell.* 2006;10(4):257-268.
24. Challen GA, Boles N, Lin KK, Goodell MA. Mouse hematopoietic stem cell identification and analysis. *Cytometry A.* 2009;75(1):14-24.
25. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.
26. Lin I, Wei A, Awamleh Z, Singh M, Ning A, Herrera A, REACH Biobank and Registry, Russell BE, Weksberg R, Arboleda VA, Arboleda VA. Multiomics of Bohring-Opitz syndrome truncating ASXL1 mutations identify canonical and noncanonical Wnt signaling dysregulation. *JCI Insight.* 2023;8(10):e167744.
27. Guenther MG, Jenner RG, Chevalier B, et al. Global and Hox-specific roles for the MLL1 methyltransferase. *Proc Natl Acad Sci U S A.* 2005;102(24):8603-8608.
28. Wilkinson AC, Ballabio E, Geng H, et al. RUNX1 is a key target in t(4;11) leukemias that contributes to gene activation through an AF4-MLL complex interaction. *Cell Rep.* 2013;3(1):116-127.
29. Zhou Y, Zhou B, Pache L, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun.* 2019;10(1):1523.
30. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012;9(4):357-359.
31. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.* 2013;29(1):15-21.
32. O'Connell RM, Balazs AB, Rao DS, Kivork C, Yang L, Baltimore D. Lentiviral vector delivery of human interleukin-7 (hIL-7) to human immune system (HIS) mice expands T lymphocyte populations. *PLoS One.* 2010;5(8):e12009.
33. Minuesa G, Albanese SK, Xie W, et al. Small-molecule targeting of MUSASHI RNA-binding activity in acute myeloid leukemia. *Nat Commun.* 2019;10(1):2691.
34. Muralidharan R, Mehta M, Ahmed R, et al. HuR-targeted small molecule inhibitor exhibits cytotoxicity towards human lung cancer cells. *Sci Rep.* 2017;7(1):9694.
35. Ahmed R, Muralidharan R, Srivastava A, et al. Molecular targeting of HuR oncoprotein suppresses MITF and induces apoptosis in melanoma cells. *Cancers.* 2021;13(2):166.
36. Wang L, Rowe RG, Jaimes A, et al. Small-molecule inhibitors disrupt let-7 oligouridylation and release the selective blockade of let-7 processing by LIN28. *Cell Rep.* 2018;23(10):3091-3101.
37. Itskovich SS, Gurunathan A, Clark J, et al. MBNL1 regulates essential alternative RNA splicing patterns in MLL-rearranged leukemia. *Nat Commun.* 2020;11(1):2369.
38. Weng H, Huang F, Yu Z, et al. The m(6)A reader IGF2BP2 regulates glutamine metabolism and represents a therapeutic target in acute myeloid leukemia. *Cancer Cell.* 2022;40(12):1566-1582.e10.
39. Costello RT, Mallet F, Gaugler B, et al. Human acute myeloid leukemia CD34+/CD38- progenitor cells have decreased sensitivity to chemotherapy and Fas-induced apoptosis, reduced immunogenicity, and impaired dendritic cell transformation capacities. *Cancer Res.* 2000;60(16):4403-4411.
40. Eppert K, Takenaka K, Lechman ER, et al. Stem cell gene expression programs influence clinical outcome in human leukemia. *Nat Med.* 2011;17(9):1086-1093.
41. Bachas C, Schuurhuis GJ, Assaraf YG, et al. The role of minor subpopulations within the leukemic blast compartment of AML patients at initial diagnosis in the development of relapse. *Leukemia.* 2012;26(6):1313-1320.
42. Shlush LI, Mitchell A, Heisler L, et al. Tracing the origins of relapse in acute myeloid leukaemia to stem cells. *Nature.* 2017;547(7661):104-108.

43. Blanco JG, Dervieux T, Edick MJ, et al. Molecular emergence of acute myeloid leukemia during treatment for acute lymphoblastic leukemia. *Proc Natl Acad Sci U S A*. 2001;98(18):10338-10343.
44. Rayes A, McMasters RL, O'Brien MM. Lineage switch in MLL-rearranged infant leukemia following CD19-directed therapy. *Pediatr Blood Cancer*. 2016; 63(6):1113-1115.
45. Ennajdaoui H, Howard JM, Sterne-Weiler T, et al. IGF2BP3 modulates the interaction of invasion-associated transcripts with RISC. *Cell Rep*. 2016; 15(9):1876-1883.
46. Xueqing H, Jun Z, Yueqiang J, et al. IGF2BP3 may contribute to lung tumorigenesis by regulating the alternative splicing of PKM. *Front Bioeng Biotechnol*. 2020;8:679.
47. Huang H, Weng H, Sun W, et al. Recognition of RNA N(6)-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. *Nat Cell Biol*. 2018;20(3):285-295.
48. Elcheva IA, Spiegelman VS. Targeting RNA-binding proteins in acute and chronic leukemia. *Leukemia*. 2021;35(2):360-376.
49. Barbieri I, Tzelepis K, Pandolfini L, et al. Promoter-bound METTL3 maintains myeloid leukaemia by m6A-dependent translation control. *Nature*. 2017; 552(7683):126-131.
50. Okuda H, Miyamoto R, Takahashi S, et al. RNA-binding proteins of KHDRBS and IGF2BP families control the oncogenic activity of MLL-AF4. *Nat Commun*. 2022;13(1):6688.
51. Velasco-Hernandez T, Säwén P, Bryder D, Cammenga J. Potential pitfalls of the Mx1-Cre system: implications for experimental modeling of normal and malignant hematopoiesis. *Stem Cell Reports*. 2016;7(1):11-18.