

Lawrence Berkeley National Laboratory

Recent Work

Title

THE VAPOR PHASE HYDROLYSIS OP THE RARE EARTH HALIDES

Permalink

<https://escholarship.org/uc/item/1ts84546>

Authors

Koch, C. W.

Cunningham, B. B.

Publication Date

1952-08-13

UNIVERSITY OF CALIFORNIA - BERKELEY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.



UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

THE VAPOR PHASE HYDROLYSIS OF THE RARE EARTH HALIDES

II. Heat and Free Energy of The Reactions:





C. W. Koch and B. B. Cunningham

August 13, 1952

THE VAPOR PHASE HYDROLYSIS OF THE RARE EARTH HALIDES

II. Heat and Free Energy of the Reactions:




C. W. Koch and B. B. Cunningham
 Department of Chemistry and Radiation Laboratory
 University of California, Berkeley, California

August 13, 1952

ABSTRACT

Measurements of equilibrium constants for the reactions:



$\text{GdCl}_3(\text{s}) + \text{H}_2\text{O}(\text{g}) = \text{GdOCl}(\text{s}) + 2\text{HCl}(\text{g})$ at various temperatures give the free energy functions



$$\Delta F^\circ (\text{kcal}) = 21.56 + \frac{6.4 T \log T}{1000} + 1.8 \times 10^{-7} T^2 - \frac{22}{T} - \frac{53.01 T}{1000}, \text{ and}$$

$$\Delta F^\circ (\text{kcal}) = 19.96 + \frac{6.4 T \log T}{1000} + 1.8 \times 10^{-7} T^2 - \frac{22}{T} - \frac{51.88 T}{1000},$$

respectively, using an estimated ΔC_p function of $-2.8 - 0.36 \times 10^{-3} T + 0.44 \times 10^5 T^{-2}$ for both reactions. Within the experimental temperature range of about 700-900° K these functions are consistent with the experimental measurements within an average deviation of about ± 0.07 kcal. Because of the possible error in the estimated ΔC_p function, however, extrapolation to 298° K involves a much larger uncertainty.

THE VAPOR PHASE HYDROLYSIS OF THE RARE EARTH HALIDES

II. Heat and Free Energy of the Reactions:



C. W. Koch and B. B. Cunningham
Department of Chemistry and Radiation Laboratory
University of California, Berkeley, California

August 13, 1952

Introduction

An experimental method of measuring equilibrium constants for the reaction: $\text{LaCl}_3(\text{s}) + \text{H}_2\text{O}(\text{g}) = \text{LaOCl}(\text{s}) + 2\text{HCl}(\text{g})$ at various temperatures has been described in a previous paper.¹ The method has now

1. C. W. Koch, A. Broido, and B. B. Cunningham, J. Am. Chem. Soc. 74, 2349 (1952).

been applied to an investigation of the corresponding reactions of samarium and gadolinium. The results are of interest because they furnish thermodynamic data not previously available for the rare earth oxychlorides and also because they provide information for a consideration of the influence of crystal geometry on the thermodynamic properties of compounds of close structural similarity. This latter subject will be considered in a subsequent paper.

Experimental

Apparatus and Technique.--A detailed description of the apparatus and experimental technique used for the hydrolysis measurements on lanthanum trichloride has been given previously.¹ No modifications of apparatus or technique were necessary for the measurements reported here. However, a platinum foil radiation shield some four inches in length has now been placed around the central portion of the reaction chamber, although it was found that the apparent temperature of the sample was not affected significantly by this change in the experimental set-up.

Materials.--Samarium was obtained as the sesquioxide from Dr. Frank Spedding of Iowa State College. Spectrographic analysis by Mr. John Conway of the Radiation Laboratory gave the following results: europium - 0.01-0.1%, calcium - 0.01-0.1%, lanthanum - 0.01-0.1%.

Gadolinium sesquioxide was purchased from Adam Hilger and Sons, London, England. The results of a spectrographic analysis of a sample of this material were as follows: iron - 0.01-0.1%, magnesium - 0.01-0.1%, and no other rare earth detected.

X-Ray Diffraction Data.--Powder patterns of rare earth trichlorides and oxychlorides prepared by the methods described in this paper were obtained by Mrs. Carol Dauben and Mrs. Helena Rubin of the x-ray diffraction group. Estimates of the crystallite size of some of our preparations were made by Professor D. H. Templeton from measurements of the line broadening of the diffraction patterns.

RESULTS

The results of the hydrolysis measurements on solid SmCl_3 and GdCl_3 are summarized in Tables I and II below. Calculated free energy values based on the function:

$$\Delta F_T^0 = \Delta H_0 + \frac{6.4 T \log T}{1000} + 1.8 \times 10^{-6} T^2 - \frac{22}{T} + I T$$

are given for comparison. This function assumes the same ΔC_p equation as that estimated for lanthanum:¹

$$\Delta C_p = -2.8 - 0.36 \times 10^{-3} T + 0.44 \times 10^5 T^{-2}.$$

The constants ΔH_0 and I were evaluated for the samarium and gadolinium reactions by taking the mean of individual values calculated from various pairs of free energy values derived from experimentally observed equilibrium constants.

The right hand columns of Tables I and II give the deviations of the experimental ΔF^0 values from the ΔF^0 values calculated from the free energy functions. Within the experimental temperature range the agreement is reasonably satisfactory.

It should be emphasized, however, that heat capacity data for the rare earth chlorides and oxychlorides are not available and the uncertainty in our estimated ΔC_p equation therefore is considerable. This leads to an uncertainty of perhaps 1.5 kcal and 2-3 entropy units in the extrapolation of our data to room temperature.

The thermodynamic constants for the hydrolysis of the solid trichlorides of lanthanum, samarium and gadolinium by water vapor are summarized in Table III.

Table I

Equilibrium Constants for the Reaction:
 $\text{SmCl}_3(\text{s}) + \text{H}_2\text{O}(\text{g}) = \text{SmOCl}(\text{s}) + 2\text{HCl}(\text{g})$

Run No.	T°K	P _{HCl} mm	P _{H₂O} mm	K _{max} K _{min}	K _{mean}	ΔF _{exp} ^o kcal	ΔF _{calc} ^o kcal	Δ kcal
7	893	418.8 406.3	1.60 1.60	144 136	140 ± 4	-8.77	-8.83	+0.06
8	885	423.7 390.8	1.77 1.74	133 116	124.5 ± 8.5	-8.48	-8.55	+0.07
9	867	454.9 442.5	2.54 2.52	107 102	104.5 ± 2.5	-8.01	-8.00	+0.01
10	848	441.9 407.8	2.99 2.89	85.9 75.7	80.8 ± 5.1	-7.40	-7.40	0.00
11	830	363.7 347.3	2.70 2.59	64.4 61.3	62.9 ± 1.6	-6.83	-6.84	+0.01
1	816	387.8 329.6	3.45 2.66	57.3 53.7	55.5 ± 1.8	-6.51	-6.40	-0.11
12	811	326.1 309.8	2.65 2.53	52.8 49.9	51.4 ± 1.5	-6.35	-6.24	-0.09
2	797	273.6 262.7	2.43 2.36	40.5 38.5	39.5 ± 1.0	-5.82	-5.81	-0.01
13	774	298.2 272.1	4.14 3.64	28.3 26.8	27.5 ± 0.8	-5.10	-5.08	-0.02
4	755	258.3 231.9	4.26 3.90	20.6 18.1	19.3 ± 1.3	-4.44	-4.48	+0.04
5	727	205.3 193.6	4.37 4.22	12.7 11.7	12.2 ± 0.5	-3.61	-3.70	+0.09
14	703	152.7 133.6	3.81 3.35	8.05 7.01	7.53 ± 0.52	-2.82	-2.84	+0.02
Average deviation								±0.05
Net deviation								+0.05

Table II

Equilibrium Constants for the Reaction:
 $\text{GdCl}_3(\text{s}) + \text{H}_2\text{O}(\text{g}) = \text{GdOCl}(\text{s}) + 2\text{HCl}(\text{g})$

Run No.	T°K	P _{HCl} mm	P _{H₂O} mm	K _{max} K _{min}	K _{mean}	ΔF° _{exp} kcal	ΔF° _{calc} kcal	Δ kcal
1	895	610 636	2.46 2.87	199 185	192 ± 7	-9.35	-9.45	+0.10
8	880	472 397	1.80 1.40	163 148	156 ± 8	-9.19	-9.00	-0.19
3	870	620 605	3.17 3.18	159 151	155 ± 4	-8.72	-8.70	-0.02
9	858	453 415	1.97 1.72	137 132	134 ± 3	-8.35	-8.34	-0.01
10	840	344 318	1.55 1.36	101 98	100 ± 2	-7.69	-7.80	+0.11
11	821	346 300	1.84 1.42	86 83	84 ± 2	-7.23	-7.23	0.00
5	800	478 387	4.36 3.22	69 60	65 ± 5	-6.64	-6.60	-0.04
12	751	322 303	4.16 4.01	32.8 30.1	31.5 ± 1.4	-5.15	-5.11	-0.04
7	701	212 218	4.08 4.66	14.4 13.5	14.0 ± 0.5	-3.68	-3.58	-0.10
Average deviation								±0.07
Net deviation								-0.19

Table III

Summary of Thermodynamic Constants For the Trichloride Hydrolysis Reactions

	ΔF_{785}°	ΔH_{785}°	ΔS_{785}°	ΔF_{298}°	ΔH_{298}°	ΔS_{298}°	ΔH_0	I
La	+1.4	25.6	30.8	17.0	27.0	33.5	27.9	-52.4
Sm	-5.4	19.2	31.4	10.4	20.6	34.0	21.6	-53.0
Gd	-6.2	17.6	30.2	9.2	19.0	32.9	20.0	-51.9

The above values are in kcal per mole and cal per degree per mole.

Neglecting electronic contributions it would be expected that ΔS for the various reactions would be very nearly the same at a given temperature.

However, since the electrical fields within the trichloride and oxychloride crystals are not of the same symmetry, the difference between the contributions of low lying energy states to the entropies of the trichloride and oxychloride of the same element, may be appreciably different for different rare earths. Such effects cannot be calculated with confidence at present, and we prefer to attribute the differences in the observed ΔS values principally to experimental error.

The most noticeable feature of the hydrolysis reactions is that their free energies become progressively more negative with increasing atomic number. A detailed consideration of this effect in terms of the crystal energies of the trichlorides and oxychlorides of lanthanum, samarium, and gadolinium will be presented in a subsequent paper.

In a previous paper¹ on the reaction of lanthanum trichloride with water vapor, we combined our thermochemical data with data given

in the Bureau of Standard's "Selected Values of Chemical Thermodynamic Properties" (Series I, Table 76-1, December 31, 1949) to calculate a heat and free energy of formation of LaOCl . It now appears^{2,3} that the

2. H. R. Lohr and B. B. Cunningham, J. Am. Chem. Soc. 73, 2025 (1951).

3. F. H. Spedding and C. F. Miller, Iowa State College Ames Laboratory Report ISC-167 (July 25, 1951).

heats of formation of the rare earth trichlorides given in the source referred to are substantially in error. It does not seem worthwhile therefore to attempt to calculate heats and free energies of formation of SmOCl and GdOCl from our data until reliable thermochemical values are available for the trichlorides.

Crystallite size estimates of some of our preparations indicated a particle size in excess of 500 \AA . It seems unlikely that the thermodynamic properties of the solid trichlorides and oxychlorides used in the equilibrium measurements were sufficiently different from those of macro crystals to have affected our results significantly.

This work was performed under the auspices of the U.S. Atomic Energy Commission.