

# UC Davis

## UC Davis Previously Published Works

### Title

LRRC10 (Leucine-Rich Repeat Containing Protein 10) and REEP5 (Receptor Accessory Protein 5) as Novel Regulators of Cardiac Excitation-Contraction Coupling Structure and Function

### Permalink

<https://escholarship.org/uc/item/1v65c20v>

### Journal

Journal of the American Heart Association, 7(3)

### ISSN

2047-9980

### Authors

Chiamvimonvat, Nipavan  
Song, Long-Sheng

### Publication Date

2018-02-06

### DOI

10.1161/jaha.117.008260

Peer reviewed

# LRRC10 (Leucine-Rich Repeat Containing Protein 10) and REEP5 (Receptor Accessory Protein 5) as Novel Regulators of Cardiac Excitation-Contraction Coupling Structure and Function

Nipavan Chiamvimonvat, MD; Long-Sheng Song, MD

In cardiomyocytes, excitation–contraction (E-C) coupling is the fundamental mechanism governing the contractile function of the heart, by which working ventricular myocytes generate sufficient contractile force to pump blood throughout the body.<sup>1</sup> In the past several decades, numerous studies have greatly increased our understanding of the structure and function of cardiac E-C coupling.<sup>2</sup> During E-C coupling, an action potential drives membrane depolarization to initiate a cascade of Ca<sup>2+</sup>-mediated processes, among which Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release is the central process controlling the efficacy and fidelity of E-C coupling. During the process of Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release, Ca<sup>2+</sup> influx through the opening of voltage-gated L-type Ca<sup>2+</sup> channels (LTCCs) located on T-tubule membrane triggers a much larger release of Ca<sup>2+</sup> from the sarcoplasmic reticulum (SR) via Ca<sup>2+</sup> release channels or ryanodine receptors residing on the terminal cisternae of the SR or junctional SR. In ventricular cardiomyocytes, T-tubules and the junctional SR closely appose each

other to form tight junctional couplings with a gap of 12 to 15 nm, dubbed cardiac dyads, which provide nanoscale ultrastructural microdomains for highly efficient and precisely controlled Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release (Figure).<sup>3</sup> It is also appreciated that the structural protein junctophilin-2 is responsible for the formation of the junctional coupling between T-tubules and the terminal cisternae of SR membranes, as well as for maintaining T-tubule integrity in a regularly arrayed organization.<sup>4,5,6</sup>

## Novel Proteins Responsible for Shaping the Junctional SR: The Receptor Expression-Enhancing Proteins Family Protein

The SR system is a network of tubules running throughout the cardiomyocytes that uptakes and stores Ca<sup>2+</sup> ions. The SR contains 3 distinct but continuous regions: the network SR, the corbular SR, and the junctional SR.<sup>7</sup> A majority of the total SR system is the network SR that is organized in an anastomosing system surrounding the myofibrils. The junctional and corbular SR are structurally specialized domains extending from the network SR. Although both contain ryanodine receptors, the junctional SR is physically connected to T-tubules or sarcolemma (as mentioned above), whereas the corbular SR is free of sarcolemmal contact. Morphologically, the junctional SR, as the terminal cisternae of the continuous SR network, changes shape drastically, from a long, randomly running tubular-like structure into a flat SR cisternae that closely apposes to the surface sarcolemma or associates along with T-tubules.<sup>8</sup> Even though much is known regarding the SR network, what remains unclear is how the junctional SR is shaped in cardiomyocytes.

In this issue of *Journal of the American Heart Association (JAHA)*, Chen and colleagues elegantly describe a novel role for REEP5 (receptor accessory protein 5) in the formation of the junctional SR.<sup>9</sup> REEP5 (also known as DP1) belongs to a family of receptor expression-enhancing proteins (REEPs) that are part of a superfamily of “morphogens” that establish

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.


From the Division of Cardiovascular Medicine, Department of Internal Medicine, School of Medicine, University of California, Davis, Davis, CA (N.C.); Department of Veterans Affairs, Northern California Health Care System, Mather, CA (N.C.); Division of Cardiovascular Medicine, Department of Internal Medicine, Abboud Cardiovascular Research Center, Carver College of Medicine, University of Iowa, Iowa City, IA (L.-S.S.); Department of Veterans Affairs, Iowa City Medical Center, Iowa City, IA (L.-S.S.).

**Correspondence to:** Nipavan Chiamvimonvat, MD, Division of Cardiovascular Medicine, Department of Internal Medicine, University of California, Davis, One Shields Ave, GBSF 6303, Davis, CA 95616; and Department of Veterans Affairs, Northern California Health Care System, 10535 Hospital Way, Mather, CA 95655. E-mail: nchiamvimonvat@ucdavis.edu

Long-Sheng Song, MD, Division of Cardiovascular Medicine, Department of Internal Medicine, Abboud Cardiovascular Research Center, Carver College of Medicine, University of Iowa, 285 Newton Rd, Iowa City, IA 52242. E-mail: long-sheng-song@uiowa.edu

*J Am Heart Assoc.* 2018;7:e008260. DOI: 10.1161/JAHA.117.008260.

© 2018 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.



**Figure.** Novel E-C coupling partners uncovered. New findings indicate that LRR10 is a novel LTCC accessory protein, and REEP5 plays an important role in shaping the morphology of junctional SR. JP2 indicates junctophilin type 2; LRR10, leucine-rich repeat-containing protein 10; LTCC, L-type  $\text{Ca}^{2+}$  channel; REEP5, receptor expression-enhancing protein 5; RyR, ryanodine receptor; SERCA, sarco/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase; SR, sarco/endoplasmic reticulum; T-tubule, transverse tubule.

the curvature of the endoplasmic reticulum.<sup>10</sup> While mutations in REEP1 and REEP6 have been associated with neurological and retinal disorders, respectively, the function of REEP proteins in the heart is completely unknown. In the article by Chen and colleagues, the authors demonstrate that REEP5 is the only REEP family protein enriched in the heart and colocalizes with ryanodine receptors in the junctional SR region. Using a novel CRISPR/Cas9-based REEP5 knockout rat model, focused-ion-beam scanning electron microscopy was used to reconstruct the ultrastructure of cardiac SR, with a focus on the junctional SR. These sophisticated and state-of-the-art techniques revealed that REEP5 deficiency alters the morphology of the junctional SR, resulting in a defect in the connections between the SR and T-tubules. At the cellular level, REEP5 deficiency reduces the triggering efficiency of L-type  $\text{Ca}^{2+}$  current and decreases  $\text{Ca}^{2+}$  release from the SR. This defect in  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release under REEP5 deficiency culminates in compromised cardiac contractility.

## Novel Auxiliary Subunit of Cardiac LTCC and Its Roles in Dilated Cardiomyopathy

Leucine-rich repeats (LRRs) are 20 to 29 amino acid motifs that are found in many proteins of diverse biological

functions and contain a conserved 11-residue segment with the consensus sequence  $\text{LxxLxLxx}^{\text{N}}/\text{c}^{\text{N}}\text{L}$  (x represents any amino acid and L positions can be occupied by valine, isoleucine, or phenylalanine).<sup>11</sup> Recent structural analyses have provided insights into the role of leucine-rich repeats as versatile structural domains critical for protein-protein interactions, including a curved overall shape with mostly helical elements on the convex side and a parallel  $\beta$  sheet on the concave side where the protein-protein interactions occur.<sup>11</sup> LRR-containing (LRR) protein 10 (LRR10) is a cardiac-specific member of the LRR superfamily with critical roles in cardiac function.<sup>12</sup> It contains 7 LRR motifs with no other known functional domains. LRR10 has been shown to be highly conserved and expressed exclusively in cardiomyocytes where it localizes to the dyad region. Previous studies in zebrafish demonstrated that *Lrrc10* plays important roles in cardiac function and that knockdown of *Lrrc10* in zebrafish results in cardiac developmental abnormalities with reduced cardiac function and death.<sup>13</sup> Subsequent studies in mice supported these earlier findings where homozygous knockout of *Lrrc10* results in dilated cardiomyopathy (DCM).<sup>14</sup> Critical roles of LRR10 in cardiac function were further solidified where mutations in *LRR10* have recently been identified in human DCM.<sup>15</sup> However, the exact mechanisms of how LRR10 may regulate cardiac function remain unknown.

In an intriguing and comprehensive study by Woon and colleagues in the current issue of *JAHA*,<sup>16</sup> the authors used whole exome sequencing to identify a homozygous recessive variant (I195T) in the LRR region of *LRR10* in a sporadic case of pediatric DCM. Comprehensive mechanistic study reveals the unique roles of LRR10 in the modulation of cardiac LTCC. Specifically, it was established that LRR10 forms multiprotein complexes with  $\text{Ca}_v1.2$  LTCC. Co-expression of LRR10 with  $\text{Ca}_v1.2$  in a heterologous expression system results in a significant increase in L-type  $\text{Ca}^{2+}$  current density while the opposite effect is observed when the mutant LRR10 (I195T) was used for the co-expression. More importantly, co-expression of LRR10 or mutant LRR10 with  $\text{Ca}_v1.2$  significantly impacts the voltage-dependent gating of the channels with hyperpolarization and depolarization shift in the steady-state activation, respectively. There is no alteration in channel trafficking to the membrane. These data strongly support the critical roles of LRR10 as an auxiliary subunit of  $\text{Ca}_v1.2$   $\text{Ca}^{2+}$  channel. Quantitative analyses show that co-expression of LRR10 results in a decrease in the late  $\text{Ca}^{2+}$  current, while co-expression of the mutant LRR10 increases the late  $\text{Ca}^{2+}$  current. Indeed, the increase in the late  $\text{Ca}^{2+}$  current may lead to increased  $\text{Ca}^{2+}$  loading and DCM. Finally, the authors corroborate their findings by taking advantage of the *Lrrc10* knockout mice.

## New Insights Into the Regulatory Mechanisms for Cardiac E-C Coupling, Cardiomyopathy, and Heart Failure

DCM and hypertrophic cardiomyopathy are the most common primary myocardial diseases with a prevalence of 1 in 2500 and 1 in 500 individuals, respectively.<sup>17</sup> Patients with either form of cardiomyopathy have increased propensity for fatal arrhythmias and sudden cardiac death. Over the past several decades, a large number of mutations in sarcomeric proteins have been shown to be linked to hypertrophic cardiomyopathy. In contrast, DCM has been associated with mutations in a wide range of genes encoding for proteins of the Z-disc, costamere, sarcolemma, nuclear lamina, and cytoskeletal proteins.<sup>17</sup> In addition, up to 50% of cases of DCM are caused by acquired pathogenesis including ischemia and viral myocarditis. However, the underlying causes in a significant number of patients with DCM remain unknown. Therefore, there is a critical need to determine new genetic determinants underlying DCM. Moreover, insights into novel proteins mechanistically linked to DCM help in further understanding the underlying pathophysiology of acquired causes of DCM.

The 2 complementary articles in this issue of *JAHA* reveal novel regulatory mechanisms for cardiac E-C coupling. Importantly, these findings help to fill our knowledge gaps about the exquisite and intricate control of cardiac E-C coupling and how the defects may result in DCM and cardiac dysfunction. Specifically, Yao et al demonstrate that REEP5 plays a critical role in shaping the SR at the junctions and that defect in REEP5 deforms the architecture of junctional SR, resulting in a decrease in cardiac contractility.<sup>9</sup> Moreover, REEP5 is downregulated in a chronic pressure-induced model of heart failure, supporting its possible contribution to cardiac dysfunction. Nonetheless, additional studies are needed to further determine the specific mechanisms of how REEP5 maintains the integrity of the junctional SR membrane. In addition, the mechanistic underpinnings of the downregulation of REEP5 in heart failure require further investigations.

Complementary findings by Woon et al provide new mechanistic insights into regulation of LTCC gating by identifying LRRC10 as a novel auxiliary subunit.<sup>16</sup> In a parallel study, LRRC26 proteins have been shown to modulate the gating of a large conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  (BK) channel by enhancing the allosteric coupling between voltage-sensor activation and the channel's closed–open transition. This results in a large hyperpolarization shift in voltage dependence and plays important roles for activation of BK channels in nonexcitable cells.<sup>18</sup> Here, Woon and coworkers demonstrated with compelling evidence that defects in LRRC10 result in a decrease in peak  $\text{Ca}^{2+}$  current but an increase in delayed current, which can predispose patients to  $\text{Ca}^{2+}$

overload as a possible pathophysiologic mechanism for DCM. Further substantiating the role of LRRC10 in normal cardiac function, the same group of investigators has previously shown that *Lrrc10* knockout mice develop severe cardiac dysfunction and ventricular dilation following mild pressure overload.<sup>19</sup> However, additional studies are required to fully decipher the interacting partners of LRRC10 within the macromolecular complex of LTCCs. LRRC10 proteins are localized within the dyad regions and may interact and regulate other component(s) of the molecular complexes involved in cardiac E-C coupling. LRRC10 was shown to interact with  $\alpha$ -actinin and  $\alpha$ -sarcomeric actin at the Z-disc in cardiomyocytes.<sup>20</sup> The biophysical effects of LRRC10 on the voltage-dependent gating of  $\text{Ca}_v1.2$  channels require additional study to fully dissect the molecular mechanisms for the observed changes. Since LRRC10 also induces a hyperpolarization shift in the voltage-dependent activation of LTCCs similar to the effects of LRRC26 on BK channels, it is tempting to hypothesize that LRRC10 may also enhance the allosteric coupling between the voltage sensor and the channel's closed–open transition similar to BK channels. However, this remains to be investigated. Structural information on the LRRC protein family may help to further determine the mechanistic interactions of LRRC10 with  $\text{Ca}_v1.2$ . It is also appealing to hypothesize that there may be a cohort of LRRC family members that facilitate the fine tuning of the voltage dependence of other ion channels. Finally, possible changes of LRRC10 in the acquired type of DCM need further investigations.

Thus, the 2 articles in this issue of *JAHA* have further peeled back and revealed the multiple layers of the intricate and precise regulation of cardiac E-C coupling. Importantly, these studies help to raise new unanswered questions: Are there additional interacting partners within the macromolecular complex of LTCCs? Do changes in the expression of LRRC10 and REEP5 lead to cardiomyopathies from different causes? How can we leverage this new information to design mechanism-based treatment for DCM and heart failure? With emerging tools such as high-resolution imaging and protein structure information as well as patient-specific pluripotent stem cell–derived cardiomyocytes, the answers to these questions may be closer than we think (perhaps even within 12–15 nm).

### Acknowledgments

The authors would like to acknowledge artistic help from Rafael Shimkunas and Shawn Roach for the figure in the editorial.

### Sources of Funding

This work is supported by NIH R01s (HL137228, HL085844, HL085727, HL090905 & HL130346), and VA Merit Awards (I01-BX000576, I01-CX001490, and I01-BX002344).

## Disclosures

None.

## References

- Bers DM. Cardiac excitation-contraction coupling. *Nature*. 2002;415:198–205.
- Eisner DA, Caldwell JL, Kistamas K, Trafford AW. Calcium and excitation-contraction coupling in the heart. *Circ Res*. 2017;121:181–195.
- Wang SQ, Song LS, Lakatta EG, Cheng H. Ca<sup>2+</sup> signalling between single L-type Ca<sup>2+</sup> channels and ryanodine receptors in heart cells. *Nature*. 2001;410:592–596.
- Takeshima H, Komazaki S, Nishi M, Iino M, Kangawa K. Junctophilins: a novel family of junctional membrane complex proteins. *Mol Cell*. 2000;6:11–22.
- Wei S, Guo A, Chen B, Kutschke W, Xie YP, Zimmerman K, Weiss RM, Anderson ME, Cheng H, Song LS. T-tubule remodeling during transition from hypertrophy to heart failure. *Circ Res*. 2010;107:520–531.
- Guo A, Zhang X, Iyer VR, Chen B, Zhang C, Kutschke WJ, Weiss RM, Franzini-Armstrong C and Song LS. Overexpression of junctophilin-2 does not enhance baseline function but attenuates heart failure development after cardiac stress. *Proc Natl Acad Sci U S A*. 2014;111:12240–12245.
- Jorgensen AO, Shen AC, Arnold W, McPherson PS, Campbell KP. The Ca<sup>2+</sup>-release channel/ryanodine receptor is localized in junctional and corbular sarcoplasmic reticulum in cardiac muscle. *J Cell Biol*. 1993;120:969–980.
- Franzini-Armstrong C, Protasi F, Tijskens P. The assembly of calcium release units in cardiac muscle. *Ann N Y Acad Sci*. 2005;1047:76–85.
- Yao L, Xie D, Geng L, Shi D, Huang J, Wu Y, Lv F, Liang D, Li L, Liu Y, Li J, Chen YH. REEP5 (Receptor Accessory Protein 5) acts as a sarcoplasmic reticulum membrane sculptor to modulate cardiac function. *J Am Heart Assoc*. 2018;7:e007205. DOI: 10.1161/JAHA.117.007205.
- Goyal U, Blackstone C. Untangling the web: mechanisms underlying ER network formation. *Biochim Biophys Acta*. 2013;1833:2492–2498.
- Kobe B, Kajava AV. The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol*. 2001;11:725–732.
- Brody MJ, Lee Y. The role of leucine-rich repeat containing protein 10 (LRRC10) in dilated cardiomyopathy. *Front Physiol*. 2016;7:337.
- Kim KH, Antkiewicz DS, Yan L, Eliceiri KW, Heideman W, Peterson RE, Lee Y. Lrrc10 is required for early heart development and function in zebrafish. *Dev Biol*. 2007;308:494–506.
- Manuylov NL, Manuylova E, Avdoshina V, Tevosian S. Serdin1/Lrrc10 is dispensable for mouse development. *Genesis*. 2008;46:441–446.
- Qu XK, Yuan F, Li RG, Xu L, Jing WF, Liu H, Xu YJ, Zhang M, Liu X, Fang WY, Yang YQ, Qiu XB. Prevalence and spectrum of LRRC10 mutations associated with idiopathic dilated cardiomyopathy. *Mol Med Rep*. 2015;12:3718–3724.
- Woon MT, Long PA, Reilly L, Evans JM, Keefe AM, Lea MR, Beglinger CJ, Balijepalli RC, Lee Y, Olson TM, Kamp TJ. Pediatric dilated cardiomyopathy-associated LRRC10 (Leucine-Rich Repeat-Containing 10) variant reveals LRRC10 as an auxiliary subunit of cardiac L-type Ca<sup>2+</sup> channels. *J Am Heart Assoc*. 2018;7:e006428. DOI: 10.1161/JAHA.117.006428.
- McNally EM, Barefield DY, Puckelwartz MJ. The genetic landscape of cardiomyopathy and its role in heart failure. *Cell Metab*. 2015;21:174–182.
- Yan J, Aldrich RW. LRRC26 auxiliary protein allows BK channel activation at resting voltage without calcium. *Nature*. 2010;466:513–516.
- Brody MJ, Feng L, Grimes AC, Hacker TA, Olson TM, Kamp TJ, Balijepalli RC, Lee Y. LRRC10 is required to maintain cardiac function in response to pressure overload. *Am J Physiol Heart Circ Physiol*. 2016;310:H269–H278.
- Brody MJ, Hacker TA, Patel JR, Feng L, Sadoshima J, Tevosian SG, Balijepalli RC, Moss RL, Lee Y. Ablation of the cardiac-specific gene leucine-rich repeat containing 10 (Lrrc10) results in dilated cardiomyopathy. *PLoS One*. 2012;7:e51621.

---

**Key Words:** Editorials • calcium channel • calcium regulation • cardiomyopathy • excitation-contraction coupling • sarcoplasmic reticulum