UC Davis

UC Davis Previously Published Works

Title

Variation in the restriction fragments of 18S-26S rRNA loci in South American Elymus (Triticeae)

Permalink

<https://escholarship.org/uc/item/1w016824>

Journal

Genome, 35(5)

ISSN

0831-2796

Authors

Dubcovsky, J
Lewis, SM
Hopp, EH

Publication Date

1992

DOI

10.1139/g92-133

Peer reviewed

Variation in the restriction fragments of 18S–26S rRNA loci in South American *Elymus* (Triticeae)

JORGE DUBCOVSKY¹ AND SILVINA M. LEWIS

*Instituto de Recursos Biológicos, INTA, Las cabañas y Los Reseros s/n, Villa Udaondo,
1712 Castelar, Buenos Aires, Argentina*

AND

ESTEBAN H. HOPP

*Instituto de Biología Molecular, INTA, Las cabañas y Los Reseros s/n, Villa Udaondo,
1712 Castelar, Buenos Aires, Argentina*

Corresponding Editor: R. Appels

Received April 6, 1992

Accepted April 23, 1992

DUBCOVSKY, J., LEWIS, S. M., and HOPP, E. H. 1992. Variation in the restriction fragments of 18S–26S rRNA loci in South American *Elymus* (Triticeae). *Genome*, **35**: 881–885.

The repeat units of the 18S–5.8S–26S ribosomal genes (rDNA) of 15 South American and 6 Asiatic and North American species of *Elymus* were shown to vary between 8.7 and 11.3 kb among species. Interspecific variation in the *Eco*RI and *Bam*HI cleavage sites was observed. *Bam*HI restriction sites were present in the subrepeats in the intergenic spacers in rDNA from diploid *Hordeum* (H genome) and absent in the diploid *Pseudoroegneria* (S genome) and in nearly all the tetraploid species of *Elymus* (SH genomes). Tetraploid *Elymus* species have only two pairs of chromosomes with secondary constrictions similar to those found in *Pseudoroegneria* and a maximum of two different repeat unit length classes. These results suggest that the rDNA in the H genome was lost during the evolution of the tetraploid *Elymus*. However, the additional H genome present in the hexaploid *Elymus* with genome formula SSHHH' H' still show active ribosomal genes. Nucleolar organizer activity was also observed in rDNA from both the S and H genomes in artificial SS × HH amphiploids, suggesting no complete nucleolar dominance (amphiplasty) between the S and H genomes.

Key words: *Elymus*, ribosomal genes, RFLP, nucleolus.

DUBCOVSKY, J., LEWIS, S. M., et HOPP, E. H. 1992. Variation in the restriction fragments of 18S–26S rRNA loci in South American *Elymus* (Triticeae). *Genome*, **35** : 881–885.

Les unités répétitives des gènes ribosomiques (ADNr) de 18S–5.8S–26S de 15 espèces d'*Elymus* sud-américaines et de 6 espèces d'*Elymus* asiatiques et nord-américaines ont présenté, parmi ces espèces, des variations de 8,7 à 11,3 kb. Des variations interspécifiques ont été observées dans les sites de clivage d'*Eco*RI et de *Bam*HI. Les sites de restriction de *Bam*HI ont été présents dans les sous-répétitions au niveau des espaces intergénomiques chez les ADNr d'*Hordeum* diploïdes (génom H), mais ils furent absents chez les *Pseudoroegneria* diploïdes (génom S) et chez presque toutes les espèces d'*Elymus* tétraploïdes (génom SH). Les espèces tétraploïdes d'*Elymus* n'ont que deux paires de chromosomes qui possèdent des constrictions secondaires semblables à celles qui sont observées chez les *Pseudoroegneria* et un maximum de deux classes différentes d'unités de longueur répétitives. Ces résultats suggèrent que l'ADNr du génome H a été perdu au cours de l'évolution des *Elymus* tétraploïdes. Toutefois, le génome H additionnel qui est présent chez les *Elymus* hexaploïdes de formule génomique SSHHH' H' montre, néanmoins, des gènes ribosomiques actifs. L'activité de l'organisateur nucléolaire a aussi été observée dans l'ADNr à la fois des génomes S et H chez des amphiploïdes SS × HH, ce qui suggère l'absence d'une dominance nucléolaire complète (amphiplastie) entre les génomes S et H.

Mots clés : *Elymus*, gènes ribosomiques, RFLP, nucléoles.

[Traduit par la rédaction]

Introduction

The genus *Elymus* L. has a worldwide distribution and is the largest among the perennial Triticeae. The majority of species (75%) are tetraploids, but hexaploids and octoploids are also found. These allopolyploids show different combinations of the SS genomes from *Pseudoroegneria*, HH from *Hordeum* and YY from an unknown diploid (Dewey 1984).

The SSHH tetraploids have four chromosomes with secondary constrictions and up to four nucleoli per nucleus (Hunziker 1966a; Dubcovsky et al. 1989; Morris and Gill 1987). This number is lower than that expected from the combination of SS and HH diploids, since four chromosomes with secondary constrictions are observed in diploid *Pseudoroegneria* and two or four are common in the diploid *Hordeum* (Hsiao et al. 1986).

Gill et al. (1988) assigned the two NOR loci in *Elymus trachycaulus* (SSHH) to the S genome. Chromosome morphology (Dubcovsky et al. 1989) and C-banding patterns (Morris and Gill 1987) showed that the chromosomes with secondary constrictions from the tetraploid *Elymus* corre-

¹Author for correspondence at the following address: Department of Agronomy and Range Science, University of California, Davis, CA 95616-8515, U.S.A.

TABLE 1. Materials, chromosome number, maximum nucleoli per nucleus and ribosomal DNA repeat unit length variants in *Elymus* species

Species	Country	Accession No.	2n	Maximum No. of nucleoli	rDNA repeat unit length (kb)	IGS <i>Bam</i> HI sites	IGS <i>Eco</i> RI site
South American spp.							
<i>E. agropyroides</i> Presl.	Argentina	657	28	4	10.1, 10.5	No	No
<i>E. andinus</i> Trin.	Chile	825	28	4	9.7, 10.3	No	No
<i>E. angulatus</i> Presl.	Argentina	H-6419	28	4	10.2, 10.6	No	No
<i>E. antarcticus</i> Hook. f.	Argentina	O-230	28	4	9.6, 10.3	No	No
<i>E. araucanus</i> (Parodi) Löve	Argentina	655	28	4	9.8	No	No
<i>E. attenuatus</i> Löve	Peru	H-6486	28	4	9.0, 9.3	No	Yes
<i>E. breviaristatus</i>							
ssp. <i>scabrifolius</i> (Döll) Löve	Argentina	B-880011	28	2	9.5	No	No
<i>E. gayanus</i> Desvaux	Chile	833	28	4	9.8, 10.4	No	No
<i>E. glaucescens</i> Seberg	Argentina	883	28	4	9.6, 10.1	No	No
<i>E. glaucescens</i> Seberg	Argentina	270	28	4	9.7, 10.3	No	No
<i>E. rigescens</i> Trin.	Argentina	630	28	4	9.7, 10.6	No	No
<i>E. tilcarensis</i> (Hunziker) Löve	Argentina	413	28	4	9.4, 9.7	No	No
<i>E. erianthus</i> Philippi	Argentina	562	42	6	9.4, 9.7, 10.1	No	Yes
<i>E. erianthus</i> Philippi	Argentina	O-213	42	6	9.4, 9.7, 10.3	No	Yes
<i>E. patagonicus</i> Speg.	Argentina	626	42	8	10.0, 10.3, 10.5	Yes	No
<i>E. patagonicus</i> Speg.	Chile	PI-286201	42	7	10.0, 10.5	Yes	No
<i>E. patagonicus</i> Speg.	Argentina	PI-297898	42	8	9.6, 10.0, 10.3, 10.5	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	H-6455	42	8	9.5, 9.8	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	412	42	8	9.5, 9.8	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	H-6466	42	8	9.2, 9.5, 9.9, 10.1	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	B-890067	42	8	9.4, 9.6, 9.8, 11.3	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	B-890073	42	8	9.4, 9.7, 9.9, 10.4	Yes	No
<i>E. scabriglumis</i> (Hackel) Löve	Argentina	PI-269646	42	8	9.5, 9.8, 10.1, 10.4	Yes	No
<i>E. mendocinus</i> (Parodi) Löve	Argentina	601	56	6	9.5, 9.8, 10.2	No	No
North American and Asiatic species							
<i>E. canadensis</i> L.	U.S.A.	PI-232249	28	4	9.8	Yes	No
<i>E. fibrosus</i> (Schrenk) Tzvelev	U.S.S.R.	PI-315493	28	4	9.0, 9.3	No	No
<i>E. glaucus</i> Buckl.	U.S.A.	PI-232263	28	4	9.6	No	No
<i>E. mutabilis</i> (Drobov) Tzvelev	U.S.S.R.	PI-314204	28	4	9.2, 9.6	No	No
<i>E. trachycaulus</i> (Link) Gould	U.S.A.	PI-232156	28	4	8.9, 9.2	No	No
<i>E. sibiricus</i> L.	P.R. China	PI-499619	28	4	8.7, 9.6	No	No
Diploids and artificial amphiploids							
<i>P. libanotica</i> (Hackel) Dewey	Iran	PI-401325	14	4	9.4, 9.9, 10.2	No	No
<i>P. spicata</i> (Pursh) Löve	Canada	PI-236670	14	4	10.4	No	No
<i>H. violaceum</i> Boiss. & Hohen.	Iran	PI-401390	14	4	9.4, 9.7	Yes	No
Hybrid 1			28	7	9.8, 10.0, 10.2	Yes	No
Hybrid 2			28	8	9.5, 10.0	Yes	No
Hybrid 17			28	7	9.4, 10.0, 10.3	Yes	No
Hybrid 14			27	7	9.4, 10.0, 10.4, 11.0	Yes	No
Hybrid 3			28	6	9.2, 10.0	Yes	No

spond to those of *Pseudoroegneria*. This suggests that the rDNA of the H genome is either not transcriptionally active in the SSHH tetraploids (amphiplasty) or that it was lost during the evolution of the tetraploid *Elymus*.

Ribosomal genes are well characterized in many cereals. They are organized as families of repeated genes in tandem arrays. Each repeat unit of rDNA contains a conserved coding region as well as variable intergenic spacer (IGS). This variation is primarily due to variation in the number of subrepeats in the IGS and to a lesser extent to the loss or gain of restriction sites within the IGS (for review see Appels and Honeycutt 1986).

The majority of the *Hordeum* species exhibit one or more *Bam*HI sites in the IGS not found in other Triticeae (Appels et al. 1980; Molnar et al. 1989; Molnar and Fedak 1989). Among the five different patterns of the distribution of

*Bam*HI sites described for rDNA in *Hordeum*, the most widely distributed, particularly among the H genome, *Hordeum* was that with additional *Bam*HI sites within the IGS subrepeats.

In the present paper we investigated the presence of these characteristic H-genome additional *Bam*HI sites in the ribosomal genes of some artificial SS × HH amphiploids (Asay et al. 1987) and in natural tetraploid, hexaploid, and octoploid *Elymus*. We also examined the maximum number of nucleoli in these species to infer the number of transcriptionally active ribosomal loci.

Materials and methods

Plant material

Species analyzed are displayed in Table 1, which also shows the collection country and the identification number.

Seeds with PI numbers were provided by Dr. Dewey and Dr. Jensen (U.S.A.), H numbers by Dr. Seberg (Denmark), B numbers by Ing. Duplancic (Experimental Agricultural Station, INTA Balcarce), and O numbers by Lic. Oliva (Experimental Agricultural Station, INTA, Río Gallegos). Identifications provided by the collectors were followed, except for *E. cf. agropyroides* PI-269646, which was included in *E. scabriglumis*. The other species were collected by the authors and identified using the keys of Nicora (1978). Löve's (1984) nomenclature was followed, but *Hordeum* was considered sensu Bothmer et al. (1991). Voucher specimens were deposited in the herbarium of the IRB-INTA Castelar (BAB).

The hybrid seeds *Pseudoroegneria spicata* PI-236670 × *Hordeum violaceum* PI-401390 sent by Dr. Dewey were obtained after open-pollination of the C1 amphiploids (Asay et al. 1987).

Chromosome numbers

Metaphase chromosomes were observed in root meristems pretreated in a saturated solution of *p*-dichlorobenzene at 4°C for 20 h and fixed in Carnoy's solution (6:3:1 ethanol – chloroform – acetic acid). After hydrolysis in 1 M HCl at 60°C for 10 min, chromosomes were stained in basic fuchsin and squashed in 2% (w/v) acetic orcein.

Nucleoli

Nucleoli were stained with a solution of 1 g of silver nitrate dissolved in 1 mL 0.008% (w/v) sodium citrate in distilled water, in a wet chamber at 60°C. Three or more root tips per plant and young microspores were analyzed and the maximum number of nucleoli was scored. A detailed analysis of nucleolar frequencies, number of chromosomes with secondary constrictions, and karyotypes of the South American *Elymus* will be published elsewhere (S.M. Lewis, J. Dubcovsky, and A.J. Martinez, unpublished).

DNA analyses

Total cellular DNA was isolated from leaves of individual plants following the procedure of Dellaporta et al. (1983). DNA (3 µg) was digested with restriction endonuclease *Bam*HI. Also, *Eco*RI and *Sst*I digestions were analyzed to estimate the total length of the repetitive unit. Double digestions with *Eco*RI–*Bam*HI were performed to map additional *Eco*RI or *Bam*HI restriction sites within the IGS. After electrophoresis in 0.7–1% agarose gels, the DNA was transferred by blotting to Nytran nylon membranes according to supplied instructions.

Plasmid pTA71, with a 8.95-kb *Eco*RI fragment of wheat rDNA (Gerlach and Bedbrook 1979) was labelled by random priming (Pharmacia, N.J.). Hybridization was performed at 65°C overnight and the final wash was for 15 min at 65°C in 0.2× SSC and 0.1% SDS buffer. *Hind*III digested bacteriophage lambda DNA was used as the molecular size standard.

Results

Digestions of genomic DNA with *Sst*I resulted in an invariant 3.9-kb fragment and variable 4.8- to 7.4-kb fragments. The invariant fragment spans the coding region of the 18S–5.8S–26S rRNA genes and the variant one includes the IGS in all the Triticeae analyzed so far (Appels et al. 1980).

Combinations of two or more different repeat unit length

FIG. 1. Restriction patterns of *Bam*HI-digested rDNA: A, *Pseudoroegneria libanotica*; B, *P. spicata*; C, *Hordeum violaceum*; D, E, and F, descendants from *P. spicata* × *H. violaceum* amphiploids. Plants numbers 1, 2, and 17; G, *Elymus agropyroides*; H, *E. andinus*; I, *E. antarcticus*; J, *E. angulatus*; K, *E. gayanus*; L, *E. rigescens*; M, *E. araucanus*; N, *E. glaucescens* (883); O, *E. glaucescens* (270); P, *E. breviaristatus* ssp. *scabrifolius*; Q, *E. scabriglumis* (H-6466); R, *E. scabriglumis* (H-6455); S, *E. scabriglumis* (H-6466); T, *E. scabriglumis* (B-890067); U, *E. scabriglumis* (B-890073); V, *E. patagonicus* (626); W, *E. erianthus* (562); X, *E. mendocinus*. Left lanes are molecular size marker bacteriophage λ digested with *Hind*III. Sizes are in kilobase pairs. Note the approximately 2-kb fragment originated by the IGS *Bam*HI sites in the H genome of *Hordeum* (C), artificial amphiploids (D–F), and SSHHH'H' hexaploids (Q–V).

classes occurred in 85% of the accessions (Table 1). Highly significant correlation ($r = 0.84$ $p < 0.01$) was observed between the maximum number of repeat unit length classes and the maximum number of nucleoli per species (Table 1),

suggesting that in most cases the different rDNA subunits are clustered at specific nucleolar organizer regions.

Only one *EcoRI* site per repeat unit was present in all species, except for *E. attenuatus* and *E. erianthus*. Though in both species this additional *EcoRI* site was within the IGS, double *EcoRI*-*BamHI* digestions showed differences in the distance between this additional site and the *BamHI* restriction site located in the 18S ribosomal gene (1.0 kb in *E. attenuatus* and 1.7 kb in *E. erianthus*).

Complex hybridization patterns were obtained with *BamHI*. All species show a conserved 3.9-kb fragment including the coding region (Fig. 1) and variable 4.8- to 7.4-kb fragments. The *BamHI* site in the 26S rRNA gene was not accessible to digestion in all repeating units (Appels and Honeycutt 1986) and this results in bands that correspond to the combined size of two adjacent *BamHI* fragments. Both species of *Pseudoroegneria* (Figs. 1A and 1B) and all tetraploid species of *Elymus*, except for *E. canadensis*, had only two *BamHI* sites. *Hordeum violaceum* (Fig. 1C) and *E. canadensis* had additional sites in the IGS. The observed *BamHI* restriction fragments did not add up to the length of the entire rDNA unit (Table 1). The most plausible explanation is that there is a *BamHI* site within the IGS subrepeats as suggested by Molnar and Fedak (1989). This hypothesis explains the ladders with approximately 0.1 kb periodicity in *H. violaceum* and *E. canadensis* observed when autoradiographs were overexposed. Additional *BamHI* sites in the IGS and ladders with the same periodicity were found in all populations of hexaploids *E. patagonicus* and *E. scabriglumis* (Fig. 1, Q-V). Inter- and intra-specific differences were found in the position of the next *BamHI* site closest to the 26S *BamHI* (Fig. 1, Q-V). No additional *BamHI* sites were found in the hexaploid *E. erianthus* and the octoploid *E. mendocinus* (Fig. 1, W and X).

Five tetraploid plants obtained from open-pollination of C1 artificial SSHH amphiploid between *H. violaceum* and *P. spicata* (Asay et al. 1987) showed the additional *BamHI* sites in the subrepeats of the IGS (Fig. 1, D-F). Chromosomes with secondary constrictions from *H. violaceum* and *P. spicata* are easy to differentiate (Hsiao et al. 1986) and both were observed in these plants. A maximum of eight nucleoli was found (Table 1).

Discussion

Natural SSHH tetraploid *Elymus* showed two pairs of satellited chromosomes corresponding to those of the SS genome (Morris and Gill 1987; Gill et al. 1988; Dubcovsky et al. 1989), a maximum of four nucleoli, and no more than two rDNA repeat unit length classes. In natural tetraploids, except for *E. canadensis*, no *BamHI* restriction sites in the IGS could be detected. These results indicate that the rDNA genes from the H genome in these allotetraploids lost its nucleolar organizer capability and are no longer detectable by DNA hybridization.

A different pattern was observed in the offspring of the artificial amphiploid between *P. spicata* (S genome) and *H. violaceum* (H genome) (Asay et al. 1987). These plants showed up to four repeat unit length classes and additional *BamHI* sites in the IGS. Chromosomes with secondary constrictions in positions similar to those found in both parents and six to eight nucleoli were observed suggesting

that H-genome rDNA genes were not only present but they were also active.

A similar situation was found in the natural tetraploid wheats (AABB genome pairs) where the number of rDNA repeat units from the A genome is so small that they are no longer active and are only detected with very sensitive in situ hybridization techniques (Mukai et al. 1991). On the other hand synthetic AABB tetraploids (Frankel et al. 1987) show no significant reduction of the A genome rDNA after 10 generations and, in addition, the A chromosome rDNA is transcriptionally active.

Genome analysis of the South American hexaploids *E. scabriglumis* and *E. patagonicus* suggested that these species had originated by hybridization of SSHH tetraploids with diploid species of *Hordeum* (Hunziker 1966b, 1967; Dewey 1972; Seberg 1991). The proposed SSHHH'H' genome formula is in accordance with the presence of additional *BamHI* restriction sites in the IGS subrepeats similar to those reported by Molnar et al. (1989) and Molnar and Fedak (1989) for nearly all the natural South American *Hordeum* species. The presence of four pairs of chromosomes with secondary constrictions and a maximum of eight nucleoli indicates that this additional H genome still possesses transcriptionally active ribosomal loci.

Nucleolar organizer activity of ribosomal loci from S and H genomes in the artificial SS × HH amphiploid and the SSHHH'H' hexaploids suggests no complete nucleolar dominance (amphiplasty) between the S and H genomes.

Acknowledgements

The authors are grateful to Dr. J. Dvořák for critically reading the manuscript. This work has been supported in part by grant PID-304520088 from the Argentine Research Council (CONICET) and concluded during a BID-CONICET fellowship at the Department of Agronomy and Range Sciences, University of California, Davis, Calif.

- Appels, R., and Honeycutt, R.L. 1986. rDNA: evolution over a billion years. In DNA systematics: plants. Edited by S.K. Dutta. CRC Press, Boca Raton, Fla. pp. 82-135.
- Appels, R., Gerlach, W.L., Dennis, E.S., et al. 1980. Molecular and chromosomal organization of the DNA sequences coding for the ribosomal RNAs in cereals. *Chromosoma*, **78**: 293-311.
- Asay, K.H., Hsiao, C., and Dewey, D.R. 1987. Intergeneric hybrids and amphiploids between *Pseudoroegneria spicata* and *Critiesion violaceum*. *Bot. Gaz.* **148**: 123-129.
- Bothmer, R. von, Jacobsen, N., Baden, C., et al. 1991. An eco-geographical study of the genus *Hordeum*. Systematic and ecogeographic studies on crop genepools. 7. International Board for Plant Genetic Resources, Rome.
- Dellaporta, S.L., Wood, J., and Hicks, J.B. 1983. A plant DNA miniprep: version II. *Plant Mol. Biol. Rep.* **1**: 19-21.
- Dewey, D.R. 1972. Genome analysis of South American *Elymus patagonicus* and its hybrids with two North American and two Asian *Agropyron* species. *Bot. Gaz.* **133**(4): 436-443.
- Dewey, D.R. 1984. The genomic system of classification as a guide to intergeneric hybridization with the perennial *Triticeae*. In Gene manipulation in plant improvement. Edited by J.P. Gustafson. Plenum Publishing Corp., New York. pp. 209-279.
- Dubcovsky, J., Soria, M.A., and Martinez, A.J. 1989. Karyotype analysis of the Patagonian *Elymus*. *Bot. Gaz.* **150**(4): 462-468.
- Frankel, O.H., Gerlach, W.L., and Peacock, W.J. 1987. The ribosomal RNA genes in synthetic tetraploids of wheat. *Theor. Appl. Genet.* **75**: 138-143.

- Gerlach, W.L., and Bedbrook, J.R. 1979. Cloning and characterization of ribosomal RNA genes from wheat and barley. *Nucleic Acids Res.* **7**: 1869-1885.
- Gill, B.S., Morris, K.L.D., and Appels, R. 1988. Assignment of the genomic affinities of chromosomes from polyploid *Elymus* species added to wheat. *Genome*, **30**: 70-82.
- Hsiao, C., Wang, R.R.-C., and Dewey, D.R. 1986. Karyotype analysis and genome relationships of 22 diploid species in the tribe Triticeae. *Can. J. Genet. Cytol.* **28**: 109-120.
- Hunziker, J.H. 1966a. Números cromosómicos y cariotipos de varias especies Sudamericanas de *Agropyron* y *Elymus* (*Gramineae*). *Kurtziana*, **3**: 151-156.
- Hunziker, J.H. 1966b. Diferencia cromosómica en el complejo hexaploide *Agropyron scabriglume*. *Kurtziana*, **3**: 127-149.
- Hunziker, J.H. 1967. Chromosome and protein differentiation in the *Agropyron scabriglume* complex. *Taxon*, **16**: 259-266.
- Löve, A. 1984. Conspectus of the *Triticeae*. *Feddes Repert.* **95**(7-8): 425-521.
- Molnar, S.J., and Fedak, G. 1989. Polymorphism in ribosomal DNA repeat units of 12 *Hordeum* species. *Genome*, **32**: 1124-1127.
- Molnar, S.J., Gupta, P.K., Fedak, G., and Wheatcroft, R. 1989. Ribosomal DNA repeat unit polymorphism in 25 *Hordeum* species. *Theor. Appl. Genet.* **78**: 387-392.
- Morris, K.L.D., and Gill, G.S. 1987. Genomic affinities of individual chromosomes based on C- and N-banding analysis of tetraploid *Elymus* species and their diploid progenitor species. *Genome*, **29**: 247-252.
- Mukai, Y., Endo, T.R., and Gill, B.S. 1991. Physical mapping of the 18S.26S rRNA multigene family in common wheat: identification of a new locus. *Chromosoma*, **100**: 71-78.
- Nicora, E.G. 1978. Gramíneas. In *Flora Patagónica*. Edited by M.N. Correa. Colección Científica del INTA, Buenos Aires.
- Seberg, O. 1991. Genome analysis of *Elymus angulatus* and *E. patagonicus* (Poaceae) and their hybrids with N. and S. American *Hordeum* spp. *Plant Syst. Evol.* **174**: 75-82.

Detection of polymorphic loci in *Arachis* germplasm using random amplified polymorphic DNAs

PATRICK G. LANHAM AND SARAH FENNELL

Cell and Molecular Genetics Department, Scottish Crop Research Institute, Invergowrie, Dundee DD2 5DA, Scotland

J. P. MOSS

International Crops Research Institute for the Semi-Arid Tropics, Patancheru, 502 324, Andhra Pradesh, India

AND

W. POWELL

Cell and Molecular Genetics Department, Scottish Crop Research Institute for the Semi-Arid Tropics, Patancheru, 502 324, Andhra Pradesh, India

Corresponding Editor: G. H. Jones

Received February 23, 1992

Accepted April 29, 1992

LANHAM, P. G., FENNELL, S., MOSS, J. P., and POWELL, W. 1992. Detection of polymorphic loci in *Arachis* germplasm using random amplified polymorphic DNAs. *Genome*, **35**: 885-889.

The development of easily scoreable genetic markers in *Arachis* will facilitate the introgression of desirable traits from wild species into adapted germplasm. We have used random amplified polymorphic DNAs (RAPDs) to identify polymorphic molecular markers in a range of wild and cultivated *Arachis* species. From a total of sixty 10-mer oligonucleotide primers, 49 polymorphic loci were identified between a cultivated *A. hypogaea* type (TMV-2) and a synthetic amphidiploid ($B \times C$)² created from a *A. batizocoi* and *A. chacoense* cross. The inheritance of polymorphic markers, both in the amphidiploid and in the F₁ progeny in a TMV-2 $\times$ ($B \times C$)² cross, has also been demonstrated. The potential exploitation of RAPD markers in groundnut improvement programs is discussed.

Key words: groundnut, *Arachis* species, RAPDs, amphidiploid.

LANHAM, P. G., FENNELL, S., MOSS, J. P., et POWELL, W. 1992. Detection of polymorphic loci in *Arachis* germplasm using random amplified polymorphic DNAs. *Genome*, **35** : 885-889.

Le développement de marqueurs génétiques pouvant être facilement inventoriés chez l'*Arachis* devrait faciliter l'introgression de traits désirables d'espèces indigènes dans le plasma germinal d'espèces cultivées. Des segments d'ADN amplifiés au hasard (RAPD) ont été utilisés pour identifier des marqueurs moléculaires polymorphes chez une gamme d'espèces indigènes et cultivées d'*Arachis*. Pour un total de 60 oligonucléotides promoteurs décimères, 49 locus polymorphes ont été identifiés chez des hybrides entre un type cultivé d'*A. hypogaea* (TMV-2) et un amphidiploïde synthétique ($B \times C$)², ce dernier étant issu d'un croisement entre l'*A. batizocoi* et l'*A. chacoense*. La transmission des marqueurs polymorphes, tant chez l'amphidiploïde que chez la descendance F₁ d'un croisement TMV-2 $\times$ ($B \times C$)², a aussi été démontrée. Le potentiel d'exploitation des marqueurs RAPD dans les programmes d'amélioration de l'arachide est discuté.

Mots clés : arachide, espèces d'*Arachis*, RAPD, amphidiploïde.

[Traduit par la rédaction]