

Lawrence Berkeley National Laboratory

Recent Work

Title

NOTE ON THE HAMILTONIAN FORM OF INTEGRAL SPIN WAVE EQUATIONS

Permalink

<https://escholarship.org/uc/item/1wp9q0rt>

Author

Case, K.M.

Publication Date

1955-07-11

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

NOTE ON THE HAMILTONIAN FORM OF INTEGRAL SPIN WAVE EQUATIONS

K.M. Gase

July 11, 1955

NOTE ON THE HAMILTONIAN FORM OF INTEGRAL SPIN WAVE EQUATIONS

K. M. Case

Radiation Laboratory, Department of Physics*
University of California, Berkeley, California

July 11, 1955

ABSTRACT

The Hamiltonian forms of the spin zero and spin one wave equations are obtained simultaneously starting from the Duffin-Kemmer form of the equations using only algebraic properties of the β matrices. From the mode of derivation the Hamiltonian forms for all integral spin equations of the Dirac-Fierz-Pauli¹ type follow immediately.

* Permanent address: Physics Department, University of Michigan, Ann Arbor, Michigan.

This work was performed under the auspices of the U. S. Atomic Energy Commission.

NOTE ON THE HAMILTONIAN FORM OF INTEGRAL SPIN WAVE EQUATIONS

K. M. Case

I. INTRODUCTION

Recently^{2,3} it has been shown that both the vector and scalar wave equations can be put in the Hamiltonian form

$$i \frac{\partial \Psi}{\partial t} = H \Psi, \tag{1}$$

where

$$H = \underline{\beta} \cdot \underline{P} + \kappa \beta_4, \tag{2}$$

and the wave function is subject to the initial condition

$$(H \beta_4 - \kappa) \Psi = 0. \tag{3}$$

The β_μ form a 10 or 5 dimensional representation of the Duffin-Kemmer algebra depending on whether it is the vector or scalar equation being considered.

The deductions of Eq. (1) were obtained starting from special representations of the β_μ . However, since it is seen to hold for both irreducible representations it would seem that it must be possible to obtain this form using only algebraic properties of the β_μ . This is done below. The present method has the advantages of handling spin 0 and 1 simultaneously, of being shorter, and of making the proofs of the properties of the β_μ in Eq. (2) almost trivial. Lastly this present method can be carried over to obtain the Hamiltonian form for all integral spin equations of the D-F-P type.

II. SPIN ZERO AND ONE

For simplicity we restrict ourselves to field free equations. The Duffin-Kemmer⁴ equations are

$$(\beta_\mu \partial_\mu + \kappa) \Psi = 0. \quad (4)$$

Here the β_μ satisfy the relations

$$\beta_\mu \beta_\nu + \beta_\nu \beta_\mu = \delta_{\mu\nu} \beta_\rho + \delta_{\rho\nu} \beta_\mu. \quad (5)$$

If we put

$$\partial_4 = -i \partial_t, \quad (6)$$

and separate the term involving β_4 , Eq. (4) becomes

$$i \beta_4 \partial_t \Psi = (\beta_\mu \cdot \partial + \kappa) \Psi. \quad (7)$$

Multiplying Eq. (7) first by β_4 and second by $(1 - \beta_4^2)$ and using Eq. (A1) we obtain the two equations

$$i \partial_t \beta_4^2 \Psi = \beta_4 (\beta_\mu \cdot \partial + \kappa) \Psi, \quad (8)$$

and

$$(1 - \beta_4^2) \Psi = - \frac{(1 - \beta_4^2)}{\kappa} (\beta_\mu \cdot \partial) \Psi. \quad (9)$$

It is seen that only the time derivative of $\beta_4^2 \Psi$ is determined by the equations. The components $(1 - \beta_4^2) \Psi$ are given in terms of these by means of the supplementary condition (9). However, if we

take the time derivative of Eq. (9) and use this as one of our equations along with Eq. (8) we need merely require that Eq. (9) hold as an initial condition. It will then hold at all later times as a consequence of the equations of motion. Thus Eq. (4) can be replaced by the Eqs. (8) and

$$i \partial_t (1 - \beta_4^2) \Psi = - \frac{(1 - \beta_4^2)}{\chi} (\beta \cdot \partial) i \partial_t \Psi. \quad (10)$$

Admissible solutions are subject to the initial condition of Eq. (9).

Eq. (10) can be simplified by writing Ψ on the right as

$$\Psi = \beta_4^2 \Psi + (1 - \beta_4^2) \Psi. \quad (11)$$

Using Eq. (A2) we see the second term in (11) does not contribute to the right hand side of (10) which is then

$$i \partial_t (1 - \beta_4^2) \Psi = - \frac{(1 - \beta_4^2)}{\chi} (\beta \cdot \partial) i \partial_t \beta_4^2 \Psi. \quad (12)$$

The time derivative of $\beta_4^2 \Psi$ may be eliminated from (12) using Eq. (8).

Hence we obtain

$$i \partial_t (1 - \beta_4^2) \Psi = - \frac{(1 - \beta_4^2)}{\chi} (\beta \cdot \partial) \beta_4 (\beta \cdot \partial + \chi) \Psi. \quad (13)$$

Equation (A3) shows the term second order in spatial derivatives in (13) vanishes. This then becomes

$$i \partial_t (1 - \beta_4^2) \Psi = - (1 - \beta_4^2) \beta \cdot \partial \beta_4 \Psi. \quad (14)$$

Adding (8) and (14) we obtain

$$i \frac{\partial \Psi}{\partial t} = H \Psi, \quad (15)$$

where

$$H = \beta_m^i P + \beta_4^i K. \quad (16)$$

Here

$$P_m = \frac{1}{i} \partial_m, \quad \beta_4^i = \beta_4, \quad (17)$$

and

$$\beta_k^i = -i(1 - \beta_4^2) \beta_k \beta_4 + i \beta_4 \beta_k, \quad (k = 1, 2, 3). \quad (18)$$

Using (A1) Eq. (17) can be simplified to

$$\beta_k^i = i(\beta_4 \beta_k - \beta_k \beta_4), \quad (k = 1, 2, 3). \quad (19)$$

Thus the Duffin-Kemmer Equation (4) has been put into the Hamiltonian form of Eq. (1) and (2). It is readily verified that the initial condition Eq. (9) is just Eq. (3) with the H given by (16).

We must still, however, show that the β_μ^i satisfy the algebraic relations (5). This is readily done by exhibiting a similarity transformation expressing the β_μ^i in terms of the β_μ . Such a transformation is

$$\beta_\mu^i = S \beta_\mu S^{-1}, \quad (\mu = 1, 2, 3, 4) \quad (20)$$

where

$$S = e^{i\pi \beta_4/2} = (1 - \beta_4^2 + i\beta_4). \quad (21)$$

We note that if the matrices β_μ are chosen hermitian, the matrix S is unitary and hence the β'_μ are also hermitian.

The situation is remarkably similar to that occurring in the case of the Dirac equation. Here we can pass from the covariant form

$$(\gamma_\mu \partial_\mu + \kappa) \Psi = 0, \quad (22)$$

to the Hamiltonian form

$$i \frac{\partial \Psi}{\partial t} = \{ \gamma'_4 \cdot \underline{p} + \gamma'_4 \kappa \} \Psi, \quad (23)$$

by multiplying by γ_4 . (Here we have denoted the conventional $(\underline{\alpha}, \beta)$ matrices by $(\underline{\gamma}, \gamma_4)$ to stress the analogy.) The γ'_μ are then given by

$$\gamma'_4 = \gamma_4, \quad (17a)$$

and

$$\gamma'_k = i \gamma_4 \gamma_k, \quad (k = 1, 2, 3). \quad (19a)$$

These γ'_μ are related to the γ_μ by the similarity transformation

$$\gamma'_\mu = S \gamma_\mu S^{-1} \quad (\mu = 1, 2, 3, 4), \quad (20a)$$

where

$$S = e^{i\pi \gamma_4 / 4} = \frac{1 + i \gamma_4}{\sqrt{2}}, \quad (21a)$$

and hence satisfy the same anticommutation relations as the γ_μ .

III. HIGHER INTEGRAL SPINS

It has been shown by Moldauer⁵ that the integral spin equations describing spin k of the D-F-P type can be written in the form

$$(\beta_p \partial_p + \chi) \Psi_{\mu_1, \mu_2 \dots \mu_{k-1}} = 0, \quad (24)$$

where the function Ψ is subject to the subsidiary condition

$$\beta_{\nu} \Psi_{\nu, \mu_2 \dots \mu_{k-1}} = 0. \quad (25)$$

Here the β matrices satisfy the relations (5) and act on a suppressed index. Ψ is symmetric and traceless in the indicated tensor indices.

Following precisely the same procedure as in Section II we obtain the Hamiltonian form of these equations

$$i \frac{\partial \Psi_{\mu_1 \dots \mu_{k-1}}}{\partial t} = \{H\} \Psi_{\mu_1 \dots \mu_{k-1}}, \quad (26)$$

with

$$H = \beta'_0 \cdot P + \beta'_4 \chi. \quad (27)$$

The admissible solutions are subject to the initial condition

$$(H \beta'_4 - \chi) \Psi_{\mu_1 \dots \mu_{k-1}} = 0, \quad (28)$$

and the supplementary condition (25). This last can be expressed in terms of the β' by multiplying by S . Thus

$$0 = S \beta_{\nu} \Psi_{\nu, \mu_2 \dots \mu_{k-1}} = S \beta_{\nu} S^{-1} S \Psi_{\nu, \mu_2 \dots \mu_{k-1}} \quad (29)$$

or

$$0 = \beta'_{\nu} S \Psi_{\nu, \mu_2 \dots \mu_{k-1}},$$

where S is given by Eq. (21).

APPENDIX

Some special consequences of the relations (5) that are used are:

$$(\beta_4)^3 = \beta_4 \quad (\text{A1})$$

$$(1 - \beta_4^2) \beta_m \partial_m (1 - \beta_4^2) = 0 \quad (\text{A2})$$

$$\beta_m \partial_m \beta_4 \beta_m \partial_m = 0 \quad (\text{A3})$$

One obtains (A1) by putting $\mu = \nu = \rho = 4$ in Eq. (3). To prove the remaining relations we will let Latin letters denote indices which run only from one to three. The identity

$$(1 - \beta_4^2) \beta_i (1 - \beta_4^2) = \beta_i - (\beta_4^2 \beta_i + \beta_i \beta_4^2) + \beta_4^2 \beta_i \beta_4^2 \quad (\text{A4})$$

becomes, on using (A1) and the relation

$$\beta_4^2 \beta_i + \beta_i \beta_4^2 = \beta_i \quad (\text{A5})$$

(obtained by putting $\mu = \nu = 4$, $\rho = i$ in Eq. (5))

$$(1 - \beta_4^2) \beta_i (1 - \beta_4^2) = 0 \quad (\text{A6})$$

Multiplying by ∂_i and summing yields (A2). Putting $\mu = 1$, $\nu = 4$, $\rho = k$ in (5) gives

$$\beta_1 \beta_4 \beta_k + \beta_k \beta_4 \beta_1 = 0 \quad (\text{A7})$$

Multiplying by $\partial_i \partial_k$ and summing yields (A3).

FOOTNOTES

1. P. A. M. Dirac, Proc. Roy. Soc. A, 155, 447 (1936); M. Fierz, Helvetica Physica Acta 12, 3 (1938); M. Fierz and W. Pauli, Proc. Roy. Soc. A, 173, 211 (1939). Hereafter this will be abbreviated as D-F-P.
2. E. Schrodinger, Proc. Roy. Soc. A, 229, 39 (1955).
3. K. M. Case, Phys. Rev. (Sept. 1) (1955).
4. N. Kemmer, Proc. Roy. Soc. A, 173, 91 (1939).
5. P. A. Moldauer, Ph.D. Thesis, University of Michigan (1955).

