UCSF

UC San Francisco Previously Published Works

Title

Marijuana, Secondhand Smoke, and Social Acceptability.

Permalink

<https://escholarship.org/uc/item/1wt0106q>

Journal

JAMA internal medicine, 178(1)

ISSN

2168-6106

Authors

Glantz, Stanton A
Halpern-Felsher, Bonnie
Springer, Matthew L

Publication Date

2018

DOI

10.1001/jamainternmed.2017.5301

Peer reviewed

Marijuana, Secondhand Smoke, and Social Acceptability

Stanton A. Glantz, PhD¹

Bonnie Halpern-Felsher, PhD²

Matthew L. Springer, PhD¹

¹Center for Tobacco Control Research and Education
Cardiovascular Research Institute
Department of Medicine (Cardiology)
University of California, San Francisco

²Division of Adolescent Medicine
Department of Pediatrics
Stanford University

Corresponding author:

Stanton A. Glantz, PhD
Professor of Medicine
Center for Tobacco Control Research and Education
530 Parnassus Suite 366
University of California, San Francisco
San Francisco, CA 94143-1390
415-476-3893
Fax 415-514-9345
Stanton.Glantz@ucsf.edu

862 words
31 references

On April 20, 2017 at 4:20 pm, 15,000 people in San Francisco's Golden Gate Park lit marijuana joints during the annual 420 Day as part of a national event. The giant cloud of secondhand marijuana smoke was visible from the University of California, San Francisco half a mile away. The cloud embodied the revelers' new freedom on this first 420 Day since November 2016 when California voters legalized recreational marijuana, and represented a growing source of air pollution.

It would have been unthinkable (and illegal) for thousands of people to congregate and smoke tobacco cigarettes in that park. The recognition that secondhand tobacco smoke causes cardiovascular disease, lung and breast cancer, and other diseases^{1,2} motivated passage of laws to protect people from secondhand smoke, including in Golden Gate Park. These laws have the beneficial side effect of stimulating voluntary smokefree home policies, discouraging initiation, supporting smoking cessation, and denormalizing tobacco use.^{3, p. 26-29,4}

Such social norm change is an effective tobacco control strategy, as exemplified by the California Tobacco Control Program, is a broad-based campaign focused on reinforcing the nonsmoking norm aimed at the population as a whole, not just smokers or youth,⁵ that included hard hitting media and support for smokefree environments.⁶ By focusing on the broader social environment, California achieved one of the lowest smoking rates – for both adults and youth – in the United States.^{6,7} Adolescents who perceive secondhand tobacco smoke as dangerous are much less likely to start smoking tobacco compared to those with lower risk perceptions of secondhand smoke.⁸

In contrast to tobacco, marijuana is widely viewed as harmless or even good for you, even something to be celebrated. Perhaps because cannabinoids are useful for treating chemotherapy-induced nausea, chronic pain, and spasms in multiple sclerosis,⁹ marijuana,

especially medical marijuana, is viewed as having positive effects. Nationally, more adolescents used marijuana than tobacco in 2016, with 16% of 10th graders and 25% of 12th graders reporting past 30-day marijuana use compared to 5% and 11% for tobacco.¹⁰ Similarly, adolescents' perceptions of marijuana risk dropped from 58% of 12th graders reporting great risk associated with marijuana in 2006 to 31% in 2016.¹⁰ California adolescents believe that, compared to tobacco, marijuana is less addictive and easier to quit, more socially acceptable, and less harmful to their own and their friends' health as well as to the environment than tobacco cigarettes.¹¹

The evidence that secondhand exposure to marijuana smoke, like the evidence for all health effects of marijuana,⁹ is more limited than for tobacco. But smoke from any source is a complex mixture of thousands of chemicals, including ultrafine particles and toxic gases. Other than nicotine and cannabinoids, tobacco and marijuana smoke are similar.¹² Indeed, the California Environmental Protection Agency identified marijuana smoke as a human carcinogen based largely on the smoke's toxicology.¹³

Cardiovascular effects of breathing secondhand smoke are particularly important because about 80% of the deaths attributed to secondhand tobacco smoke are due to heart disease, including acute myocardial infarction.¹⁴ These effects reflect the highly nonlinear dose-response curve for cardiovascular effects of secondhand smoke,¹⁵ with secondhand exposure nearly as dangerous as active smoking. Like all particulate air pollution,¹⁶ smoking and exposure to secondhand smoke both lead to impairment of endothelial function in humans as measured by arterial flow-mediated dilation¹⁷ (FMD), an effect that occurs in a few minutes in humans¹⁸⁻²⁰ and rats.²¹ Reflecting these rapid effects, implementation of comprehensive smokefree laws are followed by drops in heart attacks and other conditions.^{3, p. 435-443,22,23}

As one would expect based on this evidence, FMD drops in rats after one minute of

marijuana sidestream smoke, with impairment lasting at least 90 minutes (longer than for tobacco smoke), independent of the presence or absence of cannabinoids.²⁴ Consistent with this animal observation, a retrospective study found the risk of myocardial infarction was increased by 4.8 times in the hour after using marijuana.²⁵ Evidence on longer term marijuana use and myocardial infarction is mixed;^{9,26,27} there is consistent evidence on stroke.^{9,28,29} There is, however, already enough evidence to warrant protecting people from secondhand exposure to marijuana smoke.

While marijuana is available in many forms, smoking it remains by far the dominant mode of consumption.³⁰ The evidence that secondhand marijuana smoke is dangerous is more limited than the evidence that secondhand tobacco smoke is, yet there is already at least as much evidence concerning marijuana risk as there was for secondhand tobacco smoke in the late 1970s when the clean indoor air movement started gaining steam. In particular, we know that the cardiovascular effects of secondhand smoke are more important in terms of population impact than cancer, that these effects occur quickly, and that marijuana may be worse than tobacco smoke.²⁴ Further, misperceptions about the harms of marijuana use,¹¹ including the lack of understanding of the harms associated with secondhand smoke, point to the importance of education and policies that bring secondhand marijuana smoke to the forefront. This evidence supports maintaining and expanding clean indoor air laws to include marijuana as part of a public health framework for marijuana regulation.³¹ Stressing the right of all to breathe clean air should also be at the core of educational and legislative efforts to reinforce the marijuana smoke-free norm for everyone.

FUNDING

This work was funded in part by the National Institute of Drug Abuse (R01DA043950

[Glantz], R21DA031966 [Springer]), the California Tobacco Related Disease Research Program (25IR-0030 [Springer]) and the Elfenworks Foundation (Springer).

CONFLICT OF INTEREST

Nothing to declare.

REFERENCES

1. U.S. Department of Health and Human Services. The Health Consequences of Involuntary Exposure to Tobacco Smoke. 2006;
http://www.cdc.gov/tobacco/data_statistics/sgr/2006/.
2. California Environmental Protection Agency. *Health Effects of Exposure to Environmental Tobacco Smoke*. June 2005.
3. U.S. Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.
4. Cheng KW, Okechukwu CA, McMillen R, Glantz SA. Association between clean indoor air laws and voluntary smokefree rules in homes and cars. *Tob Control*. 2015;24(2):168-174.
5. California Department of Public Health, California Tobacco Control Program. *California Tobacco Control Update 2009: 20 Years of Tobacco Control in California*. Sacramento, CA; 2009.
6. Roeseler A, Burns D. The quarter that changed the world. *Tob Control*. 2010;19 Suppl 1:i3-15.
7. Lightwood J, Glantz SA. The effect of the California tobacco control program on smoking prevalence, cigarette consumption, and healthcare costs: 1989-2008. *PLoS One*. 2013;8(2):e47145.

8. Song AV, Glantz SA, Halpern-Felsher BL. Perceptions of second-hand smoke risks predict future adolescent smoking initiation. *J Adolesc Health*. 2009;45(6):618-625.
9. Committee on the Health Effects of Marijuana. *The Health Effects of Cannabis and Cannabinoids*. Washington, DC: National Academies Press; 2017.
10. Johnston LD, Miech RA, O'Malley PM, Bachman JG, Schulenberg JE. Teen use of any illicit drug other than marijuana at new low, same true for alcohol. 2016; <http://www.monitoringthefuture.org>. Accessed 7 Jul 2017.
11. Roditis ML, Delucchi K, Chang A, Halpern-Felsher B. Perceptions of social norms and exposure to pro-marijuana messages are associated with adolescent marijuana use. *Prev Med*. 2016;93:171-176.
12. Moir D, Rickert WS, Levasseur G, et al. A comparison of mainstream and sidestream marijuana and tobacco cigarette smoke produced under two machine smoking conditions. *Chem Res Toxicol*. 2008;21(2):494-502.
13. Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. *Evidence on the Carcinogenicity of Marijuana Smoke*. Sacramento, CA; 2009.
14. U.S. Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2006.
15. Barnoya J, Glantz SA. Cardiovascular effects of secondhand smoke: nearly as large as smoking. *Circulation*. 2005;111(20):2684-2698.

16. Pope CA, 3rd, Bhatnagar A, McCracken JP, Abplanalp W, Conklin DJ, O'Toole T. Exposure to Fine Particulate Air Pollution Is Associated With Endothelial Injury and Systemic Inflammation. *Circ Res*. 2016;119(11):1204-1214.
17. Celermajer DS, Adams MR, Clarkson P, et al. Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. *N Engl J Med*. 1996;334(3):150-154.
18. Kato T, Inoue T, Morooka T, Yoshimoto N, Node K. Short-term passive smoking causes endothelial dysfunction via oxidative stress in nonsmokers. *Can J Physiol Pharmacol*. 2006;84(5):523-529.
19. Heiss C, Amabile N, Lee AC, et al. Brief secondhand smoke exposure depresses endothelial progenitor cells activity and endothelial function: sustained vascular injury and blunted nitric oxide production. *J Am Coll Cardiol*. 2008;51(18):1760-1771.
20. Frey PF, Ganz P, Hsue PY, et al. The exposure-dependent effects of aged secondhand smoke on endothelial function. *J Am Coll Cardiol*. 2012;59(21):1908-1913.
21. Pinnamaneni K, Sievers RE, Sharma R, et al. Brief exposure to secondhand smoke reversibly impairs endothelial vasodilatory function. *Nicotine Tob Res*. 2014;16(5):584-590.
22. Tan CE, Glantz SA. Association between smoke-free legislation and hospitalizations for cardiac, cerebrovascular, and respiratory diseases: a meta-analysis. *Circulation*. 2012;126(18):2177-2183.
23. Been JV, Nurmatov UB, Cox B, Nawrot TS, van Schayck CP, Sheikh A. Effect of smoke-free legislation on perinatal and child health: a systematic review and meta-analysis. *Eur J Paediatr Dent*. 2015;16(3):210-211.

24. Wang X, Derakhshandeh R, Liu J, et al. One minute of marijuana secondhand smoke exposure substantially impairs vascular endothelial function. *J Am Heart Assoc*. 2016;5(8):e003858
25. Mittleman MA, Lewis RA, Maclure M, Sherwood JB, Muller JE. Triggering myocardial infarction by marijuana. *Circulation*. 2001;103(23):2805-2809.
26. Reis JP, Auer R, Bancks MP, et al. Cumulative Lifetime Marijuana Use and Incident Cardiovascular Disease in Middle Age: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Public Health*. 2017;107(4):601-606.
27. Chami AT, Kim C. Cannabis abuse and risk for myocardial infarction: a population based study (abstract). *J Am Coll Cardiol*. 2017;69(11 (Supl)):6
[https://doi.org/10.1016/S0735-1097\(1017\)33395-33398](https://doi.org/10.1016/S0735-1097(1017)33395-33398).
28. Kalla A, Krishnamoorthy P, Gopalakrishnan A, Garg J, Figueredo V. Cannabis use predicts risks of heart failure and cerebrovascular accidents: Results from the national inpatient sample (abstract). *J Am Coll Cardiol*. 2017;69(11 (Supl)):1784
[https://doi.org/1710.1016/S0735-1097\(1717\)35173-35172](https://doi.org/1710.1016/S0735-1097(1717)35173-35172).
29. Jouanjus E, Raymond V, Lapeyre-Mestre M, Wolff V. What is the Current Knowledge About the Cardiovascular Risk for Users of Cannabis-Based Products? A Systematic Review. *Current Atherosclerosis Reports*. 2017;19(6):26.
30. Oregon Health Authority Public Health Division. Marijuana use, attitudes and health effects in Oregon. *Marijuana Report* 2016;
<https://public.health.oregon.gov/PreventionWellness/marijuana/Documents/oha-8509-marijuana-report.pdf>.

31. Barry RA, Glantz S. A Public Health Framework for Legalized Retail Marijuana Based on the US Experience: Avoiding a New Tobacco Industry. *PLoS Med.* 2016;13(9):e1002131.