

# UC Agriculture & Natural Resources Farm

**Title**

Damping-Off Diseases

**Permalink**

<https://escholarship.org/uc/item/1xm1w425>

**Author**

Laemmlen, Franklin

**Publication Date**

2002

**DOI**

10.3733/ucanr.8041

Peer reviewed



**UNIVERSITY OF CALIFORNIA**

Agriculture  
and Natural Resources

<http://anrcatalog.ucdavis.edu>

# Damping-Off Diseases

**FRANKLIN LAEMMLEN**, University of California Cooperative Extension Farm Advisor, Santa Barbara and San Luis Obispo Counties

**D**amping-off diseases of seedlings are found worldwide and can be caused by several species of fungi under various weather conditions. The name damping-off is in standard use in the literature and usually refers to the disintegration of stem and root tissues at and below the soil line. The plant tissues become water-soaked and mushy, and the seedling wilts and falls over. Damping-off diseases, however, can have several phases. The fungi that cause these diseases can attack the seed or the seedling before it emerges above the soil surface, causing a seed rot or pre-emergent rot. When this happens, the result is a poor stand that may be mistakenly ascribed to poor seed quality or seed maggots rather than to the presence of a disease. The death of seedlings after emergence or transplanting is called postemergent damping-off and is the condition most often identified as damping-off (although this symptom may also be caused by maggots).

Most pathogens that cause damping-off diseases can also cause disease as the plant grows to maturity. Root rot, crown rot, stem lesions, basal rot, crater rot, bottom rot, and stem girdling diseases may all be associated with damping-off fungi attacking mature plants.

The two fungi that are most often associated with damping-off are *Rhizoctonia solani* and *Pythium* species. *Rhizoctonia solani* is found in most agricultural soils and survives between crops on plant residues and as microsclerotia (fig. 1). This pathogen usually attacks seedlings at or near the soil surface. Initial symptoms are stem lesions that are brick red to brown and sunken. If the disease progresses, the stem may become girdled. Stem canker, soreshin, wirestem, and damping-off are names associated with seedling and postemergent diseases caused by *R. solani*.

Damping-off diseases caused by *Pythium* species usually begin as root rot. This group of fungi survives as oospores (fig. 2) in the soil that germinate to attack root hairs and root tips, causing a progressive deterioration of the root system. The seedling may wilt before aboveground lesions are evident, or the seed may rot in the ground. *Pythium* species are often responsible for pre-emergent damping-off. These species can also infect the lower stem and cause the mushy, watery stem rot typically identified with damping-off syndrome.


Other fungi that can be associated with seedling or transplant damping-off are *Botrytis cinerea*, *Sclerotinia sclerotiorum*, *S. minor*, *Alternaria* species, *Phytophthora* species, *Fusarium* species, and *Thielaviopsis basicola*.

The environmental conditions that favor damping-off vary according to the pathogen. *R. solani* often causes injury to beans during spring germination. Cole crops transplanted in the fall are often victims to wirestem caused by this pathogen. Damping-off caused by *S. sclerotiorum*, *S. minor*, and *B. cinerea* can be severe in greenhouse seedling production and during spring transplanting and seed germination. *Pythium* species also tend to be most active during the spring months when soil temperatures are still cool and soil moisture plentiful.

Control of damping-off diseases is difficult. Damping-off must be anticipated and prevented by using seed and transplant treatments before the seed or plants are put in the field. In-furrow treatments at seeding and banded treatments over the plant line after transplanting can also be useful. Good seedbed preparation is important. Shallow planting of seed and seedlings to promote rapid germination and

thereby less stem exposure to the soil has been used to reduce losses from *R. solani* damping-off. Irrigation should be managed to enhance germination and growth and to avoid over watering.

Fungicides used for *Pythium* control, either as seed or seedling treatments, include metalaxyl, thiram, some copper compounds, and captan. *Rhizoctonia solani* damping-off may be suppressed with pentachloronitrobenzene (PCNB), chlorothalonil, benomyl, thiophanate methyl, carboxin, mancozeb, vinclozolin, and iprodione. Always refer to the product label for crop registration and dosage information.



**Figure 1.** Disease cycle of *Rhizoctonia solani* (*Thanatephorus cucumeris*).  
 (Source: Agrios, G. N. 1997. Plant pathology, 4th edition. San Diego: Academic Press.)



Visit our online catalog at <http://anrcatalog.ucdavis.edu>. You can also place orders by mail, phone, or fax, or request a printed catalog of publications, slide sets, and videos from

University of California  
Agriculture and Natural Resources  
Communication Services  
6701 San Pablo Avenue, 2nd Floor  
Oakland, CA 94608-1239

Telephone: (800) 994-8849 or (510) 642-2431

FAX: (510) 643-5470

E-mail inquiries: [danrcs@ucdavis.edu](mailto:danrcs@ucdavis.edu)

An electronic version of this publication is available on the ANR Communication Services website at <http://anrcatalog.ucdavis.edu>.

#### **Publication 8041**

©2001 by the Regents of the University of California, Division of Agriculture and Natural Resources. All rights reserved.

The University of California prohibits discrimination against or harassment of any person employed by or seeking employment with the University on the basis of race, color, national origin, religion, sex, physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized). University Policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th floor, Oakland, CA 94612-3550; (510) 987-0096. For a free catalog of other publications, telephone (800) 994-8849. For help downloading this publication, call (530) 754-5112.

To simplify information, trade names of products have been used. No endorsement of named or illustrated products is intended, nor is criticism implied of similar products that are not mentioned or illustrated.

#### **Warning on the Use of Chemicals**

Pesticides are poisonous. Always read and carefully follow all precautions and safety recommendations given on the container label. Store all chemicals in the original labeled containers in a locked cabinet or shed, away from food or feeds, and out of the reach of children, unauthorized persons, pets, and livestock. Confine chemicals to the property being treated. Avoid drift onto neighboring properties, especially gardens containing fruits or vegetables ready to be picked. Do not place containers containing pesticide in the trash nor pour pesticides down sink or toilet. Either use the pesticide according to the label or take unwanted pesticides to a Household Hazardous Waste Collection site. Contact your county agricultural commissioner for additional information on safe container disposal and for the location of the Hazardous Waste Collection site nearest you. Dispose of empty containers by following label directions. Never reuse or burn the containers or dispose of them in such a manner that they may contaminate water supplies or natural waterways.

pr-12/01-GM/VFG



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by the ANR Associate Editor for Vegetable Crops.