UC Berkeley

UC Berkeley Previously Published Works

Title

Measurement of the branching ratios $\Gamma(D_s^{*+} \rightarrow D_s + \pi^0) / \Gamma(D_s^{*+} \rightarrow D_s + \gamma)$ and $\Gamma(D^{*0} \rightarrow D^0 \pi^0) / \Gamma(D^{*0} \rightarrow D^0 \gamma)$

Permalink

<https://escholarship.org/uc/item/1xn942cv>

Journal

Physical Review D - Particles, Fields, Gravitation and Cosmology, 72(9)

ISSN

1550-7998

Authors

Aubert, B
Barate, R
Boutigny, D
et al.

Publication Date

2005-11-01

DOI

10.1103/PhysRevD.72.091101

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

**Measurement of the branching ratios $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$
and $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma)$**

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ J. S. Minamora,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ B. Spaan,²³ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaut,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negri,²⁷ L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Feroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,*} M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ G. Schott,³² W. Bhimji,³³ D. A. Bowerman,³³ P. D. Dauncey,³³ U. Egede,³³ R. L. Flack,³³ J. R. Gaillard,³³ J. A. Nash,³³ M. B. Nikolich,³³ W. Panduro Vazquez,³³ X. Chai,³⁴ M. J. Charles,³⁴ W. F. Mader,³⁴ U. Mallik,³⁴ A. K. Mohapatra,³⁴ V. Ziegler,³⁴ J. Cochran,³⁵ H. B. Crawley,³⁵ V. Eyges,³⁵ W. T. Meyer,³⁵ S. Prell,³⁵ E. I. Rosenberg,³⁵ A. E. Rubin,³⁵ J. Yi,³⁵ N. Arnaud,³⁶ M. Davier,³⁶ X. Giroux,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ F. Le Diberder,³⁶ V. Lepeltier,³⁶ A. M. Lutz,³⁶ A. Oyanguren,³⁶ T. C. Petersen,³⁶ S. Plaszczynski,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ A. Stocchi,³⁶ G. Wormser,³⁶ C. H. Cheng,³⁷ D. J. Lange,³⁷ M. C. Simani,³⁷ D. M. Wright,³⁷ A. J. Bevan,³⁸ C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ K. A. George,³⁸ D. E. Hutchcroft,³⁸ R. J. Parry,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ C. M. Cormack,³⁹ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ C. L. Brown,⁴⁰ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ M. G. Green,⁴⁰ D. A. Hopkins,⁴⁰ P. S. Jackson,⁴⁰ T. R. McMahon,⁴⁰ S. Ricciardi,⁴⁰ F. Salvatore,⁴⁰ D. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² C. L. Edgar,⁴² M. C. Hodgkinson,⁴² M. P. Kelly,⁴² G. D. Lafferty,⁴² M. T. Naisbit,⁴² J. C. Williams,⁴² C. Chen,⁴³ W. D. Hulsbergen,⁴³ A. Jawahery,⁴³ D. Kovalskyi,⁴³ C. K. Lae,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ R. Kofler,⁴⁴ V. B. Koptchev,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ S. Saremi,⁴⁴ H. Staengle,⁴⁴ S. Willocq,⁴⁴ R. Cowan,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ H. Kim,⁴⁶ P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ V. Lombardo,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ J. Reidy,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ P. Taras,⁴⁹ B. Viaud,⁴⁹ H. Nicholson,⁵⁰ N. Cavallo,^{51,†} G. De Nardo,⁵¹ F. Fabozzi,^{51,†} C. Gatto,⁵¹ L. Lista,⁵¹ D. Monorchio,⁵¹ P. Paolucci,⁵¹ D. Piccolo,⁵¹ C. Sciacca,⁵¹ M. Baak,⁵² H. Bulten,⁵² G. Raven,⁵² H. L. Snoek,⁵² L. Wilden,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ T. Allmendinger,⁵⁴ G. Benelli,⁵⁴

K. K. Gan,⁵⁴ K. Honscheid,⁵⁴ D. Hufnagel,⁵⁴ P. D. Jackson,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ T. Pulliam,⁵⁴ A. M. Rahimi,⁵⁴ R. Ter-Antonyan,⁵⁴ Q. K. Wong,⁵⁴ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ M. Lu,⁵⁵ C. T. Potter,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ F. Galeazzi,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ M. Benayoun,⁵⁷ H. Briand,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ M. J. J. John,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ L. Roos,⁵⁷ G. Therin,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ S. Pacetti,⁵⁹ M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ D. E. Wagoner,⁶¹ J. Biesiada,⁶² N. Danielson,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,⁶³ A. D'Orazio,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ H. Schröder,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Graziani,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ J. R. Wilson,⁶⁷ F. X. Yumiceva,⁶⁷ T. Abe,⁶⁸ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ J. C. Dingfelder,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va'vra,⁶⁸ N. van Bakel,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ M. Ahmed,⁷⁰ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ F. Martinez-Vidal,⁷⁶ R. S. Panvini,^{77,‡} Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ M. Graham,⁸⁰ J. J. Hollar,⁸⁰ J. R. Johnson,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰ A. Mihalys,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Dipartimento di Fisica and INFN, Università di Bari, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Institut für Experimentalphysik 1, Ruhr Universität Bochum, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸Institute for Particle Physics, University of California at Santa Cruz, Santa Cruz, California 95064, USA

¹⁹California Institute of Technology, Pasadena, California 91125, USA

²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA

- ²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany
²⁴Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Dipartimento di Fisica and INFN, Università di Ferrara, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Dipartimento di Fisica and INFN, Università di Genova, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Institut für Experimentelle Kernphysik, Universität Karlsruhe, D-76021 Karlsruhe, Germany
³³Imperial College London, London, SW7 2AZ, United Kingdom
³⁴University of Iowa, Iowa City, Iowa 52242, USA
³⁵Iowa State University, Ames, Iowa 50011-3160, USA
³⁶Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁹Queen Mary, University of London, E1 4NS, United Kingdom
⁴⁰Royal Holloway and Bedford New College, University of London, Egham, Surrey TW20 0EX, United Kingdom
⁴¹University of Louisville, Louisville, Kentucky 40292, USA
⁴²University of Manchester, Manchester M13 9PL, United Kingdom
⁴³University of Maryland, College Park, Maryland 20742, USA
⁴⁴University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁵Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
⁴⁶McGill University, Montréal, Quebec, Canada H3A 2T8
⁴⁷Dipartimento di Fisica and INFN, Università di Milano, I-20133 Milano, Italy
⁴⁸University of Mississippi, University, Mississippi 38677, USA
⁴⁹Laboratoire René J. A. Lévesque, Université de Montréal, Montréal, Quebec, Canada H3C 3J7
⁵⁰Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵¹Dipartimento di Scienze Fisiche and INFN, Università di Napoli Federico II, I-80126, Napoli, Italy
⁵²NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁴Ohio State University, Columbus, Ohio 43210, USA
⁵⁵University of Oregon, Eugene, Oregon 97403, USA
⁵⁶Dipartimento di Fisica and INFN, Università di Padova, I-35131 Padova, Italy
⁵⁷Laboratoire de Physique Nucléaire et de Hautes Energies, Universités Paris VI et VII, F-75252 Paris, France
⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁹Dipartimento di Fisica and INFN, Università di Perugia, I-06100 Perugia, Italy
⁶⁰Dipartimento di Fisica, Scuola Normale Superiore and INFN, Università di Pisa, I-56127 Pisa, Italy
⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
⁶²Princeton University, Princeton, New Jersey 08544, USA
⁶³Dipartimento di Fisica and INFN, Università di Roma La Sapienza, I-00185 Roma, Italy
⁶⁴Universität Rostock, D-18051 Rostock, Germany
⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁹Stanford University, Stanford, California 94305-4060, USA
⁷⁰State University of New York, Albany, New York 12222, USA
⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
⁷²University of Texas at Austin, Austin, Texas 78712, USA
⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
⁷⁴Dipartimento di Fisica Sperimentale and INFN, Università di Torino, I-10125 Torino, Italy
⁷⁵Dipartimento di Fisica and INFN, Università di Trieste, I-34127 Trieste, Italy
⁷⁶Universitat de Valencia-CSIC, IFIC, E-46071 Valencia, Spain
⁷⁷Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁸University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁹Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁸⁰University of Wisconsin, Madison, Wisconsin 53706, USA
⁸¹Yale University, New Haven, Connecticut 06511, USA

(Received 19 August 2005; published 8 November 2005)

Data samples corresponding to the isospin-violating decay $D_s^{*+} \rightarrow D_s^+ \pi^0$ and the decays $D_s^{*+} \rightarrow D_s^+ \gamma$, $D^{*0} \rightarrow D^0 \pi^0$ and $D^{*0} \rightarrow D^0 \gamma$ are reconstructed using 90.4 fb^{-1} of data recorded by the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- collider. The following branching ratios are extracted: $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma) = 0.062 \pm 0.005(\text{stat.}) \pm 0.006(\text{syst.})$ and $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma) = 1.74 \pm 0.02(\text{stat.}) \pm 0.13(\text{syst.})$. Both measurements represent significant improvements over present world averages.

DOI: [10.1103/PhysRevD.72.091101](https://doi.org/10.1103/PhysRevD.72.091101)

PACS numbers: 13.25.Ft, 12.39.Fe, 13.40.Hq

The decay of any higher-mass $c\bar{s}$ meson into $D_s^+ \pi^0$ [1] violates isospin conservation, thus guaranteeing a small partial width. The amount of suppression is a matter of large theoretical uncertainty according to most models of charm-meson radiative decay [2]. One such model [3] suggests that the decay $D_s^{*+} \rightarrow D_s^+ \pi^0$ may proceed via π^0 - η mixing. Even including such considerations, the radiative decay $D_s^{*+} \rightarrow D_s^+ \gamma$ is still expected to dominate. The existence of isospin-violating decay modes such as $D_s^{*+} \rightarrow D_s^+ \pi^0$ is particularly relevant given the recent observations of two narrow new D_s^+ meson states [4,5]. In particular, in contrast to the D_s^{*+} meson, there is no experimental evidence for the electromagnetic decay of the $D_{sJ}(2317)^+$; current measurements place the branching ratio to $D_s^{*+} \gamma$ at less than 18% at 90% confidence level (CL) [6].

Besides the $D_s^+ \pi^0$ and $D_s^+ \gamma$ final states, no other decay modes of the D_s^{*+} have been observed and none are expected to occur at a significant level. Only one previous observation of the decay $D_s^{*+} \rightarrow D_s^+ \pi^0$ is recorded in the literature, yielding a value of $0.062_{-0.018}^{+0.020}(\text{stat.}) \pm 0.022(\text{syst.})$ for the branching ratio $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$ [7]. The analysis presented here confirms this observation and provides a more precise measurement of this branching ratio.

The decay $D^{*0} \rightarrow D^0 \pi^0$, in contrast to $D_s^{*+} \rightarrow D_s^+ \pi^0$, does not violate isospin conservation and the world average for the branching ratio is $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma) = 1.625 \pm 0.20$ [8]. As for the D_s^{*+} meson, the π^0 and γ decay modes are expected to saturate the decay width of the D^{*0} meson.

The results presented here are based on data recorded by the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- storage rings. The data sample, corresponding to an integrated luminosity of 90.4 fb^{-1} , was recorded at and approximately 40 MeV below the $\Upsilon(4S)$ resonance. Because of the unequal beam energies, the e^+e^- center-of-mass system is boosted relative to the laboratory frame with $\beta\gamma \approx 0.55$. The *BABAR* detector and trigger are described in detail elsewhere [9]. Charged particles are

detected and their momenta measured by a silicon vertex tracker (SVT) consisting of five layers of double-sided silicon strip sensors and a cylindrical 40-layer drift chamber (DCH), both operating within a 1.5 T solenoidal magnetic field. Charged particle identification is provided by energy loss measurements in the SVT and DCH and by Cherenkov light detected in an internally reflecting ring imaging detector (DIRC). Photons are identified and their energies measured by an electromagnetic calorimeter (EMC) composed of 6580 CsI(Tl) crystals.

In the following paragraphs, the D_s^{*+} measurement is described first. The D^{*0} analysis, which uses similar procedures for signal extraction, is described afterward in less detail.

D_s^+ mesons are reconstructed via the decay sequence $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$. Kaons are identified by combining the energy deposited in the SVT and DCH with the information from the DIRC. Tracks not identified as kaons according to the particle identification criteria are considered to be pions. All $K^+ K^- \pi^+$ candidates are required to fit successfully to a common vertex. Only combinations with a $K^+ K^-$ invariant mass within $8 \text{ MeV}/c^2$ of the nominal ϕ mass [8] are retained.

In e^+e^- annihilation to charm quarks, the $c\bar{c}$ fragmentation process is characterized by the production of high-momentum (leading) charm hadrons. This property is exploited in order to reduce substantially the combinatorial background by retaining only those $\phi \pi^+$ candidates with scaled momentum x_p greater than 0.6, where x_p is defined as $x_p(D_s^+) = p^*(D_s^+)/p_{\text{max}}^*(D_s^+)$ and $p^*(D_s^+)$ is the momentum of the D_s^+ candidates in the e^+e^- center-of-mass frame with $p_{\text{max}}^*(D_s^+) = \sqrt{E_{\text{beam}}^2 - m(D_s^+)^2}$ as its maximum value.

The longitudinal polarization of the ϕ meson in the D_s^+ rest frame is used to reduce background by requiring that the absolute value of the cosine of the helicity angle, defined as the angle between the ϕ momentum direction in the D_s^+ rest frame and the momentum direction of either of the kaons in the ϕ rest frame, is 0.3 or greater.

The resulting $K^+ K^- \pi^+$ invariant mass distribution is shown in Fig. 1. This distribution can be modeled by the sum of two Gaussian functions (to represent the signal) and a third-order polynomial (to represent the background). The resulting binned χ^2 fit yields $73\,500 \pm 300$ events (statistical errors only). A D_s^+ candidate is retained if its

* Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

† Also with Università della Basilicata, Potenza, Italy

‡ Deceased

FIG. 1. The $K^+K^-\pi^+$ mass distribution. The dots represent data points with error bars corresponding to statistical uncertainties (these uncertainties are small enough that the error bars are difficult to distinguish). The solid curve shows the fitted function. The dashed curve indicates the background. D_s^+ candidates are defined by the region between the vertical dotted lines.

invariant mass is within $12 \text{ MeV}/c^2$ of the nominal D_s^+ mass [8].

A π^0 candidate is reconstructed by combining two photon candidates that fulfill the following requirements. Each photon candidate is identified by a calorimeter cluster that is not associated with a charged track and has an energy in the laboratory frame of at least 45 MeV. Additionally, to help remove the background from hadronic showers, the fractional lateral width [10], which describes the shape of the shower in the calorimeter, is required to be less than 0.55. The fiducial acceptance of photon candidates is restricted by the angular range of the EMC ($-0.92 \leq \cos\theta \leq 0.89$, where θ is the polar angle in the center-of-mass frame [9]).

A π^0 candidate is retained if it has a momentum p^* in the e^+e^- center-of-mass frame greater than $150 \text{ MeV}/c$. Furthermore, the absolute value of the cosine of the decay angle θ^* , which is defined as the angle between the direction of one of the photons in the π^0 rest frame and the direction of the π^0 candidate in the center-of-mass frame, is required to be less than 0.85. For $\pi^0 \rightarrow \gamma\gamma$ decay, the $\cos\theta^*$ distribution is uniform, while it peaks near ± 1 for random $\gamma\gamma$ combinations.

Only $\gamma\gamma$ pairs within a specified mass interval are retained. This interval is defined by the values of mass at which the π^0 signal portion of a function fitted to the $\gamma\gamma$ mass distribution falls below 0.2 times its maximum value. This requirement accommodates the asymmetric shape of the $\gamma\gamma$ mass distribution and takes into account variations in detector calibration. A kinematic fit is applied to the surviving $\gamma\gamma$ pairs to constrain their mass to the nominal π^0 mass.

After combining the D_s^{*+} and π^0 candidates in a search for the decay $D_s^{*+} \rightarrow D_s^+ \pi^0$, a fit is applied to the distribution of the mass difference $\Delta m(D_s^+ \pi^0) = m(K^+K^-\pi^+\pi^0) - m(K^+K^-\pi^+)$. The fit function is the sum of a double Gaussian function to represent the signal and the function

$$f_1(\Delta m) = N \left(1 - \exp\left(-\frac{\Delta m - m(\pi^0)}{\mu}\right) \right) \times (\Delta m^2 + a\Delta m + b), \quad (1)$$

where $m(\pi^0)$ is the π^0 mass, and N , μ , a , and b are free fit parameters to describe the background. The exponential term models the kinematic threshold; this threshold term has little influence on the background shape near the signal region. The result of this fit is shown in Fig. 2(a). A signal event yield of 560 ± 40 (statistical error only) is obtained.

For the reconstruction of the decay $D_s^{*+} \rightarrow D_s^+ \gamma$, a calorimeter cluster that is not associated with a charged track is considered a photon candidate if it fulfills the following requirements: the energy must be 50 MeV or greater in the laboratory frame and 100 MeV or greater in the e^+e^- center-of-mass frame and the fractional lateral width must be less than 0.8. To reduce the background due to photons from π^0 decay, a photon candidate is discarded if it forms a π^0 candidate with any other photon candidate in the same event. In this case, a $\gamma\gamma$ combination is considered a π^0 candidate if the invariant mass is in the range $115 < m(\gamma\gamma) < 155 \text{ MeV}/c^2$ and if the total energy is at least 200 MeV in the e^+e^- center-of-mass frame.

To obtain the $D_s^{*+} \rightarrow D_s^+ \gamma$ signal event yield, a fit is applied to the distribution of the mass difference $\Delta m(D_s^+ \gamma) = m(K^+K^-\pi^+\gamma) - m(K^+K^-\pi^+)$. The fit function is a sum of a third-order polynomial to model the background plus a function first introduced by the Crystal Ball collaboration [11] for the signal

FIG. 2. The D_s^{*+} signals: (a) $m(K^+K^-\pi^+\pi^0) - m(K^+K^-\pi^+)$; (b) $m(K^+K^-\pi^+\gamma) - m(K^+K^-\pi^+)$. The dots represent data points. The solid curve shows the fitted function. The dashed curve indicates the portion of the fit associated with background.

$$f_2(\Delta m) = N \cdot \begin{cases} A \left(B - \frac{\Delta m - \mu}{\sigma} \right)^{-n} & \text{if } (\Delta m - \mu)/\sigma \leq \alpha \\ \exp\left(-\frac{(\Delta m - \mu)^2}{2\sigma^2}\right) & \text{if } (\Delta m - \mu)/\sigma \geq \alpha, \end{cases} \quad (2)$$

where N , μ , σ , n , and α are free fit parameters and A and B are chosen such that the function and its first derivative are continuous at $(\Delta m - \mu)/\sigma = \alpha$. The fit result is shown in Fig. 2(b). A signal yield of $15\,600 \pm 200$ events (statistical error only) is obtained.

The reconstruction efficiencies are determined using a Monte Carlo simulation based on 30 000 events for each D_s^+ decay mode. The simulated events are analyzed using the same procedure as for real data. By calculating the ratio of the number of reconstructed to generated events, efficiencies of $\epsilon(D_s^+ \pi^0) = 0.041 \pm 0.002$ and $\epsilon(D_s^+ \gamma) = 0.071 \pm 0.002$ are found for the two D_s^{*+} decay modes. The efficiency ratio is $\epsilon(D_s^+ \pi^0)/\epsilon(D_s^+ \gamma) = 0.58 \pm 0.03$ (statistical error only).

Various sources of systematic uncertainties are studied. To verify that the Monte Carlo events model the data correctly, τ decays with one or two π^0 mesons in the final state are studied to obtain energy-dependent Monte Carlo efficiency corrections for π^0 mesons and photons. Although this procedure indicates that no correction is necessary, the errors on the correction functions represent uncertainties in the Monte Carlo model and contribute a systematic uncertainty of 3.6%.

To test for uncertainties in the background shape of the mass difference distributions, upper and lower sidebands in the $K^+ K^- \pi^+$ and $\gamma\gamma$ mass distributions are considered. Positive signal yields are expected in these sidebands from either misreconstructed or unassociated π^0 candidates. To measure these yields, the same fit functions used to determine the signal yields are applied to the mass difference distributions of the sideband samples. Any discrepancy in yield so obtained from data and Monte Carlo simulation is considered a systematic uncertainty (4.8%). Most of this uncertainty is attributed to the relatively large background in the $D_s^+ \pi^0$ decay mode.

The measurement of $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$ is repeated for the subsamples of candidates within various p^* intervals. By fitting either a constant function or a first-order polynomial to the branching ratio as a function of p^* , it is possible to verify that the measured branching ratios are independent of p^* (see Fig. 3). Nevertheless, it is assumed conservatively that any p^* dependence arises from unknown momentum dependencies of the efficiencies that may not cancel in the branching ratios. The difference (6.8%) between the branching ratio represented by the constant function and the integral of the first-order polynomial is therefore reported as a systematic uncertainty.

The systematic uncertainties are summarized in Table I. Combining all contributions in quadrature, a total system-

FIG. 3. The measured values of (a) $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$ and (b) $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma)$ in intervals of center-of-mass momentum p^* . The error bars indicate the associated statistical error. The solid line (dashed line) is the result of a fit to a constant (first-order polynomial). The dotted line is the result from the entire sample integrated over p^* .

atic uncertainty of 10.2% is derived for the measurement of $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$.

The ratio $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma)$, where $D^0 \rightarrow K^- \pi^+$, is measured using the same selection criteria for the π^0 and photon candidates as in the reconstruction of $D_s^{*+} \rightarrow D_s^+ \pi^0$ and $D_s^{*+} \rightarrow D_s^+ \gamma$. To be included in the $D^0 \rightarrow K^- \pi^+$ sample, a candidate K^- and π^+ combination must yield an acceptable fit to a common vertex and the scaled momentum x_p of the resulting D^0 candidate must be 0.6 or greater. Fitting the sum of a double Gaussian function and a third-order polynomial to the resulting $K^- \pi^+$ invariant mass distribution (not shown) produces $(996.0 \pm 1.5) \times 10^3$ signal events (statistical error only). A $K^- \pi^+$ combination is retained if its mass differs by less than 17 MeV/ c^2 from the nominal D^0 mass [8].

The D^0 candidates are combined with all π^0 candidates; the resulting mass difference $\Delta m(D^0 \pi^0) = m(K^- \pi^+ \pi^0) - m(K^- \pi^+)$ is shown in Fig. 4(a). A fit using a double Gaussian for the signal and the function shown in Eq. (1) for the background yields $69\,000 \pm 450$ signal events (statistical error only).

The D^0 candidates are then combined with all photon candidates producing the distribution of the mass difference $\Delta m(D^0 \gamma) = m(K^- \pi^+ \gamma) - m(K^- \pi^+)$ shown in Fig. 4(b). In this case, the peak corresponding to $D^{*0} \rightarrow D^0 \gamma$ signal is close to a large bump arising from the

TABLE I. A summary of the relative systematic uncertainties in the branching ratio measurements.

Sources	Relative Uncertainty (%)	
	$\frac{\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)}{\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)}$	$\frac{\Gamma(D^{*0} \rightarrow D^0 \pi^0)}{\Gamma(D^{*0} \rightarrow D^0 \gamma)}$
Background shape	4.8	0.1
Monte Carlo statistics	5.0	5.4
Signal model	3.6	3.8
p^* dependence	6.8	2.8
Quadrature Sum	10.2	7.2

FIG. 4. The D^{*0} signals: (a) $m(K^- \pi^+ \pi^0) - m(K^- \pi^+)$ and (b) $m(K^- \pi^+ \gamma) - m(K^- \pi^+)$. The dots represent data points. The solid curve shows the fitted function. The dashed curve indicates the portion of the fit associated with background.

reflection of $D^{*0} \rightarrow D^0 \pi^0$ in which one photon is produced by π^0 decay (the same reflection appears in D_s^{*+} decay but with a lower rate and less distinctive shape). Most of this bump is avoided by limiting the analysis to $\Delta m > 95 \text{ MeV}/c^2$. The remainder of the background is modeled using the function

$$f_3(\Delta m) = N \left(1 + \exp \left(- \frac{\Delta m - m(\pi^0)}{\mu} \right) \right) \times (\Delta m^2 + a\Delta m + b). \quad (3)$$

[Note that this function is similar to that of Eq. (1), but differs in the sign of the exponential term.] The signal is modeled by the Crystal Ball function [Eq. (2)]. The resulting fitted signal consists of $67\,880 \pm 670$ events (statistical errors only).

Efficiencies and systematic uncertainties are determined using the procedures described for $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$. Efficiencies of $\epsilon(D^0 \pi^0) = 0.037 \pm 0.002$ and $\epsilon(D^0 \gamma) = 0.064 \pm 0.002$ and an efficiency ratio of $\epsilon(D^0 \pi^0)/\epsilon(D^0 \gamma) = 0.58 \pm 0.03$ are found. The latter is consistent with the value of $\epsilon(D_s^+ \pi^0)/\epsilon(D_s^+ \gamma)$. The ratio $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma) = 1.74 \pm 0.02(\text{stat.}) \pm 0.13(\text{syst.})$ is obtained.

The branching ratio measurements are summarized in Table II. By assuming that the D_s^{*+} meson decays only to $D_s^+ \pi^0$ and $D_s^+ \gamma$, and that the D^{*0} meson decays only to $D^0 \pi^0$ and $D^0 \gamma$, it is possible to calculate the branching fractions, which are also listed in Table II.

In summary, the branching ratio $\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma) = 0.062 \pm 0.005(\text{stat.}) \pm$

TABLE II. Summary of the results. The first errors are statistical; the second represent systematic uncertainties.

$\Gamma(D_s^{*+} \rightarrow D_s^+ \pi^0)/\Gamma(D_s^{*+} \rightarrow D_s^+ \gamma)$	0.062	0.005	0.006
$\mathcal{B}(D_s^{*+} \rightarrow D_s^+ \pi^0)$	0.059	0.004	0.006
$\mathcal{B}(D_s^{*+} \rightarrow D_s^+ \gamma)$	0.942	0.004	0.006
$\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma)$	1.74	0.02	0.13
$\mathcal{B}(D^{*0} \rightarrow D^0 \pi^0)$	0.635	0.003	0.017
$\mathcal{B}(D^{*0} \rightarrow D^0 \gamma)$	0.365	0.003	0.017

$0.006(\text{syst.})$ has been measured and is consistent with the previous measurement [7], but has higher precision. Also determined is the ratio $\Gamma(D^{*0} \rightarrow D^0 \pi^0)/\Gamma(D^{*0} \rightarrow D^0 \gamma) = 1.74 \pm 0.02(\text{stat.}) \pm 0.13(\text{syst.})$. This result is in agreement with, but is more precise than, the world average [8].

It has been proposed that the decay $D_s^{*+} \rightarrow D_s^+ \pi^0$ proceeds via $\eta - \pi^0$ mixing and calculations based on Chiral perturbation theory [3] predict $\mathcal{B}(D_s^{*+} \rightarrow D_s^+ \pi^0) \approx 1\%-3\%$ based on current measurements of $\mathcal{B}(D^{*+} \rightarrow D^+ \gamma) = 1.6 \pm 0.4\%$ [8]. Newer theoretical estimates in a relativistic quark model [2] predict $\mathcal{B}(D_s^{*+} \rightarrow D_s^+ \pi^0) \approx 13\%$, somewhat larger than our measurement.

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy Physics (China), the Commissariat à l'Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, and the Particle Physics and Astronomy Research Council (United Kingdom). Individuals have received support from CONACyT (Mexico), the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

[1] The use of charge conjugate states is implied throughout this paper.

[2] J.J. Goity and W. Roberts, Phys. Rev. D **64**, 094007 (2001).

B. AUBERT *et al.*

PHYSICAL REVIEW D **72**, 091101 (2005)

- [3] P. L. Cho and M. B. Wise, Phys. Rev. D **49**, 6228 (1994).
- [4] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **90**, 242001 (2003).
- [5] D. Besson *et al.* (CLEO Collaboration), Phys. Rev. D **68**, 032002 (2003).
- [6] K. Abe *et al.* (BELLE Collaboration), Phys. Rev. Lett. **92**, 012002 (2004).
- [7] J. Gronberg *et al.* (CLEO Collaboration), Phys. Rev. Lett. **75**, 3232 (1995).
- [8] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [9] B. Aubert *et al.* (BABAR Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [10] A. Drescher *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **237**, 464 (1985).
- [11] D. Antreasyan *et al.* (Crystal Ball Collaboration), Crystal Ball Report No. 321 (1983).