

Lawrence Berkeley National Laboratory

Recent Work

Title

ORBITAL RADII, BOND LENGTH AND RESONANCE

Permalink

<https://escholarship.org/uc/item/1xt1q045>

Author

Somayajulu, G.R.

Publication Date

1961-02-01

UNIVERSITY OF
CALIFORNIA

Ernest O. Lawrence

*Radiation
Laboratory*


TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.



UCRL-9568

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405 -eng-48

ORBITAL RADII, BOND LENGTH AND RESONANCE

G.R. Somayajulu

February 1961

ORBITAL RADII, BOND LENGTH AND RESONANCE¹

G. R. Somayajulu

Department of Chemistry and
Lawrence Radiation Laboratory
University of California
Berkeley 4, California

¹ Part I: G. R. Somayajulu, J. Chem. Phys. 31, 919 (1959).

The radius of a s-p hybrid orbital has been found to be given by the expression: $r = A - B/(1 + \lambda^2)$ where A is the radius of the pure p orbital, B, a universal constant equal to 0.336 A and λ , the coefficient of mixing in the hybrid $s + \lambda p$. When radii appropriate for the orbitals that are paired together are used, bond length is additive of the orbital radii and no Schomaker-Stevenson correction is necessary. This shows that most bonds can be treated as covalent, single and localized.

1. INTRODUCTION

According to the current theories on bond length²⁻⁴, covalent bond

² L. Pauling, The Nature of the Chemical Bond, 3rd ed. (Cornell University Press, Ithaca, New York, 1960).

³ V. Schomaker and D. P. Stevenson, J. Am. Chem. Soc. 63, 37 (1941).

⁴ W. Gordy, W. V. Smith, and R. F. Trambarulo, Microwave Spectroscopy (John Wiley & Sons, Inc., New York, 1953), p. 313.

length is additive of the radii of the orbitals that are paired together. Deviations from additivity occur when the bonds are either partially ionic or partially double. The suggestion that some bonds are partially ionic is usually taken to follow from the concept of ionic-covalent resonance; the suggestion that some bonds acquire partial double bond character follows from the concept of π -electron resonance. It has been found, however, that C-C bond length in various molecules with one ground state structure is a sum^{1,5} of the orbital radii. Also the C-C bond energy has

⁵ M. G. Brown, Trans. Faraday Soc. 55, 694 (1959).

been found⁶ to be a sum of the orbital strengths. These results indicate

⁶ G. R. Somayajulu, submitted for publication.

two things; first of all they show that π -electron resonance is either negligible or absent in such molecules; secondly, they indicate the inapplicability of the Schomaker-Stevenson correction.

In many cases when hybridization is taken into account, we may speak of the molecule as composed of localized electron-pair bonds. Such bonds should be covalent. Their lengths will be simply sums of the orbital radii and the Schomaker Stevenson term becomes redundant. Similarly, Pauling's term, $(\chi)^2$, which represents the square of the difference between the electronegativities of the bonded atoms, becomes redundant when we use appropriate orbital strengths for the estimation of the bond energy. However, Mulliken's electronegativity⁷, which is conceptually

⁷ R. S. Mulliken, J. Chem. Phys. 2, 782 (1934); 3, 573 (1935).

different from that of Pauling, retains its full significance even when we take hybridization into account. Mulliken's electronegativity increases with increase in the strength (with increase in the s-character) of the hybrid and may rightly be related to the strength of the orbital. If resonance due to other schemes of pairing were to be important, we should have to seek a new relationship between the resonance energy and the electronegativities of the paired orbitals. It would seem more reasonable to expect the value of $(\Delta\chi)^2$ to vary with hybridization than having a constant value.

In this paper, we have suggested values for the orbital radii of some atoms, obtained on the assumption that the radius of a s-p hybrid is

linearly related⁸ to the percentage s-character in the hybrid. These

⁸ M. J. S. Dewar and H. N. Schmeising, Tetrahedron 5, 166 (1959).

radii prove the additive nature of the bond length. In the following paper⁶, we have estimated the energies of some bonds and showed them to be sums of the orbital strengths.

2. METHOD OF ESTIMATION OF ORBITAL RADII

The expression for the radius of a $s + \lambda p$ hybrid may be written as:

$$r = A - B/(1 + \lambda^2) \quad (1)$$

where A and B are constants characteristic of each atom and λ is the coefficient of mixing. $100/(1 + \lambda^2)$ gives the percentage s-character in the hybrid. When $\lambda = 0$, $r (= A - B)$ represents the radius of a pure s orbital. When $\lambda = \infty$, $r (= A)$ represents the radius of the pure p orbital.

The tetrahedral and digonal radii^{of carbon} are known to be^{1,5} 0.772 Å and 0.688 Å respectively. With the help of these values, the values of A and B of Eq. (1) are 0.856 Å and 0.336 Å respectively. For $\lambda^2 = 2$, i.e., for the trigonal orbital, Eq. (1) leads to a value of 0.744 Å, a value which may be deduced from the length^{9,10} of such a bond in 1,3-butadiene (1.483 Å) and

⁹ A. Almennigen, O. Bastiansen and M. Traetteberg, Acta Chem. Scand. 12, 1221 (1958).

¹⁰ A. Almennigen and O. Bastiansen, D.K.N.V.S. Skrifter, 1958, NR 4.

diphenyl (1.493 Å). A slightly lower value will result for the trigonal radius if the radius of the hybrid orbital is linearly related⁵ to λ^2 ; also it leads to a higher value for the radius of the pure s orbital. Expressions relating r and λ , suggested by Coulson¹¹ and Wilmhurst¹² lead

¹¹ C. A. Coulson, Victor Henri Commemorative Volume (Desoer, Liege, 1948), p. 15.

¹² J. K. Wilmhurst, J. Chem. Phys. 33, 813 (1960).

to radii which on addition necessitate a correction^{13,14} for the effect of

¹³ See B. Bak and L. Hansen-Nygaard, J. Chem. Phys. 33, 413 (1960).

¹⁴ See R. S. Mulliken, Tetrahedron 6, 68 (1959).

resonance or for the ionic character of the bond. When appropriate radii for the orbitals are used such corrections are unnecessary.

Estimation of radii using Eq. (1) is, however, difficult because of our poor knowledge of the valence states of atoms in various molecules. Excepting in a few simple cases, the bond angles do not seem to provide sufficient clues to the nature of bonding. However, we may take advantage of the fact that the shortenings for the double and triple bond radii of various atoms in the Periodic Table are almost the same as for carbon; one may assume, therefore, that the shortening from one hybrid to another will be approximately equal to those for the hybrids of carbon. This is equivalent to the assumption that B of Eq. (1) is constant for all atoms.

Then knowing the radius for one orbital of an atom, we may deduce the radii for the other orbitals of that atom by using Eq. (1). This procedure has been followed for the derivation of the orbital radii. When atoms use orbitals other than s and p for the formation of the hybrid orbitals, the radii of such orbitals may be worked out through the method of additivity. Such radii have not been considered in this paper. The results for the s-p hybrids are now discussed. The bond lengths in the halides of the second group elements are those obtained by Akishin et al.¹⁵ and quoted by

¹⁵ P. A. Akishin and V. P. Spiridonov, *Kristallografiya* 2, 475 (1957).

Bastiansen and Lund¹⁶. The interatomic distances in crystals with the

¹⁶ O. Bastiansen and E. W. Lund, *Ann. Rev. Phys. Chem.* 10, 31 (1959).

sphalerite and wurtzite arrangements are from Wyckoff¹⁷. If not indicated

¹⁷ R. W. G. Wyckoff, Crystal Structures, Vol. I. (Interscience Publishers, Inc., New York, 1960).

otherwise, all other bond lengths quoted in this paper are from the Tables of Interatomic Distances and Configuration of Molecules and Ions (The Chemical Society, London, 1958).

3. ORBITAL RADII OF NITROGEN

If we assume that nitrogen uses p orbital for bonding with another

nitrogen atom in hydrazine, the length (1.47 Å) of the N-N bond of hydrazine leads to the radii, 0.735 Å for p, 0.651 Å for sp^3 and 0.623 Å for sp^2 orbitals. The above values for the tetrahedral and trigonal radii of nitrogen may be supported as shown below:

The $\angle$ HNH angle of ammonia is close to the tetrahedral indicating that nitrogen atom in this molecule is in the tetrahedral state of hybridization. Additivity of the tetrahedral radius of nitrogen and the normal covalent radius of hydrogen (0.371 Å) gives 1.022 Å for the length of the N-H bond: this value compares satisfactorily with the observed value, 1.016 ± 0.01 Å.

There are sufficient grounds to believe that the nitrogen atom of formamide is hybridized in the trigonal state¹⁸. The C-N and N-H bonds

¹⁸ W. J. O. Thomas, Chem. Rev. 57, 1179 (1957).

of formamide have¹⁹ the lengths of 1.343 ± 0.007 Å and 0.995 ± 0.007 Å

¹⁹ R. J. Kurland and E. B. Wilson, Jr., J. Chem. Phys. 27, 585 (1957).

respectively. Our values for the $C(sp^2)-N(sp^2)$ and $N(sp^2)-H$ bonds are 1.367 Å and 0.994 Å respectively. These are in satisfactory agreement with the observed values. The length of the >C-NH_2 bond in p-nitraniline is 1.37 Å; this is also in good agreement with our value, 1.367 Å.

The molecule, $S_4N_4H_4$, is a puckered eight-membered ring of alternating sulphur and nitrogen atoms^{20,21}. The sulphur bond angle and the dihedral

²⁰ R. L. Saas and J. Donohue, Acta Cryst. 11, 497 (1957).

²¹ E. W. Lund and S. R. Svensen, Acta Chem. Scand. 11, 940 (1957).

angle have values close to those of S_8 . The $\hat{S}NS$ angle has been found to be close to 120° , which suggests the presence of trigonal planar $S-NH-S$ groupings. The average $S-N$ bond length is 1.674 A. Assuming the radius of sulphur atom to be 1.04 A as in S_8 , and by using the trigonal radius of nitrogen, we obtain a value of 1.663 A for the length of this bond. This is in good agreement with the observed value.

In trisilylamine, $N(SiH_3)_3$, the $\hat{Si}NSi$ angle is close to 120° and the length of the $Si-N$ bond is 1.733 A. Using our revised value for the tetrahedral radius (1.113 A) of silicon (see Section 5) and the trigonal radius of nitrogen, our value for the $Si(sp^3)-N(sp^2)$ bond is 1.741 A. This is in good agreement with the observed value. Thus these examples provide some support for our assumptions in deriving these radii. The results for the $C-N$ and $N-H$ bonds of formamide also show that one ground state structure involving a $C(sp^2)-N(sp^2)$ bond represents the structure of formamide. Slight discrepancies in the estimated lengths are probably due to the neglect of the correction for the non-equivalence of the orbitals in some of these compounds.

4. ORBITAL RADII OF HALOGENS

A good deal of information about the bonding orbitals may be obtained from a consideration of the interatomic distances in a diatomic hydride, HX , and its ion, HX^+ . Two classes of hydrides may be recognized²²; one in

²² H. A. Bent, J. Chem. Phys. 33, 1258 (1960).

which the length of the X-H bond increases when an electron is removed from X; examples for this class of hydrides are provided by CH, NH, OH, HCl, HBr, etc. In the second class of hydrides, the length of the X-H bond decreases when an electron is removed from X; examples for this class are provided by BeH, MgH, HgH, BH, AlH, etc. The reason for this may be attributed²² to a change in the s-content of the bonding orbital of X; in the first class of hydrides there may be a decrease in the s-content of the bonding orbital of X resulting in an increase in the orbital radius of X; this makes the length of the X-H bond in HX^+ greater than in HX. Just the opposite of this probably takes place in the second class of hydrides. It follows from this that the bonding orbital of C in CH is not pure p as it is usually thought to be. Likewise, in hydrogen halides, the halogen atoms do not use their pure p orbitals but hybrids of s and p. Since there are no easy methods to predict the percentage s-character in these hybrids, we will assume the applicability of Eq. (1) to these hybrids and predict the percentage s-character in the hybrid from the observed radius.

As already stated in Section 2, Eq. (1) can be used only when the radius of at least one orbital is known. It is possible to obtain the radii of the atoms in various molecules, but it is not possible in most cases to say to which orbitals they actually correspond. For example, it is customary to assume that the bonding orbitals of chlorine atoms in Cl_2 are p; what is the proof for this? In fact, there are now a number of evidences, as shown below, which suggest that the single bond in Cl_2 is probably a tetrahedral-tetrahedral bond. Pauling showed that the tetrahedral radius of chlorine in the crystal of CuCl with the sphalerite arrangement

is close to the radius of the chlorine atom in Cl_2 . Recently we have found²³

²³ G. R. Somayajulu, J. Chem. Phys. 33, 1541 (1960).

the value of $k_e r_e / D_0$ (where k_e is the force constant, r_e the internuclear distance, and D_0 , the dissociation energy) to be constant for similar diatomic molecules with constant bond type. Among the halogens, F_2 behaves anomalously indicating that the bond type in F_2 is different from that in Cl_2 , Br_2 and I_2 . However, the values of $k_e r_e / D_0$ for all the four hydrogen halides are approximately constant indicating that the bond type in the hydrogen halides is nearly the same. Further we have found the radius of the fluorine atom in F_2 is about 0.170 Å greater than its radius in HF. On the basis of Eq. (1) the difference between the p and digonal radii is also 0.168 Å. From this it may be inferred that the bonding orbital of F is p in F_2 and sp in HF. In the case of Cl, the difference between its radii in Cl_2 and HCl is 0.090 Å. On the basis of Eq. (1), the difference between either p and tetrahedral or digonal and tetrahedral is 0.084 Å. Since the bond type in all the hydrogen halides is nearly the same (as indicated by the $k_e r_e / D_0$ values) we may infer from the above that the bonding orbital of Cl is tetrahedral in Cl_2 and digonal in HCl. As expected the results for Br and I are consistent with the results for Cl. On the basis of the considerations presented above, the orbital radii of the halogen atoms have been deduced and presented in Table I. We have shown in Table II that, with the help of these radii, it is possible to estimate the lengths of the carbon-halogen bonds in several molecules. The type of bonding may be inferred from the values of the orbital

Table I

Orbital radii of the halogen atoms							
Halogen	$r(p)$	$r(sp^3)$	$r(sp^2)$	$r(sp)$	X_2	Radius in HX^+	HX
F	0.715	0.631	0.603	0.547	0.715		0.547
Cl	1.078	0.994	0.966	0.910	0.994	0.944	0.904
Br	1.225	1.141	1.113	1.057	1.141	1.088	1.043
I	1.417	1.333	1.305	1.249	1.333		1.237

Table II

Comparison of the estimated and observed interatomic distances of carbon-halogen bonds in various compounds ^a					
Molecule	Bond	r(C)	r(X)	r(C-X)	r(C-X) _{obs.}
CF ₄	C-F	0.772(te)	0.547(di)	1.319	1.325 ± 0.005
C ₂ F ₄	C-F	0.744(tr)	0.547	1.291	1.30 ± 0.01
C ₆ H ₅ F	C-F	0.744	0.547	1.291	1.30 ± 0.01
CHFO	C-F	0.744	0.603(tr)	1.343	1.351 ± 0.013
CH ₃ F	C-F	0.772	0.631(te)	1.403	1.391 ± 0.005
CCl ₄	C-Cl	0.772	0.994(te)	1.766	1.766 ± 0.005
CH ₂ =CHCl	C-Cl	0.744	0.994	1.738	1.736 ± 0.01
C ₆ H ₅ Cl	C-Cl	0.744	0.944 ^b	1.688	1.69 ± 0.01
C ₂ Cl ₂	C-Cl	0.688(di)	0.944	1.632	1.635 ± 0.004
CBr ₄	C-Br	0.772	1.141?	1.913?	1.942 ± 0.005
CH ₂ =CHBr	C-Br	0.744	1.141(te)	1.885	1.891 ± 0.01
C ₂ Br ₂	C-Br	0.688	1.141	1.829	1.80 ± 0.03
CI ₄	C-I	0.772	1.333(te)	2.105	2.12 ± 0.02
CH ₂ =CHI	C-I	0.744	1.333	2.077	2.089 ± 0.01
C ₂ I ₂	C-I	0.688	1.333	2.021	2.03 ± 0.01

- (a) The orbitals to which the radii correspond are shown in parentheses, te = tetrahedral, tr = trigonal and di = digonal.
 (b) Orbital radius of chlorine in HCl⁺.

radii required to fit the bond lengths. Further examples in support of the orbital radii of the halogens will be given elsewhere in this paper.

5. TETRAHEDRAL AND OTHER RADII OF SOME ELEMENTS

Pauling derived the tetrahedral radii of several elements from the interatomic distances in crystals with the diamond, sphalerite and wurtzite arrangements. With slight revisions most of his values are acceptable to us. In the case of silicon, however, the value, 1.110 Å that may be obtained from the interatomic distance in silicon carbide appears to be more appropriate than the value 1.17 Å obtained by Pauling from the interatomic distance in Si (c). Similarly, the value, 1.22 Å suggested by Pauling for the tetrahedral radius of Ge appears to be quite high. We suggest a value of 1.16 Å. We also suggest the values: 1.405 Å, 1.45 Å, 1.26 Å and 1.37 Å for the tetrahedral radii of Sn, Pb, Tl and Zr, respectively. These radii may be supported as shown in Table III.

From the interatomic distances in the crystals of AlN, AlAs, AlSb, GaN, GaP, GaAs, GaSb, InN, InP, InAs, and InSb, the tetrahedral radii of Al, Ga, In, N, P, As and Sb have been found to be 1.30, 1.30, 1.49, 0.65, 1.05, 1.13 and 1.33 Å respectively. From these, the p and trigonal radii of these elements have been obtained by using Eq. (1). These are presented in Table IV. We have also included in this table the orbital radii of bismuth obtained from the interatomic distances in the compounds of bismuth.

It is quite interesting to see that the trigonal radius (1.277 Å) of gallium from the length (2.50 Å) of the Ga-I bond in GaI_3 is quite close to 1.272 Å estimated by us. From the interatomic distances in AlH and InH,

Table III

Comparison of the estimated and observed interatomic distances in MX_4 and other types of compounds of Si, Ge, Sn, Pb, Ti and Zr.

Molecule	Bond	$r(\text{M})$	$r(\text{X})$	$r(\text{M-X})$	$r(\text{M-X})_{\text{Obs.}}$
SiH_4	Si-H	1.118(te)	0.371	1.489	1.480 ± 0.005
SiCl_4	Si-Cl		0.910(di)	2.028	2.03 ± 0.01
SiBr_4	Si-Br		1.057(di)	2.175	2.17 ± 0.01
SiI_4	Si-I		1.333(te)	2.451	2.435 ± 0.01
$\text{Si}(\text{CH}_3)_4$	Si-C		0.772(te)	1.890	1.888 ± 0.02
$(\text{C}_6\text{H}_5)_2\text{SiCl}_2$	Si-C		0.744(tr)	1.862	1.84 ± 0.015
Si_2Cl_6	Si-Si		1.118	2.236	2.24 ± 0.06
GeH_4	Ge-H	1.160(te)	0.371	1.531	1.53 ± 0.01
GeClF_3	Ge-F		0.547(di)	1.707	1.688 ± 0.017
	Ge-Cl		0.910	2.070	2.067 ± 0.003
GeBr_4	Ge-Br		1.141(te)	2.301	2.297 ± 0.005
GeI_4	Ge-I		1.333	2.493	2.50 ± 0.03
SnCl_4	Sn-Cl	1.405(te)	0.910	2.315	2.33 ± 0.02
SnBr_4	Sn-Br		1.057	2.462	2.46 ± 0.02
SnI_4	Sn-I		1.249(di)	2.654	2.64 ± 0.04
$\text{Sn}(\text{CH}_3)_4$	Sn-C		0.772	2.177	2.18 ± 0.03
PbCl_4	Pb-Cl	1.450(te)	0.994(te)	2.444	2.465 ± 0.02
$\text{Pb}(\text{CH}_3)_4$	Pb-C		0.772	2.222	2.203 ± 0.01
$\text{Pb}_2(\text{CH}_3)_6$	Pb-Pb		1.450	2.900	2.83 ± 0.03
TiCl_4	Ti-Cl	1.260(te)	0.910	2.170	2.18 ± 0.04
TiBr_4	Ti-Br		1.057	2.317	2.31 ± 0.02
ZrCl_4	Zr-Cl	1.370(te)	0.910	2.280	2.33 ± 0.05

Table IV

Orbital radii of Al, Ga, In, N, P, As, Sb and Bi			
Element	$r(p)$	Tetrahedral radius	Trigonal radius
Al	1.384	1.30	1.272
Ga	1.384	1.30	1.272
In	1.574	1.49	1.462
N	0.734	0.65	0.622
P	1.134	1.05	1.022
As	1.214	1.13	1.102
Sb	1.414	1.33	1.302
Bi	1.514	1.43	1.402

the radii of Al and In are 1.276 Å and 1.466 Å respectively; these are quite close to the trigonal radii obtained by us.

The values 1.214 Å and 1.414 Å obtained by us for the radii of the p orbitals of As and Sb are quite close to the values assigned by Pauling for the single bond radii of these elements. Support of these values is shown in Table V.

Pauling's values for the tetrahedral radii of S, Se and Te are 1.04 Å, 1.14 Å and 1.32 Å respectively; they are acceptable to us. Pauling's value for the tetrahedral radius of O is 0.66 Å; we recommend the value 0.651 Å, which leads to the values 0.735 Å and 0.623 Å for the p and trigonal radii of oxygen. The p orbital radii of Se and Te are 1.224 Å and 1.404 Å respectively. These radii may be supported as shown in Table VI.

The tetrahedral radii of Be, Mg, Zn, Cd and Hg have been found to be about 0.084 Å greater than the radii of these elements in their monohydrides. According to Eq. (1), the expected difference between the tetrahedral and digonal radius is 0.084 Å. Consequently, we may assume the bonding orbitals of these elements in their monohydrides to be digonal. We have, therefore, assigned the radii obtained from the interatomic distances in their monohydrides to their digonal orbitals and deduced their tetrahedral radii by using Eq.(1). The radii, thus obtained, have been compared with the tetrahedral radii obtained by Pauling in Table VII. There is a slight discrepancy in the case of Hg; we recommend the value 1.475 Å for its tetrahedral radius.

According to Eq. (1), the difference between the tetrahedral and s orbital radius is 0.252 Å. In EHg^+ , the radius of Hg is 1.223 Å; this is exactly 0.252 Å less than its tetrahedral radius 1.475 Å. On this basis,

Table V

Comparison of the estimated and observed interatomic distances
in MX_3 type compounds of P, As, Sb and Bi

Molecule	Bond	r(M)	r(X)	r(M-X)	r(M-X) _{Obs}
PH ₃	P-H	1.05(te)	0.371	1.421	1.4206 ± 0.005
PCl ₃	P-Cl	1.05	0.994(te)	2.044	2.043 ± 0.003
PBr ₃	P-Br	1.05	1.141(te)	2.191	2.18 ± 0.03
PI ₃	P-I	1.134(p)	1.333(te)	2.467	2.46 ± 0.03
AsH ₃	As-H	1.13(te)	0.371	1.501	1.519 ± 0.002
AsCl ₃	As-Cl	1.214(p)	0.994	2.208	2.17 ± 0.02
AsBr ₃	As-Br	1.214	1.141	2.355	2.36 ± 0.04
AsI ₃	As-I	1.214	1.333	2.547	2.55 ± 0.03
SbH ₃	Sb-H	1.33(te)	0.371	1.701	1.707 ± 0.0025
SbCl ₃	Sb-Cl	1.33	0.994	2.324	2.325 ± 0.005
SbBr ₃	Sb-Br	1.33	1.141	2.471	2.47 ± 0.04
SbI ₃	Sb-I	1.33	1.333	2.663	2.67 ± 0.03
BiCl ₃	Bi-Cl	1.514(p)	0.994	2.508	2.48 ± 0.02
BiBr ₃	Bi-Br	1.514	1.141	2.655	2.63 ± 0.02
(CF ₃) ₃ P	P-C	1.134	0.772(te)	1.906	1.937 ± 0.017
(CH ₃) ₃ As	As-C	1.214	0.772	1.986	1.98 ± 0.02
(CF ₃) ₃ Sb	Sb-C	1.414	0.772	2.186	2.20 ± 0.016
(C ₆ H ₅) ₃ Bi	Bi-C	1.514	0.744(tr)	2.258	2.30 ?

Table VI

Comparison of the estimated and observed lengths of some bonds
containing O, S, Se and Te atoms

Molecule	Bond	r(A)	r(B)	r(A-B)	r(A-B) _{Obs.}
CH ₃ OH	C-O	0.772(te)	0.651(te)	1.423	1.427 ± 0.007
HCOOH	C-O	0.744(tr)	0.623(tr)	1.367	1.37 ± 0.01
CH ₃ SH	C-S	0.772	1.04(te)	1.812	1.818 ± 0.005
CH ₃ COSH	C-S	0.744	1.012(tr)	1.756	1.78 ± 0.02
(CH ₃) ₂ Se	C-Se	0.772	1.224(p)	1.996	1.98 ± 0.02
(p-CH ₃ C ₆ H ₄)Te	C-Te	0.744	1.32(te)	2.064	2.05

Table VII

Comparison of the tetrahedral radii of Be, Mg, Zn, Cd and Hg obtained from the interatomic distances in the crystals with the sphalerite and wurtzite arrangements and those obtained from the digonal radius in monohydride.

Element	Digonal radius= $r(X-H)-0.371$	Tetrahedral radius= digonal radius + 0.034	Tetrahedral radius in crystals
Be	0.972	1.056	1.06
Mg	1.360	1.444	1.44 ^a
Zn	1.224	1.308	1.31
Cd	1.391	1.475	1.48
Hg	1.369	1.453	1.48

(a) Pauling's value is 1.40 Å, but subtraction of the tetrahedral radius of Te from the interatomic distance in HgTe crystal gives 1.44 Å.

Hg in HHg^+ has a s bonding orbital. From the interatomic distances in HMg^+ , HZn^+ , and HCd^+ , we obtain the values 1.278 Å for Mg, 1.144 Å for Zn and 1.296 Å for Cd. These radii correspond to about 75 % s-character. From the length of the C-Hg bond in $\text{CH}_3\text{-HgCl}$, the radius of Hg is 1.285 Å. This also corresponds to about 75% s-character. It is very strange that in dihalides (chlorides, bromides and iodides) of these elements, the radii which correspond to 75% s-character are found instead of the digonal radii. Calcium, strontium and barium, however, use their digonal radii obtained from the interatomic distances in CaH , SrH and BaH . The values for their digonal radii are 1.631 Å for Ca, 1.775 Å for Sr and 1.861 Å for Ba. The digonal radius of Be is 0.972 Å but it uses a radius of 0.85 Å in its dihalides. It is quite interesting to find that the Schomaker-Stevenson rule also leads¹⁶ to similar values for the radii of these elements. The results by using the above orbital radii are shown in Table VIII.

6. TRIGONAL RADIUS OF BORON

For the trigonal radius of boron, we propose a value of 0.795 Å. This value may be obtained from the interatomic distance in B_2 . There is, however, considerable doubt as to whether the bonding in B_2 is σ or π . The results using this value have been shown in Table IX. In borazole and BCB-trichloroborazole, both boron and nitrogen are hybridized in the trigonal state. The calculated length of the $\text{B}(\text{sp}^2)\text{-N}(\text{sp}^2)$ bond is 1.418 Å in good agreement with the observed value. Previously, several authors including the present²⁴ have thought that the structure of borazole is represented

²⁴ G. R. Somayajulu, J. Chem. Phys. 26, 822 (1958).

Table VIII

Comparison of the estimated and observed interatomic distances in the halogenides of the Group II elements				
Molecule	r(M)	r(X)	r(M-X)	r(M-X) _{obs.}
BeF ₂	0.85(?)	0.547(di)	1.397	1.40 ± 0.03
BeCl ₂		0.910(di)	1.760	1.75 ± 0.02
BeBr ₂		1.057(di)	1.907	1.91 ± 0.02
BeI ₂		1.249(di)	2.089	2.10 ± 0.02
MgCl ₂	1.278(?)	0.910	2.186	2.18 ± 0.02
MgBr ₂		1.057	2.335	2.34 ± 0.03
MgI ₂		1.249	2.527	2.52 ± 0.03
CaCl ₂	1.631(di)	0.910	2.541	2.51 ± 0.03
CaBr ₂		1.057	2.688	2.67 ± 0.03
CaI ₂		1.249	2.88	2.88 ± 0.03
SrCl ₂	1.775(di)	0.910	2.685	2.67 ± 0.03
SrBr ₂		1.057	2.832	2.82 ± 0.03
SrI ₂		1.249	3.024	3.03 ± 0.03
BaCl ₂	1.861(di)	0.994(te)	2.855	2.82 ± 0.03
BaBr ₂		1.141(te)	3.002	2.99 ± 0.03
BaI ₂		1.333(te)	3.194	3.20 ± 0.03
ZnCl ₂	1.144(?)	0.910	2.054	2.05 ± 0.01
ZnBr ₂		1.057	2.201	2.21 ± 0.01
ZnI ₂		1.249	2.393	2.38 ± 0.02
CdCl ₂	1.296(?)	0.910	2.206	2.21 ± 0.02
CdBr ₂		1.057	2.353	2.37 ± 0.02
CdI ₂		1.249	2.545	2.55 ± 0.02
HgCl ₂	1.285(?)	0.994	2.279	2.28 ± 0.02
HgBr ₂		1.141	2.426	2.41 ± 0.02
HgI ₂		1.333	2.618	2.59 ± 0.05

Table IX

Comparison of the observed and estimated lengths of B(sp ²) - X bonds. $r(B) = 0.795 \text{ \AA}$				
Molecule	Bond	$r(X)$	$r(B-X)$	$r(B-X)_{\text{Obs.}}$
BCl_3	B-Cl	0.910(di)	1.705	1.73 ± 0.02
BBr_3	B-Br	1.057(di)	1.852	1.87 ± 0.02
$\text{B}(\text{CH}_3)_3$	B-C	0.772(te)	1.567	1.56 ± 0.02
Borazole	B-N	0.623(tr)	1.418	1.44 ± 0.02
BBB-Trichloro- borazole	B-N	0.623	1.418	1.41 ± 0.02
	B-Cl	0.994(te)	1.789	1.78 ± 0.03
$\text{C}_6\text{H}_5\text{BCl}_2$	B-C	0.744(tr)	1.539	1.52 ± 0.07

by a resonance hybrid of two Kekulé-like structures, each B-N bond possessing about 50% double bond character. From the present considerations, the B-N bond of borazole appears to be a single bond.

7. CONCLUSION

From the results presented in the previous sections, we can see that this theory of bond length gives us an insight into the structure of bonding in compounds. Several examples quoted in the Tables II, III, V, VI, VIII and IX illustrate this. Our analysis is, however, qualitative. If we have to get a further insight into the structure of bonding, we shall have to consider the bond energies also. For example, the C-F bond in CF_4 is much shorter than indicated by the radii of the tetrahedral carbon of diamond and that of the F atom in F_2 . However, when the radius of the F atom in HF is used, the agreement is good. Consequently, we should also expect the F atom of CF_4 to have a much greater valence state energy than it has in F_2 , and presumably equal to that in HF. On the other hand, the C-Cl bond length in CCl_4 is given by the tetrahedral radius of carbon and the radius of the Cl atom in Cl_2 . It may be inferred from this that the valence state energy of Cl in this compound will presumably be equal to that in Cl_2 . This may be seen to be partly true. The heat of formation of CF_4 is -218 kcal/mole. Four times the heat of formation of HF is -256 kcal. On the other hand, the heat of formation of CCl_4 is -25 kcal/mole, a value which is just 9 kcal more negative than that of methane.

However, based on this theory we have failed to account for the lengths of bonds in some molecules such as CH_4 , SiF_4 , BF_3 , etc. Also

we were unable to obtain the lengths of the bonds in the halides of Be, Mg, Zn, Cd and Hg on the basis of the digonal radii of these metals. On the basis of the Pauli exclusion principle, these elements are expected to use digonal hybrids for bond formation. On the basis of Eq.(1) the radii used by them correspond to 75% s-character only. Taking these cases into consideration we think there are other factors besides hybridization which determine the bond lengths. Also Eq. (1) may not be applicable to all the elements. Probably cases which fall beyond the scope of this treatment necessitate considerations of resonance or corrections for the ionic character.

Also in some cases the bond lengths could not be estimated unambiguously. For example, we have estimated the length of the P-Br bond of PBr_3 on the basis of the tetrahedral radii of P and Br. Thus we obtain the value of 2.044 Å. This value may be obtained also by the addition of the p orbital radius of P and the digonal radius of Br. It is in such cases the bond energies will prove extremely useful.

Thus this theory suffers from several curious defects but still has several advantages over the current theories. It needs to be refined.

ACKNOWLEDGEMENTS

I am very greatly indebted to Professor Leo Brewer and Professor Kenneth S. Pitzer for helpful discussions on this subject, and to the U.S. Atomic Energy Commission for financial assistance.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

