UCSF

UC San Francisco Previously Published Works

Title

Sleep and Mood: Chicken or Egg?

Permalink

<https://escholarship.org/uc/item/1zj3s69s>

Journal

Biological psychiatry, 80(11)

ISSN

0006-3223

Authors

Ptáček, Louis J
Fu, Ying-Hui
Krystal, Andrew D

Publication Date

2016-12-01

DOI

10.1016/j.biopsych.2016.09.012

Peer reviewed

AUTHOR QUERY FORM

Journal: BPS Article Number: 12999	Please e-mail your responses and any corrections to: E-mail: a.norwitz@elsevier.com
---	---	--

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the [Q](#) link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
Q1	Please confirm that given names and surnames have been identified correctly and are presented in the desired order.
Q2	Author: If there are any drug dosages in your article, please verify them and indicate that you have done so by initialing this query.
Q3	Please review and confirm the accuracy and completeness of any affiliations.
Q4	AUTHOR: Please confirm that gene symbols are formatted in the correct style throughout your Commentary (human genes in all caps and italicized and mouse/rat genes in initial cap and italicized). Thank you.

Thank you for your assistance.

Please check this box or indicate your approval
if you have no corrections to make to the PDF file

Sleep and Mood: Chicken or Egg?

Q1 Louis J. Ptáček, Ying-Hui Fu, and Andrew D. Krystal

It has long been recognized that there is an association between sleep and mood. Both of these phenomena are light-sensitive; light entrains the circadian clock, and too little light predisposes a significant portion of the population to seasonal affective disorder. Sleep deprivation can precipitate mania in patients with bipolar disease (1) but can also be an effective treatment for “breaking” a bout of pharmacologically refractory depression (2). Glycogen synthase kinase 3 beta has been shown to be a circadian gene in flies (3) and mammals (4) and is a target of lithium therapy, which is useful in many patients with bipolar disease. Some mice with mutations causing circadian phenotypes show mood-like behaviors. However, people with mood disorders frequently experience difficulty falling and staying asleep and can manifest circadian phenotypes (e.g., early morning awakening and diurnal mood variation of severe depression) (5). Genetic variants in the circadian gene *PER3* were recently shown to cause a human circadian phenotype associated with seasonal affective disorder (6). Given this remarkable list of associations between sleep/circadian function and mood, one is left asking: “Which came first, the chicken or the egg?”

Evidence from previous studies suggests that sleep/circadian system modifications have mood effects. This includes alterations of light schedules or lesioning of the suprachiasmatic nuclei (SCN, the master circadian regulator in the brain), which can lead to phenotypes that are thought to mimic psychiatric disorders seen in humans. However, altering light schedules and lesioning of the SCN may have pleiotropic effects separate from central circadian dysregulation that complicate interpretations of these results.

In the article by Landgraf *et al.* (7), the authors show that circadian disruption resulting from knocking down expression of a gene (*BMAL1*, an essential positive regulator of circadian rhythmicity) in SCN leads to helplessness-like, behavioral despair-like, and depression-like behavior in mice. This is the best evidence to date suggesting that disturbance of the circadian system may, indeed, precipitate mood dysfunction.

Still, the issue of causality remains far from resolved. It is worth noting that there are complex interactions between behavioral phenotypes, circadian function, and sleep. It is also important to note that circadian function (i.e., rhythmicity) is distinct from sleep homeostasis. Clock genes in cells lead to cell autonomous oscillatory behavior of transcription, translation, and the posttranslational function of many genes and proteins. Therefore, circadian disruption itself may inevitably have behavioral consequences, and some behavioral phenotypes might result from altered sleep. Shift workers often do not fully synchronize to the “new” time schedule and this, in turn, can lead to sleep deprivation. We must therefore

consider separate contributions of circadian alterations and resulting sleep deprivation in thinking about behavioral consequences in humans where it is impossible to regulate the light environment and manipulate circadian function as easily as we can in model organisms. The fact that disruption of rhythms can lead to mice with features of human behavioral disorders is not mutually exclusive from the converse.

It is important to exploit opportunities that are possible in each system (including *Homo sapiens*!) while being cognizant of limitations that exist. Many researchers use *Mus musculus* and other model organisms because they afford genetic and other tools that are not possible in humans. Landgraf *et al.* (7) inject a viral vector encoding a short hairpin RNA to knock down expression of the clock gene *BMAL1*. This is a powerful tool that leads to circadian disruption without changing the lighting schedule or creating an ablation that leads to neuronal loss in SCN and disruption of light-sensing inputs and physiological outputs of the master clock. However, there are also inherent challenges in behavioral phenotyping of mice. It is impossible to really determine whether a mouse is depressed or anxious. The authors use available assays for measuring several mouse behaviors and show differences between experimental and control mice. For example, the mice with disrupted circadian function show increased depression-like behavior. They interpret this to suggest that depression can result from circadian disruption, but are careful to note important caveats in extending behavioral assessments from mice to humans.

Another important question is whether the converse is true. Is it possible for a primary behavioral phenotype like depression to lead to circadian dysregulation? Although it is clear that affect is (in part) genetically determined, genes or mutations causing affective disorders have been elusive. Once we have better understanding of the genetics underlying affective disorder, it will be possible to begin thinking about how to test this question. However, an inherent problem is the fact that affect is almost certainly a distributed network property of the brain. While it is possible to disrupt circadian function by knocking down an essential circadian gene in the SCN, there is no recognized locus in the central nervous system whose disruption will give rise to depression. Therefore, expression of a “depression mutation” may have systemic effects that could preclude strong conclusions regarding depression causing a circadian phenotype. Given the complex interactions of circadian function/sleep, mood, and other behaviors, it would not be surprising if the causal relationship between circadian/sleep and mood phenotypes is bidirectional. Several observations in humans suggest—but do not prove—that this may be the case (8–10). In the end, it may not be a question of “chicken versus egg” as much as one of “chicken AND egg.”

Acknowledgments and Disclosures

This work was supported by funding from the Howard Hughes Medical Institute and the National Institutes of Health (to LJP and Y-HF). ADK has received grant funding from the National Institutes of Health, Teva, Sunovion, Janssen, and Jazz Pharmaceuticals, and has also served as a consultant for Attentiv, Flamel, Teva, Jazz, Janssen, Merck, Neurocrine, Otsuka, Pfizer, Lundbeck, Sunovion, and Pernix.

Article Information

From the Department of Neurology (LJP, Y-HF, ~~ADK~~), Weill Neuroscience Institute (LJP, Y-HF, ADK), Kavli Institute for Fundamental Neuroscience (LJP, Y-HF), Howard Hughes Medical Institute (LJP), and the Department of Psychiatry (ADK), University of California, San Francisco, San Francisco, California.

Address correspondence to Louis J. Ptáček, M.D., HHMI/UCSF, Neurology, 548F Rock Hall, MC-2922, 1550 4th Street, San Francisco, CA 94143; E-mail: ljp@ucsf.edu.

Received Sep 16, 2016; revised Sep 21, 2016; accepted Sep 22, 2016.

References

1. Wehr TA (1991): Sleep-loss as a possible mediator of diverse causes of mania. *Br J Psychiatry* 159:576–578.
2. Post RM, Kotin J, Goodwin FK (1976): Effects of sleep deprivation on mood and central amine metabolism in depressed patients. *Arch Gen Psychiatry* 33:627–632.
3. Martinek S, Inonog S, Manoukian AS, Young MW (2001): A role for the segment polarity gene *shaggy/GSK-3* in the *Drosophila* circadian clock. *Cell* 105:769–779.
4. Kaasik K, Kivimae S, Allen JJ, Chalkley RJ, Huang Y, Baer K, *et al.* (2013): Glucose sensor O-GlcNAcylation coordinates with phosphorylation to regulate circadian clock. *Cell Metab* 17:291–302.
5. Schotte CK, Maes M, Cluydts R, Cosyns P (1997): Cluster analytic validation of the DSM melancholic depression. The threshold model: Integration of quantitative and qualitative distinctions between unipolar depressive subtypes. *Psychiatry Res* 71:181–195.
6. Zhang L, Hirano A, Hsu PK, Jones CR, Sakai N, Okuro M, *et al.* (2016): A *PERIOD3* variant causes a circadian phenotype and is associated with a seasonal mood trait. *Proc Natl Acad Sci U S A* 113: E1536–E1544.
7. Landgraf DLJ, Proulx CD, Barandas R, Malinow R, Welsh DK (2016): Genetic disruption of circadian rhythms in the suprachiasmatic nucleus causes helplessness, behavioral despair, and anxiety-like behavior in mice. *Biol Psychiatry* ●●:●●●–●●●.
8. Baglioni C, Battagliese G, Feige B, Spiegelhalter K, Nissen C, Voderholzer U, *et al.* (2011): Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord* 135:10–19.
9. Johnson EO, Roth T, Breslau N (2006): The association of insomnia with anxiety disorders and depression: exploration of the direction of risk. *J Psychiatr Res* 40:700–708.
10. Szklo-Coxe M, Young T, Peppard PE, Finn LA, Benca RM (2010): Prospective associations of insomnia markers and symptoms with depression. *Am J Epidemiol* 171:709–720.