

UC Berkeley

UC Berkeley Previously Published Works

Title

Contribution of GnIH Research to the Progress of Reproductive Neuroendocrinology

Permalink

<https://escholarship.org/uc/item/1zx9r8t8>

Journal

Frontiers in Endocrinology, 6(NOV)

ISSN

1664-2392

Authors

Tsutsui, Kazuyoshi
Ubuka, Takayoshi
Son, You Lee
[et al.](#)

Publication Date

2015

DOI

10.3389/fendo.2015.00179

Peer reviewed



Contribution of GnIH Research to the Progress of Reproductive Neuroendocrinology

Kazuyoshi Tsutsui^{1*}, Takayoshi Ubuka^{1,2}, You Lee Son¹, George E. Bentley³ and Lance J. Kriegsfeld⁴

¹Laboratory of Integrative Brain Sciences, Department of Biology and Center for Medical Life Science, Waseda University, Tokyo, Japan, ²Brain Research Institute Monash Sunway of the Jeffrey Cheah School of Medicine and Health Sciences, Monash University Malaysia, Petaling Jaya, Malaysia, ³Department of Integrative Biology, Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, CA, USA, ⁴Department of Psychology, Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, CA, USA

OPEN ACCESS

Edited by:

Yong Zhu,
East Carolina University, USA

Reviewed by:

T. John Wu,
Uniformed Services University of the
Health Sciences, USA
Pierre J. Deviche,
Arizona State University, USA

Yonathan Zohar,
University of Maryland, USA

*Correspondence:

Kazuyoshi Tsutsui
k-tsutsui@waseda.jp

Specialty section:

This article was submitted to
Experimental Endocrinology,
a section of the journal
Frontiers in Endocrinology

Received: 18 August 2015

Accepted: 05 November 2015

Published: 23 November 2015

Citation:

Tsutsui K, Ubuka T, Son YL,
Bentley GE and Kriegsfeld LJ (2015)
Contribution of GnIH Research to the
Progress of Reproductive
Neuroendocrinology.
Front. Endocrinol. 6:179.
doi: 10.3389/fendo.2015.00179

Since the discovery of gonadotropin-releasing hormone (GnRH) in mammals at the beginning of the 1970s, it was generally accepted that GnRH is the only hypothalamic neuropeptide regulating gonadotropin release in mammals and other vertebrates. In 2000, however, gonadotropin-inhibitory hormone (GnIH), a novel hypothalamic neuropeptide that actively inhibits gonadotropin release, was discovered in quail. Numerous studies over the past decade and a half have demonstrated that GnIH serves as a key player regulating reproduction across vertebrates, acting on the brain and pituitary to modulate reproductive physiology and behavior. In the latter case, recent evidence indicates that GnIH can regulate reproductive behavior through changes in neurosteroid, such as neuroestrogen, biosynthesis in the brain. This review summarizes the discovery of GnIH, and the contributions to GnIH research focused on its mode of action, regulation of biosynthesis, and how these findings advance our understanding of reproductive neuroendocrinology.

Keywords: gonadotropin-inhibitory hormone, gonadotropin-releasing hormone, gonadotropins, reproduction, reproductive behavior, melatonin, stress, social environment

HISTORY OF NEUROENDOCRINOLOGY AS A FIELD

The discovery of “neurosecretion” in the first half of the twentieth century led to the creation of neuroendocrinology, a new research field of endocrinology. In the 1920s, Ernst and Berta Scharrer first proposed the new concept of neurosecretion, suggesting that hypothalamic neurons terminating in the neurohypophysis secrete neurohormones analogous to those of endocrine glands. This seminal idea was not accepted easily by the scientific community, leading to marked criticism, including the comments like: “We have just heard some very interesting things, ... and also a great deal of nonsense.” In 1949, however, Bargmann established the concept of neurosecretion proposed by Ernst and Berta Scharrer. Subsequently, hypothalamic neuropeptides, oxytocin (1) and vasopressin (2), were identified to be neurohormones that are secreted from the neurohypophysis.

The morphology of hypothalamic neurons that terminate at the median eminence (ME) made Harris (3) to hypothesize that these hypothalamic neurons may secrete neurohormones from the ME into the hypophysial portal system to regulate the secretion of anterior pituitary hormones. Harris further summarized the first map showing different areas in the hypothalamus responsible for various

pituitary hormone release, identified by contemporary lesions and electrical stimulation studies [(4), reviewed in Ref. (5)]. Halasz et al. (6) also contributed to the identification of the hypophysiotrophic area in the hypothalamus by pituitary transplantation method (5). McCann and Ramirez were the first to demonstrate the biological existence of luteinizing hormone (LH)-releasing factor (LHRF) in the basal middle hypothalamus [see Ref. (7) for a review]. Subsequently, the groups of Schally and Guillemin identified the structure of several neurohormones that regulate anterior pituitary hormone release, including thyrotropin-releasing hormone (TRH) (8, 9), gonadotropin-releasing hormone (GnRH) (10, 11), and growth hormone-inhibiting hormone (somatostatin) (12), in the brain of mammals. Schally and Guillemin were awarded a Nobel Prize in 1977 for the discoveries of these neurohormones. At the same time, Yalow also received this prize for her role in the development of radioimmunoassay that permitted the measurement of these neurohormones.

As described above, Schally's (10) and Guillemin's (11) groups discovered GnRH, a hypothalamic neuropeptide that stimulates the release of both LH and follicle-stimulating hormone (FSH) from gonadotropes in the anterior pituitary, in mammals. Subsequently, several GnRHs have been identified in other vertebrates (13–16). Although McCann et al. (17) suggested differential hypothalamic control mechanism of FSH secretion, it had been generally accepted that GnRH is the only hypothalamic neuropeptide regulating gonadotropin release in mammals and other vertebrates.

In 2000, however, Tsutsui and colleagues challenged this notion with the discovery of gonadotropin-inhibitory hormone (GnIH), a hypothalamic neuropeptide that actively inhibits gonadotropin release, in quail (18). Subsequent studies conducted by Tsutsui and colleagues over the past decade and a half demonstrated that GnIH is highly conserved among vertebrates, from agnathans to humans, acting as a key player regulating reproduction [for reviews, see Ref. (19–30)]. Recent studies by Tsutsui's group have demonstrated that GnIH has important functions beyond the control of reproduction (31, 32). Based on these findings, it now appears that GnIH not only acts on the pituitary but in the brain to affect a number of behaviors, including reproductive behavior through changes in neurosteroid, such as neuroestrogen, biosynthesis in the brain [(32), for a review, see Ref. (33)]. Thus, the following 15 years of GnIH research in collaboration with world's leading laboratories have permitted a more complete understanding of the neuroendocrine control of reproductive behavior and physiology [for reviews, see Ref. (19–22, 24–29, 34)].

Herein, this review summarizes the discovery of GnIH and the contribution of GnIH research over the past decade and a half, focusing on its mode of action, regulation of biosynthesis, and the ways that such contributions have contributed to the field of reproductive neuroendocrinology. This review also highlights the commonalities and diversity of GnIH structures and actions as well as the evolutionary origin of GnIH in chordates.

DISCOVERY OF GnIH AND ITS ROLE IN REPRODUCTION

Gonadotropin-inhibitory hormone was discovered in the brain of the Japanese quail while searching for a novel peptide

possessing a C-terminal sequence Arg-Phe-NH₂ (RFamide peptide) in vertebrates (18). The first identification of an RFamide peptide dates back to the late 1970s, when Price and Greenberg purified a peptide Phe-Met-Arg-Phe-NH₂ (FMRFamide) from the ganglia of the venus clam (35). Since this initial discovery, various RFamide peptides that act as neurotransmitters, neuromodulators, and peripheral hormones had been identified in invertebrates. However, immunohistochemical studies in vertebrates suggested the presence of an unknown hypothalamic RFamide peptide(s) that may regulate the secretion of anterior pituitary hormones (36, 37). Based on this information, Tsutsui's laboratory searched for novel RFamide peptide(s) in the brain of the Japanese quail.

A breakthrough occurred in 2000 when Tsutsui and colleagues discovered a novel RFamide peptide Ser-Ile-Lys-Pro-Ser-Ala-Tyr-Leu-Pro-Leu-Arg-Phe-NH₂ (SIKPSAYLPLRFamide) that actively inhibited gonadotropin release in quail pituitary, providing the first demonstration of a hypothalamic neuropeptide inhibiting gonadotropin release in any vertebrate (18) (**Table 1**). Given its functional role, this neuropeptide was named GnIH (18) (**Figure 1**). In birds, cell bodies and terminals of GnIH neurons are located in the paraventricular nucleus (PVN) and ME, respectively (18). The C-terminal structure of GnIH is identical to chicken LPLRFamide which is the first reported RFamide peptide in vertebrates (38), but this peptide is likely to be a degraded fragment of chicken GnIH [(39), for reviews, see Ref. (20, 26, 27)] (**Table 1**). Subsequently, a cDNA encoding the precursor polypeptide for GnIH was identified in quail (40) and other avian species, such as chickens, sparrows, starlings, and zebra finches [for reviews, see Ref. (20, 26, 27)]. The GnIH precursor encompasses one GnIH and two GnIH-related peptides (GnIH-RP-1 and GnIH-RP-2) possessing an LPXRFamide (X = L or Q) motif at their C-terminus in all avian species studied. GnIH was further identified as a mature peptide in starlings (41) and zebra finches (42), and GnIH-RP-2 was identified in quail (40) (**Table 1**).

Gonadotropin-inhibitory hormone is considered to be a key neuropeptide inhibiting avian reproduction because GnIH inhibits gonadotropin release in most avian species studied [for reviews, see Ref. (20, 26, 27)] (**Figure 1**). To demonstrate the biological mode of action of GnIH *in vivo*, Ubuka et al. (57) treated mature male quail with chronic GnIH. Chronic GnIH treatment decreases plasma LH concentration and the expressions of the common α , LH β , and FSH β subunit mRNAs. Additionally, Ubuka et al. (57) established that GnIH treatment induces testicular apoptosis and decreases the diameter of seminiferous tubules and testosterone concentration in mature birds. In immature birds, GnIH treatment suppresses normal testicular growth and testosterone concentration (57). Thus, GnIH appears to suppress gonadal development and maintenance by decreasing gonadotropin synthesis and release in birds (**Figure 1**).

To determine if these findings extend to mammals, including humans, Tsutsui and colleagues sought to identify GnIH in the hypothalamus of mammals and primates (43, 44, 47, 48, 58). All the identified and putative mammalian and primate GnIH peptides possess a common C-terminal LPXRFamide (X = L or Q) motif, like avian GnIH and GnIH-RPs [for reviews, see Ref. (20,

TABLE 1 | Molecular structure of mature GnIH peptides in chordates.

Chordates	Peptide name	Molecular structure	Reference
Mammals	<i>human RFRP-1</i>	MPHSFANLPLRFa	Ubuka et al. (43)
	<i>human RFRP-3</i>	VPNLPQRFa	Ubuka et al. (43)
	<i>macaque RFRP-3</i>	SGRNMEVSLVRQVLNLPQRFa	Ubuka et al. (44)
	<i>bovine RFRP-1</i>	SLTFEEVKDWAPKIKMNKPVWNKMPPSAANLPLRFa	Fukusumi et al. (45)
	<i>bovine RFRP-3</i>	AMAHLPRLGKNGREDSLSRWVNPQRFa	Yoshida et al. (46)
	<i>rat RFRP-3</i>	ANMEAGTMSHFPSLPQRFa	Ukena et al. (47)
	<i>Siberian hamster RFRP-1</i>	SPAPANKVPHSAANLPLRFa	Ubuka et al. (48)
	<i>Siberian hamster RFRP-3</i>	TLSRVPSLPQRFa	Ubuka et al. (48)
Birds	<i>quail GnIH</i>	SIKPSAYLPLRFa	Tsutsui et al. (18)
	<i>quail GnIH-RP-2</i>	SSIQSLNLPQRFa	Satake et al. (40)
	<i>chicken GnIH</i>	SIRPSAYLPLRFa	McConn et al. (39)
	<i>European starling GnIH</i>	SIKPFANLPLRFa	Ubuka et al. (41)
	<i>zebra finch GnIH</i>	SIKPFANLPLRFa	Tobari et al. (42)
Reptiles	<i>red-eared slider GnIH</i>	SIKPVANLPLRFa	Ukena et al. (49)
	<i>red-eared slider GnIH-RP-1</i>	STPTVNKMPNSLANLPLRFa	Ukena et al. (49)
	<i>red-eared slider GnIH-RP-2</i>	SSIQSLANLPQRFa	Ukena et al. (49)
Amphibians	<i>bullfrog GRP/R-RFa</i>	SLKPAANLPLRFa	Koda et al. (50) and Chartrel et al. (51)
	<i>bullfrog GRP-RP-1</i>	SIPNLPQRFa	Ukena et al. (52)
	<i>bullfrog GRP-RP-2</i>	YLSGKTKVQSMANLPQRFa	Ukena et al. (52)
	<i>bullfrog GRP-RP-3</i>	AQYTNHFVHSLDTLPLRFa	Ukena et al. (52)
	<i>red-bellied newt LPXRFa-1</i>	SVPNLPQRFa	Chowdhury et al. (53)
	<i>red-bellied newt LPXRFa-2</i>	MPHASANLPLRFa	Chowdhury et al. (53)
	<i>red-bellied newt LPXRFa-3</i>	SIQPLANLPQRFa	Chowdhury et al. (53)
	<i>red-bellied newt LPXRFa-4</i>	APSAGQFIQTLANLPQRFa	Chowdhury et al. (53)
Teleost fishes	<i>goldfish LPXRFa-3</i>	SGTGLSATLPQRFa	Sawada et al. (54)
Agnathans	<i>sea lamprey LPXRFa-1a</i>	SGVGQGRSSKTLFQPQRFa	Osugi et al. (55)
	<i>sea lamprey LPXRFa-1b</i>	AALRSGVGQGRSSKTLFQPQRFa	Osugi et al. (55)
	<i>sea lamprey LPXRFa-2</i>	SEPFWHRTRPQRFa	Osugi et al. (55)
Protochordates	<i>amphioxus PQRFa-1</i>	WDEAWRPQRFa	Osugi et al. (56)
	<i>amphioxus PQRFa-2</i>	GDHTKDGWRPQRFa	Osugi et al. (56)
	<i>amphioxus PQRFa-3</i>	GRDQGWPRQRFa	Osugi et al. (56)


Only structurally determined mature endogenous peptides are shown.

21, 26–29)] (**Table 1**). Therefore, these GnIH peptides were designated as LPXRFamide peptides on the basis of their structures. Mammalian and primate GnIH peptides are also called RFamide-related peptide 1 and 3 (RFRP-1 and -3) (**Table 1**). Kriegsfeld et al. (58) found that *in vivo* administration of avian GnIH centrally or peripherally to female Syrian hamsters inhibits LH release (58). Central administration of hamster GnIHs (RFRP-1 and -3) also inhibits LH release in Siberian hamsters (48). Central administration of rat GnIH (RFRP-3) to male rats also inhibits LH release (59) and GnRH-elicited gonadotropin release (60, 61). Reduction in LH pulse amplitude and inhibition of GnRH-elicited gonadotropin release and synthesis by mammalian GnIH (RFRP-3) have also been reported in ovine (62, 63) and bovine (64). Because the structure of human GnIH (RFRP-3) is the same as ovine GnIH (RFRP-3) (43), the hypophysiotropic action of human/ovine GnIH (RFRP-3) was assessed in ovine pituitary in collaboration with the Clarke laboratory. Human/ovine GnIH (RFRP-3) inhibits GnRH-stimulated secretion of both LH and FSH (62) demonstrating that, as with avian GnIH, mammalian and primate GnIH inhibit gonadotropin release and synthesis and GnRH-elicited gonadotropin secretion [for reviews, see Ref. (19–21, 26–29)] (**Figure 1**).

COMMONALITIES AND DIVERSITY OF GnIH STRUCTURES AND ACTIONS

To place these findings into a broader perspective, Tsutsui and colleagues further identified GnIH peptides in the brain of reptiles, amphibians, and fish. All of the identified and putative GnIHs in these vertebrates also possess a common C-terminal LPXRFamide (X = L or Q) motif, like avian, mammalian, and primate GnIHs (49–54, 65, 66) (**Table 1**). Thus, GnIH peptides exist in the brain of vertebrates from fish to humans [for reviews, see Ref. (20–30)]. Goldfish GnIH precursor cDNA encodes three GnIHs, gflPXRFa-1, -2, and -3 (54). These goldfish GnIH peptides (gflPXRFa-1, -2, and -3) have both inhibitory and stimulatory effects on gonadotropin synthesis and release, possibly depending on reproductive conditions (67–70). Zebrafish GnIH peptide, zflPXRF-3, also has an inhibitory effect on gonadotropin release (71).

As mentioned above, GnIH peptides were identified in representative species of gnathostomes. However, GnIH peptide had not been identified in agnathans, the most ancient lineage of vertebrates (72). Therefore, Tsutsui and colleagues searched for agnathan GnIH in collaboration with the Sower and Nozaki



laboratories (55). Synteny analysis showed the existence of the gene for GnIH in sea lamprey, and Osugi et al. (55) cloned lamprey GnIH precursor cDNA that encodes GnIH peptides. Subsequently, three mature GnIH peptides were identified from the brain of sea lamprey by immunoaffinity purification and mass spectrometry (55) (Table 1). The identified lamprey GnIHs share a common C-terminal PQRfamamide motif (55), unlike GnIHs identified of gnathostomes. However, phylogenetic analysis showed that the identified lamprey GnIH precursor groups with the LPXRfa peptide precursors of vertebrates, whereas the previously identified lamprey PQRfa peptide precursor (73) groups with the PQRfa peptide precursors of vertebrates. Accordingly, we concluded that the lamprey GnIH precursor gene is the ortholog of LPXRfa peptide gene (55).

Lamprey GnIH neurons are located in the hypothalamus (55) with immunoreactive fibers projecting to GnRH3 neurons (55). Few lamprey GnIH immunoreactive fibers were observed in the neurohypophysis compared to abundant lamprey GnRH3

immunoreactive fibers (55). Osugi et al. (55) then analyzed the effects of lamprey GnIH peptides on the expressions of lamprey GnRHs and the gonadotropin β subunit. It was found that one of the lamprey GnIH peptides increases the expressions of lamprey GnRH3 and gonadotropin β RNA (55). Thus, GnIH is present in the brain of lamprey, the oldest lineage of vertebrates, and may act on GnRH3 neurons to stimulate the expression of gonadotropin β in the pituitary (55). Based on these findings, it is speculated that GnIH emerged in agnathans as a stimulatory neuropeptide and changed into an inhibitory neuropeptide during vertebrate evolution.

EVOLUTIONARY ORIGIN OF GnIH

The C-terminal structure of most GnIH peptides is LPXRfamamide (X = L or Q), making them a member of the RFamide peptide family [for reviews, see Ref. (20–30)]. Four more groups, i.e., the neuropeptide FF (NPFF; PQRfamamide peptide) group, the kisspeptin group, the pyroglutamylated RFamide peptide (QRFP)/26RFamide group, and the prolactin-releasing peptide (PrRP) group, have been documented in vertebrates [for reviews, see Ref. (20, 21, 23, 30)]. Because the C-terminal structure of GnIH peptides is similar to NPFF peptides that have a C-terminal PQRfamamide motif, further clarification of the NPFF peptide gene in agnathans was warranted. NPFF is a neuropeptide involved in pain modulation [for a review, see Ref. (74)]. Accordingly, Tsutsui and colleagues sought to identify the cDNAs of NPFF peptides in the brain of lamprey and hagfish (73, 75). Phylogenetic analysis established that agnathans possess both GnIH and NPFF precursor genes. Agnathan NPFF peptides were further identified in sea lamprey and hagfish. The identified agnathan NPFF peptides had the same C-terminal PQRfamamide motif as agnathan GnIH peptides (73, 75).

The findings that agnathans have both GnIH and NPFF genes and their mature peptides have the same C-terminal PQRfamamide motif (55, 73, 75) suggest that the GnIH and NPFF genes were derived from a common ancestral gene in protochordates. To test this hypothesis, Tsutsui and colleagues identified an amphioxus PQRfamamide peptide precursor cDNA that encodes three putative PQRfamamide peptides (56). Subsequently, three endogenous amphioxus PQRfamamide peptides were identified by immunoaffinity purification and mass spectrometry (56) (Table 1). Phylogenetic analysis showed that the amphioxus PQRfamamide peptide precursor occurs before the divergence between the GnIH and NPFF groups in vertebrates (56). Synteny analysis showed that the conserved synteny region exists around the loci of the amphioxus PQRfamamide peptide gene, GnIH gene, and NPFF gene in vertebrates (56). The amphioxus PQRfamamide peptide gene is located near the HOX cluster, whereas the GnIH and NPFF genes in vertebrates are located near the HOXA and HOXC clusters, respectively, suggesting that the GnIH and NPFF genes may have duplicated through whole-genome duplications (56). Accordingly, the amphioxus PQRfamamide peptide gene is considered to be close to the ancestor of the GnIH and NPFF genes (56, 76). Thus, the GnIH and NPFF genes may have diverged from a common ancestral gene in the protochordate through whole-genome duplication event during vertebrate evolution.

MODE OF GnIH ACTION ON GONADOTROPIN SECRETION

Identification of GnIH Receptor

To investigate the mode of action of GnIH on gonadotropin secretion in birds, Tsutsui and colleagues identified the receptor for GnIH in quail. They identified GnIH receptor as GPR147, a member of the G-protein-coupled receptor (GPCR) superfamily (77), which is also called NPFF receptor 1 (NPFF1). Membrane fraction of COS-7 cells that are transfected with GnIH receptor cDNA binds with high affinity to GnIH and GnIH-RPs (77). Since GnIH receptor is expressed in gonadotropes in the anterior pituitary, GnIH can act directly on gonadotropes to reduce gonadotropin release in birds [for reviews, see Ref. (19–21, 26–29, 34)] (**Figure 1**). In addition to acting on the anterior pituitary, GnIH neurons project to GnRH1 neurons (41, 78) that express GnIH receptor (41) (**Figure 1**). Thus, GnIH not only acts on gonadotropes but also acts on GnRH1 neurons to inhibit gonadotropin synthesis and release in birds [for reviews, see Ref. (19–21, 26–29, 34)] (**Figure 1**).

In mammals, Hinuma et al. (79) identified a specific receptor for mammalian GnIH, RFRP, which is identical to GPR147 and named it OT7T022 by the reverse pharmacological approach. In the human genome, there are approximately 700 GPCR genes, and the receptors whose ligands are still unknown are called orphan receptors. Hinuma et al. (79) searched for specific receptors for ligands by testing whether (1) increases in calcium ions, (2) increases in cAMP, or (3) decreases in cAMP happens in the cells transfected with the receptor by ligand stimulation [for a review, see Ref. (80)]. In the same year, Bonini et al. (81) reported two GPCRs for NPFF and designated them as NPFF1 (identical to GPR147) and NPFF2 (identical to GPR74). Bonini et al. (81) cloned these receptors by GPCR-targeted degenerate PCR using rat genomic DNA. As mentioned previously, it is thought that the GnIH (LPXRFamide peptide) and NPFF (PQRFamide peptide) genes have diverged from a common ancestral gene through gene duplication (55, 56, 76). It is also thought that GPR147 and GPR74 are paralogous from synteny analysis (82) and phylogenetic analysis (83). Analyses of binding affinities of GnIH and NPFF for GPR147 and GPR74 and their signal transduction pathways reveal that GnIH has a higher affinity for GPR147, whereas NPFF has potent agonistic activity for GPR74 (46, 81, 84), suggesting that GPR147 (NPFF1, OT7T022) is the primary receptor for GnIH.

GnIH Cell Signaling

To further understand the cellular cascade by which GnIH impacts gonadotropes in the anterior pituitary, Tsutsui and colleagues investigated GnIH signaling pathways in the mouse gonadotrope cell line, L β T2. First, the expression of GnIH receptor mRNA in L β T2 cells was shown by RT-PCR (85). Subsequently, the inhibitory effects of GnIH on GnRH-induced signaling pathways were demonstrated; mouse GnIHs effectively reduce GnRH-induced cAMP production and extracellular signal-regulated kinase (ERK) phosphorylation (85). Furthermore, mouse GnIHs reduce GnRH-induced LH β expression and LH release (85). The

stimulatory effect of GnRH on gonadotropin expression is suppressed by adenylate cyclase (AC) and protein kinase A (PKA) inhibitors but not by protein kinase C (PKC) inhibitor (85). Accordingly, mouse GnIH reduces GnRH-stimulated gonadotropin secretion by specifically interfering with GnRH actions via a AC/cAMP/PKA-dependent ERK pathway (85).

Following the discovery of GnIH, kisspeptin was discovered in mammals. Opposite to GnIH, kisspeptin has a stimulatory effect on GnRH neurons and the hypothalamic–pituitary–gonadal axis (HPG axis) in mammals (86–89). GnIH neurons project not only to GnRH1 neurons in the preoptic area (POA) but also to kisspeptin neurons in the hypothalamus and may regulate their activities [for reviews, see Ref. (19–21, 26–29, 90)] (**Figure 1**). GnIH neurons project to GnRH2 neurons and many other neurons suggesting multiple actions of GnIH [for reviews, see Ref. (19–21, 26–29)] (**Figure 1**).

REGULATION OF GnIH BIOSYNTHESIS IN THE BRAIN BY ENVIRONMENTAL AND INTERNAL FACTORS

Influence of Photoperiod Mediated by Melatonin

Investigating the regulatory mechanisms of GnIH expression in the brain has important implications for understanding the physiological role of GnIH. Photoperiodic mammals regulate reproductive activities according to the annual cycle of changes in the nocturnal secretion of melatonin (91). There is also evidence in birds that melatonin is involved in the regulation of several seasonal processes including gonadotropin secretion and gonadal activity (92–95), despite the accepted dogma that birds do not use seasonal changes in melatonin secretion to time their reproductive effort (96, 97).

To explore whether or not GnIH is part of the mechanism driving melatonin-induced seasonal changes in reproduction, Tsutsui and colleagues investigated the action of melatonin on the expression of GnIH in quail, a highly photoperiodic bird species. Initial findings demonstrated that melatonin induces GnIH expression in birds (**Figure 2**). More specifically, Ubuka et al. (98) first found that melatonin removal by pinealectomy, combined with orbital enucleation (Px plus Ex), decreases the expressions of GnIH mRNA and GnIH peptide labeling in the brain of quail (98). Ubuka et al. (98) further found that melatonin administration increases the expressions of GnIH mRNA and GnIH peptide in the brain of quail (98). Importantly, they found that Mel_{1c}, a melatonin receptor subtype, is expressed in GnIH neurons in quail (98). These findings established that melatonin acts directly on GnIH neurons to induce GnIH expression in this species (**Figure 2**). Chowdhury et al. (99) further demonstrated that melatonin not only increases GnIH expression but also increases GnIH release in quail (**Figure 2**). Interestingly, GnIH release is photoperiodically controlled in quail with diurnal changes that are negatively correlated with plasma LH concentration (99). As one would expect, GnIH release increases under short day (SD) photoperiods, when the duration of nocturnal secretion of melatonin increases (99). Together, these findings indicate that




FIGURE 2 | A schematic model of the neuroendocrine integration of environmental factors and internal factors to control GnIH expression and release. The neuroendocrine integration of environmental factors, such as photoperiod, stress, and social interaction, and internal factors, such as GnIH, melatonin, glucocorticoid, and norepinephrine (NE), is important for the control of reproduction and reproductive behaviors. GnIH inhibits the expression and release of gonadotropins and the expression of reproductive behaviors in birds and mammals. GnIH expression and release are photoperiodically modulated via a melatonin-dependent process in birds and mammals. Melatonin increases GnIH expression in quail and rats, whereas melatonin decreases GnIH expression in hamsters and sheep. Stress increases GnIH expression mediated by the actions of glucocorticoids in birds and mammals. GnIH may be a mediator of stress-induced reproductive disruption. The social environment also changes GnIH expression and release mediated by the action of NE. See the text for details.

melatonin derived from the pineal gland and eyes acts directly on GnIH neurons via Mel_{1c} to induce GnIH expression and release in birds (29, 98–101) (**Figure 2**).

In contrast to the results seen in quail, melatonin reduces GnIH expression in Syrian and Siberian hamsters, both photoperiodic mammals (48, 102, 103) (**Figure 2**). Specifically, GnIH mRNA levels and the number of GnIH immunoreactive cell bodies are reduced in sexually quiescent Syrian and Siberian hamsters exposed to SD photoperiods, compared to sexually active animals maintained under long day (LD) photoperiods. These photoperiodic changes in GnIH expression are abolished in Px hamsters, and injections of LD hamsters with melatonin reduce the expression of GnIH to SD levels (48, 102). Analogous seasonal patterns of GnIH expression have been observed in European (104) and Turkish (105) hamsters as well as the semi-desert rodent, Jerboa (106). Although these results are suggestive of a role for GnIH in seasonal breeding, they are inconsistent with a straightforward model of seasonal reproductive control by this peptide. One possibility is that hamsters require enhanced GnIH expression to suppress GnRH during the initial period of regression, whereas this level of inhibition is not necessary in hamsters with a fully regressed reproductive axis and low testosterone (T) concentrations. Another possibility emerged from our findings

in male Siberian hamsters. In this species, GnIH administration suppresses HPG axis function in LD, reproductive-competent animals but stimulates gonadotropin secretion in SD, reproductively quiescent animals (48), indicating that the impact of GnIH may depend on reproductive status or season. There are also reports showing that the expression of GnIH is regulated by melatonin and season in sheep (107, 108) and rats (109). Thus, as in quail, the expression of GnIH is photoperiodically modulated via a melatonin-dependent process in mammals (**Figure 2**).

We still do not have data as to why melatonin stimulates GnIH expression in quail (98) and inhibits GnIH expression in hamsters (48, 102) and other mammals. Although expression of melatonin receptor in GnIH neurons is still not demonstrated in mammals, one possibility is that melatonin triggers different intracellular signals in GnIH neurons in quail because mammals do not have Mel_{1c} melatonin receptor subtype (110). Another possibility is that melatonin indirectly regulates GnIH expression in mammals unlike quail. These possibilities should be tested in future studies.

Influence of Stress Mediated by Glucocorticoids

It is well known that stress can reduce reproduction across vertebrates (111). To explore whether stress changes GnIH expression in birds, Calisi et al. (112) investigated the effect of capture-handling stress on GnIH expression in male and female adult house sparrows. Calisi et al. (112) found that more GnIH-positive neurons are observed in fall birds versus those sampled in the spring, and GnIH-positive neurons are increased by capture-handling stress in spring birds. These observations indicate that stress influences GnIH during the breeding season (112). These findings suggested that stress may act through GnIH neurons to inhibit reproductive function in birds.

In mammals, Kirby et al. (113) showed that both acute and chronic immobilization stress lead to an upregulation of the expression of GnIH in the dorsomedial hypothalamic area (DMH) of male rats associated with inhibition of downstream hypothalamic–pituitary–testicular activity (**Figure 2**). Adrenalectomy blocks this stress-induced increase in GnIH expression. Immunohistochemistry revealed that GnIH neurons express glucocorticoid receptor (GR), suggesting that adrenal glucocorticoids act directly on GnIH neurons to increase GnIH expression (**Figure 2**). Together, these observations indicate that GnIH is an important integrator of stress-induced suppression of reproductive function in mammals (113).

Recently, Son et al. (114) demonstrated that GR mRNA is expressed in GnIH neurons in the PVN of quail, suggesting that glucocorticoids can directly regulate GnIH transcription (**Figure 2**). It was also found that treatment with corticosterone (CORT) increases GnIH mRNA expression in the quail diencephalon (114) (**Figure 2**). Subsequently, Son et al. (114) investigated the mechanism of activation of GnIH transcription by CORT using a GnIH-expressing neuronal cell line, rHypoE-23, derived from rat hypothalamus. Importantly, GR mRNA is expressed in rHypoE-23 cells, and CORT treatment increases GnIH mRNA expression (114). Son et al. (114) further found that CORT stimulates GR recruitment to the GC-response element (GRE)

present in the rat GnIH promoter region, providing further support that CORT induces GnIH expression via GR in GnIH neurons (Figure 2). Taken together, it appears that stress reduces gonadotropin release, at least in part, through an increase in GnIH expression. More recent evidence also indicates that GnIH might itself regulate the stress response in mice (115).

Influence of Social Interactions Mediated by Norepinephrine

In addition to the regulation of GnIH expression by environmental factors, photoperiod, and stress, the social environment may influence the GnIH system (Figure 2). To examine this possibility, Calisi et al. (116) investigated the impact of mating competition on GnIH. Nesting opportunities for pairs of European starlings were manipulated, and GnIH mRNA and GnIH content as well as GnRH content in the brain were examined. Birds with nest boxes have fewer numbers of GnIH-producing cells than those without nest boxes. However, GnRH content does not vary with nest box ownership. These observations suggest that GnIH may serve as a modulator of reproductive function in response to social environment (116).

Reproductive physiology and behavior are variable, both within and between individuals. It is known that the presence of a female bird as well as copulation rapidly decrease plasma T concentrations in male quail (117, 118). Based on these earlier observations, Tsutsui and colleagues investigated the neurochemical mechanism by which social stimuli alter reproductive physiology and behavior (Figure 2). Tobari et al. (31) first found that norepinephrine (NE) release increases rapidly in the PVN of quail when viewing a female conspecific (Figure 2). Likewise, GnIH mRNA expression increases in the PVN, with associated decreases in LH concentrations in plasma, when males view a female (Figure 2). Tobari et al. (31) then established a link between these two events by showing that NE application stimulates GnIH release from diencephalic tissue blocks *in vitro*. Double-label immunohistochemistry revealed that GnIH neurons are innervated by noradrenergic fibers and immunohistochemistry combined with *in situ* hybridization demonstrated that GnIH neurons expressed α 2A-adrenergic receptor mRNA. Together, these observations indicate that female presence increases NE release in the PVN and stimulates GnIH release, resulting in the suppression of LH release in quail (31) (Figure 2).

MULTIPLE ACTIONS OF GnIH

Direct Regulation of Gonadal Activity

The aforementioned findings indicate that GnIH is a key player in the regulation of reproduction across vertebrates, reducing gonadotropin synthesis and release by decreasing the activity of GnRH1 neurons and decreasing the activity of pituitary gonadotropes, inevitably suppressing gonadal steroid secretion and spermatogenesis (Figure 1). In addition to these central actions of GnIH, several lines of evidence point to direct, local regulation of gonadal activity [for reviews, see Ref. (19, 20, 26–29, 32, 34)] (Figure 1). GnIH and GnIH receptor are expressed in steroidogenic cells and germ cells in the gonads of birds and mammals

(119–125), with GnIH possibly acting in an autocrine/paracrine manner to suppress gonadal steroid production and germ cell differentiation and maturation (119–125) (Figure 1). There is also evidence in songbirds that gonadal GnIH responds directly to melatonin, metabolic challenge, and cues of stress in a seasonal manner (126–128)

Regulation of Feeding Behavior

It is becoming clear that GnIH participates not only in neuroendocrine functions but also in behavioral control. In environments where energy availability fluctuates, animals use photoperiod to phase breeding with anticipated times of maximal food availability (129). Should food become scarce during the breeding season, reproduction is temporarily inhibited (130, 131). Food deprivation and other metabolic challenges inhibit reproductive axis functioning and sexual motivation (132–136). GnIH may relay metabolic information to the HPG axis and regulate neural feeding circuits [for a review, see Ref. (19)].

Tachibana et al. (137) showed that intracerebroventricular (ICV) injections of GnIH, GnIH-RP-1, and GnIH-RP-2 stimulate food intake in chicks (137). In further support of a stimulatory role for GnIH in feeding, anti-GnIH antiserum suppresses appetite induced by fasting but does not modify feeding under *ad libitum* conditions (137). Similarly, Fraley et al. (138) reported that ICV injection of GnIH, but not of GnIH-RP1, suppresses plasma LH and stimulates feeding in adult Pekin ducks. To establish the neurochemical cascade underlying the actions of GnIH on feeding, Tachibana et al. (139) explored the possibility that the orexigenic effect of GnIH occurs via actions on the opioid and nitric oxide (NO) systems. It was found that the orexigenic effect of ICV injected GnIH is attenuated by coinjection of β -funaltrexamine (an opioid μ -receptor antagonist) but not ICI-174,864 (an opioid δ -receptor antagonist) and nor-binaltorphimine (an opioid κ -receptor antagonist) in chicks. It was also found that coinjection of a non-selective NO synthase inhibitor does not affect GnIH-induced feeding behavior (139). More recently, McConn et al. (39) investigated the central mechanism of the GnIH orexigenic response in chicks. It was found that neuropeptide Y (NPY) mRNA is increased, while pro-opiomelanocortin (POMC) mRNA is decreased in the hypothalamus following ICV injection of chicken GnIH. Additionally, ICV GnIH injections increase c-fos immunoreactive cells in the lateral hypothalamic area (LHA). McConn et al. (39) further found that in isolated LHA, melanin-concentrating hormone (MCH) mRNA is increased by ICV administration of GnIH. Together, these observations suggest that opioid μ -receptor-positive, NPY, POMC, and MCH neurons are likely involved in the GnIH orexigenic response.

In mammals, there are several reports indicating that ICV administration of GnIH increases food intake in rats (59) and sheep (140). Fu and van den Pol (141) showed that chicken GnIH and human GnIH inhibit POMC neurons and attenuate kisspeptin cell excitation by a mechanism based on opening potassium channels in mouse brain slices. Jacobi et al. (142) reported that GnIH inhibits the firing rate in POMC neurons and has a predominantly inhibitory effect on action potential activity in NPY neurons in mice. Jacobi et al. (142) also reported that NPY neurons have close contacts from GnIH fibers.

Together, these observations indicate that GnIH participates not only in reproduction but also in feeding behavior in birds and mammals.

Regulation of Reproductive Behaviors

Gonadotropin-inhibitory hormone also acts on the brain to regulate reproductive behaviors, such as sexual and aggressive behaviors (32, 143, 144) (Figure 1). For example, Bentley et al. (143) showed that a centrally administered physiological dose of GnIH inhibits copulation solicitation in estrogen-primed female white-crowned sparrows exposed to the song of males. It is known that GnRH2 enhances copulation solicitation in estrogen-primed female white-crowned sparrows exposed to the song of males (145). Because GnIH neurons terminate in close proximity of GnRH2 neurons and GnRH2 neurons express GnIH receptor in songbirds (41), GnIH may inhibit copulation solicitation by inhibiting GnRH2 neurons in female songbirds (143) (Figure 1). Ubuka et al. (144) directly investigated this possibility by applying RNA interference (RNAi) to the GnIH gene and examining the behavior of male and female white-crowned sparrows in collaboration with the Wingfield laboratory. GnIH RNAi reduces resting time, spontaneous production of complex vocalizations, and stimulates agonistic vocalizations. Additionally, GnIH RNAi enhances song production of short duration in male birds when they are challenged by playbacks of novel male songs. These observations indicate that GnIH gene silencing induces arousal. Ubuka et al. (144) further showed that the activities of male and female birds are negatively correlated with GnIH mRNA expression in the PVN. The density of GnIH neuronal fibers in the ventral tegmental area is decreased by GnIH RNAi in female birds, and the number of GnRH1 and GnRH2 neurons that receive close appositions of GnIH neuronal fiber terminals is negatively correlated with the activity of male birds (144) (Figure 1). Recently, Ubuka et al. (32) have demonstrated that GnIH also inhibits aggressive behavior in male quail. Thus, it is considered that GnIH decreases sexual and aggressive behaviors in birds [for reviews, see Ref. (19, 29, 34)] (Figure 1).

Similar results have been observed in mammals. For example, Johnson et al. (59) reported that ICV administration of GnIH suppresses male sex behavior in rats. Piekarski et al. (146) found that ICV administration of GnIH reduces sexual motivation and vaginal scent marking but not lordosis behavior in female hamsters. According to Piekarski et al. (146), GnIH administration alters fos expression in key neural loci implicated in female sexual behavior, including the medial POA, medial amygdala, and bed nucleus of the stria terminalis. These observations suggest that GnIH is an important modulator of female proceptive sexual behavior and motivation (Figure 1). Thus, as in birds, GnIH is not only in a position to regulate the HPG axis but may also act as a neuromodulator to drive the neural circuitry underlying socially motivated behavior.

Regulation of Neurosteroidogenesis

It is becoming increasingly clear that the interaction of neuropeptides and neurosteroids plays a role in the regulations of some brain functions [for a review, see Ref. (147)]. Recently, Ubuka

et al. (32) found that GnIH activates cytochrome P450 aromatase (P450arom) and increases neuroestrogen synthesis in the avian brain (32) (Figures 1 and 3). Importantly, the actions of GnIH on neuroestrogen synthesis change the expression of aggressive behavior in birds (32) (Figures 1 and 3), providing a new concept that GnIH modifies the neurosteroid milieu in the brain to modulate aggressive behavior.

Further exploring the ability of GnIH to alter neurosteroid production has led to important insight into the control of aggressive behavior in quail. Sexually mature male quail frequently fight with intense aggressiveness, by using stereotyped actions (148,




FIGURE 3 | Model of the intracellular mechanism of GnIH and its receptor (GPR147) that control male aggressive behavior by regulating the activity of aromatase and neuroestrogen synthesis in the brain. GPR147 is expressed in aromatase immunoreactive cells in the brain. GPR147 is coupled to G α_i protein that inhibits the activity of adenylate cyclase (AC) and decreases cAMP production and the activity of protein kinase A (PKA). Inhibition of AC/cAMP/PKA pathway may decrease phosphorylated aromatase and increase dephosphorylated aromatase. 17 β -Estradiol (E2) synthesized from testosterone (T) by aromatase in the brain especially in the preoptic area (POA) regulates male aggression. The administration of GnIH activates aromatase by decreasing phosphorylated aromatase and stimulates neuroestrogen synthesis in the brain. GnIH may inhibit aggressive behavior by directly activating aromatase and increasing neuroestrogen synthesis in the brain beyond its optimum concentration for the expression of aggressive behavior. Partially adapted from Ubuka and Tsutsui (33).

149). Aggressive behavior of male quail is thought to be androgen dependent (148–150), but there is generally no correlation between aggressiveness and peripheral T concentration (150). It is also known that aggression in males is activated by aromatizable androgens, such as T and androstenedione (AD), but not by non-aromatizable androgens, such as dihydrotestosterone (DHT), and that administration of P450arom inhibitors blocks T-induced aggression (150, 151). Thus, the action of gonadal androgen on aggressive behavior requires its aromatization into estrogen (neuroestrogen) in the brain (152–154).

GnIH neurons project to the ME and other brain areas, such as the POA (18, 155, 156) and the periaqueductal central gray [PAG (41)] in birds. GnIH receptor is also expressed in the POA (41, 77) and PAG (41). These brain areas are known to regulate aggressive behavior (157, 158). The POA is also known to be the most critical site of aromatization of gonadal androgen by P450arom and neuroestrogen action for the activation of aggressive behavior in male quail (159, 160). Because GnIH decreases aggressive behavior in male birds (32, 144), GnIH may decrease this behavior by regulating P450arom activity and neuroestrogen synthesis in the brain (Figures 1 and 3). Ubuka et al. (32) therefore investigated whether GnIH neuronal fibers innervate P450arom cells and whether P450arom cells express GnIH receptor in the POA of male quail. It was found that abundant GnIH immunoreactive neuronal fibers are distributed in the vicinity of P450arom immunoreactive cells in the POA (32). It was also found that GnIH receptor is expressed in P450arom immunoreactive cells in the POA (32). Furthermore, GnIH stimulates P450arom activity and increases neuroestrogen synthesis in the POA through GnIH receptor (32) (Figures 1 and 3). These studies provided the first evidence that a hypothalamic neuropeptide can regulate neuroestrogen synthesis in the brain.

Importantly, the increase in neuroestrogen concentrations in the POA was associated with a decrease in aggressive behavior (32). Therefore, the effect of central administration of various doses of estradiol-17 β (E2) on aggressive behavior was tested in male quail. Ubuka et al. (32) found that central administration of E2 at higher doses decreases aggressive behavior unlike E2 at lower doses (32). This observation indicates that the action of neuroestrogen is essential for the expression of aggressive behavior, but higher concentrations of neuroestrogen in the brain decrease this behavior. Taken together, GnIH decreases aggressive behavior by activating P450arom and increasing neuroestrogen synthesis in the brain beyond its optimum concentration for the expression of aggressive behavior of male birds (32) (Figures 1 and 3).

Ubuka et al. (32) further investigated the mode of action of GnIH on the stimulation of P450arom activity. There is important evidence that P450arom activity is not only controlled in the long term by transcription of the P450arom gene *Cyp19* by steroids, but also in the short term by phosphorylation by neurotransmitters, such as glutamate (161). Balthazart's group demonstrated that P450arom activity in the hypothalamus of male quail is rapidly downregulated by phosphorylation (161–165). Therefore, GnIH may activate P450arom by dephosphorylation of phosphorylated P450arom. Ubuka et al. (32) found that ICV administration of GnIH reduces phosphorylated P450arom in the POA in the short term compared with control birds (32). Ubuka et al. (32)

also found that the action of GnIH on neuroestrogen synthesis in the POA is abolished by concomitant administration of RF9 (166, 167), a potent antagonist of GnIH receptor, or fadrozole (168, 169), an inhibitor of P450arom. Based on these findings, it is apparent that GnIH stimulates neuroestrogen synthesis in the POA by activating P450arom through dephosphorylation after binding to GnIH receptor in P450arom cells (Figure 3).

CONCLUSION AND FUTURE DIRECTIONS

The discovery of GnIH in 2000 and the contributions aimed at understanding its evolutionary history and functions have markedly advanced the progress of reproductive neuroendocrinology. Studies on GnIH over the past decade and a half have demonstrated that GnIH is a key player in the regulation of reproduction across vertebrates. It now appears that GnIH acts on the pituitary and the brain to modulate the reproductive axis and reproductive behaviors (Figure 1). In this review, the commonalities and diversity of GnIH structures and actions and the evolutionary origin of GnIH in chordates were also highlighted. The discovery of GnIH has changed our understanding of the regulation of reproductive physiology and behavior. As a result, more than 50 laboratories worldwide are now working on GnIH.

Following the discovery of GnIH, kisspeptin was also discovered in mammals. GnIH and kisspeptin are both comparatively new members of the RFamide peptide family and act on the HPG axis to downregulate and upregulate the reproductive system, respectively. Thus, we now know that GnRH is not the only hypothalamic neuropeptide regulating reproduction. Importantly, GnIH neurons project not only to GnRH neurons but also to kisspeptin neurons in the hypothalamus (Figure 1). GnRH neurons and kisspeptin neurons express GnIH receptor. We expect that future studies will reveal previously unknown interactions among GnIH, GnRH, and kisspeptin (Figure 1). The fact that GnIH neurons project to many other neurons in the brain suggests multiple actions of GnIH that have not yet been discovered (Figure 1).

GnIH activates P450arom activity in the brain (Figure 1) and may change the formation of other neurosteroids by activating or inactivating their steroidogenic enzymes. Furthermore, steroidogenic enzymes are expressed not only in the brain but also in the pineal gland (170–173). Future studies are needed to further develop the emerging concept that hypothalamic neuropeptides may modify the neurosteroid milieu in the brain and pineal gland to impact their function.

ACKNOWLEDGMENTS

This review is dedicated to Tsutsui's beloved wife, Rieko Tsutsui. The works described in this review were supported at least partially by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture, Japan (15207007, 16086206, 18107002, 22132004, and 22227002 to KT). I am grateful to the following collaborators, J. C. Wingfield, P. J. Sharp, I. J. Clarke, R. P. Millar, S. A. Sower, S. Haraguchi, Y. Muneoka, K. Ukena, T. Osugi, V. S. Chowdhury, J. F. Cockrem, E. Saigoh, Y. Tobar, H. Yin, K. Inoue, H. Teranishi, Y. Fukuda, T. Mizuno, M. Narihito, K.

Ishikawa, S. Ishii, O. Koizumi, M. Ueno, H. Minakata, H. Satake, E. Iwakoshi, D. Daukss, K. Gazda, T. Kosugi, G. Bedecarrats, M. Hisada, T. Kawada, N. L. McGuire, R. Calisi, N. Perfito, S. O'Brien, I. T. Moore, J. P. Jensen, G. J. Kaur, D. W. Wacker, N. A. Ciccone, I. C. Dunn, T. Boswell, S. Kim, Y. C. Huang, J. Reid, J. Jiang, P. Deviche, T. W. Small, M. A. Ottinger, T. Tachibana, M. Furuse, M. A. Cline, D. F. Mei, A. Mason, E. M. Gibson, S. A. Humber, S. Jain, W. P. Williams III, S. Zhao, I. P. Sari, Y. Qi, J. T. Smith, H. C. Parkington, J. Iqbal, Q. Li, A. Tilbrook, K. Morgan, A. J. Pawson, M. Murakami, T. Matsuzaki, T. Iwasa, T. Yasui, M. Irahara, M. A. Johnson, G. S. Fraley, H. Oishi, C. Klausen, C. B. Gilks, T. Yano, P. C. K. Leung, M. Binns, P. A. Cadigan, H. Lai, M. Kitani, A. Suzuuchi, V. Pham, S. Kikuyama, K. Yamamoto, A. Koda, I. Hasumuma, F. Toyoda, K. Sawada, K. Tsunekawa, P. Singh, S. Anjum, A. Krishna, R. Sridaran, S. Sethi, C. M. Chaturvedi, M. R. Jafarzadeh Shirazi,

M. J. Zamiri, A. Tamadon, M. Amano, S. Moriyama, M. Iigo, K. Uchida, M. Nozaki, H. Kawauchi, M. Shahjahan, T. Ikegami, H. Doi, A. Hattori, H. Ando, S. Honda, N. Harada, M. Hatori, T. Hirota, M. Iitsuka, N. Kurabayashi, K. Kokame, R. Sato, A. Nakai, T. Miyata, Y. Fukada, M. Sato, K. Tashiro, T. Tokita, J. Y. Seong, J. L. Do-Rego, J. Leprince, G. Pelletier, and H. Vaudry.

FUNDING

Grants-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture, Japan (15207007, 16086206, 18107002, 22132004, and 22227002 to KT). NSF IOS 1122044 (to GEB), NIH grant HD-050470 and NSF grant IOS-1257638 (to LJK), Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan (23570091 to TU).

REFERENCES

- Livermore AH, Du Vigneaud V. Preparation of high potency oxytocic material by the use of counter-current distribution. *J Biol Chem* (1949) **180**:365–73.
- Turner RA, Pierce JG, Du Vigneaud V. The purification and the amino acid content of vasopressin preparations. *J Biol Chem* (1951) **191**:21–8.
- Harris GW. Neuronal control of the pituitary gland. *Physiol Rev* (1948) **28**:139–79.
- Harris GW. The function of the pituitary stalk. *Bull Johns Hopkins Hosp* (1955) **97**:358–75.
- Watts AG. 60 years of neuroendocrinology: the structure of the neuroendocrine hypothalamus: the neuroanatomical legacy of Geoffrey Harris. *J Endocrinol* (2015) **226**:T25–39. doi:10.1530/JOE-15-0157
- Halasz B, Pupp L, Uhlarik S. Hypophysiotropic area in the hypothalamus. *J Endocrinol* (1962) **25**:147–54. doi:10.1677/joe.0.0250147
- McCann SM, Ramirez VD. The neuroendocrine regulation of hypophysial luteinizing hormone secretion. *Recent Prog Horm Res* (1964) **20**:131–81.
- Burgus R, Dunn TF, Desiderio D, Guillemin R. Molecular structure of the hypothalamic hypophysiotropic TRF factor of ovine origin: mass spectrometry demonstration of the PCA-His-Pro-NH₂ sequence. *C R Acad Sci (Paris)* (1969) **269**:1870–3.
- Boler J, Enzmann F, Folkers K, Bowers CY, Schally AV. The identity of chemical and hormonal properties of the thyrotropin releasing hormone and pyroglutamyl-histidyl-proline amide. *Biochem Biophys Res Commun* (1969) **37**:705–10. doi:10.1016/0006-291X(69)90868-7
- Matsuo H, Baba Y, Nair RM, Arimura A, Schally AV. Structure of the porcine LH- and FSH-releasing hormone. I. The proposed amino acid sequence. *Biochem Biophys Res Commun* (1971) **43**:1334–9. doi:10.1016/S0006-291X(71)80019-0
- Burgus R, Butcher M, Amoss M, Ling N, Monahan M, Rivier J, et al. Primary structure of the ovine hypothalamic luteinizing hormone-releasing factor (LRF) (LH-hypothalamus-LRF-gas chromatography-mass spectrometry-decapeptide-Edman degradation). *Proc Natl Acad Sci U S A* (1972) **69**:278–82. doi:10.1073/pnas.69.1.278
- Brazeau P, Vale W, Burgus R, Ling N, Butcher M, Rivier J, et al. Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science* (1973) **179**:77–9. doi:10.1126/science.179.4068.77
- King JA, Millar RP. Structure of chicken hypothalamic luteinizing hormone-releasing hormone. I. Structural determination on partially purified material. *J Biol Chem* (1982) **257**:10722–8.
- Miyamoto K, Hasegawa Y, Minegishi T, Nomura M, Takahashi Y, Igarashi M, et al. Isolation and characterization of chicken hypothalamic luteinizing hormone-releasing hormone. *Biochem Biophys Res Commun* (1982) **107**:820–7. doi:10.1016/0006-291X(82)90596-4
- Miyamoto K, Hasegawa Y, Nomura M, Igarashi M, Kangawa K, Matsuo H. Identification of the second gonadotropin-releasing hormone in chicken hypothalamus: evidence that gonadotropin secretion is probably controlled by two distinct gonadotropin-releasing hormones in avian species. *Proc Natl Acad Sci U S A* (1984) **81**:3874–8. doi:10.1073/pnas.81.12.3874
- Sherwood NM, Sower SA, Marshak DR, Fraser BA, Brownstein MJ. Primary structure of gonadotropin-releasing hormone in lamprey brain. *J Biol Chem* (1986) **15**:4812–9.
- McCann SM, Mizunuma H, Samson WK, Lumpkin MD. Differential hypothalamic control of FSH secretion: a review. *Psychoneuroendocrinology* (1983) **8**:299–308. doi:10.1016/0306-4530(83)90004-5
- Tsutsui K, Saigoh E, Ukena K, Teranishi H, Fujisawa Y, Kikuchi M, et al. A novel avian hypothalamic peptide inhibiting gonadotropin release. *Biochem Biophys Res Commun* (2000) **275**:661–7. doi:10.1006/bbrc.2000.3350
- Kriegsfeld LJ, Ubuka T, Bentley GE, Tsutsui K. Seasonal control of gonadotropin-inhibitory hormone (GnIH) in birds and mammals. *Front Neuroendocrinol* (2015) **37**:65–75. doi:10.1016/j.yfrne.2014.12.001
- Tsutsui K. Review: a new key neurohormone controlling reproduction, gonadotropin-inhibitory hormone (GnIH): Biosynthesis, mode of action and functional significance. *Prog Neurobiol* (2009) **88**:76–88. doi:10.1016/j.pneurobio.2009.02.003
- Tsutsui K. Review: how to contribute to the progress of neuroendocrinology: new insights from discovering novel neuropeptides and neurosteroids regulating pituitary and brain functions. *Gen Comp Endocrinol* (2015). doi:10.1016/j.ygcen.2015.05.019
- Tsutsui K, Ubuka T. Gonadotropin-inhibitory hormone. In: Kastin AJ, Vaudry H, editors. *Handbook of Biologically Active Peptides*. London: Academic Press (2012). p. 802–11.
- Tsutsui K, Ukena K. Review: hypothalamic LPXRF-amide peptides in vertebrates: identification, localization and hypophysiotropic activity. *Peptides* (2006) **27**:1121–9. doi:10.1016/j.peptides.2005.06.036
- Tsutsui K, Ubuka T, Yin H, Osugi T, Ukena K, Bentley GE, et al. Review: mode of action and functional significance of avian gonadotropin-inhibitory hormone (GnIH). *J Exp Zool* (2006) **305A**:801–6. doi:10.1002/jez.a.305
- Tsutsui K, Bentley GE, Ubuka T, Saigoh E, Yin H, Osugi T, et al. Review: the general and comparative biology of gonadotropin-inhibitory hormone (GnIH). *Gen Comp Endocrinol* (2007) **153**:365–70. doi:10.1016/j.ygcen.2006.10.005
- Tsutsui K, Bentley GE, Bedecarrats GT, Osugi T, Ubuka T, Kriegsfeld LJ. Review: gonadotropin-inhibitory hormone (GnIH) and its control of central and peripheral reproductive function. *Front Neuroendocrinol* (2010) **31**:284–95. doi:10.1016/j.yfrne.2010.03.001
- Tsutsui K, Bentley GE, Kriegsfeld LJ, Osugi T, Seong JY, Vaudry H. Review: discovery and evolutionary history of gonadotropin-inhibitory hormone and kisspeptin: new key neuropeptides controlling reproduction. *J Neuroendocrinol* (2010) **22**:716–27. doi:10.1111/j.1365-2826.2010.02018.x
- Tsutsui K, Ubuka T, Bentley GE, Kriegsfeld LJ. Review: gonadotropin-inhibitory hormone (GnIH): discovery, progress and prospect. *Gen Comp Endocrinol* (2012) **177**:305–14. doi:10.1016/j.ygcen.2012.02.013
- Tsutsui K, Ubuka T, Bentley GE, Kriegsfeld LJ. Review: regulatory mechanisms of gonadotropin-inhibitory hormone (GnIH) synthesis and release in photoperiodic animals. *Front Neurosci* (2013) **7**:60. doi:10.3389/fnins.2013.00060

30. Ukena K, Tsutsui K. A new member of the hypothalamic RF-amide peptide family, LPXRF-amide peptides: structure, localization, and function. *Mass Spectrom Rev* (2005) **24**:469–86. doi:10.1002/mas.20031
31. Tobari Y, Son YL, Ubuka T, Hasegawa Y, Tsutsui K. A new pathway mediating social effects on the endocrine system: female presence acting via norepinephrine release stimulates gonadotropin-inhibitory hormone in the paraventricular nucleus and suppresses luteinizing hormone in quail. *J Neurosci* (2014) **34**:9803–11. doi:10.1523/JNEUROSCI.3706-13.2014
32. Ubuka T, Haraguchi S, Tobari Y, Narihiro M, Ishikawa K, Hayashi T, et al. Hypothalamic inhibition of socio-sexual behaviors by increasing neuroestrogen synthesis. *Nat Commun* (2014) **5**:3061. doi:10.1038/ncomms4061
33. Ubuka T, Tsutsui K. Review: neuroestrogen regulation of socio-sexual behavior of males. *Front Neurosci* (2014) **8**:323. doi:10.3389/fnins.2014.00323
34. Ubuka T, Son YL, Bentley GE, Millar RP, Tsutsui K. Gonadotropin-inhibitory hormone (GnIH), GnIH receptor and cell signaling. *Gen Comp Endocrinol* (2013) **190**:10–7. doi:10.1016/j.ygcen.2013.02.030
35. Price DA, Greenberg MJ. Structure of a molluscan cardioexcitatory neuro-peptide. *Science* (1977) **197**:670–1. doi:10.1126/science.877582
36. Raffia RB. The action of FMRFamide (Phe-Met-Arg-Phe-NH₂) and related peptides on mammals. *Peptides* (1988) **9**:915–22. doi:10.1016/0196-9781(88)90141-6
37. Rastogi RK, D'Aniello B, Pinelli C, Fiorentino M, Di Fiore MM, Di Meglio M, et al. FMRFamide in the amphibian brain: a comprehensive survey. *Microsc Res Tech* (2001) **54**:158–72. doi:10.1002/jemt.1130
38. Dockray GJ, Reeve JR Jr, Shively J, Gayton RJ, Barnard CS. A novel active pentapeptide from chicken brain identified by antibodies to FMRFamide. *Nature* (1983) **305**:328–30. doi:10.1038/305328a0
39. McConn B, Wang G, Yi J, Gilbert ER, Osugi T, Ubuka T, et al. Gonadotropin-inhibitory hormone-stimulation of food intake is mediated by hypothalamic effects in chicks. *Neuropeptides* (2014) **48**:327–34. doi:10.1016/j.npep.2014.09.001
40. Satake H, Hisada M, Kawada T, Minakata H, Ukena K, Tsutsui K. Characterization of a cDNA encoding a novel avian hypothalamic neuropeptide exerting an inhibitory effect on gonadotropin release. *Biochem J* (2001) **354**:379–85. doi:10.1042/bj3540379
41. Ubuka T, Kim S, Huang YC, Reid J, Jiang J, Osugi T, et al. Gonadotropin-inhibitory hormone neurons interact directly with gonadotropin-releasing hormone-I and -II neurons in European starling brain. *Endocrinology* (2008) **149**:268–78. doi:10.1210/en.2007-0983
42. Tobari Y, Iijima N, Tsunekawa K, Osugi T, Okanoya K, Tsutsui K, et al. Identification of gonadotropin-inhibitory hormone in the zebra finch (*Taeniopygia guttata*): peptide isolation, cDNA cloning and brain distribution. *Peptides* (2010) **31**:816–26. doi:10.1016/j.peptides.2010.01.015
43. Ubuka T, Morgan K, Pawson AJ, Osugi T, Chowdhury VS, Minakata H, et al. Identification of human GnIH homologs, RFRP-1 and RFRP-3, and the cognate receptor, GPR147 in the human hypothalamic pituitary axis. *PLoS One* (2009) **4**:e8400. doi:10.1371/journal.pone.0008400
44. Ubuka T, Lai H, Kitani M, Suzuuchi A, Pham V, Cadigan PA, et al. Gonadotropin-inhibitory hormone identification, cDNA cloning, and distribution in rhesus macaque brain. *J Comp Neurol* (2009) **517**:841–55. doi:10.1002/cne.22191
45. Fukusumi S, Habata Y, Yoshida H, Iijima N, Kawamata Y, Hosoya M, et al. Characteristics and distribution of endogenous RFamide-related peptide-1. *Biochim Biophys Acta* (2001) **1540**:221–32. doi:10.1016/S0167-4889(01)00135-5
46. Yoshida H, Habata Y, Hosoya M, Kawamata Y, Kitada C, Hinuma S. Molecular properties of endogenous RFamide-related peptide-3 and its interaction with receptors. *Biochim Biophys Acta* (2003) **1593**:151–7. doi:10.1016/S0167-4889(02)00389-0
47. Ukena K, Iwakoshi E, Minakata H, Tsutsui K. A novel rat hypothalamic RFamide-related peptide identified by immunoaffinity chromatography and mass spectrometry. *FEBS Lett* (2002) **512**:255–8. doi:10.1016/S0014-5793(02)02275-5
48. Ubuka T, Inoue K, Fukuda Y, Mizuno T, Ukena K, Kriegsfeld LJ, et al. Identification, expression, and physiological functions of Siberian hamster gonadotropin-inhibitory hormone. *Endocrinology* (2012) **153**:373–85. doi:10.1210/en.2011-1110
49. Ukena K, Iwakoshi-Ukena E, Osugi T, Tsutsui K. Identification and localization of gonadotropin-inhibitory hormone (GnIH) orthologs in the hypothalamus of the red-eared slider turtle, *Trachemys scripta elegans*. *Gen Comp Endocrinol* (2015). doi:10.1016/j.ygcen.2015.06.009
50. Koda A, Ukena K, Teranishi H, Ohta S, Yamamoto K, Kikuyama S, et al. A novel amphibian hypothalamic neuropeptide: isolation, localization, and biological activity. *Endocrinology* (2002) **143**:411–9. doi:10.1210/endo.143.2.8630
51. Chartrel N, Dujardin C, Leprince J, Desrués L, Tonon MC, Cellier E, et al. Isolation, characterization, and distribution of a novel neuropeptide, Rana RFamide (R-RFa), in the brain of the European green frog *Rana esculenta*. *J Comp Neurol* (2002) **448**:111–27. doi:10.1002/cne.10253
52. Ukena K, Koda A, Yamamoto K, Kobayashi T, Iwakoshi-Ukena E, Minakata H, et al. Novel neuropeptides related to frog growth hormone-releasing peptide: isolation, sequence, and functional analysis. *Endocrinology* (2003) **144**:3879–84. doi:10.1210/en.2003-0359
53. Chowdhury VS, Ubuka T, Osugi T, Shimura T, Tsutsui K. Identification, localization and expression of LPXRFamide peptides, and melatonin-dependent induction of their precursor mRNA in the newt brain. *J Endocrinol* (2011) **209**:211–20. doi:10.1530/JOE-10-0494
54. Sawada K, Ukena K, Satake H, Iwakoshi E, Minakata H, Tsutsui K. Novel fish hypothalamic neuropeptide. *Eur J Biochem* (2002) **269**:6000–8. doi:10.1046/j.1432-1033.2002.03351.x
55. Osugi T, Dauks D, Gazda K, Ubuka T, Kosugi T, Nozaki M, et al. Evolutionary origin of the structure and function of gonadotropin-inhibitory hormone: insights from lampreys. *Endocrinology* (2012) **153**:2362–74. doi:10.1210/en.2011-2046
56. Osugi T, Okamura T, Son YL, Ohkubo M, Ubuka T, Henmi Y, et al. Evolutionary origin of GnIH and NPDF in chordates: insights from novel amphioxus RFamide peptides. *PLoS One* (2014) **9**:e100962. doi:10.1371/journal.pone.0100962
57. Ubuka T, Ukena K, Sharp PJ, Bentley GE, Tsutsui K. Gonadotropin-inhibitory hormone inhibits gonadal development and maintenance by decreasing gonadotropin synthesis and release in male quail. *Endocrinology* (2006) **147**:1187–94. doi:10.1210/en.2005-1178
58. Kriegsfeld LJ, Mei DF, Bentley GE, Ubuka T, Mason AO, Inoue K, et al. Identification and characterization of a gonadotropin-inhibitory system in the brains of mammals. *Proc Natl Acad Sci U S A* (2006) **103**:2410–5. doi:10.1073/pnas.0511003103
59. Johnson MA, Tsutsui K, Fraley GS. Rat RFamide-related peptide-3 stimulates GH secretion, inhibits LH secretion, and has variable effects on sex behavior in the adult male rat. *Horm Behav* (2007) **51**:171–80. doi:10.1016/j.yhbeh.2006.09.009
60. Murakami M, Matsuzaki T, Iwasa T, Yasui T, Irahara M, Osugi T, et al. Hypophysiotropic role of RFamide-related peptide-3 in the inhibition of LH secretion in female rats. *J Endocrinol* (2008) **199**:105–12. doi:10.1677/JOE-08-0197
61. Rizwan MZ, Porteous R, Herbison AE, Anderson GM. Cells expressing RFamide-related peptide-1/3, the mammalian gonadotropin-inhibitory hormone orthologs, are not hypophysiotropic neuroendocrine neurons in the rat. *Endocrinology* (2009) **150**:1413–20. doi:10.1210/en.2008-1287
62. Clarke IJ, Sari IP, Qi Y, Smith JT, Parkington HC, Ubuka T, et al. Potent action of RFamide-related peptide-3 on pituitary gonadotropes indicative of a hypophysiotropic role in the negative regulation of gonadotropin secretion. *Endocrinology* (2008) **149**:5811–21. doi:10.1210/en.2008-0575
63. Sari IP, Rao A, Smith JT, Tilbrook AJ, Clarke IJ. Effect of RF-amide-related peptide-3 on luteinizing hormone and follicle-stimulating hormone synthesis and secretion in ovine pituitary gonadotropes. *Endocrinology* (2009) **150**:5549–56. doi:10.1210/en.2009-0775
64. Kadokawa H, Shibata M, Tanaka Y, Kojima T, Matsumoto K, Oshima K, et al. Bovine C-terminal octapeptide of RFamide-related peptide-3 suppresses luteinizing hormone (LH) secretion from the pituitary as well as pulsatile LH secretion in bovines. *Domest Anim Endocrinol* (2009) **36**:219–24. doi:10.1016/j.domaniend.2009.02.001
65. Sawada K, Ukena K, Kikuyama S, Tsutsui K. Identification of a cDNA encoding a novel amphibian growth hormone-releasing peptide and localization of its transcript. *J Endocrinol* (2002) **174**:395–402. doi:10.1677/joe.0.1740395
66. Shahjahan M, Ikegami T, Osugi T, Ukena K, Doi H, Hattori A, et al. Synchronised expressions of LPXRFamide peptide and its receptor genes: seasonal, diurnal and circadian changes during spawning period in grass puffer. *J Neuroendocrinol* (2011) **23**:39–51. doi:10.1111/j.1365-2826.2010.02081.x

67. Amano M, Moriyama S, Iigo M, Kitamura S, Amiya N, Yamamori K, et al. Novel fish hypothalamic neuropeptides stimulate the release of gonadotrophins and growth hormone from the pituitary of sockeye salmon. *J Endocrinol* (2006) **188**:417–23. doi:10.1677/joe.1.06494
68. Moussavi M, Wlasichuk M, Chang JP, Habibi HR. Seasonal effect of GnIH on gonadotrope functions in the pituitary of goldfish. *Mol Cell Endocrinol* (2012) **350**:53–60. doi:10.1016/j.mce.2011.11.020
69. Moussavi M, Wlasichuk M, Chang JP, Habibi HR. Seasonal effect of gonadotrophin inhibitory hormone on gonadotrophin-releasing hormone-induced gonadotroph functions in the goldfish pituitary. *J Neuroendocrinol* (2013) **25**:506–16. doi:10.1111/jne.12024
70. Qi X, Zhou W, Li S, Lu D, Yi S, Xie R, et al. Evidences for the regulation of GnRH and GTH expression by GnIH in the goldfish, *Carassius auratus*. *Mol Cell Endocrinol* (2013) **366**:9–20. doi:10.1016/j.mce.2012.11.001
71. Zhang Y, Li S, Liu Y, Lu D, Chen H, Huang X, et al. Structural diversity of the GnIH/GnIH receptor system in teleost: its involvement in early development and the negative control of LH release. *Peptides* (2010) **31**:1034–43. doi:10.1016/j.peptides.2010.03.003
72. Janvier P. Modern look for ancient lamprey. *Nature* (2006) **443**:921–4. doi:10.1038/443921a
73. Osugi T, Ukena K, Sower SA, Kawauchi H, Tsutsui K. Evolutionary origin and divergence of PQRamide peptides and LPXRFamide peptides in the RFamide peptide family. Insights from novel lamprey RFamide peptides. *FEBS J* (2006) **273**:1731–43. doi:10.1111/j.1742-4658.2006.01587.x
74. Moulédous L, Mollereau C, Zajac JM. Opioid-modulating properties of the neuropeptide FF system. *Biofactors* (2010) **36**:423–9. doi:10.1002/biof.116
75. Osugi T, Uchida K, Nozaki M, Tsutsui K. Characterization of novel RFamide peptides in the central nervous system of the brown hagfish: isolation, localization, and functional analysis. *Endocrinology* (2011) **152**:4252–64. doi:10.1210/en.2011-1375
76. Osugi T, Ubuka T, Tsutsui K. An evolutionary scenario for GnIH in chordates. *J Neuroendocrinol* (2015) **27**:556–66. doi:10.1111/jne.12246
77. Yin H, Ukena K, Ubuka T, Tsutsui K. A novel G protein-coupled receptor for gonadotropin-inhibitory hormone in the Japanese quail (*Coturnix japonica*): identification, expression and binding activity. *J Endocrinol* (2005) **184**:257–66. doi:10.1677/joe.1.05926
78. Bentley GE, Perfito N, Ukena K, Tsutsui K, Wingfield JC. Gonadotropin-inhibitory peptide in song sparrows (*Melospiza melodia*) in different reproductive conditions, and in house sparrows (*Passer domesticus*) relative to chicken-gonadotropin-releasing hormone. *J Neuroendocrinol* (2003) **15**:794–802. doi:10.1046/j.1365-2826.2003.01062.x
79. Hinuma S, Shintani Y, Fukusumi S, Iijima N, Matsumoto Y, Hosoya M, et al. New neuropeptides containing carboxy-terminal RFamide and their receptor in mammals. *Nat Cell Biol* (2000) **2**:703–8. doi:10.1038/35036326
80. Fukusumi S, Fujii R, Hinuma S. Recent advances in mammalian RFamide peptides: the discovery and functional analyses of PrRP, RFRPs and QRFP. *Peptides* (2006) **27**:1073–86. doi:10.1016/j.peptides.2005.06.031
81. Bonini JA, Jones KA, Adham N, Forray C, Artymyshyn R, Durkin MM, et al. Identification and characterization of two G protein-coupled receptors for neuropeptide FF. *J Biol Chem* (2000) **275**:39324–31. doi:10.1074/jbc.M004385200
82. Fredriksson R, Lagerström MC, Lundin LG, Schiöth HB. The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol* (2003) **63**:1256–72. doi:10.1124/mol.63.6.1256
83. Ubuka T, Tsutsui K. Evolution of gonadotropin-inhibitory hormone receptor and its ligand. *Gen Comp Endocrinol* (2014) **209**:148–61. doi:10.1016/j.yggen.2014.09.002
84. Liu Q, Guan XM, Martin WJ, McDonald TP, Clements MK, Jiang Q, et al. Identification and characterization of novel mammalian neuropeptide FF-like peptides that attenuate morphine-induced antinociception. *J Biol Chem* (2001) **276**:36961–9. doi:10.1074/jbc.M105308200
85. Son YL, Ubuka T, Millar RP, Kanasaki H, Tsutsui K. Gonadotropin-inhibitory hormone inhibits GnRH-induced gonadotropin subunit gene transcriptions by inhibiting AC/cAMP/PKA-dependent ERK pathway in LβT2 cells. *Endocrinology* (2012) **153**:2332–43. doi:10.1210/en.2011-1904
86. de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, Milgrom E. Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. *Proc Natl Acad Sci U S A* (2003) **100**:10972–6. doi:10.1073/pnas.1834399100
87. Kauffman AS, Clifton DK, Steiner RA. Emerging ideas about kisspeptin-GPR54 signaling in the neuroendocrine regulation of reproduction. *Trends Neurosci* (2007) **30**:504–11. doi:10.1016/j.tins.2007.08.001
88. Pinilla L, Aguilar E, Dieguez C, Millar RP, Tena-Sempere M. Kisspeptins and reproduction: physiological roles and regulatory mechanisms. *Physiol Rev* (2012) **92**:1235–316. doi:10.1152/physrev.00037.2010
89. Seminara SB, Messenger S, Chatzidakis EE, Thresher RR, Acierno JS Jr, Shagoury JK, et al. The GPR54 gene as a regulator of puberty. *N Engl J Med* (2003) **349**:1614–27. doi:10.1056/NEJMoa035322
90. Poling MC, Quennell JH, Anderson GM, Kauffman AS. Kisspeptin neurons do not directly signal to RFRP-3 neurones but RFRP-3 may directly modulate a subset of hypothalamic kisspeptin cells in mice. *J Neuroendocrinol* (2013) **25**:876–86. doi:10.1111/jne.12084
91. Bronson FH. *Mammalian Reproductive Biology*. Chicago, IL: University of Chicago Press (1990).
92. Ohta M, Kadota C, Konishi H. A role of melatonin in the initial stage of photoperiodism in the Japanese quail. *Biol Reprod* (1989) **40**:935–41. doi:10.1095/biolreprod40.5.935
93. Guymarc'h C, Lumineau S, Vivien-Roels B, Richard J, Deregnacourt S. Effect of melatonin supplementation on the sexual development in European quail (*Coturnix coturnix*). *Behav Processes* (2001) **53**:121–30. doi:10.1016/S0376-6357(01)00133-4
94. Rozenboim I, Aharony T, Yahav S. The effect of melatonin administration on circulating plasma luteinizing hormone concentration in castrated white leghorn roosters. *Poult Sci* (2002) **81**:1354–9. doi:10.1093/ps/81.9.1354
95. Greives TJ, Kingma SA, Beltrami G, Hau M. Melatonin delays clutch initiation in a wild songbird. *Biol Lett* (2012) **8**:330–2. doi:10.1098/rsbl.2011.1100
96. Wilson FE. Neither retinal nor pineal photoreceptors mediate photoperiodic control of seasonal reproduction in American tree sparrows (*Spizella arborea*). *J Exp Zool* (1991) **259**:117–27. doi:10.1002/jez.1402590114
97. Juss TS, Meddle SL, Servant RS, King VM. Melatonin and photoperiodic time measurement in Japanese quail (*Coturnix coturnix japonica*). *Proc Biol Sci* (1993) **254**:21–8. doi:10.1098/rspb.1993.0121
98. Ubuka T, Bentley GE, Ukena K, Wingfield JC, Tsutsui K. Melatonin induces the expression of gonadotropin-inhibitory hormone in the avian brain. *Proc Natl Acad Sci U S A* (2005) **102**:3052–7. doi:10.1073/pnas.0403840102
99. Chowdhury VS, Yamamoto K, Ubuka T, Bentley GE, Hattori A, Tsutsui K. Melatonin stimulates the release of gonadotropin-inhibitory hormone by the avian hypothalamus. *Endocrinology* (2010) **151**:271–80. doi:10.1210/en.2009-0908
100. Chowdhury VS, Ubuka T, Tsutsui K. Review: Melatonin stimulates the synthesis and release of gonadotropin-inhibitory hormone in birds. *Gen Comp Endocrinol* (2013) **181**:175–8. doi:10.1016/j.yggen.2012.08.005
101. Ubuka T, Bentley GE, Tsutsui K. Neuroendocrine regulation of gonadotropin secretion in seasonally breeding birds. *Front Neurosci* (2013) **7**:38. doi:10.3389/fnins.2013.00038
102. Revel FG, Saboureau M, Pévet P, Simonneaux V, Mikkelsen JD. RFamide-related peptide gene is a melatonin-driven photoperiodic gene. *Endocrinology* (2008) **149**:902–12. doi:10.1210/en.2007-0848
103. Mason AO, Duffy S, Zhao S, Ubuka T, Bentley GE, Tsutsui K, et al. Photoperiod and reproductive condition are associated with changes in RFamide-related peptide (RFRP) expression in Syrian hamsters (*Mesocricetus auratus*). *J Biol Rhythms* (2010) **25**:176–85. doi:10.1177/0748730410368821
104. Simonneaux V, Ancel C, Poirel VJ, Gauer F. Kisspeptins and RFRP-3 act in concert to synchronize rodent reproduction with seasons. *Front Neurosci* (2013) **7**:22. doi:10.3389/fnins.2013.00022
105. Piekarski DJ, Jarjisan SG, Perez L, Ahmad H, Dhawan N, Zucker I, et al. Effects of pinealectomy and short day lengths on reproduction and neuronal RFRP-3, kisspeptin, and GnRH in female Turkish hamsters. *J Biol Rhythms* (2014) **29**:181–91. doi:10.1177/0748730414532423
106. Janati A, Talbi R, Klosen P, Mikkelsen JD, Magoul R, Simonneaux V, et al. Distribution and seasonal variation in hypothalamic RF-amide peptides in a semi-desert rodent, the jerboa. *J Neuroendocrinol* (2013) **25**:402–11. doi:10.1111/jne.12015
107. Dardente H, Birnie M, Lincoln GA, Hazlerigg DG. RFamide-related peptide and its cognate receptor in the sheep: cDNA cloning, mRNA distribution in

- the hypothalamus and the effect of photoperiod. *J Neuroendocrinol* (2008) **20**:1252–9. doi:10.1111/j.1365-2826.2008.01784.x
108. Smith JT, Coolen LM, Kriegsfeld LJ, Sari IP, Jaafarzadehshirazi MR, Maltby M, et al. Variation in kisspeptin and RFamide-related peptide (RFRP) expression and terminal connections to gonadotropin-releasing hormone neurons in the brain: a novel medium for seasonal breeding in the sheep. *Endocrinology* (2008) **149**:5770–82. doi:10.1210/en.2008-0581
 109. Gingerich S, Wang X, Lee PK, Dhillon SS, Chalmers JA, Koletar MM, et al. The generation of an array of clonal, immortalized cell models from the rat hypothalamus: analysis of melatonin effects on kisspeptin and gonadotropin-inhibitory hormone neurons. *Neuroscience* (2009) **162**:1134–40. doi:10.1016/j.neuroscience.2009.05.026
 110. Vanecek J. Cellular mechanisms of melatonin action. *Physiol Rev* (1998) **78**:687–721.
 111. Chand D, Lovejoy DA. Stress and reproduction: controversies and challenges. *Gen Comp Endocrinol* (2011) **171**:253–7. doi:10.1016/j.yggen.2011.02.022
 112. Calisi RM, Rizzo NO, Bentley GE. Seasonal differences in hypothalamic EGR-1 and GnIH expression following capture-handling stress in house sparrows (*Passer domesticus*). *Gen Comp Endocrinol* (2008) **157**:283–7. doi:10.1016/j.yggen.2008.05.010
 113. Kirby ED, Geraghty AC, Ubuka T, Bentley GE, Kaufer D. Stress increases putative gonadotropin inhibitory hormone and decreases luteinizing hormone in male rats. *Proc Natl Acad Sci USA* (2009) **106**:11324–9. doi:10.1073/pnas.0901176106
 114. Son YL, Ubuka T, Narihiro M, Fukuda Y, Hasunuma I, Yamamoto K, et al. Molecular basis for the activation of gonadotropin-inhibitory hormone gene transcription by corticosterone. *Endocrinology* (2014) **155**:1817–26. doi:10.1210/en.2013-2076
 115. Kim JS, Brownjohn PW, Dyer BS, Beltramo M, Walker CS, Hay DL, et al. Anxiogenic and stressor effects of the hypothalamic neuropeptide RFRP-3 are overcome by a novel antagonist. *Endocrinology* (2015) **156**:4152–62. doi:10.1210/en.2015-1532
 116. Calisi RM, Diaz-Muñoz SL, Wingfield JC, Bentley GE. Social and breeding status are associated with the expression of GnIH. *Genes Brain Behav* (2011) **10**:557–64. doi:10.1111/j.1601-183X.2011.00693.x
 117. Delville Y, Sulon J, Hendrick JC, Balthazart J. Effect of the presence of females on the pituitary-testicular activity in male Japanese quail (*Coturnix coturnix japonica*). *Gen Comp Endocrinol* (1984) **55**:295–305. doi:10.1016/0016-6480(84)90115-1
 118. Cornil CA, Stevenson TJ, Ball GF. Are rapid changes in gonadal testosterone release involved in the fast modulation of brain estrogen effects? *Gen Comp Endocrinol* (2009) **163**:298–305. doi:10.1016/j.yggen.2009.04.029
 119. Anjum S, Krishna A, Sridaran R, Tsutsui K. Inhibitory roles of the mammalian GnIH ortholog RFRP3 in testicular activities in adult mice. *J Endocrinol* (2014) **223**:79–91. doi:10.1530/JOE-14-0333
 120. Bentley GE, Ubuka T, McGuire NL, Chowdhury VS, Morita Y, Yano T, et al. Gonadotropin-inhibitory hormone and its receptor in the avian reproductive system. *Gen Comp Endocrinol* (2008) **156**:34–43. doi:10.1016/j.yggen.2007.10.003
 121. Oishi H, Klausen C, Bentley GE, Osugi T, Tsutsui K, Gilks CB, et al. The human gonadotropin-inhibitory hormone ortholog RFamide-related peptide-3 suppresses gonadotropin-induced progesterone production in human granulosa cells. *Endocrinology* (2012) **153**:3435–45. doi:10.1210/en.2012-1066
 122. Singh P, Krishna A, Sridaran R, Tsutsui K. Changes in GnRH I, bradykinin and their receptors and GnIH in the ovary of *Calotes versicolor* during reproductive cycle. *Gen Comp Endocrinol* (2008) **159**:158–69. doi:10.1016/j.yggen.2008.08.016
 123. Singh P, Krishna A, Sridaran R, Tsutsui K. Immunohistochemical localization of GnRH and RFamide-related peptide-3 in the ovaries of mice during the estrous cycle. *J Mol Histol* (2011) **42**:371–81. doi:10.1007/s10735-011-9340-8
 124. Singh P, Krishna A, Tsutsui K. Effects of gonadotropin-inhibitory hormone on folliculogenesis and steroidogenesis of cyclic mice. *Fertil Steril* (2011) **95**:1397–404. doi:10.1016/j.fertnstert.2010.03.052
 125. Zhao S, Zhu E, Yang C, Bentley GE, Tsutsui K, Kriegsfeld LJ. RFamide-related peptide and messenger ribonucleic acid expression in mammalian testis: association with the spermatogenic cycle. *Endocrinology* (2010) **151**:617–27. doi:10.1210/en.2009-0978
 126. McGuire NL, Bentley GE. A functional neuropeptide system in vertebrate gonads: gonadotropin-inhibitory hormone and its receptor in testes of field-caught house sparrow (*Passer domesticus*). *Gen Comp Endocrinol* (2010) **166**:565–72. doi:10.1016/j.yggen.2010.01.010
 127. McGuire NL, Kangas K, Bentley GE. Effects of melatonin on peripheral reproductive function: regulation of testicular GnIH and testosterone. *Endocrinology* (2011) **152**:3461–70. doi:10.1210/en.2011-1053
 128. McGuire NL, Koh A, Bentley GE. The direct response of the gonads to cues of stress in a temperate songbird species is season-dependent. *PeerJ* (2013) **1**:e139. doi:10.7717/peerj.139
 129. Wingfield JC, Farner DS. Control of seasonal reproduction in temperate-zone birds. In: Reiter RJ, Follett BK, editors. *Progress in Reproductive Biology*. Vol. 5, New York: Karger (1980). p. 62–101.
 130. Bronson FH. Food-restricted, prepubertal, female rats: rapid recovery of luteinizing hormone pulsing with excess food, and full recovery of pubertal development with gonadotropin-releasing hormone. *Endocrinology* (1986) **118**:2483–7. doi:10.1210/endo-118-6-2483
 131. Schneider JE, Wise JD, Benton NA, Brozek JM, Keen-Rhinehart E. When do we eat? Ingestive behavior, survival, and reproductive success. *Horm Behav* (2013) **64**:702–28. doi:10.1016/j.yhbeh.2013.07.005
 132. Bronson FH. Effect of food manipulation on the GnRH-LH-estradiol axis of young female rats. *Am J Physiol* (1988) **254**:R616–21.
 133. Bronson FH, Marsteller FA. Effect of short-term food deprivation on reproduction in female mice. *Biol Reprod* (1985) **33**:660–7. doi:10.1095/biolreprod33.3.660
 134. Schneider JE, Wade GN. Availability of metabolic fuels controls estrous cyclicity of Syrian hamsters. *Science* (1989) **244**:1326–8. doi:10.1126/science.2734610
 135. Shahab M, Zaman W, Bashir K, Arslan M. Fasting-induced suppression of hypothalamic-pituitary-gonadal axis in the adult rhesus monkey: evidence for involvement of excitatory amino acid neurotransmitters. *Life Sci* (1997) **61**:1293–300. doi:10.1016/S0024-3205(97)00674-7
 136. Sisk CL, Bronson FH. Effects of food restriction and restoration on gonadotropin and growth hormone secretion in immature male rats. *Biol Reprod* (1986) **35**:554–61. doi:10.1095/biolreprod35.3.554
 137. Tachibana T, Sato M, Takahashi H, Ukena K, Tsutsui K, Furuse M. Gonadotropin-inhibiting hormone stimulates feeding behavior in chicks. *Brain Res* (2005) **1050**:94–100. doi:10.1016/j.brainres.2005.05.035
 138. Fraley GS, Coombs E, Gerometta E, Colton S, Sharp PJ, Li Q, et al. Distribution and sequence of gonadotropin-inhibitory hormone and its potential role as a molecular link between feeding and reproductive systems in the Pekin duck (*Anas platyrhynchos domestica*). *Gen Comp Endocrinol* (2013) **184**:103–10. doi:10.1016/j.yggen.2012.11.026
 139. Tachibana T, Masuda N, Tsutsui K, Ukena K, Ueda H. The orexigenic effect of GnIH is mediated by central opioid receptors in chicks. *Comp Biochem Physiol A Mol Integr Physiol* (2008) **150**:21–5. doi:10.1016/j.cbpa.2008.02.018
 140. Clarke IJ, Smith JT, Henry BA, Oldfield BJ, Stefanidis A, Millar RP, et al. Gonadotropin-inhibitory hormone is a hypothalamic peptide that provides a molecular switch between reproduction and feeding. *Neuroendocrinology* (2012) **95**:305–16. doi:10.1159/000332822
 141. Fu LY, van den Pol AN. Kisspeptin directly excites anorexigenic proopiomelanocortin neurons but inhibits orexigenic neuropeptide Y cells by an indirect synaptic mechanism. *J Neurosci* (2010) **30**:10205–19. doi:10.1523/JNEUROSCI.2098-10.2010
 142. Jacobi JS, Coleman HA, Enriori PJ, Parkinson HC, Li Q, Pereira A, et al. Paradoxical effect of gonadotropin-inhibiting hormone to negatively regulate neuropeptide Y neurons in mouse arcuate nucleus. *J Neuroendocrinol* (2013) **25**:1308–17. doi:10.1111/jne.12114
 143. Bentley GE, Jensen JP, Kaur GJ, Wacker DW, Tsutsui K, Wingfield JC. Rapid inhibition of female sexual behavior by gonadotropin-inhibitory hormone (GnIH). *Horm Behav* (2006) **49**:550–5. doi:10.1016/j.yhbeh.2005.12.005
 144. Ubuka T, Mukai M, Wolfe J, Beverly R, Clegg S, Wang A, et al. RNA interference of gonadotropin-inhibitory hormone gene induces arousal in songbirds. *PLoS One* (2012) **7**:e30202. doi:10.1371/journal.pone.0030202
 145. Maney DL, Richardson RD, Wingfield JC. Central administration of chicken gonadotropin-releasing hormone-II enhances courtship behavior in a female sparrow. *Horm Behav* (1997) **32**:11–8. doi:10.1006/hbeh.1997.1399

146. Piekarski DJ, Zhao S, Jennings KJ, Iwasa T, Legan SJ, Mikkelsen JD, et al. Gonadotropin-inhibitory hormone reduces sexual motivation but not lordosis behavior in female Syrian hamsters (*Mesocricetus auratus*). *Horm Behav* (2013) **64**:501–10. doi:10.1016/j.yhbeh.2013.06.006
147. Do-Rego JL, Seong JY, Burel D, Leprince J, Luu-The V, Tsutsui K, et al. Review: neurosteroid biosynthesis: enzymatic pathways and neuroendocrine regulation by neurotransmitters and neuropeptides. *Front Neuroendocrinol* (2009) **30**:259–301. doi:10.1016/j.yfrne.2009.05.006
148. Mills AD, Crawford LL, Domjan M, Faure JM. The behavior of the Japanese or domestic quail *Coturnix japonica*. *Neurosci Biobehav Rev* (1997) **21**:261–81. doi:10.1016/S0149-7634(96)00028-0
149. Selinger HE, Bermant G. Hormonal control of aggressive behavior in Japanese quail (*Coturnix coturnix japonica*). *Behaviour* (1967) **28**:255–68. doi:10.1163/156853967X00037
150. Tsutsui K, Ishii S. Effects of sex steroids on aggressive behavior of adult male Japanese quail. *Gen Comp Endocrinol* (1981) **44**:480–6. doi:10.1016/0016-6480(81)90336-1
151. Schlinger BA, Callard GV. Aromatization mediates aggressive behavior in quail. *Gen Comp Endocrinol* (1990) **79**:39–53. doi:10.1016/0016-6480(90)90086-2
152. Balthazart J, Cornil CA, Charlier TD, Taziaux M, Ball GF. Estradiol, a key endocrine signal in the sexual differentiation and activation of reproductive behavior in quail. *J Exp Zool A Ecol Genet Physiol* (2009) **311**:323–45. doi:10.1002/jez.464
153. Balthazart J, Charlier TD, Cornil CA, Dickens MJ, Harada N, Konkle AT, et al. Sex differences in brain aromatase activity: genomic and non-genomic controls. *Front Endocrinol* (2011) **2**:34. doi:10.3389/fendo.2011.00034
154. Yahr P. Data and hypotheses in tales of dihydrotestosterone. *Horm Behav* (1979) **13**:92–6. doi:10.1016/0018-506X(79)90037-0
155. Ubuka T, Ueno M, Ukena K, Tsutsui K. Developmental changes in gonadotropin-inhibitory hormone in the Japanese quail (*Coturnix japonica*) hypothalamo-hypophysial system. *J Endocrinol* (2003) **178**:311–8. doi:10.1677/joe.0.1780311
156. Ukena K, Ubuka T, Tsutsui K. Distribution of a novel avian gonadotropin-inhibitory hormone in the quail brain. *Cell Tissue Res* (2003) **312**:73–9.
157. Absil P, Ritters LV, Balthazart J. Preoptic aromatase cells project to the mesencephalic central gray in the male Japanese quail (*Coturnix japonica*). *Horm Behav* (2001) **40**:369–83. doi:10.1006/hbeh.2001.1702
158. Cornil CA, Ball GF, Balthazart J. Rapid control of male typical behaviors by brain-derived estrogens. *Front Neuroendocrinol* (2012) **33**:425–46. doi:10.1016/j.yfrne.2012.08.003
159. Balthazart J, Surlmont C. Androgen and estrogen action in the preoptic area and activation of copulatory behavior in quail. *Physiol Behav* (1990) **48**:599–609. doi:10.1016/0031-9384(90)90198-D
160. Panzica GC, Viglietti-Panzica C, Balthazart J. The sexually dimorphic medial preoptic nucleus of quail: a key brain area mediating steroid action on male sexual behavior. *Front Neuroendocrinol* (1996) **17**:51–125. doi:10.1006/frne.1996.0002
161. Balthazart J, Baillien M, Ball GF. Rapid control of brain aromatase activity by glutamatergic inputs. *Endocrinology* (2006) **147**:359–66. doi:10.1210/en.2005-0845
162. Balthazart J, Baillien M, Ball GF. Rapid and reversible inhibition of brain aromatase activity. *J Neuroendocrinol* (2001) **13**:63–73. doi:10.1046/j.1365-2826.2001.00598.x
163. Balthazart J, Baillien M, Ball GF. Phosphorylation processes mediate rapid changes of brain aromatase activity. *J Steroid Biochem Mol Biol* (2001) **79**:261–77. doi:10.1016/S0960-0760(01)00143-1
164. Balthazart J, Baillien M, Charlier TD, Ball GF. Calcium-dependent phosphorylation processes control brain aromatase in quail. *Eur J Neurosci* (2003) **17**:1591–606. doi:10.1046/j.1460-9568.2003.02598.x
165. Charlier TD, Harada N, Balthazart J, Cornil CA. Human and quail aromatase activity is rapidly and reversibly inhibited by phosphorylating conditions. *Endocrinology* (2011) **152**:4199–210. doi:10.1210/en.2011-0119
166. Pineda R, Garcia-Galiano D, Sanchez-Garrido MA, Romero M, Ruiz-Pino F, Aguilar E, et al. Characterization of the inhibitory roles of RFRP3, the mammalian ortholog of GnIH, in the control of gonadotropin secretion in the rat: in vivo and in vitro studies. *Am J Physiol Endocrinol Metab* (2010) **299**:E39–46. doi:10.1152/ajpendo.00108.2010
167. Simonin F, Schmitt M, Laulin JP, Laboureyras E, Jhamandas JH, MacTavish D, et al. RF9, a potent and selective neuropeptide FF receptor antagonist, prevents opioid-induced tolerance associated with hyperalgesia. *Proc Natl Acad Sci U S A* (2006) **103**:466–71. doi:10.1073/pnas.0502090103
168. Steele RE, Mellor LB, Sawyer WK, Wasvary JM, Browne LJ. *In vitro* and *in vivo* studies demonstrating potent and selective estrogen inhibition with the nonsteroidal aromatase inhibitor CGS 16949A. *Steroids* (1987) **50**:147–61. doi:10.1016/0039-128X(83)90068-5
169. Wade J, Schlinger BA, Hodges L, Arnold AP. Fadrozole: a potent and specific inhibitor of aromatase in the zebra finch brain. *Gen Comp Endocrinol* (1994) **94**:53–61. doi:10.1006/gcen.1994.1059
170. Haraguchi S, Hara S, Ubuka T, Mita M, Tsutsui K. Possible role of pineal allopregnanolone in Purkinje cell survival. *Proc Natl Acad Sci U S A* (2012) **109**:21110–5. doi:10.1073/pnas.1210804109
171. Hatori M, Hirota T, Iitsuka M, Kurabayashi N, Haraguchi S, Kokame K, et al. Light-dependent and circadian clock-regulated activation of sterol regulatory element-binding protein, X-box-binding protein 1, and heat shock factor pathways. *Proc Natl Acad Sci U S A* (2011) **108**:4864–9. doi:10.1073/pnas.1015959108
172. Tsutsui K, Haraguchi S, Fukada Y, Vaudry H. Brain and pineal 7 α -hydroxypregnenolone stimulating locomotor activity: identification, mode of action and regulation of biosynthesis. *Front Neuroendocrinol* (2013) **34**:179–89. doi:10.1016/j.yfrne.2013.05.002
173. Tsutsui K, Haraguchi S, Hatori M, Hirota T, Fukada Y. Biosynthesis and biological actions of pineal neurosteroids in domestic birds. *Neuroendocrinology* (2013) **98**:97–105. doi:10.1159/000353782

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Tsutsui, Ubuka, Son, Bentley and Kriegsfeld. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.