

UCSF

UC San Francisco Previously Published Works

Title

Longitudinal stability in cigarette smokers of urinary biomarkers of exposure to the toxicants acrylonitrile and acrolein

Permalink

<https://escholarship.org/uc/item/2034023n>

Journal

PLOS ONE, 14(1)

ISSN

1932-6203

Authors

Chen, Menglan
Carmella, Steven G
Sipe, Christopher
[et al.](#)

Publication Date

2019

DOI

10.1371/journal.pone.0210104

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-ShareAlike License, available at <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Peer reviewed

RESEARCH ARTICLE

Longitudinal stability in cigarette smokers of urinary biomarkers of exposure to the toxicants acrylonitrile and acrolein

Menglan Chen¹, Steven G. Carmella¹, Christopher Sipe¹, Joni Jensen¹, Xianghua Luo^{1,2}, Chap T. Le^{1,2}, Sharon E. Murphy¹, Neal L. Benowitz³, F. Joseph McClernon⁴, Ryan Vandrey⁵, Sharon S. Allen⁶, Rachel Denlinger-Apte⁷, Paul M. Cinciripini⁸, Andrew A. Strasser⁹, Mustafa al'Absi¹⁰, Jason D. Robinson⁸, Eric C. Donny¹¹, Dorothy Hatsukami¹, Stephen S. Hecht¹*

1 Masonic Cancer Center, University of Minnesota, Minneapolis, Minnesota, United States of America, **2** Division of Biostatistics, School of Public Health, University of Minnesota, Minneapolis, Minnesota, United States of America, **3** Department of Medicine, University of California, San Francisco, California, United States of America, **4** Department of Psychiatry and Behavioral Sciences, Duke University, Durham, North Carolina, United States of America, **5** Department of Psychiatry and Behavioral Sciences, Johns Hopkins University, Baltimore, Maryland, United States of America, **6** Department of Family Medicine and Community Health, University of Minnesota Medical School, Minneapolis, Minnesota, United States of America, **7** Department of Behavioral and Social Sciences, Brown University, Providence, Rhode Island, United States of America, **8** Department of Behavioral Science, University of Texas MD Anderson Cancer Center, Houston, Texas, United States of America, **9** Department of Psychiatry, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America, **10** Behavioral Medicine Laboratories, University of Minnesota Medical School, Duluth, Minnesota, United States of America, **11** Department of Physiology and Pharmacology, Wake Forest School of Medicine, Winston-Salem, North Carolina, United States of America

* hecht002@umn.edu



OPEN ACCESS

Citation: Chen M, Carmella SG, Sipe C, Jensen J, Luo X, Le CT, et al. (2019) Longitudinal stability in cigarette smokers of urinary biomarkers of exposure to the toxicants acrylonitrile and acrolein. *PLoS ONE* 14(1): e0210104. <https://doi.org/10.1371/journal.pone.0210104>

Editor: Yu Ru Kou, National Yang-Ming University, TAIWAN

Received: October 17, 2018

Accepted: December 17, 2018

Published: January 4, 2019

Copyright: © 2019 Chen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This work was supported by National Institute on Drug Abuse and Food and Drug Administration grant U54 DA-031659 (ECD, DKH, CTL, and SSH). The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the

Abstract

The urinary metabolites cyanoethyl mercapturic acid (CEMA) and 3-hydroxypropyl mercapturic acid (3-HPMA) have been widely used as biomarkers of exposure to acrylonitrile and acrolein, respectively, but there are no published data on their consistency over time in the urine of cigarette smokers. We provided, free of charge over a 20 week period, Spectrum NRC600/601 research cigarettes to cigarette smokers in the control arm of a randomized clinical trial of the reduced nicotine cigarette. Urine samples were collected at weeks 4, 8, 12, 16, and 20 and analyzed for CEMA and 3-HPMA, and total nicotine equivalents (TNE) using validated methods. Creatinine-corrected intra-class correlation coefficients for CEMA, 3-HPMA, and TNE were 0.67, 0.46, and 0.68, respectively, indicating good longitudinal consistency for CEMA, while that of 3-HPMA was fair. A strong correlation between CEMA and TNE values was observed. These data support the use of CEMA as a reliable biomarker of tobacco smoke exposure. This is the first report of the longitudinal stability of the biomarkers of acrylonitrile and acrolein exposure in smokers. The data indicate that CEMA, the biomarker of acrylonitrile exposure, is consistent over time in cigarette smokers, supporting its use. While 3-HPMA levels were less stable over time, this biomarker is nevertheless a useful monitor of human acrolein exposure because of its specificity to this toxicant.

manuscript; and decision to submit the manuscript for publication.

Competing interests: The authors have declared that no competing interests exist.

Introduction

The U.S. National Toxicology Program 14th Report on Carcinogens classifies acrylonitrile as “reasonably anticipated to be a human carcinogen” based on sufficient evidence of carcinogenicity from studies in experimental animals [1]. Acrylonitrile causes a variety of tumors in rats including cancer of the central nervous system and Zymbal gland. However, data from epidemiological studies are considered inadequate to evaluate specific exposures to acrylonitrile as causes of human cancer. The International Agency for Research on Cancer concluded that acrylonitrile is “possibly carcinogenic to humans.” [2] Tobacco smoke is a major source of human exposure to acrylonitrile, with a mean value of 28.4 µg per cigarette in mainstream smoke of U.S. brand cigarettes using the Health Canada smoking regimen [3]. Cyanoethyl mercapturic acid (CEMA, Fig 1) is a urinary metabolite of acrylonitrile formed by its initial Michael addition reaction with glutathione followed by metabolic removal of the glutamic acid and glycine residues by peptidases and *N*-acetylation of the remaining cysteine residue [4].

Jakubowski *et al* were the first to measure CEMA in humans exposed to acrylonitrile. They used gas chromatography with flame ionization detection [5]. Schettgen *et al* developed a column-switching liquid chromatography tandem mass spectrometry (LC-MS/MS) method for CEMA in human urine and found a relationship to cigarette smoking as well as passive smoke exposure [6]. Scherer *et al* reported a method for CEMA in human urine using derivatization with pentafluorobenzyl bromide followed by LC-MS/MS analysis. They observed a significant decrease in CEMA levels upon smoking cessation [7]. Minet *et al* showed that smokers excreted from 75–165 times more urinary CEMA than non-smokers and that CEMA levels in smokers were significantly correlated with total nicotine equivalents (TNE) and other biomarkers [8]. Multiple other investigators have confirmed and extended these results, including application to studies of marijuana smokers, waterpipe users, and e-cigarette users [9–21].

The second toxicant investigated in this study is acrolein, an intense eye and respiratory tract irritant which causes a variety of adverse effects including irritation, inflammation, and cell proliferation, although there is scant evidence for its carcinogenicity [22]. It is considered to be one of the most toxic compounds in tobacco smoke [23, 24]. Its respiratory toxicity results from its chemical reactivity with respiratory tract proteins and other macromolecules, depletion of glutathione, and necrotic cell death among other effects [23, 24]. Tobacco smoke is an important source of acrolein exposure, with a mean level of 177 µg per cigarette in mainstream smoke [3]. 3-Hydroxypropyl mercapturic acid (3-HPMA, Fig 1) is an accepted biomarker of acrolein exposure. This urinary metabolite forms by Michael addition of glutathione followed by endogenous reduction of the aldehyde group and metabolic processing to the *N*-acetylcysteine conjugate 3-HPMA.

Mascher *et al* were the first to develop an LC-MS/MS method for 3-HPMA, applying it to analysis of urine from smokers and non-smokers [25]. Scherer *et al* demonstrated a borderline significant decrease in the amount of 3-HPMA in the urine of smokers who used charcoal filtered cigarettes compared to cellulose acetate filtered cigarettes [26]. We demonstrated a




Fig 1. Structures of CEMA and 3-HPMA.

<https://doi.org/10.1371/journal.pone.0210104.g001>

significant decrease in urinary 3-HPMA when cigarette smokers stopped smoking [27, 28]. Yan *et al* developed a rapid LC-MS/MS method for analysis of 3-HPMA in urine [29]. Morin *et al* and Shepperd *et al* demonstrated significant correlations of mouth level exposure to acrolein in smokers and 3-HPMA in urine [30, 31]. Minet *et al* carried out an inter-laboratory comparison of 3-HPMA levels in urine and found good inter-laboratory reproducibility [32]. We developed a high throughput LC-MS/MS method for 3-HPMA in cigarette smokers' urine [33]. and demonstrated significantly lower levels of this acrolein metabolite in the urine of e-cigarette users than in cigarette smokers [34]. Alwis *et al* determined 3-HPMA levels in the NHANES study, 2005–2006, and found five times higher levels among tobacco smokers than non-tobacco users [35]. Kassem *et al* found that hookah smoking was a significant source of acrolein exposure based on urinary 3-HPMA measurements [36]. We observed that urinary levels of 3-HPMA were significantly different among five ethnic groups of cigarette smokers in the Multiethnic Cohort Study [37].

Thus, multiple reports in the literature have described the development and application of CEMA and 3-HPMA as biomarkers of acrylonitrile and acrolein exposure, respectively, focusing mainly on exposure to these significant toxicants from cigarette smoking. However, to our knowledge, there have been no reports on the longitudinal consistency of these two important biomarkers of volatile toxicant uptake in cigarette smokers. It is critical to know whether measurement of a urinary biomarker at a given point in time is representative of its levels over time in cigarette smokers. We addressed this question here by examining the consistency over time of CEMA and 3-HPMA using urine samples obtained at 5 time points over 20 weeks from smokers participating in the control arm of a clinical trial of reduced nicotine content cigarettes [38]. These smokers were given, free of charge, "Spectrum" research cigarettes containing 15.5 mg nicotine per gram tobacco.

Materials and methods

Study design

The data reported here are from the control group of a 10-site randomized clinical trial that studied the effect of immediate vs. gradual reduction in nicotine content of cigarettes on smoking-related behavior and biomarkers of smoke exposure. Details of this multi-site trial have been reported [38]. The study protocol was approved by the University of Minnesota Institutional Review Board (IRB Code Number 1106M01561, FWA Number 00000312). Written consent was obtained from all subjects. Briefly, the control group consisted of 249 cigarette smokers (43% female) randomly assigned to smoke Spectrum NRC600/601 research cigarettes, which are non-menthol and mentholated, respectively, containing 15.5 mg nicotine/g tobacco, similar to commercially available cigarettes. The NRC600 cigarette mainstream smoke contained 54.9 μg /cigarette acrolein (ISO conditions) and 104 μg /cigarette acrolein under Canadian Intense conditions. The corresponding values for NRC601 cigarette smoke were 54.1 and 101 [39]. Data for acrylonitrile levels in mainstream smoke of these cigarettes was not available. The cigarettes were provided free of charge to participants over a period of 20 weeks. Participants attended a weekly clinic visit for the first 4 weeks and then bi-weekly visits for the next 16 weeks to complete questionnaires and other measures and receive supplies of additional cigarettes, approximately twice their self-reported baseline level of consumption. At all visits, smokers were advised on the importance of smoking only the study cigarettes, as long as they continued to smoke. Support was given for attempts to quit smoking if interest were expressed. At weeks 4, 8, 12, 16, and 20, first morning urine voids were collected for measurement of biomarkers.

Analysis of CEMA and 3-HPMA in urine

The analysis was performed essentially as described [33] with slight modifications for analysis of CEMA. Briefly, $[CD_3]_3$ -HPMA and $[CD_3]$ CEMA were added to 200 μ l urine. An Oasis Max 60 mg mixed mode anion exchange 96 well plate (Waters Corp.) was preconditioned with methanol and 2% aqueous NH_4OH . The sample was applied, and eluted with 30% methanol containing 2% formic acid, to give the fraction containing 3-HPMA and CEMA. The fraction was concentrated to dryness, and the residue dissolved in 60 μ l MeOH/ NH_4OAc :1/3, pH 6.8, and analyzed by atmospheric pressure chemical ionization (APCI)-LC-MS/MS-SRM using an Agilent 1100 HPLC system (Agilent Technologies) coupled to a TSQ Quantum Discovery Max instrument (Thermo Scientific) equipped with a 50 x 3.0 mm, 2.5-micron Sunfire C18 LC column (Waters Corp.). Solvent A was 15 mM NH_4OAc , pH 6.8, and solvent B was methanol. The column temperature was 50 °C and the flow rate was 0.4 ml/min. Three μ l of the sample were injected. The LC solvent system was 98% A and 2% B held for 2 min, then ramped to 12% B in 6.5 min, then to 70% B in 2.5 min, and then to 2% B in 0.5 min and held for 6 min. The MS/MS system was run in the negative APCI mode using the following ion mass transitions for detection: 3-HPMA, m/z 220 \rightarrow m/z 91; $[CD_3]_3$ -HPMA, m/z 223 \rightarrow m/z 91; CEMA, m/z 215 \rightarrow m/z 162, $[CD_3]$ CEMA, m/z 218 \rightarrow m/z 165. Other MS parameters were as follows: collision energy (V), 3-HPMA, 14; CEMA, 10; peak width, Q1 and Q3 (FWHM) 0.7; scan width (m/z), 0.40; scan time (sec), 0.2; vaporizer temp (°C), 450; capillary temperature (°C), 200; N_2 sheath gas pressure, psi, 30; N_2 auxiliary gas pressure, psi, 5.

Method validation sample

Aliquots of urine from 9 cigarette smokers who smoked an average of 16 cigarettes per day were pooled together and spiked with various amounts of CEMA ranging from 50 to 600 ng/ml.

Validation parameters

Intra-day precision was determined by repetitive analysis ($N = 6$) of the pooled smokers' urine sample on a single day. Inter-day precision was determined by including the pooled smokers' urine sample in each set of samples analyzed during the study ($N = 129$). Accuracy was determined by adding known amounts of CEMA to a pooled smokers' urine sample and then carrying out the analysis.

Analysis of total nicotine equivalents (TNE) and creatinine in urine

TNE (sum of nicotine, cotinine, 3'-hydroxycotinine and their glucuronides and nicotine-*N*-oxide) and creatinine were quantified as described previously [40].

Statistical analyses

Summary statistics, including mean, median, and interquartile ranges, for creatinine-corrected 3-HPMA, CEMA, and TNE at weeks 4, 8, 12, 16, and 20 are presented using boxplots. The intraclass correlation coefficient (ICC) and 95% confidence intervals (CI) were estimated by fitting a linear mixed model for repeatedly measured, log-transformed biomarkers with a random intercept using the SAS macro%ICC9 [41] where missing visits were assumed to be at random. The coefficient of variation (CV) for each participant who had two or more measurements was calculated as the sample standard deviation (SD) divided by the sample mean; and then the mean CV was calculated for all such participants. Note that higher ICC

and lower CV correspond to better longitudinal stability. The repeated-measure correlation (r_{rm}) between two log-transformed biomarkers and 95% CI were estimated by using the R package `rmcorr`; [42] the scatterplot of raw data was presented together with parallel lines which were fitted to the data of each subject. Note that the repeated-measure correlation is invariant to creatinine correction (or any other biomarker corrections or no correction) because of the log-transformation.

Results

The study sample consisted of 249 smokers (43% female), mean age 45.0 ± 13.4 years. They were 61% Caucasian, 30% African American, and 9% other races. They smoked a mean of 20.6 ± 11.4 cigarettes per day during the 20 week study period; 97.7% of these were the Spectrum research cigarettes, based on self-report. Further characteristics of the group have been reported [38].

The analytical method has been previously validated for 3-HPMA [33]. In this study, validation of the method was extended to CEMA. An LC-MS/MS chromatogram of CEMA and the internal standard [CD_3]CEMA is illustrated in Fig 2. As shown in Fig 3, there was excellent agreement between amounts of CEMA quantified in a pooled smokers' urine sample, and the amounts added to that sample ($R^2 = 0.997$). The expected value for CEMA in the unspiked urine sample was 117 ± 7.50 ng/ml (S.D.) and the observed y-intercept value was 120 ng/ml (0.56 nmol/ml). Intra-day precision ($N = 6$, CV x 100%) was 6.40% and inter-day precision ($N = 129$) was 6.60%. The limit of quantitation was 0.18 pmol/ml.

Means and other summary statistics for 3-HPMA, CEMA, and TNE over the course of the 20 week period are shown in Fig 4. Both 3-HPMA and CEMA were significantly correlated with TNE; the correlation of log transformed CEMA with TNE ($r_{rm}^2 = 0.58$) was stronger than that of 3-HPMA ($r_{rm}^2 = 0.38$) (Fig 5).

For participants who provided 2 or more urine samples ($N = 224$), the mean CV for creatinine corrected 3-HPMA, CEMA, and TNE were 48%, 41%, and 33%, respectively. The ICC, expressed per mg creatinine and per ml urine (e.g., non-corrected), are summarized in Table 1. For the creatinine-corrected data, the ICC of CEMA was similar to that of TNE and significantly superior to that of 3-HPMA. The ICC of the non-corrected values shared the same pattern as that of the creatinine-corrected values (see Table 1).

Discussion

Our data demonstrate that the longitudinal consistency of CEMA (ICC, 0.67) is similar to that of TNE (ICC, 0.68) and superior to that of 3-HPMA (ICC, 0.46). Quantitation of CEMA and 3-HPMA as urinary biomarkers of exposure to acrylonitrile and acrolein, respectively, has been reported extensively, but to our knowledge this is the first study to investigate the longitudinal consistency of these biomarkers in humans. Longitudinal consistency is an important parameter as it provides an indication of the generalizability of a measurement taken at a particular point in time. The control group in a randomized clinical trial of reduced nicotine content cigarettes, as reported here, provided an ideal opportunity to test biomarker longitudinal consistency. This control group of daily cigarette smokers was provided with Spectrum NRC600/601 research cigarettes, free of charge, over a period of 20 weeks. They were asked to smoke only these cigarettes, thus providing a relatively stable source of smoke exposure to acrylonitrile and acrolein. According to Cicchetti [43], an ICC between 0.75 and 1.00 is "excellent," one between 0.64 and 0.74 is "good", while a value between 0.40 and 0.59 is "fair." The similar ICC values of CEMA and TNE are clearly related to their relatively strong correlation, as indicated in Fig 5, which has also been noted by others [6, 8, 10, 14, 15]. We have previously




Fig 2. LC-MS/MS chromatograms of CEMA and the internal standard [CD₃]CEMA.

<https://doi.org/10.1371/journal.pone.0210104.g002>

determined ICC values for total urinary cotinine and total urinary 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) of 0.58 and 0.76, respectively, in smokers over a period of one year [44].

The better longitudinal consistency of CEMA as opposed to 3-HPMA undoubtedly results from the fact that exposure to acrolein from sources other than cigarette smoke is more likely than it is for acrylonitrile. Published studies [7, 8, 10–12] and our own unpublished data indicate that the relative levels of CEMA in smokers vs. non-smokers are far greater than those of 3-HPMA. In one recently completed unpublished study we observed a mean value of 0.31 ng/ml CEMA in the urine of 328 non-smokers analyzed by the method reported here, while the mean value in the current study was 300 ng/ml (1.38 nmol/ml), about 1000 times greater than in non-smokers. Others reported differences of 10–100 fold in CEMA values between smokers and non-smokers [6, 8, 10, 11, 15]. In contrast, data for 3-HPMA in our studies noted above were 443 ng/ml in the 328 non-smokers and 2190 ng/ml (9.9 nmol/ml) in smokers, a 6.7-fold difference, which is generally consistent with the literature [45]. These contrasting differences undoubtedly result from the fact that human environmental and endogenous exposure to




Fig 3. Relationship between added and measured CEMA in urine. MW of CEMA = 216.

<https://doi.org/10.1371/journal.pone.0210104.g003>

acrylonitrile from sources other than tobacco smoke is relatively rare while acrolein has multiple sources, most notably endogenous formation resulting from lipid peroxidation, polyamine metabolism and related processes, in addition to multiple exogenous combustion processes [46]. The formation of acrylonitrile from tobacco nitrate and nitrite has been postulated based on pyrolysis experiments [47]. Other potential sources of acrylonitrile in cigarette smoke may include pyrolysis of nicotine and anatabine.

The use of first morning void urine samples in this study may also have contributed to the relatively poor longitudinal stability of 3-HPMA. A study by Sarkar *et al* [48] evaluated spot urine samples at three time points (early morning, post-lunch and evening) along with 24 h urine collections in cigarette smokers. They observed diurnal variation in 3-HPMA levels, with lower levels in morning spot urine collections than in post lunch or evening collections, in contrast to tobacco-specific biomarkers such as TNE.

The method reported here for quantitation of CEMA uses APCI as the MS ionization technique. Most, if not all, other methods for CEMA analysis report the use of electrospray ionization techniques. Our choice of APCI was based to some extent on our analytical method for 3-HPMA which followed descriptions of APCI techniques used for other mercapturic acids [27, 49]. Furthermore, some studies indicate that APCI is less susceptible to ion suppression than electrospray ionization [50]. Our data for CEMA in smokers are similar to those of other reports that used electrospray ionization and the validation parameters are also similar [6–8, 10, 11, 15]. Thus, both ionization techniques appear to be acceptable for CEMA quantitation, when coupled with appropriate LC conditions and MS instrument parameters.

In summary, we report data which strongly support the use of CEMA as a biomarker of acrylonitrile exposure in cigarette smokers. Consistent with published studies, we find that LC-MS analysis of CEMA is accurate, precise, and dependable, and that the data from smokers correlate with TNE. Unique to this study, we also demonstrate good longitudinal consistency of urinary CEMA in cigarette smokers sample over a 20 week period. While the longitudinal stability of 3-HPMA was lower than that of CEMA, it remains a valid and specific quantitative biomarker of acrolein exposure in humans.




Fig 4. Mean values of 3-HPMA, CEMA, TNE, and CPD over the 20 week period in which subjects smoked the Spectrum cigarettes. Horizontal line inside the box: median; black diamond: mean; bottom and top edge of the box: 1st and 3rd quartile (interquartile range [IQR]); the upper whisker extends from the top of the box to the largest value no further than 1.5 times IQR and the bottom whisker extends from the bottom of the box to the smallest value no further than 1.5 times IQR; the y-axis is in natural log scale.

<https://doi.org/10.1371/journal.pone.0210104.g004>




Fig 5. Correlation of CEMA and 3-HPMA with TNE; each individual is represented by a different color.

<https://doi.org/10.1371/journal.pone.0210104.g005>

Table 1. Estimated intra-class correlation coefficients (ICC) for 3-HPMA, CEMA, and TNE.

Variable	N	Creatinine-corrected ICC (95% CI)	Non-corrected ICC (95% CI)
3-HPMA	233	0.46 (0.39, 0.52)	0.38 (0.32, 0.45)
CEMA	234	0.67 (0.62, 0.72)	0.55 (0.49, 0.61)
TNE	236	0.68 (0.63, 0.72)	0.58 (0.52, 0.64)

Creatinine-corrected values were calculated using nmol/mg creatinine. Non-corrected values were calculated using nmol/ml urine.

<https://doi.org/10.1371/journal.pone.0210104.t001>

Supporting information

S1 Table. Biomarker values.

(XLS)

Acknowledgments

The authors are indebted to the contributions of the Masonic Cancer Center Analytical Biochemistry Shared Resource, which provided mass spectrometry support essential for this study.

Author Contributions

Conceptualization: Sharon E. Murphy, Neal L. Benowitz, F. Joseph McClernon, Ryan Vandrey, Sharon S. Allen, Rachel Denlinger-Apte, Paul M. Cinciripini, Andrew A. Strasser, Mustafa al'Absi, Jason D. Robinson, Eric C. Donny, Dorothy Hatsukami, Stephen S. Hecht.

Data curation: Xianghua Luo, Chap T. Le.

Formal analysis: Xianghua Luo, Chap T. Le, Dorothy Hatsukami, Stephen S. Hecht.

Funding acquisition: Sharon E. Murphy, Neal L. Benowitz, F. Joseph McClernon, Sharon S. Allen, Rachel Denlinger-Apte, Paul M. Cinciripini, Andrew A. Strasser, Mustafa al'Absi, Jason D. Robinson, Eric C. Donny, Dorothy Hatsukami.

Investigation: Menglan Chen, Steven G. Carmella, Christopher Sipe, Joni Jensen, Xianghua Luo, Chap T. Le, Sharon E. Murphy, Neal L. Benowitz, F. Joseph McClernon, Ryan Vandrey, Sharon S. Allen, Rachel Denlinger-Apte, Paul M. Cinciripini, Andrew A. Strasser, Mustafa al'Absi, Jason D. Robinson, Eric C. Donny, Dorothy Hatsukami, Stephen S. Hecht.

Methodology: Menglan Chen, Steven G. Carmella, Christopher Sipe, Joni Jensen, Xianghua Luo, Chap T. Le, Sharon E. Murphy, Ryan Vandrey.

Project administration: Eric C. Donny, Dorothy Hatsukami, Stephen S. Hecht.

Supervision: Menglan Chen, Steven G. Carmella, Joni Jensen, Stephen S. Hecht.

Writing – original draft: Stephen S. Hecht.

Writing – review & editing: Menglan Chen, Steven G. Carmella, Christopher Sipe, Joni Jensen, Xianghua Luo, Chap T. Le, Sharon E. Murphy, Neal L. Benowitz, F. Joseph McClernon, Ryan Vandrey, Sharon S. Allen, Rachel Denlinger-Apte, Paul M. Cinciripini, Andrew A. Strasser, Mustafa al'Absi, Jason D. Robinson, Eric C. Donny, Dorothy Hatsukami.

References

1. NTP (National Toxicology Program). Report on Carcinogens, Fourteenth Edition. Research Triangle Park: U.S. Department of Health and Human Services; 2016.
2. International Agency for Research on Cancer. Re-evaluation of some organic chemicals, hydrazine and hydrogen peroxide (part one). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 71. Lyon, FR: IARC; 1999. p. 43–108
3. Bodnar JA, Morgan WT, Murphy PA, Ogden MW. Mainstream smoke chemistry analysis of samples from the 2009 US cigarette market. *Regul Toxicol Pharmacol.* 2012; 64(1):35–42. <https://doi.org/10.1016/j.yrtph.2012.05.011> PMID: 22683394
4. Sumner SC, Selvaraj L, Nauhaus SK, Fennell TR. Urinary metabolites from F344 rats and B6C3F1 mice coadministered acrylamide and acrylonitrile for 1 or 5 days. *Chem Res Toxicol.* 1997; 10(10):1152–60. <https://doi.org/10.1021/tx9602123> PMID: 9348438
5. Jakubowski M, Linhart I, Pielas G, Kopecky J. 2-Cyanoethylmercapturic acid (CEMA) in the urine as a possible indicator of exposure to acrylonitrile. *Br J Ind Med.* 1987; 44(12):834–40. PMID: 3689720
6. Schettgen T, Musiol A, Alt A, Ochsmann E, Kraus T. A method for the quantification of biomarkers of exposure to acrylonitrile and 1,3-butadiene in human urine by column-switching liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem.* 2009; 393(3):969–81. <https://doi.org/10.1007/s00216-008-2510-1> PMID: 19018522

7. Scherer G, Urban M, Hagedorn HW, Serafin R, Feng S, Kapur S, et al. Determination of methyl-, 2-hydroxyethyl- and 2-cyanoethylmercapturic acids as biomarkers of exposure to alkylating agents in cigarette smoke. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2010; 878(27):2520–8. <https://doi.org/10.1016/j.jchromb.2010.02.023> PMID: 20227354
8. Minet E, Cheung F, Errington G, Sterz K, Scherer G. Urinary excretion of the acrylonitrile metabolite 2-cyanoethylmercapturic acid is correlated with a variety of biomarkers of tobacco smoke exposure and consumption. *Biomarkers*. 2011; 16(1):89–96. <https://doi.org/10.3109/1354750X.2010.533287> PMID: 21108560
9. Wu CF, Uang SN, Chiang SY, Shih WC, Huang YF, Wu KY. Simultaneous quantitation of urinary cotinine and acrylonitrile-derived mercapturic acids with ultraperformance liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem*. 2012; 402(6):2113–20. <https://doi.org/10.1007/s00216-011-5661-4> PMID: 22231508
10. Hou H, Xiong W, Gao N, Zhang X, Tang G, Hu Q. A column-switching liquid chromatography-tandem mass spectrometry method for quantitation of 2-cyanoethylmercapturic acid and 2-hydroxyethylmercapturic acid in Chinese smokers. *Anal Biochem*. 2012; 430(1):75–82. <https://doi.org/10.1016/j.ab.2012.07.026> PMID: 22868354
11. Schettgen T, Bertram J, Kraus T. Accurate quantification of the mercapturic acids of acrylonitrile and its genotoxic metabolite cyanoethylene-epoxide in human urine by isotope-dilution LC-ESI/MS/MS. *Talanta*. 2012; 98:211–9. <https://doi.org/10.1016/j.talanta.2012.06.074> PMID: 22939149
12. Xiaotao Z, Hongwei H, Wei X, Qingyuan H. Simultaneous measurement of N-Acetyl-S-(2-cyanoethyl)-cysteine and N-acetyl-S-(2-hydroxyethyl)-cysteine in human urine by liquid chromatography-tandem mass spectrometry. *J Chromatogr Sci*. 2014; 52(7):719–24. <https://doi.org/10.1093/chromsci/bmt106> PMID: 23902664
13. Jacob P, 3rd, Abu Raddaha AH, Dempsey D, Havel C, Peng M, Yu L, et al. Comparison of nicotine and carcinogen exposure with water pipe and cigarette smoking. *Cancer Epidemiol Biomarkers Prev*. 2013; 22(5):765–72. <https://doi.org/10.1158/1055-9965.EPI-12-1422> PMID: 23462922
14. Zhang X, Xiong W, Shi L, Hou H, Hu Q. Simultaneous determination of five mercapturic acid derived from volatile organic compounds in human urine by LC-MS/MS and its application to relationship study. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2014; 967:102–9. <https://doi.org/10.1016/j.jchromb.2014.07.013> PMID: 25086756
15. Pluym N, Gilch G, Scherer G, Scherer M. Analysis of 18 urinary mercapturic acids by two high-throughput multiplex-LC-MS/MS methods. *Anal Bioanal Chem*. 2015; 407(18):5463–76. <https://doi.org/10.1007/s00216-015-8719-x> PMID: 25935678
16. Wei B, Alwis KU, Li Z, Wang L, Valentin-Blasini L, Sosnoff CS, et al. Urinary concentrations of PAH and VOC metabolites in marijuana users. *Environ Int*. 2016; 88:1–8. <https://doi.org/10.1016/j.envint.2015.12.003> PMID: 26690539
17. Boyle EB, Viet SM, Wright DJ, Merrill LS, Alwis KU, Blount BC, et al. Assessment of exposure to VOCs among pregnant women in the National Children's Study. *Int J Environ Res Public Health*. 2016; 13(4):376. <https://doi.org/10.3390/ijerph13040376> PMID: 27043585
18. Goniewicz ML, Gawron M, Smith DM, Peng M, Jacob P, 3rd, Benowitz NL. Exposure to nicotine and selected toxicants in cigarette smokers who switched to electronic cigarettes: a longitudinal within-subjects observational study. *Nicotine Tob Res*. 2017; 19(2):160–7. <https://doi.org/10.1093/ntr/ntw160> PMID: 27613896
19. Lin CY, Lee HL, Sung FC, Su TC. Investigating the association between urinary levels of acrylonitrile metabolite N-acetyl-S-(2-cyanoethyl)-L-cysteine and the oxidative stress product 8-hydroxydeoxyguanosine in adolescents and young adults. *Environ Pollut*. 2018; 239:493–8. <https://doi.org/10.1016/j.envpol.2018.04.010> PMID: 29684876
20. Lorkiewicz P, Riggs DW, Keith RJ, Conklin DJ, Xie Z, Sutaria S, et al. Comparison of urinary biomarkers of exposure in humans using electronic cigarettes, combustible cigarettes, and smokeless tobacco. *Nicotine Tob Res*. 2018.
21. Pulvers K, Emami AS, Nollen NL, Romero DR, Strong DR, Benowitz NL, et al. Tobacco consumption and toxicant exposure of cigarette smokers using electronic cigarettes. *Nicotine Tob Res*. 2018; 20(2):206–14. <https://doi.org/10.1093/ntr/ntw333> PMID: 28003511
22. International Agency for Research on Cancer. Acrolein. *IARC Monogr Eval Carcinog Risks Hum*. 63. Lyon, FR: IARC; 1995. p. 337–72. PMID: 9097100
23. Haussmann HJ. Use of hazard indices for a theoretical evaluation of cigarette smoke composition. *Chem Res Toxicol*. 2012; 25(4):794–810. <https://doi.org/10.1021/tx200536w> PMID: 22352345
24. Yeager RP, Kushman M, Chemerynski S, Weil R, Fu X, White M, et al. Proposed mode of action for acrolein respiratory toxicity associated with inhaled tobacco smoke. *Toxicol Sci*. 2016; 151(2):347–64. <https://doi.org/10.1093/toxsci/kfw051> PMID: 26969371

25. Mascher DG, Mascher HJ, Scherer G, Schmid ER. High-performance liquid chromatographic-tandem mass spectrometric determination of 3-hydroxypropylmercapturic acid in human urine. *J Chromatogr B Biomed Sci Appl.* 2001; 750(1):163–9. PMID: [11204217](https://pubmed.ncbi.nlm.nih.gov/11204217/)
26. Scherer G, Urban M, Engl J, Hagedorn HW, Riedel K. Influence of smoking charcoal filter tipped cigarettes on various biomarkers of exposure. *Inhal Toxicol.* 2006; 18(10):821–9. <https://doi.org/10.1080/08958370600747945> PMID: [16774872](https://pubmed.ncbi.nlm.nih.gov/16774872/)
27. Carmella SG, Chen M, Zhang Y, Zhang S, Hatsukami DK, Hecht SS. Quantitation of acrolein-derived 3-hydroxypropylmercapturic acid in human urine by liquid chromatography-atmospheric pressure chemical ionization-tandem mass spectrometry: effects of cigarette smoking. *Chem Res Toxicol.* 2007; 20:986–90. <https://doi.org/10.1021/tx700075y> PMID: [17559234](https://pubmed.ncbi.nlm.nih.gov/17559234/)
28. Carmella SG, Chen M, Han S, Briggs A, Jensen J, Hatsukami DK, et al. Effects of smoking cessation on eight urinary tobacco carcinogen and toxicant biomarkers. *Chem Res Toxicol.* 2009; 22(4):734–41. <https://doi.org/10.1021/tx800479s> PMID: [19317515](https://pubmed.ncbi.nlm.nih.gov/19317515/)
29. Yan W, Byrd GD, Brown BG, Borgerding MF. Development and validation of a direct LC-MS-MS method to determine the acrolein metabolite 3-HPMA in urine. *J Chromatogr Sci.* 2010; 48(3):194–9. PMID: [20223085](https://pubmed.ncbi.nlm.nih.gov/20223085/)
30. Morin A, Shepperd CJ, Eldridge AC, Poirier N, Voisine R. Estimation and correlation of cigarette smoke exposure in Canadian smokers as determined by filter analysis and biomarkers of exposure. *Regul Toxicol Pharmacol.* 2011; 61(3 Suppl):S3–12. <https://doi.org/10.1016/j.yrtph.2010.09.020> PMID: [20937342](https://pubmed.ncbi.nlm.nih.gov/20937342/)
31. Shepperd CJ, Eldridge AC, Mariner DC, McEwan M, Errington G, Dixon M. A study to estimate and correlate cigarette smoke exposure in smokers in Germany as determined by filter analysis and biomarkers of exposure. *Regul Toxicol Pharmacol.* 2009; 55(1):97–109. <https://doi.org/10.1016/j.yrtph.2009.06.006> PMID: [19539004](https://pubmed.ncbi.nlm.nih.gov/19539004/)
32. Minet E, Errington G, Scherer G, Newland K, Sharifi M, Bailey B, et al. An inter-laboratory comparison of urinary 3-hydroxypropylmercapturic acid measurement demonstrates good reproducibility between laboratories. *BMC Res Notes.* 2011; 4:391. <https://doi.org/10.1186/1756-0500-4-391> PMID: [21985092](https://pubmed.ncbi.nlm.nih.gov/21985092/)
33. Carmella SG, Chen M, Zarth A, Hecht SS. High throughput liquid chromatography-tandem mass spectrometry assay for mercapturic acids of acrolein and crotonaldehyde in cigarette smokers' urine. *J Chromatogr B.* 2013; 935:36–40.
34. Hecht SS, Carmella SG, Kotandeniya D, Pillsbury ME, Chen M, Ransom BW, et al. Evaluation of toxicant and carcinogen metabolites in the urine of e-cigarette users versus cigarette smokers. *Nicotine Tob Res.* 2015; 17(6):704–9. <https://doi.org/10.1093/ntr/ntu218> PMID: [25335945](https://pubmed.ncbi.nlm.nih.gov/25335945/)
35. Alwis KU, deCastro BR, Morrow JC, Blount BC. Acrolein exposure in U.S. tobacco smokers and non-tobacco users: NHANES 2005–2006. *Environ Health Perspect.* 2015; 123(12):1302–8. <https://doi.org/10.1289/ehp.1409251> PMID: [26024353](https://pubmed.ncbi.nlm.nih.gov/26024353/)
36. Kassem NOF, Kassem NO, Liles S, Zarth AT, Jackson SR, Daffa RM, et al. Acrolein exposure in hookah smokers and non-smokers exposed to hookah tobacco secondhand smoke: implications for regulating hookah tobacco products. *Nicotine Tob Res.* 2018; 20(4):492–501. <https://doi.org/10.1093/ntr/ntx133> PMID: [28591850](https://pubmed.ncbi.nlm.nih.gov/28591850/)
37. Park SL, Carmella SG, Chen M, Patel Y, Stram DO, Haiman CA, et al. Mercapturic acids derived from the toxicants acrolein and crotonaldehyde in the urine of cigarettes smokers from five ethnic groups with differing risks for lung cancer. *PLoS One.* 2015; 10:e0124841. <https://doi.org/10.1371/journal.pone.0124841> PMID: [26053186](https://pubmed.ncbi.nlm.nih.gov/26053186/)
38. Hatsukami DK, Luo X, Jensen JA, al'Absi M, Allen SS, Carmella SG, et al. Effect of immediate vs gradual reduction in nicotine content of cigarettes on biomarkers of smoke exposure: a randomized clinical trial. *JAMA.* 2018; 320(9):880–91. <https://doi.org/10.1001/jama.2018.11473> PMID: [30193275](https://pubmed.ncbi.nlm.nih.gov/30193275/)
39. Ding YS, Richter P, Hearn B, Zhang L, Bravo R, Yan X, et al. Chemical characterization of mainstream smoke from SPECTRUM variable nicotine research cigarettes. *Tob Regul Sci.* 2017; 3(1):81–94. PMID: [28220149](https://pubmed.ncbi.nlm.nih.gov/28220149/)
40. Murphy SE, Park SS, Thompson EF, Wilkens LR, Patel Y, Stram DO, et al. Nicotine N-glucuronidation relative to N-oxidation and C-oxidation and UGT2B10 genotype in five ethnic/racial groups. *Carcinogenesis.* 2014; 35(11):2526–33. <https://doi.org/10.1093/carcin/bgu191> PMID: [25233931](https://pubmed.ncbi.nlm.nih.gov/25233931/)
41. Hertzmark E, Spiegelman D. The SAS ICC9 macro. Harvard T.H. Chan School of Public Health; 2010.
42. Bakdash JZ, Marusich LR. Repeated measures correlation. *Front Psychol.* 2017; 8:456. <https://doi.org/10.3389/fpsyg.2017.00456> PMID: [28439244](https://pubmed.ncbi.nlm.nih.gov/28439244/)
43. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment.* 1994; 6:284–90.

44. Church TR, Anderson KE, Le C, Zhang Y, Kampa DM, Benoit AR, et al. Temporal stability of urinary and plasma biomarkers of tobacco smoke exposure among cigarette smokers. *Biomarkers*. 2010; 15:345–52. <https://doi.org/10.3109/13547501003753881> PMID: 20374068
45. Hecht SS, Yuan JM, Hatsukami DK. Applying tobacco carcinogen and toxicant biomarkers in product regulation and cancer prevention. *Chem Res Toxicol*. 2010; 23:1001–8. <https://doi.org/10.1021/tx100056m> PMID: 20408564
46. Stevens JF, Maier CS. Acrolein: sources, metabolism, and biomolecular interactions relevant to human health and disease. *Mol Nutr Food Res*. 2008; 52(1):7–25. <https://doi.org/10.1002/mnfr.200700412> PMID: 18203133
47. Johnson WR, Hale RW, Clough SC, Chen PH. Chemistry of the conversion of nitrate nitrogen to smoke products. *Nature*. 1973; 243:223–5. PMID: 4706291
48. Sarkar M, Muhammad-Kah R, Liang Q, Kapur S, Feng S, Roethig H. Evaluation of spot urine as an alternative to 24h urine collection for determination of biomarkers of exposure to cigarette smoke in adult smokers. *Environ Toxicol Pharmacol*. 2013; 36(1):108–14. <https://doi.org/10.1016/j.etap.2013.03.001> PMID: 23603463
49. Urban M, Gilch G, Schepers G, van Miert E, Scherer G. Determination of the major mercapturic acids of 1,3-butadiene in human and rat urine using liquid chromatography with tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2003; 796(1):131–40. PMID: 14552824
50. Boyd B, Basic C, Bethem R. Trace quantitative analysis by mass spectrometry. Chichester, West Sussex, England; Hoboken, N.J.: John Wiley & Sons; 2008.