

# UC Irvine

## UC Irvine Previously Published Works

### Title

Real-Time Subcellular Imaging of Breast Cancer Cell Attachment in Blood Vessels Using GFP-Labeled Paxillin in Live Mice.

### Permalink

<https://escholarship.org/uc/item/206379qw>

### Journal

CANCER RESEARCH, 71

### ISSN

0008-5472

### Authors

Suetsugu, A  
Digman, M  
Gratton, E  
[et al.](#)

### Publication Date

2011-12-15

### DOI

10.1158/0008-5472.SABCS11-P5-02-05

### License

<https://creativecommons.org/licenses/by/4.0/> 4.0

Peer reviewed

# Cancer Research

Poster Session Abstracts

## P5-02-05: Real-Time Subcellular Imaging of Breast Cancer Cell Attachment in Blood Vessels Using GFP-Labeled Paxillin in Live Mice.

A Suetsugu, M Digman, E Gratton, H Moriwaki, S Saji, M Bouvet, and RM Hoffman

DOI: 10.1158/0008-5472.SABCS11-P5-02-05 Published December 2011

[Article](#)

[Info & Metrics](#)

Abstracts: Thirty-Fourth Annual CTRC-AACR San Antonio Breast Cancer Symposium-- Dec 6-10, 2011; San Antonio, TX

### Abstract

**Background:** Paxillin is involved in the assembly of focal adhesions. We wish to visualize paxillin behavior in breast cancer cells *in vivo*, as well as *in vitro*.

**Materials and Methods:** Dual-photon confocal microscopy was used to image paxillin behavior by linking it to GFP. MDA-MB-231 human breast cancer cells expressing paxillin-GFP were imaged *in vitro* and *in vivo* adhering and trafficking in blood vessels in mice.  $10^6$  paxillin-GFP expressing breast cancer cells were injected in the epigastric cranialis vein.

**Results:** *In vitro*, round breast cancer cells had greater paxillin movement than stretched cancer cells as seen by fluorescence imaging. Paxillin-GFP breast cancer cells in the epigastric cranialis vein were initially rounded and had GFP-expressing protrusions. At later timepoints, many paxillin-GFP-expressing breast cancer cells stretched. The breast cancer cells then extravasated and subsequently grew around the outer surface of the blood vessel after one week. Two weeks after injection, paxillin-GFP expressing breast cancer cells were imaged migrating along the vessel wall. Most of the paxillin-GFP expressing breast cancer cells were stretched and were not mobile. With anti-VEGF treatment, paxillin was observed in round structures within the cells rather than stretched structures and paxillin movement within the cell was arrested.

Discussion: These results demonstrate that breast cancer cells brightly expressing paxillin-GFP and two-photon confocal microscopy can allow the visualization of the behavior of paxillin within breast cancer cells during adhesion and migration along the walls of blood vessels, as well as during anti-angiogenesis therapy.

**Citation Information:** Cancer Res 2011;71(24 Suppl):Abstract nr P5-02-05.

[^ Back to top](#)



December 2011  
Volume 71, Issue 24 Supplement  
[Table of Contents](#)

Search this issue



[Sign up for alerts](#)

[© Request Permissions](#)

[! Article Alerts](#)

[✉ Email Article](#)

[🌐 Citation Tools](#)

[↪ Share](#)

Tweet

Like 0

Advertisement