UCLA

UCLA Previously Published Works

Title

Dexmedetomidine Prevents Cognitive Decline by Enhancing Resolution of High Mobility Group Box 1 Protein-induced Inflammation through a Vagomimetic Action in Mice

Permalink

<https://escholarship.org/uc/item/20q30538>

Journal

ANESTHESIOLOGY, 128(5)

ISSN

0003-3022

Authors

Hu, Jun
Vacas, Susana
Feng, Xiaomei
et al.

Publication Date

2018

DOI

10.1097/ALN.0000000000002038

Peer reviewed

Dexmedetomidine Prevents Cognitive Decline by Enhancing Resolution of High Molecular Group Box 1 Protein–induced Inflammation through a Vagomimetic Action in Mice

Jun Hu, M.D., Susana Vacas, M.D., Ph.D., Xiaomei Feng, M.D., Ph.D., David Lutrin, M.D., Yosuke Uchida, M.D., Ph.D., Ieng Kit Lai, B.S., Mervyn Maze, M.B.Ch.B.

ABSTRACT

Background: Inflammation initiated by damage-associated molecular patterns has been implicated for the cognitive decline associated with surgical trauma and serious illness. We determined whether resolution of inflammation mediates dexmedetomidine-induced reduction of damage-associated molecular pattern–induced cognitive decline.

Methods: Cognitive decline (assessed by trace fear conditioning) was induced with high molecular group box 1 protein, a damage-associated molecular pattern, in mice that also received blockers of neural (vagal) and humoral inflammation-resolving pathways. Systemic and neuroinflammation was assessed by proinflammatory cytokines.

Results: Damage-associated molecular pattern–induced cognitive decline and inflammation (mean $\pm$ SD) was reversed by dexmedetomidine (trace fear conditioning: $58.77 \pm 8.69\%$ vs. $41.45 \pm 7.64\%$, $P < 0.0001$; plasma interleukin [IL]-1 β : 7.0 ± 2.2 pg/ml vs. 49.8 ± 6.0 pg/ml, $P < 0.0001$; plasma IL-6: 3.2 ± 1.6 pg/ml vs. 19.5 ± 1.7 pg/ml, $P < 0.0001$; hippocampal IL-1 β : 4.1 ± 3.0 pg/mg vs. 41.6 ± 8.0 pg/mg, $P < 0.0001$; hippocampal IL-6: 3.4 ± 1.3 pg/mg vs. 16.2 ± 2.7 pg/mg, $P < 0.0001$). Reversal by dexmedetomidine was prevented by blockade of vagomimetic imidazoline and α_7 nicotinic acetylcholine receptors but not by α_2 adrenoceptor blockade. Netrin-1, the orchestrator of inflammation–resolution, was upregulated (fold-change) by dexmedetomidine (lung: 1.5 ± 0.1 vs. 0.7 ± 0.1 , $P < 0.0001$; spleen: 1.5 ± 0.2 vs. 0.6 ± 0.2 , $P < 0.0001$), resulting in upregulation of proresolving (lipoxin-A $_4$: 1.7 ± 0.2 vs. 0.9 ± 0.2 , $P < 0.0001$) and downregulation of proinflammatory (leukotriene-B $_4$: 1.0 ± 0.2 vs. 3.0 ± 0.3 , $P < 0.0001$) humoral mediators that was prevented by α_7 nicotinic acetylcholine receptor blockade.

Conclusions: Dexmedetomidine resolves inflammation through vagomimetic (neural) and humoral pathways, thereby preventing damage-associated molecular pattern–mediated cognitive decline. (ANESTHESIOLOGY 2018; XXX:00-00)

OVER the last decade we have generated several lines of evidence that implicate surgery-initiated systemic inflammation and neuroinflammation in the development of postoperative cognitive decline both in preclinical^{1–3} and clinical settings.⁴ The engagement of the innate immune system, which triggers the inflammatory response, resulting in postoperative cognitive decline, is due to the damage-associated molecular pattern, high molecular group box 1 protein (HMGB1), which is passively released from traumatized tissue.^{5,6}

While postoperative cognitive decline was first reported in the setting of general anesthesia,⁷ no difference has been reported in the incidence of postoperative cognitive decline in patients randomized to receive general *versus* regional anesthetic techniques.⁸ These previous studies may have been underpowered, and larger trials are progressing to understand whether regional or general anesthesia is less likely to result in postoperative cognitive decline (Regional versus General Anesthesia

What We Already Know about This Topic

- Inflammation has been associated with cognitive disorders following surgery and medical illness
- Preclinical evidence suggests a role for high molecular group box 1 protein in cognitive disorders after an inflammatory insult

What This Article Tells Us That Is New

- In a preclinical model, dexmedetomidine prevented cognitive deficits resulting from administration of high molecular group box 1 protein via both vagomimetic and humoral pathways
- The results are consistent with the notion that the cognitive deficits noted after surgery or medical illness may be prevented by the administration of dexmedetomidine

for Promoting Independence after Hip Fracture [REGAIN] Trial; NCT02507505).⁹ Notwithstanding the outcome of the REGAIN trial, many patients will require either sedatives and/or general anesthesia combined with a regional technique.

Submitted for publication August 27, 2017. Accepted for publication November 21, 2017. From the Department of Anesthesia and Perioperative Care and Center for Cerebrovascular Research, University of California, San Francisco, San Francisco, California (J.H., X.F., D.L., Y.U., I.K.L., M.M.); the Department of Anesthesia, Tongling People's Hospital, Tongling, People's Republic of China (J.H.); and the Department of Anesthesia and Perioperative Medicine, University of California, Los Angeles, Los Angeles, California (S.V.).

Copyright © 2017, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2018; XXX:00-00

While some comparative studies have suggested that the frequency and/or severity of postoperative cognitive decline may be affected by the choice and dose of the anesthetic/sedative agent that is used for the surgical procedure,¹⁰ and that exposure to deeper anesthetic stages results in a higher incidence of delirium,¹¹ larger, appropriately powered studies are needed to avoid a type 2 statistical error, as was suspected in the Hip Fracture in Elderly Patients study (NCT01199276).¹²

It is notable that animal cohorts receiving anesthesia and analgesia alone did not differ from the control group (no anesthesia/analgesia, no surgery).^{1,2} Ethical considerations prevented us from using a “surgery only” (*i.e.*, without anesthesia/analgesia) cohort to which we could compare the neuroinflammatory and cognitive effects provoked by the addition of anesthetic and analgesic drugs. Now that we have demonstrated that systemically administered HMGB1 reproduces the surgical phenotype with high fidelity,⁵ we are ethically able to study a surrogate of the “surgery only” cohort without use of anesthesia/hypnotics to which we can compare the effects of adding these drugs.

To determine whether anesthetic/hypnotics have a modulating effect on the surgical phenotype, we have chosen to first study dexmedetomidine because it reduces cognitive decline^{13,14} and inflammation,¹⁵ including that associated with acute neurologic injury,¹⁶ by an as-yet-undetermined mechanism. Because of the challenge that there is “questionable biologic plausibility” for the beneficial effects that were noted postoperatively with dexmedetomidine,¹⁴ we addressed the hypothesis that the ameliorative effect of dexmedetomidine on cognitive decline is due to resolution of inflammation through neural and humoral processes.

Materials and Methods

Animals

All experimental procedures involving animals were approved by the Institutional Animal Care and Use Committee of the University of California, San Francisco (AN 167062) and conformed to National Institutes of Health (Bethesda, Maryland) guidelines. Twelve- to 14-week-old wild-type male C57BL/6J mice (Jackson Laboratory, USA) were used for this study. All animals were fed standard rodent food and water and were housed (five mice per cage) in a controlled environment with 12-h light/dark cycles. Mice were tagged and randomly allocated to each group before any treatment or procedure. Researchers were blinded to the group assignment, which was revealed only after completing the analysis. Mice did not experience unexpected lethality in the study, and animals were euthanized according to our institutional animal care and use committee guidelines.

Drug Administration

Recombinant HMGB1 (R&D Systems, USA) was dissolved in phosphate-buffered saline and administered intraperitoneally 50 µg/kg, a dose that we had previously reported to

produce a similar inflammatory and cognitive response as that seen after surgery.⁵

Dexmedetomidine (Sigma-Aldrich, USA) was dissolved in 0.9% sterile saline and administered 50 µg/kg intraperitoneally every 2 h for three doses immediately after HMGB1 (fig. 1, A and B) or at 20 µg/kg intraperitoneally for four doses in the surgical model (fig. 1, C and D). These doses were selected to simulate perioperative sedation either with (fig. 1, C and D) or without (fig. 1, A and B) isoflurane anesthesia.

Yohimbine (Sigma-Aldrich) was dissolved in 0.9% sterile saline, and 1.5 mg/kg was administered intraperitoneally, a dose that effectively blocks α_2 adrenoceptor-mediated responses.¹⁷ Atipamezole (Sigma-Aldrich) was dissolved in 5% dimethyl sulfoxide in saline, and 3 mg/kg was administered intraperitoneally, a dose that effectively blocks both imidazole receptor and α_2 adrenoceptor-mediated responses. Methlylcaconitine (Sigma-Aldrich) was dissolved in 0.9% sterile saline and 4 mg/kg was administered intraperitoneally, a dose that blocks α_7 nicotinic acetylcholine receptor-mediated response.²

Aseptic Surgical Trauma

Under aseptic conditions, groups of mice were subjected to an open tibia fracture of the left hind paw with an intramedullary fixation as previously described.⁵ Briefly, mice received general anesthesia with 2% isoflurane, and analgesia was achieved with buprenorphine 0.1 mg/kg administered subcutaneously, immediately after anesthetic induction. Warming pads and temperature-controlled lights were used to maintain body temperature. The entire procedure from the induction of anesthesia to the end of surgery lasted 12 ± 5 min.

Cognitive Testing

Trace fear conditioning was used to assess learning and memory in rodents as previously described.^{2,3} Briefly, mice are trained to associate a conditional stimulus, such as a tone, with an aversive, unconditional stimulus, such as a foot shock. Aversive memory is associated with freezing behavior when the rodent is reexposed to the same context. The behavioral study was conducted using a conditioning chamber (Med Associates Inc., USA) and an unconditional stimulus (two periods of 2-s foot shock of 0.75 mA). Behavior was captured with an infrared video camera (Video Freeze; Med Associates Inc.). Mice underwent a context test 72 h after training, during which no tones or foot shocks were delivered. Lack of movement, indicating freezing behavior, was analyzed by the software of the video recordings. With this model, perturbations of the hippocampus that are associated with memory impairment result in disruption of recall of the fear responses to the presentation of the same context, resulting in a reduction in freezing behavior.⁵

Blood and Tissue Sample Harvesting

Twenty-four hours after the specific intervention, blood was collected transcardially after thoracotomy under isoflurane

Fig. 1. Study design. (A) Mice were randomly allocated to 10 groups ($n = 15/\text{group}$) and were pretreated intraperitoneally (ip) with antagonists (yohimbine/atipamezole/methyllycaconitine). Thirty minutes later, mice were trained in the trace fear conditioning paradigm. After the training session, high mobility group box 1 protein (HMGB1) or vehicle (phosphate-buffered saline) was administered ip. Dexmedetomidine was administered every 2 h, for a total of three times. Seventy-two hours after HMGB1, testing was performed in the trace fear conditioning. (B) Mice were randomly allocated to 10 groups ($n = 8/\text{group}$) and were pretreated ip with antagonists (yohimbine/atipamezole/methyllycaconitine), and 30 min later HMGB1 was administered. Dexmedetomidine was administered every 2 h for a total of three times. Blood and tissue were collected 24 h later. (C) Mice were randomly allocated to three groups ($n = 15/\text{group}$): control (vehicle only), surgery/anesthesia, and surgery/anesthesia + dexmedetomidine. Mice were trained in the trace fear conditioning paradigm. After the training session, animals were anesthetized with isoflurane and subjected to aseptic trauma. Dexmedetomidine was administered and the mice were tested in the trace fear conditioning three days later. (D) Mice were randomly allocated to three groups ($n = 5\text{--}6/\text{group}$): control (vehicle only), surgery/anesthesia, and surgery/anesthesia + dexmedetomidine. Mice were anesthetized with isoflurane and subjected to aseptic trauma. Dexmedetomidine was administered and blood and tissue were collected 24 h later.

anesthesia and placed in heparin-coated syringes. After collection of blood, mice were immediately perfused with saline, and the hippocampus, brain, lung, and spleen were then rapidly extracted and stored at -80°C for further testing. After centrifugation of the blood sample at 3,400 rotations per minute for 10 min at 4°C , plasma was collected and stored at -80°C until these were assayed.

Circulating Cytokines

Plasma interleukin (IL)-6 and IL- 1β were quantified using commercially available enzyme-linked immunosorbent kits, according to the manufacturer's instructions (R&D Systems, USA).

Hippocampal Inflammatory Markers

Two different techniques were used to assess hippocampal inflammation. For experiments described in figure 1B, the hippocampus was homogenized and sonicated in cell lysis buffer (Cell Signaling Technology, USA) plus protease inhibitor (Halt Protease Inhibitor Single-Use Cocktail, Thermo Fisher Scientific, USA) and phenylmethanesulfonyl fluoride (Cell Signaling Technology). Protein concentration was assayed with a Pierce BCA Protein Assay kit (Thermo Fisher Scientific). IL-6 and IL- 1β were measured using commercially available enzyme-linked immunosorbent kits, according to the manufacturer's instructions (R&D Systems). For experiments described in figure 1D, after mice were perfused

with saline, the hippocampus was rapidly extracted, placed in RNeasy lysis solution (Qiagen, The Netherlands), and stored at 4°C overnight. Total RNA was extracted using RNeasy Lipid Tissue Kit (Qiagen). Extracted RNA was treated with recombinant DNase I by using a RNase-Free DNase set (Qiagen). Messenger RNA (mRNA) concentrations were determined with a ND-1000 Spectrophotometer (NanoDrop; Thermo Fisher Scientific), and mRNA was reverse-transcribed to complementary DNA with a High Capacity RNA to-cDNA Kit (Applied Biosystems, USA). TaqMan Fast Advanced Master Mix (Applied Biosystems) and specific gene-expression assays were used for quantitative polymerase chain reaction, actin- β (NM_007393.1), and IL-6 (Mm00446190_m1). Quantitative polymerase chain reaction was performed using StepOnePlus (Applied Biosystems). Each sample was run in triplicate, and relative gene expression was calculated using the comparative threshold cycle $\Delta\Delta\text{Ct}$ and normalized to β -actin. Results are expressed as fold increases relative to controls.

Measurement of Circulating Leukotriene B_4 and Lipoxin A_4

Plasma leukotriene B_4 (LTB $_4$) and lipoxin A_4 (LXA $_4$) were quantified using commercially available enzyme-linked immunosorbent kits, according to the manufacturer's instructions (Biomatik USA, LLC, USA). Results are expressed as fold-change compared with that measured in control mice that did not receive any intervention.

Measurement of Netrin-1 (in the Lung and Spleen) and Albumin (in the Brain)

Tissues were homogenized with RIPA Lysis Buffer (Cell Signaling Technology) plus protease inhibitor (Halt Protease Inhibitor Single-Use Cocktail, Thermo Fisher Scientific) and phenylmethanesulfonyl fluoride (Cell Signaling Technology) and sonicated. Protein concentration was assayed with Pierce BCA Protein Assay kit (Thermo Prod). For immunoblotting, the buffer for the samples was prepared by adding 950 μ l of 2 $\times$ Laemmli Sample Buffer to 50 μ l of 2-mercaptoethanol (Bio-Rad, USA). The protein samples were mixed in a 1:1 ratio with the sample buffer. After boiling for 5 min, 20 μ g of protein was loaded onto 12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. After electrophoresis, the analytes were transferred onto nitrocellulose transfer membranes. After the membranes were incubated with blocking buffer (LICOR Biosciences, USA) for 1 h at room temperature, these were incubated with primary antibodies at 1:1,000 dilution overnight at 4°C. The primary antibodies that were used for immunoblotting were rabbit monoclonal antibodies directed against murine netrin-1 (ab126729, Abcam, USA) and murine albumin (ab207327, Abcam). For the loading control, a rabbit monoclonal antibody directed at murine glyceraldehyde-3-phosphate dehydrogenase (ab181602, Abcam) was used. After washing four times with Tween buffered saline containing 0.1% Tween, membranes were incubated with a 1:10,000 dilution of IRDye 680RD- or 800RD-labeled goat antirabbit antibody (LICOR Biosciences) for 1 h at room temperature. Membranes were washed three times with TBS containing 0.1% Tween and once with TBS, and images were captured and quantified using a LICOR Imager (LICOR Biosciences).

Statistical Analysis

All data in this study were analyzed using Prism 6.0 (GraphPad Software, USA) and were expressed as mean $\pm$ SD. Statistical comparison was performed by a one-way ANOVA followed by Tukey test for *post hoc* analysis. Significance was set at *P* value less than 0.05. No statistical power calculation was conducted before our study, and the sample size selected was based on our previous experience using this design.^{3,18}

Results

Dexmedetomidine Prevents Cognitive Decline Induced by HMGB1 in an Imidazoline- and α_7 Nicotinic Acetylcholine Receptor-dependent Mechanism

Consistent with our previous report,⁵ HMGB1 significantly decreased freezing time (%) compared to the control group ($41.45 \pm 7.64\%$ vs. $58.94 \pm 7.22\%$; $P < 0.0001$; fig. 2). Administration of dexmedetomidine prevented HMGB1-induced cognitive decline ($58.77 \pm 8.69\%$ vs. $41.45 \pm 7.64\%$; $P < 0.0001$). In contrast to the lack of effect by an α_2 adrenoceptor-blocking dose of yohimbine, atipamezole, which has blocking activity at both the α_2 adrenergic receptor and the imidazoline receptor, prevented dexmedetomidine-induced

Fig. 2. Dexmedetomidine prevents high mobility group box 1 protein (HMGB1)-induced decrement in freezing behavior in an atipamezole- and methyllycaconitine-sensitive manner. Ten groups of randomly assigned mice ($n = 15$ /group) were administered antagonists (methyllycaconitine, atipamezole, yohimbine) before HMGB1 and subjected to trace fear conditioning training with and without dexmedetomidine exposure. Testing for freezing behavior in the trace fear conditioning context was undertaken 72 h later. Freezing time data are expressed as means $\pm$ SD and were analyzed by one-way ANOVA and Tukey *post hoc* test. * $P < 0.0001$ for comparisons shown.

reversal of HMGB1-mediated cognitive decline ($40.87 \pm 6.60\%$ vs. $58.77 \pm 8.69\%$; $P < 0.0001$). The reversal effect by dexmedetomidine of HMGB1-induced cognitive decline was prevented by the α_7 nicotinic acetylcholine receptor antagonist, methyllycaconitine ($37.79 \pm 9.51\%$ vs. $58.77 \pm 8.69\%$; $P < 0.0001$).

Dexmedetomidine Prevents HMGB1-induced Systemic Inflammation through an Imidazoline- and α_7 Nicotinic Acetylcholine Receptor-dependent Mechanism

Twenty-four hours after HMGB1 administration, plasma IL-1 β (fig. 3A) and IL-6 (fig. 3B) were significantly increased ninefold (IL-1 β : 49.8 ± 6.0 pg/ml vs. 5.2 ± 2.5 pg/ml, $P < 0.0001$) and sixfold (IL-6: 19.5 ± 1.7 pg/ml vs. 3.1 ± 1.7 pg/ml, $P < 0.0001$), respectively. Exposure to dexmedetomidine reduced the plasma concentration of both proinflammatory cytokines to normal levels (IL-1 β : 7.0 ± 2.2 pg/ml vs. 49.8 ± 6.0 pg/ml, $P < 0.0001$; IL-6: 3.2 ± 1.6 pg/ml vs. 19.5 ± 1.7 pg/ml, $P < 0.0001$). Among the α_2 adrenoceptor antagonists, only atipamezole, which also has activity at the imidazole receptor, abolished the antiinflammatory response of dexmedetomidine (IL-1 β : 47.8 ± 7.2 pg/ml vs. 7.0 ± 2.2 pg/ml, $P < 0.0001$). While methyllycaconitine, the α_7 nicotinic acetylcholine receptor antagonist, also prevented inhibition by dexmedetomidine of the peripheral inflammatory response to HMGB1 (IL-1 β : 65.0 ± 9.0 pg/ml vs. 7.0 ± 2.2 pg/ml, $P < 0.0001$), it is notable that methyllycaconitine, alone, significantly enhanced

Fig. 3. Dexmedetomidine prevents high mobility group box 1 protein (HMGB1)-induced peripheral inflammation in an atipamezole- and methyllycaconitine-sensitive manner. Ten groups of randomly assigned mice ($n = 8/\text{group}$) were administered antagonists (methyllycaconitine, atipamezole, yohimbine) before HMGB1 in the presence or absence of dexmedetomidine. Twenty-four hours after HMGB1, mice were euthanized, and the blood was harvested and assayed by enzyme-linked immunosorbent assay for circulating interleukin (IL)-1 β (A) and IL-6 (B). Data are expressed as means $\pm$ SD and analyzed by one-way ANOVA and Tukey *post hoc* test. $*P < 0.0001$, $\#P < 0.01$ for comparisons shown.

the inflammatory response to HMGB1 (IL-1 β : 68.4 ± 5.8 pg/ml *vs.* 49.8 ± 6.0 pg/ml, $P = 0.006$).

Dexmedetomidine Prevents HMGB1-induced Hippocampal Inflammation through an Imidazoline- and α_7 Nicotinic Acetylcholine Receptor-dependent Mechanism

At 24 h after HMGB1, hippocampal IL-1 β (fig. 4A) and IL-6 (fig. 4B) were significantly increased eightfold (41.6 ± 8.0 pg/mg *vs.* 5.5 ± 2.0 pg/mg, $P < 0.0001$) and sixfold (16.2 ± 2.7 pg/mg *vs.* 2.7 ± 1.5 pg/mg, $P < 0.0001$), respectively. Exposure to dexmedetomidine reduced the hippocampal concentration of both proinflammatory cytokines to normal levels for IL-1 β (4.1 ± 3.0 pg/mg *vs.* 41.6 ± 8.0 pg/mg, $P < 0.0001$) and for IL-6 (3.4 ± 1.3 pg/mg *vs.* 16.2 ± 2.7 pg/mg, $P < 0.0001$). Among the α_2 adrenoceptor antagonists, only atipamezole abolished dexmedetomidine's antiinflammatory response of IL-1 β (43.7 ± 7.7 pg/mg *vs.* 4.1 ± 3.0 pg/mg, $P < 0.0001$) and IL-6 (14.4 ± 2.8 pg/mg *vs.* 3.4 ± 1.3 pg/mg, $P < 0.0001$). Methyllycaconitine, the α_7 nicotinic acetylcholine receptor antagonist, also prevented inhibition by dexmedetomidine of the HMGB1-induced peripheral inflammatory response as reflected by IL-1 β (45.8 ± 10.1 pg/mg *vs.* 4.1 ± 3.0 pg/mg, $P < 0.0001$) and IL-6 (16.5 ± 2.9 pg/mg *vs.* 3.4 ± 1.3 pg/mg, $P < 0.0001$). Unlike systemic inflammation (fig. 3), methyllycaconitine did not enhance the hippocampal inflammatory response to HMGB1 (fig. 4).

Dexmedetomidine Reverses HMGB1-induced Downregulation of Netrin-1 Expression in an α_7 Nicotinic Acetylcholine Receptor-sensitive Manner

Accompanying the sterile inflammation induced by HMGB1 (figs. 3 and 4), netrin-1 expression is significantly decreased in the lung (0.7 ± 0.1 *vs.* 1.0 ± 0.2 , $P = 0.0173$, fig. 5A) and spleen (0.6 ± 0.2 *vs.* 1.0 ± 0.2 , $P = 0.0220$, fig. 5B), organs that are vagally innervated. Exposure to dexmedetomidine reverses HMGB1-induced netrin-1 downregulation in both

organs (1.5 ± 0.1 *vs.* 0.7 ± 0.1 , $P < 0.0001$, fig. 5A; 1.4 ± 0.2 *vs.* 0.7 ± 0.2 , $P < 0.0001$, fig. 5B), and pretreatment with methyllycaconitine, the α_7 nicotinic acetylcholine receptor antagonist, prevented reversal of netrin expression by dexmedetomidine (0.7 ± 0.1 *vs.* 1.5 ± 0.1 , $P < 0.0001$, fig. 5A; 0.6 ± 0.2 *vs.* 1.4 ± 0.2 , $P < 0.0001$, fig. 5B).

Dexmedetomidine Reverses HMGB1-induced Changes in the Expression of Circulating LTB $_4$ and LXA $_4$ in an α_7 Nicotinic Acetylcholine Receptor-sensitive Mechanism

Accompanying the sterile inflammation induced by HMGB1 (figs. 3 and 4) and the downregulation of netrin-1 expression (fig. 5), there is a threefold upregulation in the relative expression of LTB $_4$, the proinflammatory lipid mediator (2.7 ± 0.4 *vs.* 1.0 ± 0.2 , $P < 0.0001$, fig. 6A). Dexmedetomidine reverses HMGB1-induced upregulation of LTB $_4$ (1.0 ± 0.2 *vs.* 2.7 ± 0.4 , $P < 0.0001$, fig. 6A), and this reversal is prevented by methyllycaconitine (3.0 ± 0.3 *vs.* 1.0 ± 0.2 , $P < 0.0001$, fig. 6A). Conversely, dexmedetomidine upregulated the expression of LXA $_4$ (1.7 ± 0.2 *vs.* 1.2 ± 0.2 , $P < 0.0001$, fig. 6B), the specific proresolving mediator, an effect that was reversed by methyllycaconitine (0.9 ± 0.2 *vs.* 1.7 ± 0.2 , $P < 0.0001$, fig. 6B).

Dexmedetomidine Reverses HMGB1-induced Leakage of the Blood-Brain Barrier

Accompanying the HMGB1-induced inflammation (figs. 3 and 4), the blood-brain barrier is disrupted, as evidenced by a significant upregulation of albumin expression in the brain assessed by immunoblotting. The upregulation in brain albumin expression (1.0 ± 0.2 *vs.* 1.7 ± 0.3 , $P = 0.0019$) is suppressed by dexmedetomidine (1.7 ± 0.3 *vs.* 1.0 ± 0.2 , $P = 0.002$; fig. 7).

Dexmedetomidine Reverses Surgery-induced Cognitive Decline and Inflammation

Because isoflurane had no effect on HMGB1-induced cognitive decline (data not shown), the effect of dexmedetomidine

Fig. 4. Dexmedetomidine prevents high mobility group box 1 protein (HMGB1)–induced hippocampal inflammation in an atipamezole- and methyllycaconitine-sensitive manner. Ten groups of randomly assigned mice ($n = 8/\text{group}$) were administered antagonists (methyllycaconitine, atipamezole, yohimbine) before HMGB1 in the presence or absence of dexmedetomidine. Twenty-four hours after HMGB1, mice were euthanized, and the hippocampus was harvested and assayed by ELISA for interleukin (IL)-1 β (A) and IL-6 (B). Data are expressed as means $\pm$ SD and were analyzed by one-way ANOVA and Tukey *post hoc* test. * $P < 0.0001$ for comparisons shown.

Fig. 5. Dexmedetomidine prevents high mobility group box 1 protein (HMGB1)–induced downregulation of netrin-1 expression in the lung (A) and spleen (B) in an α_7 nicotinic acetylcholine receptor–dependent manner. Four groups of randomly assigned mice ($n = 5/\text{group}$) were administered saline vehicle (control), HMGB1 alone, HMGB1 + dexmedetomidine, or HMGB1 + dexmedetomidine + methyllycaconitine. Twenty-four hours later, mice were euthanized, and lung (A) and spleen (B) were harvested for expression of netrin-1 by immunoblotting. Data are expressed as means $\pm$ SD fold-change relative to control and were analyzed by one-way ANOVA and Tukey *post hoc* test. # $P < 0.05$ and * $P < 0.0001$ for comparisons shown. GAPDH = glyceraldehyde-3-phosphate dehydrogenase.

on surgery plus isoflurane-induced cognitive decline was investigated. Surgery-induced cognitive decline was reversed by dexmedetomidine ($35.91 \pm 5.03\%$ vs. $54.62 \pm 8.82\%$, $P < 0.0001$, fig. 8A). Twenty-four hours after surgery, we observed a significant decrease in circulating IL-6 (83.2 ± 60.2 pg/ml vs. 30.1 ± 13.7 pg/ml, $P = 0.013$, fig. 8B). The change of hippocampal mRNA expression of IL-6 after surgery was reversed by dexmedetomidine (8.6 ± 1.3 vs. 10.7 ± 1.0 , $P = 0.012$, fig. 8C).

Discussion

Recapitulation of Main Findings

Using HMGB1 to generate a surrogate of the surgical phenotype without the confounding influence of general anesthesia, dexmedetomidine reversed HMGB1-induced cognitive decline (fig. 2), as well as systemic (fig. 3) and hippocampal (fig. 4) inflammation. In each case, reversal by dexmedetomidine of the HMGB1-induced surgical phenotype

Fig. 6. Dexmedetomidine downregulates the circulating proinflammatory mediator leukotriene B₄ (LTB₄; A) and upregulates the circulating proresolving mediator lipoxin A₄ (LXA₄; B). Four groups of randomly assigned mice (n = 8/group) were administered saline vehicle (control), high mobility group box 1 protein (HMGB1) alone, HMGB1+ dexmedetomidine, or HMGB1 + dexmedetomidine + methyllycaconitine. Twenty-four hours later, mice were euthanized and the blood was harvested and assayed by ELISA for plasma LTB₄ (A) and LXA₄ (B). Data are expressed as means ± SD fold-change relative to control and were analyzed by one-way ANOVA and Tukey *post hoc* test. **P* < 0.0001 for comparisons shown.

Fig. 7. Dexmedetomidine reverses high mobility group box 1 protein (HMGB1)-induced leakage of blood–brain barrier. Three groups of randomly assigned mice (n = 5/group) were treated with vehicle (control), HMGB1, or HMGB1 + dexmedetomidine. Twenty-four hours after treatment, mice were euthanized, and the brains were harvested for immunoblotting of albumin expression. Data are expressed as means ± SD relative to control and were analyzed by one-way ANOVA and Tukey *post hoc* test. **P* < 0.01 for comparisons shown. GAPDH = glyceraldehyde-3-phosphate dehydrogenase.

was blocked by atipamezole (figs. 2–4), an α_2 adrenoceptor antagonist that also has activity at the imidazole receptor,¹⁹ but not by yohimbine, an α_2 adrenoceptor antagonist that has no activity at the imidazole receptor. Reversal by

dexmedetomidine of the surgical phenotype was also blocked by methyllycaconitine (figs. 2–4), the α_7 nAChR antagonist. Dexmedetomidine reversed the HMGB1-induced downregulation of netrin-1 in the vagally innervated lung (fig. 5A) and spleen (fig. 5B); reversal by dexmedetomidine was attenuated by antagonism of signaling at the vagal termini by α_7 nicotinic acetylcholine receptor blockade with methyllycaconitine. Upregulation of LTB₄, the proinflammatory humoral response to HMGB1, was prevented by dexmedetomidine (fig. 6A), while the proresolving mediator, LXA₄, was upregulated by dexmedetomidine (fig. 6B). In each case, the effects of dexmedetomidine were negated by pretreatment with methyllycaconitine, the α_7 nicotinic acetylcholine receptor blocker (fig. 6). Disruption of the blood–brain barrier by HMGB1 was reversed by dexmedetomidine (fig. 7). Finally, perioperative exposure to dexmedetomidine prevented surgery-induced cognitive decline and peripheral and hippocampal inflammation (fig. 8).

Justification for the Use of HMGB1 to Produce the Surgical Phenotype

HMGB1 regulates transcription of nuclear factor kappa-light-chain-enhancer of activated B cells,²⁰ establishing it as an extracellular orchestrator of the systemic inflammatory response.²¹ Previously, we showed the causal role of HMGB1 in mediating postoperative cognitive decline after passive release from traumatized tissues.⁵ After binding to pattern recognition receptors on circulating immunocytes, HMGB1 initiates the innate immune response to aseptic surgical trauma. Exogenously administered HMGB1 reproduces the surgical phenotype if myeloid-derived circulating monocytes are present,⁵ a feature also noted after surgery. Other cardinal features of the surgical phenotype, including peripheral (fig. 3) and neuroinflammation (fig. 4) and disruption of the blood–brain barrier (fig. 7), are reproduced after HMGB1 administration.²

Fig. 8. Dexmedetomidine reverses surgery-induced cognitive decline (A) and peripheral (B) and neuroinflammation (C). Three groups of randomly assigned mice ($n = 15/\text{group}$) were treated with (1) vehicle, (2) tibia fracture during anesthesia (Sx/Anesth) + vehicle, and (3) Sx/Anesth + dexmedetomidine before training in the trace fear conditioning paradigm (A). Testing for freezing behavior in the trace fear conditioning context was undertaken 72 h later. Freezing time data are expressed as means $\pm$ SD and were analyzed by one-way ANOVA and Tukey *post hoc* test. Three groups of randomly assigned mice ($n = 6/\text{group}$) were treated with (1) vehicle, (2) tibia fracture during anesthesia (Sx/Anesth) + vehicle, and (3) Sx/Anesth + dexmedetomidine. Mice were euthanized at 24 h, and blood and brain were harvested. Plasma interleukin (IL)-6 was assayed by enzyme-linked immunosorbent assay (B), and hippocampal IL-6 was assayed by quantitative polymerase chain reaction (C). The means $\pm$ SD of the mean for expression of IL-6 protein (B) and mRNA (C) were analyzed by one-way ANOVA and Tukey *post hoc* test. # $P < 0.05$; * $P < 0.01$; ** $P < 0.0001$.

Use of Antagonists with Which to Probe the Mechanism for Dexmedetomidine's Effects in Reversing the Surgical Phenotype

α_2 Adrenoceptor agonists such as dexmedetomidine and clonidine have an imidazole ring structure facilitating binding to and activation of the imidazoline receptor.²² Because atipamezole, which has antagonist activity at the imidazoline receptor,²⁰ was able to block dexmedetomidine's reversal of the surgical phenotype while yohimbine, which has no activity at the imidazoline receptor, was ineffective, we invoke an action mediated by the imidazoline receptor for the reversal of dexmedetomidine's effect. An important property of imidazoline receptor agonists is its negative chronotropic property,²³ which is prevented by vagotomy.²⁴ As enhancement of vagal activity by α_2 agonists with an imidazole ring has been well documented²⁵ and, as the bradycardic effect of α_2 agonists with an imidazole ring structure is unrelated to its binding affinity to α_2 adrenoceptors,¹⁹ we propose that this imidazoline receptor-mediated vagomimetic action may be the mechanism whereby dexmedetomidine reverses the surgical phenotype.

Increase in vagal activity has been shown to be an important mechanism for resolving inflammation.²⁶ The α_7 nicotinic acetylcholine receptor transduces the inflammation-resolving vagomimetic effect.²⁷ A supporting finding that the enhanced vagal activity is the likely explanation for dexmedetomidine's reversal of the surgical phenotype is provided by the fact that α_7 nicotinic acetylcholine receptor blockade by methyllycaconitine eliminated the ameliorative effect produced by dexmedetomidine (figs. 2–4).

Role of Vagal Stimulation on Neural- and Humoral-mediated Resolution of Inflammation

Neural²⁸ and humoral²³ pathways have been implicated in the resolution of acute inflammation. Neural mechanisms

involve efferent vagal fibers in which release of acetylcholine activates α_7 nicotinic acetylcholine receptor on immunocompetent cells,²⁹ thereby inhibiting NF- κ B and downregulating synthesis of proinflammatory cytokines.³⁰ For the humoral pathway, biotransformation of free fatty acids (arachidonic acid, eicosapentaenoic acid, and docosahexaenoic acid) elaborates specific proresolving lipid mediators.³¹ During synthesis of eicosanoid hormones, there are juncture points at which a precursor, such as leukotriene A_4 , can elaborate two different products (LTB_4 and LXA_4) with diametrically opposite effects, namely, proinflammation and proresolution, respectively.³² Dexmedetomidine appears to shunt the biotransformation pathway to the synthesis of LXA_4 at the expense of LTB_4 (fig. 6).

Humoral and neural pathways have been recently linked to vagally mediated expression of netrin-1.³³ Netrin-1, originally identified as a neuronal guidance protein, also limits inflammation.³⁴ Interestingly, netrin-1 expression is downregulated in vagally innervated lung and spleen after HMGB1 administration, an effect that is reversed by dexmedetomidine (fig. 5). A similar downregulation of netrin-1 expression has been noted in other inflammation-inducing experimental settings.³³

Integrating Results into Existing Molecular Mechanism Model for Postoperative Cognitive Decline

Previously, we had shown that trauma-induced release of HMGB1 engages circulating monocytes⁵ to stimulate the synthesis of proinflammatory cytokines that are capable of disrupting the blood–brain barrier,² enabling the passage of bone marrow–derived monocytes to enter into the brain, attracted by upregulation of the chemokine MCP-1 by microglia in the hippocampus.³ Within the hippocampus, the bone marrow–derived monocytes activate microglia, resulting in release of proinflammatory cytokines that disrupt

long-term potentiation,³⁵ the neurobiologic correlate of learning and memory. Within 7 days, the inflammation usually resolves, as does postoperative cognitive decline, except in vulnerable animals, in which inflammation and cognitive decline can both be exaggerated³ and more persistent.¹⁸

Dexmedetomidine appears to reverse postoperative cognitive decline by enhancing the inflammation-resolving pathways. Dexmedetomidine stimulates the vagus through imidazoline receptor activation, resulting in inhibition of the NF- κ B–dependent synthesis of proinflammatory cytokines (figs. 3 and 4) through activation of α_7 nicotinic acetylcholine receptor (figs. 3 and 4). Netrin-1 is downregulated by HMGB1 through a vagal pathway, and this is reversed by dexmedetomidine (fig. 5), resulting in a change in the elaboration of specific proresolving mediators (increase in LXA₄ and decrease in LTB₄) that resolves inflammation (fig. 6).

Caveats

Methyllycaconitine-induced Upregulation of Circulating Proinflammatory Cytokines

Methyllycaconitine was used in a dose that blocks the α_7 nicotinic acetylcholine receptor–mediated inhibition of synthesis of proinflammatory cytokines.² However, at this dose there was an increase in HMGB1-induced proinflammatory cytokines (fig. 3), and it may be argued that the reversal of the anti-inflammatory effect of dexmedetomidine could be due to the nonspecific enhancement of proinflammatory cytokines independent of any specific action that dexmedetomidine exerts on the α_7 nicotinic acetylcholine receptor through vagal stimulation. It is notable that in the hippocampus, where cognitive decline is produced, there was not a similar methyllycaconitine-induced enhancement of proinflammatory cytokines.

Doses of Dexmedetomidine Used in HMGB1- and Surgery-induced Cognitive Decline

In the HMGB1-induced surrogate of the surgical phenotype, we used a dexmedetomidine dose of 50 μ g/kg, while in the trauma-induced model, a dexmedetomidine dose of 20 μ g/kg was used. The reason that these doses differ was to prevent a significant increase in the sedative effect of dexmedetomidine in the presence of isoflurane³⁶ that may increase postoperative cognitive decline.¹¹ The lower dexmedetomidine dose in the surgical model (fig. 8) was as effective as the higher dose in the HMGB1 model (figs. 2–4) at reversing cognitive decline and inflammation.

Relevance of Findings in Young Mice to Vulnerable Animal Models

In the current study, we only tested the efficacy and putative mechanisms whereby dexmedetomidine prevents inflammation and cognitive decline in young mice. Whether these observed cognitive decline–reducing properties of dexmedetomidine occurs in vulnerable models (including aging, obesity,³ and metabolic syndrome¹⁸) remains to be determined.

Relevance of Preclinical Mechanistic Findings in Light of Dexmedetomidine's Ability to Reverse the Surgical Phenotype in Clinical Studies

Sedation with dexmedetomidine reduces the likelihood of delirium in the intensive care unit when compared to benzodiazepines^{13,37} and when dexmedetomidine was administered during the first postoperative night to elderly surgical patients.¹⁴ Because dexmedetomidine changes the activity in neuronal pathways in a similar manner to those altered during natural sleep³⁸ and produces similar electroencephalographic changes to those seen during natural sleep,³⁹ and because both sleep deprivation⁴⁰ and sleep fragmentation⁴¹ induce neuroinflammation and cognitive decline, we had conjectured that the cognition-enhancing effects of dexmedetomidine were due to its unique sedative profile. However, as dexmedetomidine-induced sedation is mediated by the α_{2A} adrenoceptors⁴² that are antagonized by yohimbine,⁴³ our finding that reversal by dexmedetomidine of the surgical phenotype is insensitive to yohimbine challenges that explanation.

We recognize that the reversal of postoperative cognitive decline by dexmedetomidine is not universal. Recently, Deiner *et al.*⁴⁴ reported that dexmedetomidine lacked efficacy in preventing the onset of postoperative delirium.⁴⁴ Further studies are needed to define the patient subgroups that may be resistant to the cognitive decline–reducing properties of dexmedetomidine.

Future Applications

These preclinical studies were performed in mice that do not exhibit exaggerated and persistent cognitive decline. The effectiveness of dexmedetomidine will need to be established in reagents that have abnormalities in their inflammation-resolving pathways including in advanced age with its “inflammaging” processes⁴⁵ and the metabolic syndrome with its precocious aging phenotype.⁴⁶

While we have already shown that dexmedetomidine is effective at decreasing delirium in mechanically ventilated medical/surgical intensive care patients¹³ as well as in non-ventilated surgical patients,¹⁴ it will be important to demonstrate that wound healing and the ability to combat infection are not jeopardized by a dexmedetomidine intervention that perturbs the innate immune system. Regarding infection, it is notable that survival is enhanced by dexmedetomidine in a preclinical model of sepsis,⁴⁷ a finding that was also noted in a *post hoc* analysis of the Maximizing Efficacy of Targeted Sedation and Reducing Neurological Dysfunction (MENDS) trial.⁴⁸ However, dexmedetomidine was not shown to improve outcome in a recently reported trial of septic patients; this question is being further addressed in the MENDS II trial (NCT01739933) comparing outcomes in septic patients sedated with dexmedetomidine *versus* propofol. Whether the existing role of dexmedetomidine as a perioperative and procedural sedative agent can be supplemented by indications in which resolution of inflammation requires bolstering will need to be studied further. Furthermore, if the

activation of the imidazoline receptor and subsequent vagal outflow are confirmed to be the reason for dexmedetomidine's antiinflammatory properties, it may be possible to reproduce these effects with a selective imidazoline receptor agonist and avoid α_2 adrenoceptor properties such as sedation.

Acknowledgments

The authors wish to thank Jinbao Chen, M.D. (Department of Anesthesia, Tongling People's Hospital, Tongling Anhui, People's Republic of China), for providing postdoctoral support to Dr. Hu and to Christopher Chang, B.A. (University of California, Santa Cruz, California), for help with the preparation of the manuscript.

Research Support

Supported by grant No. R01GM104194 from the National Institutes of Health, Bethesda, Maryland (to Dr. Maze).

Competing Interests

Dr. Maze is a coinventor on a patent for the use of dexmedetomidine for sedation. Between 1987 and 1991, Dr. Maze's laboratory at Stanford University (Stanford, California) received \$250,000 for the assignment of the patent to Farnos (Espoo, Finland), the company that synthesized dexmedetomidine. Between 1995 and 2008, Dr. Maze was intermittently paid as a consultant by Orion-Farnos (Espoo, Finland), Abbott Labs (Chicago, Illinois), and Hospira (Lake Forest, Illinois) for advising on the pivotal phase III clinical trials, approval of the new drug application, and subsequent marketing of the product. Dr. Maze has not received any payments for at least the last 5 yr. Dr. Maze has not and will not receive royalty payments for sales of dexmedetomidine. The other authors declare no competing interests.

Correspondence

Address correspondence to Dr. Maze: Zuckerberg San Francisco General, 1001 Potrero Avenue, Building 10, Room 1206, San Francisco, California 94110-3518. mervyn.maze@ucsf.edu. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

1. Terrando N, Monaco C, Ma D, Foxwell BM, Feldmann M, Maze M: Tumor necrosis factor- α triggers a cytokine cascade yielding postoperative cognitive decline. *Proc Natl Acad Sci USA* 2010; 107:20518–22
2. Terrando N, Eriksson LI, Ryu JK, Yang T, Monaco C, Feldmann M, Jonsson Fagerlund M, Charo IF, Akassoglou K, Maze M: Resolving postoperative neuroinflammation and cognitive decline. *Ann Neurol* 2011; 70:986–95
3. Feng X, Valdearcos M, Uchida Y, Lutrind D, Maze M, Koliwad SK: Microglia mediate postoperative hippocampal inflammation and cognitive decline in mice. *JCI Insight* 2017; 2:e91229
4. Forsberg A, Cervenka S, Jonsson Fagerlund M, Rasmussen LS, Zetterberg H, Erlandsson Harris H, Stridh P, Christensson E, Granström A, Schenning A, Dymmel K, Knave N, Terrando N, Maze M, Borg J, Varrone A, Halldin C, Blennow K, Farde L, Eriksson LI: The immune response of the human brain to abdominal surgery. *Ann Neurol* 2017; 81:572–82
5. Vacas S, Degos V, Tracey KJ, Maze M: High-mobility group box 1 protein initiates postoperative cognitive decline by engaging bone marrow-derived macrophages. *ANESTHESIOLOGY* 2014; 120:1160–7
6. Terrando N, Yang T, Wang X, Fang J, Cao M, Andersson U, Erlandsson HH, Ouyang W, Tong J: Systemic HMGB1 neutralization prevents postoperative neurocognitive dysfunction in aged rats. *Front Immunol* 2016; 7:441
7. Bedford PD: Adverse cerebral effects of anaesthesia on old people. *Lancet* 1955; 269:259–63
8. Williams-Russo P, Sharrock NE, Mattis S, Szatrowski TP, Charlson ME: Cognitive effects after epidural vs general anesthesia in older adults. A randomized trial. *JAMA* 1995; 274:44–50
9. Neuman MD, Ellenberg SS, Sieber FE, Magaziner JS, Feng R, Carson JL; and the REGAIN Investigators: Regional *versus* General Anesthesia for Promoting Independence after Hip Fracture (REGAIN): Protocol for a pragmatic, international multicentre trial. *BMJ Open* 2016; 6:e013473
10. Radtke FM, Franck M, Lendner J, Krüger S, Wernecke KD, Spies CD: Monitoring depth of anaesthesia in a randomized trial decreases the rate of postoperative delirium but not postoperative cognitive dysfunction. *Br J Anaesth* 2013; 110 (Suppl 1):i98–105
11. Siddiqi N, Harrison JK, Clegg A, Teale EA, Young J, Taylor J, Simpkins SA: Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database Syst Rev* 2016; 3:CD005563
12. Coburn M, Sanders RD, Maze M, Rossaint R; HIPELD Investigators: The Hip Fracture Surgery in Elderly Patients (HIPELD) study: Protocol for a randomized, multicenter controlled trial evaluating the effect of xenon on postoperative delirium in older patients undergoing hip fracture surgery. *Trials* 2012; 13:180
13. Pandharipande PP, Pun BT, Herr DL, Maze M, Girard TD, Miller RR, Shintani AK, Thompson JL, Jackson JC, Deppen SA, Stiles RA, Dittus RS, Bernard GR, Ely EW: Effect of sedation with dexmedetomidine vs lorazepam on acute brain dysfunction in mechanically ventilated patients: The MENDS randomized controlled trial. *JAMA* 2007; 298:2644–53
14. Su X, Meng ZT, Wu XH, Cui F, Li HL, Wang DX, Zhu X, Zhu SN, Maze M, Ma D: Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: A randomised, double-blind, placebo-controlled trial. *Lancet* 2016; 388:1893–902
15. Chen Y, Miao L, Yao Y, Wu W, Wu X, Gong C, Qiu L, Chen J: Dexmedetomidine ameliorate CLP-induced rat intestinal injury via inhibition of inflammation. *Mediators Inflamm* 2015; 2015:918361
16. Rong H, Zhao Z, Feng J, Lei Y, Wu H, Sun R, Zhang Z, Hou B, Zhang W, Sun Y, Gu X, Ma Z, Liu Y: The effects of dexmedetomidine pretreatment on the pro- and anti-inflammation systems after spinal cord injury in rats. *Brain Behav Immun* 2017; 64:195–207
17. Hsing CH, Lin CF, So E, Sun DP, Chen TC, Li CF, Yeh CH: α_2 -Adrenoceptor agonist dexmedetomidine protects septic acute kidney injury through increasing BMP-7 and inhibiting HDAC2 and HDAC5. *Am J Physiol Renal Physiol* 2012; 303:F1443–53
18. Feng X, Degos V, Koch LG, Britton SL, Zhu Y, Vacas S, Terrando N, Nelson J, Su X, Maze M: Surgery results in exaggerated and persistent cognitive decline in a rat model of the metabolic syndrome. *ANESTHESIOLOGY* 2013; 118:1098–105
19. Ernsberger P, Giuliano R, Willette RN, Reis DJ: Role of imidazole receptors in the vasodepressor response to clonidine analogs in the rostral ventrolateral medulla. *J Pharmacol Exp Ther* 1990; 253:408–18
20. Tang D, Kang R, Zeh HJ III, Lotze MT: High-mobility group box 1, oxidative stress, and disease. *Antioxid Redox Signal* 2011; 14:1315–35

21. Wang H, Liao H, Ochani M, Justiniani M, Lin X, Yang L, Al-Abed Y, Wang H, Metz C, Miller EJ, Tracey KJ, Ulloa L: Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nat Med* 2004; 10:1216–21
22. Khan ZP, Ferguson CN, Jones RM: Alpha-2 and imidazoline receptor agonists. Their pharmacology and therapeutic role. *Anaesthesia* 1999; 54:146–65
23. Serhan CN: Treating inflammation and infection in the 21st century: New hints from decoding resolution mediators and mechanisms. *FASEB J* 2017; 31:1273–88
24. Boblewski K, Lehmann A, Sączewski F, Kornicka A, Rybczyńska A: Vagotomy reveals the importance of the imidazoline receptors in the cardiovascular effects of marsanidine and 7-ME-marsanidine in rats. *Pharmacol Rep* 2014; 66:874–9
25. Philbin KE, Bateman RJ, Mendelowitz D: Clonidine, an alpha2-receptor agonist, diminishes GABAergic neurotransmission to cardiac vagal neurons in the nucleus ambiguus. *Brain Res* 2010; 1347: 65–70
26. Chavan SS, Tracey KJ: Essential neuroscience in immunology. *J Immunol* 2017; 198: 3389–97
27. Gallowitsch-Puerta M, Tracey KJ: Immunologic role of the cholinergic anti-inflammatory pathway and the nicotinic acetylcholine alpha 7 receptor. *Ann NY Acad Sci* 2005; 1062:209–19
28. Pavlov VA, Tracey KJ: Neural regulation of immunity: Molecular mechanisms and clinical translation. *Nat Neurosci* 2017; 20:156–66
29. Rosas-Ballina M, Olofsson PS, Ochani M, Valdes-Ferrer SI, Levine YA, Reardon C, Tusche MW, Pavlov VA, Andersson U, Chavan S, Mak TW, Tracey KJ: Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit. *Science* 2011; 334:98–101
30. van Maanen MA, Lebre MC, van der Poll T, LaRosa GJ, Elbaum D, Vervoordeldonk MJ, Tak PP: Stimulation of nicotinic acetylcholine receptors attenuates collagen-induced arthritis in mice. *Arthritis Rheum* 2009; 60:114–22
31. Dalli J, Serhan CN: Specific lipid mediator signatures of human phagocytes: Microparticles stimulate macrophage efferocytosis and pro-resolving mediators. *Blood* 2012; 120:e60–72
32. Levy BD, Clish CB, Schmidt B, Gronert K, Serhan CN: Lipid mediator class switching during acute inflammation: Signals in resolution. *Nat Immunol* 2001; 2:612–9
33. Mirakaj V, Dalli J, Granja T, Rosenberger P, Serhan CN: Vagus nerve controls resolution and pro-resolving mediators of inflammation. *J Exp Med* 2014; 211:1037–48
34. Aherne CM, Collins CB, Masterson JC, Tizzano M, Boyle TA, Westrich JA, Parnes JA, Furuta GT, Rivera-Nieves J, Eltzschig HK: Neuronal guidance molecule netrin-1 attenuates inflammatory cell trafficking during acute experimental colitis. *Gut* 2012; 61:695–705
35. Terrando N, Gómez-Galán M, Yang T, Carlström M, Gustavsson D, Harding RE, Lindskog M, Eriksson LI: Aspirin-triggered resolvin D1 prevents surgery-induced cognitive decline. *FASEB J* 2013; 27:3564–71
36. Savola MK, MacIver MB, Doze VA, Kendig JJ, Maze M: The alpha 2-adrenoceptor agonist dexmedetomidine increases the apparent potency of the volatile anesthetic isoflurane in rats *in vivo* and in hippocampal slice *in vitro*. *Brain Res* 1991; 548:23–8
37. Riker RR, Shehabi Y, Bokesch PM, Ceraso D, Wisemandle W, Koura F, Whitten P, Margolis BD, Byrne DW, Ely EW, Rocha MG; SEDCOM (Safety and Efficacy of Dexmedetomidine Compared With Midazolam) Study Group: Dexmedetomidine vs midazolam for sedation of critically ill patients: A randomized trial. *JAMA* 2009; 301:489–99
38. Nelson LE, Lu J, Guo T, Saper CB, Franks NP, Maze M: The alpha2-adrenoceptor agonist dexmedetomidine converges on an endogenous sleep-promoting pathway to exert its sedative effects. *ANESTHESIOLOGY* 2003; 98:428–36
39. Akeju O, Kim SE, Vazquez R, Rhee J, Pavone KJ, Hobbs LE, Purdon PL, Brown EN: Spatiotemporal dynamics of dexmedetomidine-induced electroencephalogram oscillations. *PLoS One* 2016; 11:e0163431
40. Zhu B, Dong Y, Xu Z, Gompf HS, Ward SA, Xue Z, Miao C, Zhang Y, Chamberlin NL, Xie Z: Sleep disturbance induces neuroinflammation and impairment of learning and memory. *Neurobiol Dis* 2012; 48:348–55
41. Vacas S, Degos V, Maze M: Fragmented sleep enhances post-operative neuroinflammation but not cognitive dysfunction. *Anesth Analg* 2017; 124:270–6
42. Lakhilani PP, MacMillan LB, Guo TZ, McCool BA, Lovinger DM, Maze M, Limbird LE: Substitution of a mutant alpha2a-adrenergic receptor via “hit and run” gene targeting reveals the role of this subtype in sedative, analgesic, and anesthetic-sparing responses *in vivo*. *Proc Natl Acad Sci USA* 1997; 94:9950–5
43. Tang Y, Axelsson AS, Spégl P, Andersson LE, Mulder H, Groop LC, Renström E, Rosengren AH: Genotype-based treatment of type 2 diabetes with an α 2A-adrenergic receptor antagonist. *Sci Transl Med* 2014; 6:257ra139
44. Deiner S, Luo X, Lin HM, Sessler DI, Saager L, Sieber FE, Lee HB, Sano M, Jankowski C, Bergese SD, Candiotti K, Flaherty JH, Arora H, Shander A, Rock P; and the Dexlirium Writing Group: Intraoperative Infusion of Dexmedetomidine for Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly Patients Undergoing Major Elective Noncardiac Surgery: A randomized clinical trial. *JAMA Surg* 2017; 152:e171505
45. Franceschi C, Garagnani P, Vitale G, Capri M, Salvioli S: Inflammaging and ‘Garb-aging.’ *Trends Endocrinol Metab* 2017; 28:199–212
46. Su X, Feng X, Terrando N, Yan Y, Chawla A, Koch LG, Britton SL, Matthay MA, Maze M: Dysfunction of inflammation-resolving pathways is associated with exaggerated postoperative cognitive decline in a rat model of the metabolic syndrome. *Mol Med* 2013; 18:1481–90
47. Qiao H, Sanders RD, Ma D, Wu X, Maze M: Sedation improves early outcome in severely septic Sprague Dawley rats. *Crit Care* 2009; 13:R136
48. Pandharipande PP, Sanders RD, Girard TD, McGrane S, Thompson JL, Shintani AK, Herr DL, Maze M, Ely EW; MENDS investigators: Effect of dexmedetomidine *versus* lorazepam on outcome in patients with sepsis: An a priori-designed analysis of the MENDS randomized controlled trial. *Crit Care* 2010; 14:R38